



Sample Information

Patient Name: 江欽思  
Gender: Male  
ID No.: F104017141  
History No.: 48438239  
Age: 70  
  
Ordering Doctor: DOC1050C 劉嘉仁  
Ordering REQ.: 0CLCTNF  
Signing in Date: 2023/05/25

Path No.: M112-00113  
MP No.: MY23028  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marro  
Bone Marrow Aspirating Date: 2023/05/19

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	4 Therapies Available
Relevant Therapy Summary	5	0 Clinical Trials
Relevant Therapy Details	5	
Diagnostic Details	8	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	<b>CEBPA p.(G123Efs*37) c.368delG</b>	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>CEBPA p.(G123Efs*37) c.368delG</b> CCAAT enhancer binding protein alpha Allele Frequency: 45.04%  <b>Diagnostic significance:</b> Acute Myeloid Leukemia	cytarabine + daunorubicin cytarabine + idarubicin cytarabine + mitoxantrone gemtuzumab ozogamicin + chemotherapy	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO  
Public data sources included in diagnostic significance: NCCN, ESMO

### Prevalent cancer biomarkers without relevant evidence based on included data sources

*BCOR p.(R810\*) c.2428C>T, NRAS p.(G12D) c.35G>A, EZH2 c.2110+1G>A, KRAS p.(K117N) c.351A>C, TET2 p.(N281\*) c.840\_841insT, KRAS p.(A146T) c.436G>A, NRAS p.(G12S) c.34G>A, NRAS p.(G13C) c.37G>T, TET2 p.(S1087\*) c.3260\_3269delCTTCAGAAAA*

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
NRAS	p.(G13C)	c.37G>T	COSM570	chr1:115258745	4.46%	NM_002524.5	missense	1996
NRAS	p.(G12D)	c.35G>A	COSM564	chr1:115258747	12.30%	NM_002524.5	missense	2000
NRAS	p.(G12S)	c.34G>A	COSM563	chr1:115258748	5.20%	NM_002524.5	missense	2000
TET2	p.(N281*)	c.840_841insT	.	chr4:106155938	47.17%	NM_001127208.2	nonsense	1993
TET2	p.(S1087*)	c.3260_3269delCTTCAGAAAA	.	chr4:106158358	45.15%	NM_001127208.2	nonsense	1980
EZH2	p.(?)	c.2110+1G>A	.	chr7:148506401	46.65%	NM_004456.5	unknown	2000
KRAS	p.(A146T)	c.436G>A	COSM19404	chr12:25378562	8.30%	NM_033360.4	missense	2000
KRAS	p.(K117N)	c.351A>C	COSM19940	chr12:25378647	7.45%	NM_033360.4	missense	2000
CEBPA	p.(G123Efs*37)	c.368delG	.	chr19:33792952	45.04%	NM_004364.4	frameshift Deletion	897
BCOR	p.(R810*)	c.2428C>T	.	chrX:39932171	88.40%	NM_001123385.2	nonsense	1810
EZH2	p.(R732T)	c.2195G>C	.	chr7:148506163	47.90%	NM_004456.5	missense	2000
SH2B3	p.(H541Y)	c.1621C>T	.	chr12:111885999	48.72%	NM_005475.3	missense	1995

## Biomarker Descriptions

### BCOR (BCL6 corepressor)

Background: The BCOR gene encodes the B-cell CLL/lymphoma 6 (BCL6) corepressor protein which potentiates transcriptional repression by BCL6<sup>1,2</sup>. BCOR also associates with class I and II histone deacetylases (HDACs) suggesting an alternate mechanism for BCOR mediated transcriptional repression independent of BCL6<sup>2</sup>. Genetic alterations in BCOR result in protein dysfunction which suggests BCOR functions as a tumor suppressor gene<sup>3,4,5</sup>.

## Biomarker Descriptions (continued)

**Alterations and prevalence:** Genetic alterations in BCOR include missense, nonsense, and frameshift mutations that result in loss of function and have been observed in up to 5% of myelodysplastic syndromes (MDS), 5-10% of chronic myelomonocytic leukemia (CMML), and 1-5% of acute myeloid leukemia (AML)<sup>6,7,8,9</sup>. Higher mutational frequencies are reported in some solid tumors, including up to 15% of uterine cancer and 5-10% of colorectal cancer, stomach cancer, cholangiocarcinoma, and melanoma. Although less common, BCOR fusions and internal tandem duplications (ITDs) have been reported in certain rare cancer types<sup>10,11,12</sup>. Specifically, BCOR-CCNB3 rearrangements define a particular subset of sarcomas with Ewing sarcoma-like morphology known as BCOR-CCNB3 sarcomas (BCS)<sup>13,14</sup>. In clear cell carcinoma of the kidney, a rare pediatric renal malignant tumor, one study described the presence of BCOR ITDs in more than 90% of cases<sup>10</sup>.

**Potential relevance:** BCOR rearrangement, including inv(X)(p11.4p11.22) resulting in BCOR-CCNB3 fusion, is diagnostic of sarcoma with BCOR genetic alterations, a subset of undifferentiated round cell sarcomas<sup>15,16</sup>. Additionally, translocation t(x;22)(p11;q13) resulting in ZC3H7B-BCOR fusion is a useful ancillary diagnostic marker of high-grade endometrial stromal sarcoma<sup>15</sup>. Nonsense, frameshift, and splice site mutations in BCOR are associated with poor prognosis in myelodysplastic syndromes<sup>6,7</sup>. In FLT3-ITD negative AML patients under 65 with intermediate cytogenetic prognosis, mutations in BCOR confer inferior overall survival (OS) as well as relapse free survival (RFS) compared to those without BCOR abnormalities (OS= 13.6% vs. 55%; RFS= 14.3% vs. 44.5%)<sup>9</sup>.

### CEBPA (CCAAT enhancer binding protein alpha)

**Background:** The CEBPA gene encodes the enhancer binding protein alpha, a member of the basic region leucine zipper family of transcription factors that recognizes the CCAAT promoter<sup>17</sup>. CEBPA gives rise to two protein isoforms— p42 and p30, where p30 is the shorter isoform lacking the N-terminal 117 amino acids that is present in p42. Both isoforms contain the basic leucine zipper (bZip) domain involved in hetero/homo-dimerization with other CEBP family members and are required for DNA binding<sup>17</sup>. CEBPA is a tumor suppressor gene that plays a critical role in the development of granulocytes<sup>17</sup>. Specifically, CEBPA can influence the expression of granulocyte colony-stimulating factor (G-CSF) and interleukin 6 (IL-6), which are required for neutrophil maturation<sup>18,19</sup>. CEBPA also directly interacts and inhibits cell cycle kinases, including CDK2 and CDK4, thereby hindering cell proliferation<sup>20</sup>. CEBPA is the target of monoallelic or biallelic mutations leading to a loss of function, which can promote the development of cancers such as acute myeloid leukemia (AML)<sup>21</sup>. Germline mutations in CEBPA are also frequent among AML patients and are associated with predisposition to the disease<sup>22,23</sup>.

**Alterations and prevalence:** Mutations in CEBPA are reported in 6-18% of all AML cases<sup>8,24,25,26</sup>. In AML, CEBPA mutations are observed to occur as either monoallelic (single mutant) or bi-allelic (double mutant)<sup>26,27,28</sup>. Biallelic CEBPA mutations are heterozygous and occur as a specific combination of an N-terminal frameshift on one allele and a C-terminal in frame mutation on the other, referred to as an N/C mutant<sup>28,29</sup>. Frameshift mutations result in the N-terminal truncation of approximately 120 amino acids while preserving the remaining 300 amino acids that are initiated further downstream<sup>29</sup>. C-terminal in-frame mutations disrupt the bZip domain which interferes with DNA binding and hetero/homo-dimerization with other CEBP family members. Specifically, N/C mutants possess one N-terminal truncated allele coding for the p30 isoform while the other allele codes for either p30 or p42 isoforms harboring C-terminal mutations<sup>28</sup>.

**Potential relevance:** Single mutations located in the basic leucine zipper (bZIP) region of the gene (smbZIP-CEBPA) as well as biallelic CEBPA mutations are recognized as a diagnostic entity for AML with CEBPA mutation by the World Health Organization (WHO)<sup>30</sup>. Both smbZIP-CEBPA and biallelic CEBPA mutations are associated with a favorable prognosis in AML<sup>31</sup>.

### EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit)

**Background:** The EZH2 gene encodes the enhancer of zeste homolog 2 protein, a histone methyltransferase that functions as both a transcriptional suppressor and co-activator<sup>32</sup>. EZH2 mediates methylation of histone H3 at Lys 27 (H3K27me3) and promotes tumor growth and metastasis through regulation of the cell cycle<sup>32,33</sup>. Since EZH2 loss-of-function is associated with the development of cancer, it is considered a tumor suppressor. EZH2 is overexpressed in various cancer types, consequently, it can also function as an oncogene<sup>32</sup>.

**Alterations and prevalence:** Diverse EZH2 alterations including missense, nonsense, frameshift mutations, and inactivating deletions are observed in 18-25% of T-cell acute lymphocytic leukemia (T-ALL), 3-13% of myeloproliferative neoplasms (MPN), 8-12% of myelodysplastic/myeloproliferative neoplasms overlap disorders (MDS/MPN), and 6% of diverse MDS<sup>33,34</sup>. Heterozygous gain-of-function mutations at tyrosine 641 (Y641) are observed in 22% of germinal center B-cell (GBC) and diffuse large B-cell lymphoma (DLBCL), and 7-17% of follicular lymphoma (FL)<sup>33,35</sup>. In solid tumors, EZH2 mutations are observed in up to 8% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma, and 3% of cholangiocarcinoma<sup>8,36</sup>. Amplifications are observed in up to 7% of ovarian carcinoma<sup>8,36</sup>. Increased EZH2 copy number corresponds with enhanced protein expression and is observed in over 50% of hormone-refractory prostate cancers<sup>37</sup>.

**Potential relevance:** The methyltransferase inhibitor tazemetostat<sup>38</sup> was FDA approved (2020) for EZH2 mutated relapsed or refractory follicular lymphoma after at least 2 prior systemic therapies. Tazemetostat was also granted FDA fast track designation in 2016

## Biomarker Descriptions (continued)

for DLBCL harboring EZH2 activating mutations<sup>39</sup>. EZH2 nonsense or frameshift mutations are independently associated with poor prognosis in MDS and MDS/MPN<sup>6</sup>. EZH2 mutations confer poor prognosis in essential thrombocythemia (ET) and primary myelofibrosis (PMF)<sup>40</sup>. EZH2 overexpression correlates with malignancy, poor prognosis, and poor survival, and has been detected in MDS and acute myeloid leukemia (AML)<sup>32,41</sup>. Several studies have shown EZH2 overexpression enhances chemoresistance in solid tumor types<sup>42,43</sup>.

### KRAS (KRAS proto-oncogene, GTPase)

**Background:** The KRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes NRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>44,45,46</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. KRAS mutations are observed in up to 10-20% of uterine cancer, 30-35% of lung adenocarcinoma and colorectal cancer, and about 60% of pancreatic cancer<sup>8</sup>. The majority of KRAS mutations consist of point mutations occurring at G12, G13, and Q61<sup>8,47,48</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>36,49</sup>.

**Potential relevance:** The FDA has approved the small molecule inhibitors, sotorasib<sup>50</sup> (2021) and adagrasib<sup>51</sup> (2022), for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC). The FDA has also granted breakthrough therapy designation (2022) to the KRAS G12C inhibitor, GDC-6036<sup>52</sup>, for KRAS G12C mutation in non-small cell lung cancer. The small molecular inhibitor, RO-5126766, was granted breakthrough designation (2021) alone for KRAS G12V mutant non-small cell lung cancer or in combination with defactinib, for KRAS mutant endometrial carcinoma and KRAS G12V mutant non-small cell lung cancer<sup>53</sup>. The PLK1 inhibitor, onvansertib<sup>54</sup>, was granted fast track designation (2020) in combination with bevacizumab and FOLFIRI for second-line treatment of patients with KRAS-mutated metastatic colorectal cancer (mCRC). Additionally, the SHP2 inhibitor, BBP-398<sup>55</sup> was granted fast track designation (2022) in combination with sotorasib for previously treated patients with KRAS G12C-mutated metastatic NSCLC. The EGFR antagonists, cetuximab<sup>56</sup> and panitumumab<sup>57</sup>, are contraindicated for treatment of colorectal cancer patients with KRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>49</sup>. Additionally, KRAS mutations are associated with poor prognosis in NSCLC<sup>58</sup>.

### NRAS (NRAS proto-oncogene, GTPase)

**Background:** The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>44,45,46</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers<sup>8,59</sup>. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61<sup>8,60</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>36,49</sup>.

**Potential relevance:** Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab<sup>56</sup> and panitumumab<sup>57</sup>, are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>49</sup>. The FDA has granted fast track designation to the pan-RAF inhibitor, KIN-2787<sup>61</sup>, for the treatment of NRAS mutation positive metastatic or unresectable melanoma. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome<sup>6</sup> as well as melanoma<sup>62</sup>. In a phase III clinical trial in patients with advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively<sup>63</sup>.

### TET2 (tet methylcytosine dioxygenase 2)

**Background:** TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3<sup>64</sup>. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to 5-hydroxymethylcytosine<sup>65,66</sup>. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded  $\beta$ -helix domain (DSBH)<sup>67</sup>. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies<sup>64,65,66</sup>.

**Alterations and prevalence:** Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)<sup>6</sup>. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies<sup>65,68</sup>. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

## Biomarker Descriptions (continued)

Potential relevance: The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations<sup>40,69</sup>. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia<sup>40,70</sup>

## Relevant Therapy Summary

☒ In this cancer type   ☐ In other cancer type   ☒ In this cancer type and other cancer types   ☒ No evidence

### CEBPA p.(G123Efs\*37) c.368delG

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cytarabine + daunorubicin	×	●	×	×	×
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×

## Relevant Therapy Details

### Current NCCN Information

☒ In this cancer type   ☐ In other cancer type   ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-03-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### CEBPA p.(G123Efs\*37) c.368delG

#### ● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**CEBPA p.(G123Efs\*37) c.368delG (continued)****● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**CEBPA p.(G123Efs\*37) c.368delG (continued)****● gemtuzumab ozogamicin + cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

Other criteria: CD33 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## Diagnostic Details

### Current ESMO Information

ESMO information is current as of 2023-03-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

**CEBPA p.(G123Efs\*37) c.368delG**

### Diagnostic significance: Acute Myeloid Leukemia

**Variant class:** CEBPA mutation

**Diagnostic notes:**

- CEBPA biallelic mutations; AML with recurrent genetic abnormalities; WHO classification of AML

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]



## References

1. Gearhart et al. Polycomb group and SCF ubiquitin ligases are found in a novel BCOR complex that is recruited to BCL6 targets. *Mol. Cell. Biol.* 2006 Sep;26(18):6880-9. PMID: 16943429
2. Huynh et al. BCoR, a novel corepressor involved in BCL-6 repression. *Genes Dev.* 2000 Jul 15;14(14):1810-23. PMID: 10898795
3. Kelly et al. Bcor loss perturbs myeloid differentiation and promotes leukaemogenesis. *Nat Commun.* 2019 Mar 22;10(1):1347. PMID: 30902969
4. Cao et al. BCOR regulates myeloid cell proliferation and differentiation. *Leukemia.* 2016 May;30(5):1155-65. PMID: 26847029
5. Yamamoto et al. Clarifying the impact of polycomb complex component disruption in human cancers. *Mol. Cancer Res.* 2014 Apr;12(4):479-84. PMID: 24515802
6. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
7. Damm et al. BCOR and BCORL1 mutations in myelodysplastic syndromes and related disorders. *Blood.* 2013 Oct 31;122(18):3169-77. PMID: 24047651
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Terada et al. Usefulness of BCOR gene mutation as a prognostic factor in acute myeloid leukemia with intermediate cytogenetic prognosis. *Genes Chromosomes Cancer.* 2018 Aug;57(8):401-408. PMID: 29663558
10. Wong et al. Clear cell sarcomas of the kidney are characterised by BCOR gene abnormalities, including exon 15 internal tandem duplications and BCOR-CCNB3 gene fusion. *Histopathology.* 2018 Jan;72(2):320-329. PMID: 28833375
11. Cramer et al. Successful Treatment of Recurrent Primitive Myxoid Mesenchymal Tumor of Infancy With BCOR Internal Tandem Duplication. *J Natl Compr Canc Netw.* 2017 Jul;15(7):868-871. PMID: 28687574
12. Peters et al. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. *Mod. Pathol.* 2015 Apr;28(4):575-86. PMID: 25360585
13. Puls et al. BCOR-CCNB3 (Ewing-like) sarcoma: a clinicopathologic analysis of 10 cases, in comparison with conventional Ewing sarcoma. *Am. J. Surg. Pathol.* 2014 Oct;38(10):1307-18. PMID: 24805859
14. Kao et al. BCOR-CCNB3 Fusion Positive Sarcomas: A Clinicopathologic and Molecular Analysis of 36 Cases With Comparison to Morphologic Spectrum and Clinical Behavior of Other Round Cell Sarcomas. *Am. J. Surg. Pathol.* 2018 May;42(5):604-615. PMID: 29300189
15. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2022]
16. NCCN Guidelines® - NCCN-Bone Cancer [Version 2.2023]
17. Pabst et al. Complexity of CEBPA dysregulation in human acute myeloid leukemia. *Clin. Cancer Res.* 2009 Sep 1;15(17):5303-7. PMID: 19706798
18. Zhang et al. Upregulation of interleukin 6 and granulocyte colony-stimulating factor receptors by transcription factor CCAAT enhancer binding protein alpha (C/EBP alpha) is critical for granulopoiesis. *J. Exp. Med.* 1998 Sep 21;188(6):1173-84. PMID: 9743535
19. Heath et al. C/EBPalpha deficiency results in hyperproliferation of hematopoietic progenitor cells and disrupts macrophage development in vitro and in vivo. *Blood.* 2004 Sep 15;104(6):1639-47. PMID: 15073037
20. Wang et al. C/EBPalpha arrests cell proliferation through direct inhibition of Cdk2 and Cdk4. *Mol. Cell.* 2001 Oct;8(4):817-28. PMID: 11684017
21. Lourenço et al. A tumor suppressor role for C/EBPa in solid tumors: more than fat and blood. *Oncogene.* 2017 Sep 14;36(37):5221-5230. PMID: 28504718
22. Pabst et al. Somatic CEBPA mutations are a frequent second event in families with germline CEBPA mutations and familial acute myeloid leukemia. *J. Clin. Oncol.* 2008 Nov 1;26(31):5088-93. PMID: 18768433
23. Tawana et al. Disease evolution and outcomes in familial AML with germline CEBPA mutations. *Blood.* 2015 Sep 3;126(10):1214-23. PMID: 26162409
24. Benthous et al. Rapid and sensitive screening for CEBPA mutations in acute myeloid leukaemia. *Br. J. Haematol.* 2008 Oct;143(2):230-9. PMID: 18752591
25. Su et al. Mutational spectrum of acute myeloid leukemia patients with double CEBPA mutations based on next-generation sequencing and its prognostic significance. *Oncotarget.* 2018 May 18;9(38):24970-24979. PMID: 29861846
26. Dufour et al. Acute myeloid leukemia with biallelic CEBPA gene mutations and normal karyotype represents a distinct genetic entity associated with a favorable clinical outcome. *J. Clin. Oncol.* 2010 Feb 1;28(4):570-7. PMID: 20038735
27. Green et al. Prognostic significance of CEBPA mutations in a large cohort of younger adult patients with acute myeloid leukemia: impact of double CEBPA mutations and the interaction with FLT3 and NPM1 mutations. *J. Clin. Oncol.* 2010 Jun 1;28(16):2739-47. PMID: 20439648

## References (continued)

28. Avellino et al. Expression and regulation of C/EBPα in normal myelopoiesis and in malignant transformation. *Blood*. 2017 Apr 13;129(15):2083-2091. PMID: 28179278
29. Pabst et al. Transcriptional dysregulation during myeloid transformation in AML. *Oncogene*. 2007 Oct 15;26(47):6829-37. PMID: 17934489
30. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022 Jul;36(7):1703-1719. PMID: 35732831
31. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
32. Gan et al. Epigenetic regulation of cancer progression by EZH2: from biological insights to therapeutic potential. 6:10. PMID: 29556394
33. Yamagishi et al. Targeting EZH2 in cancer therapy. *Curr Opin Oncol*. 2017 Sep;29(5):375-381. PMID: 28665819
34. Li. EZH2: novel therapeutic target for human cancer. 4:1. PMID: 25520914
35. Morin et al. Somatic mutations altering EZH2 (Tyr641) in follicular and diffuse large B-cell lymphomas of germinal-center origin. *Nat. Genet*. 2010 Feb;42(2):181-5. PMID: 20081860
36. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
37. Saramäki et al. The gene for polycomb group protein enhancer of zeste homolog 2 (EZH2) is amplified in late-stage prostate cancer. *Genes Chromosomes Cancer*. 2006 Jul;45(7):639-45. PMID: 16575874
38. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/213400s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/213400s000lbl.pdf)
39. <https://www.healio.com/news/hematology-oncology/20161129/fda-grants-fast-track-designation-to-tazemetostat-for-diffuse-large-bcell-lymphoma>
40. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
41. Sauvageau et al. Polycomb group proteins: multi-faceted regulators of somatic stem cells and cancer. *Cell Stem Cell*. 2010 Sep 3;7(3):299-313. PMID: 20804967
42. Gardner et al. Chemosensitive Relapse in Small Cell Lung Cancer Proceeds through an EZH2-SLFN11 Axis. *Cancer Cell*. 2017 Feb 13;31(2):286-299. PMID: 28196596
43. Fan et al. Inhibition of EZH2 reverses chemotherapeutic drug TMZ chemosensitivity in glioblastoma. *Int J Clin Exp Pathol*. 2014;7(10):6662-70. PMID: 25400745
44. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer*. 2011 Oct 13;11(11):761-74. PMID: 21993244
45. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol*. 2008 Jul;9(7):517-31. PMID: 18568040
46. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J*. 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
47. Román et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. *Mol Cancer*. 2018 Feb 19;17(1):33. doi: 10.1186/s12943-018-0789-x. PMID: 29455666
48. Dinu et al. Prognostic significance of KRAS gene mutations in colorectal cancer—preliminary study. *J Med Life*. 2014 Oct-Dec;7(4):581-7. PMID: 25713627
49. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol*. 2016 Jan 10;34(2):179-85. PMID: 26438111
50. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/214665s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/214665s003lbl.pdf)
51. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/216340Orig1s000Corrected\\_lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/216340Orig1s000Corrected_lbl.pdf)
52. <https://assets.cwp.roche.com/f/126832/x/5738a7538b/irp230202.pdf>
53. <https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-reports-third-quarter-2022-financial-results>
54. <https://cardiffoncology.investorroom.com/2020-05-28-Cardiff-Oncology-Announces-Fast-Track-Designation-Granted-by-the-FDA-to-Onvansertib-for-Second-Line-Treatment-of-KRAS-Mutated-Colorectal-Cancer>
55. <https://bridgebio.com/news/bridgebio-pharma-announces-first-lung-cancer-patient-dosed-in-phase-1-2-trial-and-us-fda-fast-track-designation-for-shp2-inhibitor-bbp-398-in-combination-with-amgens-lumakras-sotorasib/>
56. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)
57. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125147s210lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf)

## References (continued)

58. Slebos et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *N. Engl. J. Med.* 1990 Aug 30;323(9):561-5. PMID: 2199829
59. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE.* 2011;6(7):e22769. PMID: 21829508
60. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res.* 2013 May 1;19(9):2584-91. PMID: 23515407
61. <https://investors.kinnate.com/news-releases/news-release-details/kinnate-biopharma-inc-receives-fast-track-designation-us-food>
62. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol.* 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
63. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2017 Apr;18(4):435-445. PMID: 28284557
64. Pan et al. The TET2 interactors and their links to hematological malignancies. *IUBMB Life.* 2015 Jun;67(6):438-45. PMID: 26099018
65. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature.* 2010 Dec 9;468(7325):839-43. PMID: 21057493
66. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. *Leukemia.* 2014 Mar;28(3):485-96. PMID: 24220273
67. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. *Exp. Mol. Med.* 2017 Apr 28;49(4):e323. PMID: 28450733
68. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood.* 2009 Oct 8;114(15):3285-91. PMID: 19666869
69. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
70. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood.* 2014 Apr 3;123(14):2220-8. PMID: 24478400