



Sample Information

Patient Name: 陳雪美
Gender: Female
ID No.: Q201104496
History No.: 18096770
Age: 84

Ordering Doctor: DOC1751J 蕭樑材
Ordering REQ.: H45NC3L
Signing in Date: 2023/03/03

Path No.: M112-00035
MP No.: MY23011
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/02/21

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	3 Relevant Biomarkers
Biomarker Descriptions	3	13 Therapies Available
Relevant Therapy Summary	4	1 Clinical Trials
Relevant Therapy Details	5	
Prognostic Details	13	
Clinical Trials Summary	13	

Relevant Myelodysplastic Syndrome Variants

Gene	Finding	Gene	Finding
ASXL1	None detected	NPM1	None detected
BCOR	None detected	NRAS	None detected
CBL	None detected	NUP214	None detected
CREBBP	None detected	RUNX1	RUNX1 p.(E88*) c.262G>T
ETV6	None detected	SF3B1	None detected
EZH2	None detected	SRSF2	None detected
FLT3	None detected	STAG2	STAG2 p.(S174Nfs*11) c.518_519insAAATC
GATA2	None detected	TP53	None detected
IDH2	IDH2 p.(R140Q) c.419G>A	U2AF1	None detected
KIT	None detected	WT1	None detected
KMT2A	None detected	ZRSR2	None detected

Relevant Myelodysplastic Syndrome Variants (continued)

Gene	Finding	Gene	Finding
MECOM	None detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>IDH2</i> p.(R140Q) c.419G>A isocitrate dehydrogenase (NADP(+)) 2 Allele Frequency: 32.80% Prognostic significance: NCCN: Poor	None	enasidenib ¹ azacitidine decitabine venetoclax + chemotherapy	1
IA	<i>RUNX1</i> p.(E88*) c.262G>T RUNX family transcription factor 1 Allele Frequency: 32.75% Prognostic significance: NCCN: Poor	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine gemtuzumab ozogamicin + chemotherapy venetoclax + chemotherapy	0
IA	<i>STAG2</i> p.(S174Nfs*11) c.518_519insAAATC stromal antigen 2 Allele Frequency: 35.49% Prognostic significance: NCCN: Poor	None	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
IDH2	p.(R140Q)	c.419G>A	COSM41590	chr15:90631934	32.80%	NM_002168.4	missense	2000
RUNX1	p.(E88*)	c.262G>T	.	chr21:36259229	32.75%	NM_001754.4	nonsense	2000
STAG2	p.(S174Nfs*11)	c.518_519insAAATC	.	chrX:123179062	35.49%	NM_001042749.2	frameshift Insertion	1995
CEBPA	p.(A303P)	c.907G>C	.	chr19:33792414	36.82%	NM_004364.4	missense	1999
STAG2	p.(S173=)	c.519C>T	.	chrX:123179070	36.49%	NM_001042749.2	synonymous	1998

Biomarker Descriptions

IDH2 (isocitrate dehydrogenase (NADP(+)) 2)

Background: The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to α -ketoglutarate (α -KG)¹. The IDH1 gene encodes the NADP⁺ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

Alterations and prevalence: Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)². Recurrent IDH2 variants include predominately R140Q and R172K plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity³. Although wild-type enzymatic activity is ablated, recurrent IDH2 variants catalyze the conversion of α -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair^{1,4}. Recurrent IDH2 mutations are present in 10-20% of patients with AML and 5% of patients with MDS^{5,6,7}.

Potential relevance: Enasidenib⁸ is FDA approved (2017) for the treatment of AML patients with IDH2 R140G/L/Q/W and R172G/K/M/S/W mutations. In AML, acquired resistance to enasidenib has been associated with the emergence of Q316E or I319M mutations⁹. IDH2 R172 and R140Q variants are associated with poor prognosis in MDS but have been shown to confer improved prognosis in lower grade gliomas^{10,11,12}. Additionally, IDH2 mutations are associated with inferior overall survival in polycythemia vera (PV) and essential thrombocythemia (ET) as well as inferior leukemia-free survival in primary myelofibrosis (PMF)¹³.

RUNX1 (RUNX family transcription factor 1)

Background: The RUNX1 gene encodes the runt-related transcription factor (RUNX) 1, part of the RUNX family of transcription factors which also includes RUNX2 and RUNX3¹⁴. All RUNX proteins share several conserved regions with similar functionality including a highly conserved N-terminal 'runt' domain responsible for binding DNA, a C-terminal region composed of an activation domain, inhibitory domain, protein interacting motifs, and a nuclear matrix targeting signal¹⁵. Each of these proteins are capable of interacting with core binding factor beta (CBF β) to form the core binding factor (CBF) complex. Consequently, RUNX1, RUNX2, and RUNX3 are collectively known as core binding factor alpha (CBF α) since they can each function as the alpha subunit of CBF. Specifically, CBF β binds to the 'runt' domain of RUNX1 leading to RUNX1 stabilization and increased affinity of the CBF complex for promoters involved in hematopoietic differentiation and cell cycle regulation^{16,17}. RUNX1 is frequently mutated in various hematological malignancies¹⁷. Germline mutations in RUNX1 result in a rare autosomal dominant condition known as familial platelet disorder, with predisposition to acute myeloid leukemia (FPD/AML)^{18,19}. Somatic mutations and chromosomal translocations in RUNX1 are often observed in myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), and chronic myelomonocytic leukemia (CMML)¹⁷.

Alterations and prevalence: RUNX1 is frequently rearranged in hematological malignancies with over 50 different observed translocations²⁰. The most recurrent translocation, t(12;21)(q34;q11), results in ETV6-RUNX1 fusion and is observed in 20-25% of childhood ALL^{21,22,23}. This translocation is also observed in adult ALL at a lower frequency (2%)^{22,23}. Another recurrent translocation, t(8;21)(q22;q22), results in RUNX1-RUNX1T1 fusion and is observed in 5-10% of AML²⁴. The RUNX1-RUNX1T1 fusion, consists of the RHD domain of RUNX1 and the majority of RUNX1T1, which promotes oncogenesis by altering transcriptional regulation of RUNX1 target genes^{17,24}. Somatic mutations in RUNX1 include missense, nonsense, and frameshift mutations resulting in loss of function or dominant negative effects¹⁷. RUNX1 mutations are reported in approximately 10% of de novo AML as well as 10-15% of MDS^{10,17,25,26}.

Potential relevance: AML with RUNX1-RUNX1T1 fusions is considered a distinct molecular subtype by the World Health Organization (WHO)^{25,27}. Translocations involving RUNX1, specifically t(8;21)(q22;q22)/RUNX1-RUNX1T1 in AML and t(12;21)(q34;q11)/ETV6-RUNX1 in ALL, are associated with favorable risk^{25,28}. On the other hand, mutations in RUNX1 confer poor prognosis in AML, MDS, and systemic mastocytosis (SM)^{10,25,29}.

STAG2 (stromal antigen 2)

Background: The STAG2 gene encodes the stromal antigen 2 protein, one of the core proteins in the cohesin complex, which regulates the separation of sister chromatids during cell division^{30,31}. Components of the cohesin complex include SMC1A, SMC3, and RAD21, which bind to STAG1/STAG2 paralogs^{32,33}. Inactivating mutations in STAG2 contribute to X-linked neurodevelopmental disorders, aneuploidy, and chromosomal instability in cancer^{32,34}.

Alterations and prevalence: Somatic mutations in STAG2 include nonsense, frameshift, splice site variants¹⁰. Somatic mutations in STAG2 are observed in various solid tumors including 14% of bladder cancer, 10% of uterine cancer, 3% of stomach cancer, and 4% of lung adenocarcinoma³⁵. In addition, mutations in STAG2 are observed in 5-10% of myelodysplastic syndrome (MDS), 3% of acute myeloid leukemia, and 2% of diffuse large B-cell lymphoma^{10,35}.

Biomarker Descriptions (continued)

Potential relevance: Nonsense, frameshift, and splice site STAG2 mutations are associated with poor prognosis in MDS¹⁰. Truncating mutations in STAG2 lead to a loss of function in bladder cancer and are often identified as an early event associated with low grade and stage tumors³⁶.

Relevant Therapy Summary

● In this cancer type
 ○ In other cancer type
 ● In this cancer type and other cancer types
 ✕ No evidence

IDH2 p.(R140Q) c.419G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
enasidenib	○	○	✕	○	✕
azacitidine	✕	○	✕	✕	✕
decitabine	✕	○	✕	✕	✕
venetoclax + azacitidine	✕	○	✕	✕	✕
venetoclax + cytarabine	✕	○	✕	✕	✕
venetoclax + decitabine	✕	○	✕	✕	✕
LY-3410738	✕	✕	✕	✕	● (I)

RUNX1 p.(E88*) c.262G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✕	○	✕	✕	✕
azacitidine	✕	○	✕	✕	✕
cytarabine	✕	○	✕	✕	✕
cytarabine + daunorubicin	✕	○	✕	✕	✕
cytarabine + daunorubicin + etoposide	✕	○	✕	✕	✕
cytarabine + etoposide + idarubicin	✕	○	✕	✕	✕
cytarabine + fludarabine + idarubicin + filgrastim	✕	○	✕	✕	✕
cytarabine + idarubicin	✕	○	✕	✕	✕
cytarabine + mitoxantrone	✕	○	✕	✕	✕
decitabine	✕	○	✕	✕	✕
gemtuzumab ozogamicin + cytarabine + daunorubicin	✕	○	✕	✕	✕
venetoclax + azacitidine	✕	○	✕	✕	✕
venetoclax + cytarabine	✕	○	✕	✕	✕
venetoclax + decitabine	✕	○	✕	✕	✕

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

- ☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

FDA information is current as of 2023-01-18. For the most up-to-date information, search www.fda.gov.

IDH2 p.(R140Q) c.419G>A

☐ enasidenib

Cancer type: Acute Myeloid Leukemia

Label as of: 2020-11-24

Variant class: IDH2 R140Q mutation

Indications and usage:

IDHIFA® is an isocitrate dehydrogenase-2 inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA-approved test.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf

Current NCCN Information

- ☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-01-03. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

IDH2 p.(R140Q) c.419G>A

☐ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

☐ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

IDH2 p.(R140Q) c.419G>A (continued)**○ venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

RUNX1 p.(E88*) c.262G>T**○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

RUNX1 p.(E88*) c.262G>T (continued)**○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

RUNX1 p.(E88*) c.262G>T (continued)**○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

RUNX1 p.(E88*) c.262G>T (continued)**○ venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

RUNX1 p.(E88*) c.262G>T (continued)**○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

○ cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Current ESMO Information

☒ In this cancer type ☐ In other cancer type ☐ In this cancer type and other cancer types

ESMO information is current as of 2023-01-03. For the most up-to-date information, search www.esmo.org.

IDH2 p.(R140Q) c.419G>A

☐ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2023-01-03. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

IDH2 p.(R140Q) c.419G>A

Prognostic significance: NCCN: Poor

Cancer type: Myelodysplastic Syndrome

Variant class: IDH2 R140Q mutation

NCCN Recommendation category: 2A

Summary:

- NCCN Guidelines® associate the biomarker with poor prognosis

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

RUNX1 p.(E88*) c.262G>T

Prognostic significance: NCCN: Poor

Cancer type: Myelodysplastic Syndrome

Variant class: RUNX1 truncating mutation

NCCN Recommendation category: 2A

Summary:

- NCCN Guidelines® independently associate the biomarker with poor prognosis

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

STAG2 p.(S174Nfs*11) c.518_519insAAATC

Prognostic significance: NCCN: Poor

Cancer type: Myelodysplastic Syndrome

Variant class: STAG2 truncating mutation

NCCN Recommendation category: 2A

Summary:

- NCCN Guidelines® associate the biomarker with poor prognosis

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

Clinical Trials Summary

IDH2 p.(R140Q) c.419G>A

NCT ID	Title	Phase
NCT04603001	A Phase I Study of Oral LY3410738 in Patients With Advanced Hematologic Malignancies With IDH1 or IDH2 Mutations.	I

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene*. 2018 Apr;37(15):1949-1960. PMID: 29367755
2. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
3. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature*. 2009 Dec 10;462(7274):739-44. PMID: 19935646
4. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell*. 2010 Mar 16;17(3):225-34. PMID: 20171147
5. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
6. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia*. 2011 Feb;25(2):246-53. PMID: 21079611
7. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
8. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf
9. Intlekofer et al. Acquired resistance to IDH inhibition through trans or cis dimer-interface mutations. *Nature*. 2018 Jul;559(7712):125-129. PMID: 29950729
10. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
11. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med*. 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
12. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology*. 2010 Oct 26;75(17):1560-6. PMID: 20975057
13. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
14. de et al. Runx transcription factors in the development and function of the definitive hematopoietic system. *Blood*. 2017 Apr 13;129(15):2061-2069. PMID: 28179276
15. Chuang et al. RUNX family: Regulation and diversification of roles through interacting proteins. *Int. J. Cancer*. 2013 Mar 15;132(6):1260-71. PMID: 23180629
16. Jung et al. Prognostic factor analysis in core-binding factor-positive acute myeloid leukemia. *Anticancer Res*. 2014 Feb;34(2):1037-45. PMID: 24511052
17. Sood et al. Role of RUNX1 in hematological malignancies. *Blood*. 2017 Apr 13;129(15):2070-2082. PMID: 28179279
18. Béri-Dexheimer et al. Clinical phenotype of germline RUNX1 haploinsufficiency: from point mutations to large genomic deletions. *Eur. J. Hum. Genet.* 2008 Aug;16(8):1014-8. PMID: 18478040
19. Hayashi et al. Myeloid neoplasms with germ line RUNX1 mutation. *Int. J. Hematol.* 2017 Aug;106(2):183-188. PMID: 28534116
20. De et al. RUNX1 translocations and fusion genes in malignant hemopathies. *Future Oncol.* 2011 Jan;7(1):77-91. PMID: 21174539
21. De et al. ETV6 fusion genes in hematological malignancies: a review. *Leuk. Res.* 2012 Aug;36(8):945-61. PMID: 22578774
22. Pui et al. Acute lymphoblastic leukemia. *N. Engl. J. Med.* 2004 Apr 8;350(15):1535-48. PMID: 15071128
23. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
24. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D920-4. PMID: 23161685
25. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
26. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
27. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016 May 19;127(20):2391-405. PMID: 27069254
28. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 1.2022]
29. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
30. Mehta et al. Cohesin: functions beyond sister chromatid cohesion. *FEBS Lett.* 2013 Aug 2;587(15):2299-312. PMID: 23831059
31. Aquila et al. The role of STAG2 in bladder cancer. *Pharmacol. Res.* 2018 May;131:143-149. PMID: 29501732
32. Mullegama et al. De novo loss-of-function variants in STAG2 are associated with developmental delay, microcephaly, and congenital anomalies. *Am. J. Med. Genet. A.* 2017 May;173(5):1319-1327. PMID: 28296084

References (continued)

33. van et al. Synthetic lethality between the cohesin subunits STAG1 and STAG2 in diverse cancer contexts. *Elife*. 2017 Jul 10;6. PMID: 28691904
34. Solomon et al. Mutational inactivation of STAG2 causes aneuploidy in human cancer. *Science*. 2011 Aug 19;333(6045):1039-43. PMID: 21852505
35. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
36. Solomon et al. Frequent truncating mutations of STAG2 in bladder cancer. *Nat. Genet*. 2013 Dec;45(12):1428-30. PMID: 24121789