



Sample Information

Patient Name: 賴蘇美慧
Gender: Female
ID No.: A202003913
History No.: 47994752
Age: 81

Ordering Doctor: DOC6266E 徐千富
Ordering REQ.: 0CJTEHR
Signing in Date: 0CJSKKW

Path No.: M112-00077
MP No.: MY23024
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/04/17

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	2 Relevant Biomarkers
Biomarker Descriptions	2	8 Therapies Available
Relevant Therapy Summary	3	1 Clinical Trials
Relevant Therapy Details	4	
Prognostic Details	10	
Diagnostic Details	11	
Clinical Trials Summary	11	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	CEBPA p.(S190*) c.569C>A	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	IDH2 p.(R140Q) c.419G>A	RUNX1	None detected
KMT2A	None detected	TP53	None detected

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>IDH2</i> p.(R140Q) c.419G>A isocitrate dehydrogenase (NADP(+)) 2 Allele Frequency: 20.40%	enasidenib ¹ azacitidine decitabine venetoclax + chemotherapy	None	1
IA	<i>CEBPA</i> p.(S190*) c.569C>A CCAAT enhancer binding protein alpha Allele Frequency: 25.42% Prognostic significance: ELN 2017: Favorable Diagnostic significance: Acute Myeloid Leukemia	cytarabine + daunorubicin cytarabine + idarubicin cytarabine + mitoxantrone gemtuzumab ozogamicin + chemotherapy	None	0

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

STAG2 p.(W102*) c.305G>A

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
IDH2	p.(R140Q)	c.419G>A	COSM41590	chr15:90631934	20.40%	NM_002168.4	missense	2000
CEBPA	p.(S190*)	c.569C>A	.	chr19:33792752	25.42%	NM_004364.4	nonsense	531
STAG2	p.(W102*)	c.305G>A	.	chrX:123171393	46.42%	NM_001042749.2	nonsense	1999
IKZF1	p.(S88=)	c.264G>A	.	chr7:50444334	47.60%	NM_006060.6	synonymous	1998

Biomarker Descriptions

CEBPA (CCAAT enhancer binding protein alpha)

Background: The CEBPA gene encodes the enhancer binding protein alpha, a member of the basic region leucine zipper family of transcription factors that recognizes the CCAAT promoter¹. CEBPA gives rise to two protein isoforms— p42 and p30, where p30 is the shorter isoform lacking the N-terminal 117 amino acids that is present in p42. Both isoforms contain the basic leucine zipper (bZip) domain involved in hetero/homo-dimerization with other CEBP family members and are required for DNA binding¹. CEBPA is a tumor suppressor gene that plays a critical role in the development of granulocytes¹. Specifically, CEBPA can influence the expression of granulocyte colony-stimulating factor (G-CSF) and interleukin 6 (IL-6), which are required for neutrophil maturation^{2,3}. CEBPA also directly interacts and inhibits cell cycle kinases, including CDK2 and CDK4, thereby hindering cell proliferation⁴. CEBPA is the target of monoallelic or biallelic mutations leading to a loss of function, which can promote the development of cancers such as acute myeloid leukemia (AML)⁵. Germline mutations in CEBPA are also frequent among AML patients and are associated with predisposition to the disease^{6,7}.

Alterations and prevalence: Mutations in CEBPA are reported in 6-18% of all AML cases^{8,9,10,11}. In AML, CEBPA mutations are observed to occur as either monoallelic (single mutant) or bi-allelic (double mutant)^{11,12,13}. Biallelic CEBPA mutations are heterozygous and occur as a specific combination of an N-terminal frameshift on one allele and a C-terminal in frame mutation on the other, referred to as an N/C mutant^{13,14}. Frameshift mutations result in the N-terminal truncation of approximately 120 amino acids while preserving the remaining 300 amino acids that are initiated further downstream¹⁴. C-terminal in-frame mutations disrupt the bZip domain which interferes with DNA binding and hetero/homo-dimerization with other CEBP family members. Specifically, N/C mutants possess one

Biomarker Descriptions (continued)

N-terminal truncated allele coding for the p30 isoform while the other allele codes for either p30 or p42 isoforms harboring C-terminal mutations¹³.

Potential relevance: Biallelic CEBPA mutations are recognized as a diagnostic entity for AML by the World Health Organization (WHO)¹⁵. Biallelic CEBPA mutations are associated with favorable risk and improved prognosis in AML^{11,15,16,17,18}.

IDH2 (isocitrate dehydrogenase (NADP(+)) 2)

Background: The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to α -ketoglutarate (α -KG)¹⁹. The IDH1 gene encodes the NADP+ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

Alterations and prevalence: Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)²⁰. Recurrent IDH2 variants include predominately R140Q and R172K plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity²¹. Although wild-type enzymatic activity is ablated, recurrent IDH2 variants catalyze the conversion of α -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair^{19,22}. Recurrent IDH2 mutations are present in 10-20% of patients with AML and 5% of patients with MDS^{23,24,25}.

Potential relevance: Enasidenib²⁶ is FDA approved (2017) for the treatment of AML patients with IDH2 R140G/L/Q/W and R172G/K/M/S/W mutations. In AML, acquired resistance to enasidenib has been associated with the emergence of Q316E or I319M mutations²⁷. IDH2 R172 and R140Q variants are associated with poor prognosis in MDS but have been shown to confer improved prognosis in lower grade gliomas^{28,29,30}. Additionally, IDH2 mutations are associated with inferior overall survival in polycythemia vera (PV) and essential thrombocythemia (ET) as well as inferior leukemia-free survival in primary myelofibrosis (PMF)³¹.

STAG2 (stromal antigen 2)

Background: The STAG2 gene encodes the stromal antigen 2 protein, one of the core proteins in the cohesin complex, which regulates the separation of sister chromatids during cell division^{32,33}. Components of the cohesion complex include SMC1A, SMC3, and RAD21, which bind to STAG1/STAG2 paralogs^{34,35}. Inactivating mutations in STAG2 contribute to X-linked neurodevelopmental disorders, aneuploidy, and chromosomal instability in cancer^{34,36}.

Alterations and prevalence: Somatic mutations in STAG2 include nonsense, frameshift, splice site variants²⁸. Somatic mutations in STAG2 are observed in various solid tumors including 14% of bladder cancer, 10% of uterine cancer, 3% of stomach cancer, and 4% of lung adenocarcinoma³⁷. In addition, mutations in STAG2 are observed in 5-10% of myelodysplastic syndrome (MDS), 3% of acute myeloid leukemia, and 2% of diffuse large B-cell lymphoma^{28,37}.

Potential relevance: Nonsense, frameshift, and splice site STAG2 mutations are associated with poor prognosis in MDS²⁸. Truncating mutations in STAG2 lead to a loss of function in bladder cancer and are often identified as an early event associated with low grade and stage tumors³⁸.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

IDH2 p.(R140Q) c.419G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
enasidenib	●	●	×	●	×
azacitidine	×	●	×	×	×
decitabine	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

IDH2 p.(R140Q) c.419G>A (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + azacitidine	×	●	×	×	×
venetoclax + cytarabine	×	●	×	×	×
venetoclax + decitabine	×	●	×	×	×
LY-3410738	×	×	×	×	● (I)

CEBPA p.(S190*) c.569C>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cytarabine + daunorubicin	×	●	×	×	×
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2023-01-18. For the most up-to-date information, search www.fda.gov.

IDH2 p.(R140Q) c.419G>A

● enasidenib

Cancer type: Acute Myeloid Leukemia

Label as of: 2020-11-24

Variant class: IDH2 R140Q mutation

Indications and usage:

IDHIFA® is an isocitrate dehydrogenase-2 inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA-approved test.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-01-03. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

IDH2 p.(R140Q) c.419G>A

● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

IDH2 p.(R140Q) c.419G>A (continued)**● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

CEBPA p.(S190*) c.569C>A**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

CEBPA p.(S190*) c.569C>A (continued)**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

● gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

CEBPA p.(S190*) c.569C>A (continued)**● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Current ESMO Information

- ☒ In this cancer type ☐ In other cancer type ☐ In this cancer type and other cancer types

ESMO information is current as of 2023-01-03. For the most up-to-date information, search www.esmo.org.

IDH2 p.(R140Q) c.419G>A

☒ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2023-01-03. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

CEBPA p.(S190*) c.569C>A

Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

NCCN Recommendation category: 2A

Summary:

- Genetic Abnormality: Biallelic mutated CEBPA.

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

Current ESMO Information

ESMO information is current as of 2023-01-03. For the most up-to-date information, search www.esmo.org.

CEBPA p.(S190*) c.569C>A

Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: CEBPA mutation

Summary:

- Double-mutant CEBPA

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Diagnostic Details

Current ESMO Information

ESMO information is current as of 2023-01-03. For the most up-to-date information, search www.esmo.org.

CEBPA p.(S190*) c.569C>A

Diagnostic significance: Acute Myeloid Leukemia

Variant class: CEBPA mutation

Diagnostic notes:

- CEBPA biallelic mutations; AML with recurrent genetic abnormalities; WHO classification of AML

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Clinical Trials Summary

IDH2 p.(R140Q) c.419G>A

NCT ID	Title	Phase
NCT04603001	A Phase I Study of Oral LY3410738 in Patients With Advanced Hematologic Malignancies With IDH1 or IDH2 Mutations.	I

References

1. Pabst et al. Complexity of CEBPA dysregulation in human acute myeloid leukemia. *Clin. Cancer Res.* 2009 Sep 1;15(17):5303-7. PMID: 19706798
2. Zhang et al. Upregulation of interleukin 6 and granulocyte colony-stimulating factor receptors by transcription factor CCAAT enhancer binding protein alpha (C/EBP alpha) is critical for granulopoiesis. *J. Exp. Med.* 1998 Sep 21;188(6):1173-84. PMID: 9743535
3. Heath et al. C/EBPalph deficiency results in hyperproliferation of hematopoietic progenitor cells and disrupts macrophage development in vitro and in vivo. *Blood.* 2004 Sep 15;104(6):1639-47. PMID: 15073037
4. Wang et al. C/EBPalph arrests cell proliferation through direct inhibition of Cdk2 and Cdk4. *Mol. Cell.* 2001 Oct;8(4):817-28. PMID: 11684017
5. Lourenço et al. A tumor suppressor role for C/EBPa in solid tumors: more than fat and blood. *Oncogene.* 2017 Sep 14;36(37):5221-5230. PMID: 28504718
6. Pabst et al. Somatic CEBPA mutations are a frequent second event in families with germline CEBPA mutations and familial acute myeloid leukemia. *J. Clin. Oncol.* 2008 Nov 1;26(31):5088-93. PMID: 18768433
7. Tawana et al. Disease evolution and outcomes in familial AML with germline CEBPA mutations. *Blood.* 2015 Sep 3;126(10):1214-23. PMID: 26162409
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Benthous et al. Rapid and sensitive screening for CEBPA mutations in acute myeloid leukaemia. *Br. J. Haematol.* 2008 Oct;143(2):230-9. PMID: 18752591
10. Su et al. Mutational spectrum of acute myeloid leukemia patients with double CEBPA mutations based on next-generation sequencing and its prognostic significance. *Oncotarget.* 2018 May 18;9(38):24970-24979. PMID: 29861846
11. Dufour et al. Acute myeloid leukemia with biallelic CEBPA gene mutations and normal karyotype represents a distinct genetic entity associated with a favorable clinical outcome. *J. Clin. Oncol.* 2010 Feb 1;28(4):570-7. PMID: 20038735
12. Green et al. Prognostic significance of CEBPA mutations in a large cohort of younger adult patients with acute myeloid leukemia: impact of double CEBPA mutations and the interaction with FLT3 and NPM1 mutations. *J. Clin. Oncol.* 2010 Jun 1;28(16):2739-47. PMID: 20439648
13. Avellino et al. Expression and regulation of C/EBPa in normal myelopoiesis and in malignant transformation. *Blood.* 2017 Apr 13;129(15):2083-2091. PMID: 28179278
14. Pabst et al. Transcriptional dysregulation during myeloid transformation in AML. *Oncogene.* 2007 Oct 15;26(47):6829-37. PMID: 17934489
15. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
16. Kühnl et al. Molecular markers in acute myeloid leukaemia. *Int. J. Hematol.* 2012 Aug;96(2):153-63. PMID: 22791509
17. Barjesteh et al. Biallelic mutations in the CEBPA gene and low CEBPA expression levels as prognostic markers in intermediate-risk AML. *Hematol. J.* 2003;4(1):31-40. PMID: 12692518
18. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
19. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene.* 2018 Apr;37(15):1949-1960. PMID: 29367755
20. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
21. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature.* 2009 Dec 10;462(7274):739-44. PMID: 19935646
22. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell.* 2010 Mar 16;17(3):225-34. PMID: 20171147
23. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
24. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia.* 2011 Feb;25(2):246-53. PMID: 21079611
25. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
26. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf

References (continued)

27. Intlekofer et al. Acquired resistance to IDH inhibition through trans or cis dimer-interface mutations. *Nature*. 2018 Jul;559(7712):125-129. PMID: 29950729
28. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
29. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med*. 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
30. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology*. 2010 Oct 26;75(17):1560-6. PMID: 20975057
31. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
32. Mehta et al. Cohesin: functions beyond sister chromatid cohesion. *FEBS Lett*. 2013 Aug 2;587(15):2299-312. PMID: 23831059
33. Aquila et al. The role of STAG2 in bladder cancer. *Pharmacol. Res*. 2018 May;131:143-149. PMID: 29501732
34. Mullegama et al. De novo loss-of-function variants in STAG2 are associated with developmental delay, microcephaly, and congenital anomalies. *Am. J. Med. Genet. A*. 2017 May;173(5):1319-1327. PMID: 28296084
35. van et al. Synthetic lethality between the cohesin subunits STAG1 and STAG2 in diverse cancer contexts. *Elife*. 2017 Jul 10;6. PMID: 28691904
36. Solomon et al. Mutational inactivation of STAG2 causes aneuploidy in human cancer. *Science*. 2011 Aug 19;333(6045):1039-43. PMID: 21852505
37. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
38. Solomon et al. Frequent truncating mutations of STAG2 in bladder cancer. *Nat. Genet*. 2013 Dec;45(12):1428-30. PMID: 24121789