

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE	Brain glioblastoma (GBM)	PHYSICIAN	ORDERING PHYSICIAN	Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE	Brain
	NAME	Li, Yu-Ching		MEDICAL FACILITY	Taipei Veterans General Hospital		SPECIMEN ID	S111-20614 C
	DATE OF BIRTH	31 December 1984		ADDITIONAL RECIPIENT	None		SPECIMEN TYPE	Slide Deck
	SEX	Male		MEDICAL FACILITY ID	205872		DATE OF COLLECTION	24 May 2022
	MEDICAL RECORD #	48535291		PATHOLOGIST	Not Provided		SPECIMEN RECEIVED	08 June 2022

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 0 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

IDH1 R132H

ARID1A splice site 3539+1G>A

ATM rearrangement intron 58

ATRX L764fs*2, Q193* - subclonal[†]

CIC R215W, S146fs*1, R1214* - subclonal[†]

MLL2 Q3910_Q3911del

NOTCH2 NOTCH2- ALG6 rearrangement

SETD2 K583fs*17 - subclonal[†]

TP53 M237I

2 Disease relevant genes with no reportable alterations: **EGFR**, **PDGFRA**

[†] See About the Test in appendix for details.

Report Highlights

- Variants with **diagnostic implications** that may indicate a specific cancer type: **ATRX** L764fs*2, Q193* (p. 7), **IDH1** R132H (p. 4)
- Targeted therapies with potential clinical benefit **approved in another tumor type**: Ivosidenib (p. 11)
- Variants that may inform **nontargeted treatment approaches** (e.g., chemotherapy) in this tumor type: **IDH1** R132H (p. 4)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 12)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **IDH1** R132H (p. 4)
- Variants that may represent **clonal hematopoiesis** and may originate from non-tumor sources: **MLL2** Q3910_Q3911del (p. 8)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 0 Muts/Mb

GENOMIC FINDINGS

IDH1 - R132H

10 Trials see p. 16

ARID1A - splice site 3539+1G>A

8 Trials see p. 12

ATM - rearrangement intron 58

10 Trials see p. 14

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
none	Ivosidenib
none	none
none	none

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

MLL2 - Q3910_Q3911del p. [8](#)

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

ATRX - L764fs*2, Q193* - subclonal	p. 7	NOTCH2 - NOTCH2- ALG6 rearrangement	p. 8
CIC - R215W, S146fs*1, R1214* - subclonal	p. 7	SETD2 - K583fs*17 - subclonal	p. 9
MLL2 - Q3910_Q3911del	p. 8	TP53 - M237I	p. 10

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

Low-level MSI has been reported in 5-9% of glioblastoma (GBM) samples⁶⁻⁸. A large-scale study did not find high-level microsatellite instability (MSI-H) in any of 129 GBM samples⁶, although a small-scale study reported MSI-H in 4 of 15 pediatric GBMs and 1 of 12 adult GBMs⁹. The frequency of MSI has been reported to be increased in relapsed compared to primary GBM⁶, in GBMs with a previous lower grade astrocytoma⁷, and in giant cell GBM compared to classic GBM⁸.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁰⁻¹². This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹³⁻¹⁵. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{10,12,14-15}.

BIOMARKER

Tumor Mutational Burden

RESULT

0 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁶⁻¹⁸, anti-PD-1 therapies¹⁶⁻¹⁹, and combination nivolumab and ipilimumab²⁰⁻²⁵. In glioma, a lack of association between TMB and clinical benefit from immune checkpoint inhibitors has been reported^{16,26-27}. However, multiple case studies have reported that patients with ultramutated gliomas driven by POLE

mutations have benefited from treatment with anti-PD-1²⁸⁻²⁹ or anti-PD-L1³⁰ therapies. Therefore, although increased TMB alone may not be a strong biomarker for PD-1 or PD-L1 inhibitors in this cancer type, these agents may have efficacy for patients with glioma harboring both high TMB and POLE mutation.

FREQUENCY & PROGNOSIS

Glioblastoma (GBM) harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 4.2% of cases have high TMB (>20 mut/Mb)³¹. For pediatric patients, high TMB has been reported in a subset of high-grade gliomas, frequently in association with mutations in mismatch repair or proofreading genes and in TP53, whereas BRAF alterations or other oncogene fusions were observed more frequently in brain tumors harboring low TMB³²⁻³³. Increased TMB has been reported to correlate with higher tumor grade in glioma³⁴ and glioblastoma (GBM) tissue samples with biallelic mismatch repair deficiency

(bMMRD)²⁸, as well as with shorter OS of patients with diffuse glioma³⁵.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁶⁻³⁷ and cigarette smoke in lung cancer³⁸⁻³⁹, treatment with temozolomide-based chemotherapy in glioma⁴⁰⁻⁴¹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁴²⁻⁴⁶, and microsatellite instability (MSI)^{42,45-46}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{16,26-30}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

IDH1

ALTERATION

R132H

TRANSCRIPT ID

NM_005896

CODING SEQUENCE EFFECT

395G>A

VARIANT ALLELE FREQUENCY (% VAF)

39.6%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

IDH1 mutations that lead to production of 2-HG, most commonly R132 alterations, may predict sensitivity to IDH1-mutation-specific inhibitors such as ivosidenib⁴⁷. A Phase 1b/2 study of the IDH1 inhibitor olutasidenib for patients with IDH1-mutated glioma reported a DCR of 50% (n=24) with 1 PR⁴⁸. A Phase 1 study of the pan-IDH1/IDH2 inhibitor vorasidenib for patients with IDH1- or IDH2-mutated glioma reported an ORR of 18.2% (4/22; RANO criteria) and median PFS of 31.4 months for non-enhancing cases and median PFS of 7.5 months for the overall glioma population (n=52)⁴⁹. Preclinical studies suggested that IDH1 neomorphic mutations may also confer sensitivity to PARP inhibitors⁵⁰⁻⁵³. In a Phase 1 trial of the PD-L1 inhibitor atezolizumab in

patients with glioblastoma (GBM), 2 of the 3 patients with IDH1-mutant tumors experienced clinical benefit (1 PR and 1 long-term SD; the third patient experienced short-term SD), whereas none of the 8 patients with IDH1-wild-type GBM experienced benefit (8/8 PD); significantly longer PFS and a trend toward longer OS were observed in the patients with IDH1-mutated tumors compared to the patients with IDH1-wild-type tumors³⁰. Preclinical data indicate that IDH1-mutated glioma may be sensitive to the glutaminase inhibitor telaglenastat in combination with radiotherapy⁵⁴.

— Nontargeted Approaches —

IDH1/2 mutations are associated with improved survival outcomes for patients with glioma treated with radiation or alkylating chemotherapy (NCCN CNS Cancers Guidelines, v2.2021).

FREQUENCY & PROGNOSIS

IDH1 mutation is characteristic of low-grade gliomas and secondary glioblastoma, and is relatively rare in primary glioblastoma⁵⁵⁻⁵⁹. In the TCGA datasets, IDH1 mutation has been found in 77% of lower grade glioma cases and in 5% of glioblastoma cases⁶⁰⁻⁶¹. IDH1/2 mutations are a strong favorable prognostic marker for OS in Grade 2-3 glioma, particularly in combination with 1p/19q codeletion (NCCN CNS Cancers Guidelines, v2.2021). Several studies have found IDH1 mutations to be associated with improved

prognosis and longer PFS and OS in patients with various types of glioma including anaplastic astrocytoma and GBM^{59,62-68}.

FINDING SUMMARY

The isocitrate dehydrogenases IDH1 and IDH2 encode highly homologous enzymes that are involved in the citric acid (TCA) cycle and other metabolic processes, playing roles in normal cellular metabolism and in protection against oxidative stress and apoptosis⁶⁹. R132 is located within the active site of IDH1 and is a hotspot for mutations in cancer⁶⁹⁻⁷³. Substitutions at IDH1 R132 alter the enzymatic activity of IDH1, resulting in the production of the oncometabolite, D-2-hydroxyglutarate (2-HG)⁷¹⁻⁷⁵, which promotes tumorigenesis^{71,76-79}.

POTENTIAL DIAGNOSTIC IMPLICATIONS

IDH mutation in the absence of TERT mutation is suggestive of astrocytoma (NCCN CNS Cancers Guidelines, v2.2021)⁸⁰. IDH1/2 mutation is associated with Grade 2 and 3 astrocytomas and oligodendrogliomas, and distinguishes secondary glioblastoma (GBM) from primary GBM (NCCN CNS Cancers Guidelines, v2.2021). ATRX mutations often co-occur with IDH1/2 mutations and may be indicative of Grade 2-3 astrocytoma or secondary glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v2.2021)⁸⁰⁻⁸¹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

ARID1A

ALTERATION

splice site 3539+1G>A

TRANSCRIPT ID

NM_006015

CODING SEQUENCE EFFECT

3539+1G>A

VARIANT ALLELE FREQUENCY (% VAF)

34.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies approved to address the mutation or loss of ARID1A in cancer. However, on the basis of limited clinical and preclinical evidence, ARID1A inactivating mutations may lead to sensitivity to ATR inhibitors such as M6620 and ceralasertib⁸². In a Phase 2 study of ceralasertib in solid tumors, 2 patients with endometrial carcinoma in the cohort with loss of ARID1A expression achieved CRs on ceralasertib monotherapy; at least 1 of these 2 patients carried an inactivating ARID1A mutation. In contrast, no responses were observed for patients with normal ARID1A expression treated with ceralasertib combined with olaparib⁸³. One patient with small cell lung cancer harboring an ARID1A mutation

experienced a PR when treated with M6620 combined with topotecan⁸⁴. In a Phase 1 trial, a patient with metastatic colorectal cancer harboring both an ARID1A mutation and ATM loss treated with single-agent M6620 achieved a CR that was ongoing at 29 months⁸⁵. On the basis of limited preclinical evidence from studies in ovarian cancer, ARID1A inactivation may predict sensitivity to EZH2 inhibitors⁸⁶⁻⁸⁷, which are under investigation in clinical trials. Other studies have reported that the loss of ARID1A may activate the PI3K-AKT pathway and be linked with sensitivity to inhibitors of this pathway⁸⁸⁻⁹⁰. Patients with ARID1A alterations in advanced or metastatic solid tumors may derive benefit from treatment with anti-PD-1 or anti-PD-L1 immunotherapy⁹¹. Loss of ARID1A expression has been associated with chemoresistance to platinum-based therapy for patients with ovarian clear cell carcinoma⁹²⁻⁹³ and to 5-fluorouracil in colorectal cancer cell lines⁹⁴.

FREQUENCY & PROGNOSIS

ARID1A alterations are particularly prevalent in ovarian clear cell carcinoma (46-50%), ovarian and uterine endometrioid carcinomas (24-44%), and cholangiocarcinoma (27%); they are also reported in up to 27% of gastric carcinoma, esophageal adenocarcinoma, Waldenstrom macroglobulinemia, pediatric Burkitt lymphoma, hepatocellular carcinoma, colorectal carcinoma,

and urothelial carcinoma samples analyzed (COSMIC, cBioPortal, Jan 2022)⁹⁵⁻¹⁰³. ARID1A loss is associated with microsatellite instability in ovarian and endometrial endometrioid adenocarcinomas^{91,104-107}, CRC^{91,108-110}, and gastric cancer^{91,111-115}. ARID1A protein loss is associated with tumors of poor histological grade for many tumor types, including colorectal cancer (CRC)¹⁰⁸⁻¹¹⁰, cervical cancer¹¹⁶⁻¹¹⁷, gastric cancer¹¹¹⁻¹¹⁵, urothelial carcinoma¹¹⁸⁻¹²⁰, ovarian and endometrial cancers^{93,104-107,121-125}, breast carcinoma¹²⁶⁻¹²⁸, and clear cell renal cell carcinoma¹²⁹; ARID1A mutation has been associated with poor outcomes for patients with cholangiocarcinoma¹³⁰⁻¹³³. However, prognostic data regarding patient survival are often mixed and conflicting.

FINDING SUMMARY

ARID1A encodes the AT-rich interactive domain-containing protein 1A, also known as Baf250a, a member of the SWI/SNF chromatin remodeling complex. Mutation, loss, or inactivation of ARID1A has been reported in many cancers, and the gene is considered a tumor suppressor^{99,114,127,134-139}. ARID1A mutations, which are mostly truncating, have been identified along the entire gene and often correlate with ARID1A protein loss^{99,112,135-136,140}, whereas ARID1A missense mutations are mostly uncharacterized.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

ATM

ALTERATION

rearrangement intron 58

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Loss of functional ATM results in a defective DNA damage response and homologous recombination-mediated DNA repair and may predict sensitivity to PARP inhibitors¹⁴¹⁻¹⁴². Clinical responses have been reported for patients with ATM-mutated prostate cancer treated with PARP inhibitors¹⁴³⁻¹⁴⁵ and PARP inhibitors have shown limited clinical benefit for patients with other ATM-mutated solid tumors including pancreatic cancer¹⁴⁶⁻¹⁴⁷, colorectal cancer¹⁴⁸, papillary renal cell carcinoma¹⁴⁹, ovarian cancer¹⁵⁰, small cell bowel cancer,¹⁴⁷ and biliary tract cancer¹⁵¹. In Phase 1 trials of ATR inhibitors, a heavily pretreated patient with colorectal cancer who achieved a CR to berzosertib⁴⁹ and 4 out of 4 patients with diverse solid tumors who achieved PRs to BAY1895344¹⁵² harbored ATM inactivation or protein loss; studies showing reduced cell viability

and increased DNA damage in preclinical models of solid tumors¹⁵³⁻¹⁵⁵ and hematologic malignancies^{153,156} also support the increased sensitivity of ATM-deficient cells to ATR inhibitors. Preclinical experiments also indicate that loss of ATM causes dependency on DNA-PKcs in cancer cells; DNA-PKcs inhibitors promoted apoptosis in ATM-deficient cells and were active in a lymphoma mouse model lacking ATM activity¹⁵⁷.

FREQUENCY & PROGNOSIS

ATM mutations have been reported in 1% of glioblastoma cases^{61,158}. An analysis of Grades 1-4 astrocytomas detected upregulation of ATM mRNA expression¹⁵⁹. Higher expression of the ATM protein correlated with longer overall and progression-free survival in a study of 69 glioblastoma cases¹⁶⁰. However, ATM gene and protein expression in glioblastoma cells has also been correlated with resistance to radiation therapy; glioma cells with lower ATM expression have been found to be more sensitive to radiation than cells with higher ATM expression¹⁶¹⁻¹⁶³. In addition, one study reported that the ATM kinase inhibitor KU-60019 inhibited glioma cell growth and invasion and sensitized cells to radiation¹⁶³. Preclinical studies have also shown that ATM

kinase inactivation and ATM inhibitors such as LY294002 and KU-55933 increase the sensitivity of glioblastoma cells to cisplatin and temozolomide¹⁶⁴⁻¹⁶⁶.

FINDING SUMMARY

ATM encodes the protein ataxia telangiectasia mutated, which is a serine/threonine protein kinase that plays a key role in the DNA damage response¹⁶⁷. Loss of functional ATM promotes tumorigenesis¹⁶⁸. Alterations such as seen here may disrupt ATM function or expression¹⁶⁹⁻¹⁷¹.

POTENTIAL GERMLINE IMPLICATIONS

ATM mutation carriers have increased cancer risk, with female carriers displaying a 38% lifetime risk of breast cancer¹⁷². Biallelic mutations in ATM underlie the rare autosomal-recessive inherited disorder ataxia-telangiectasia (A-T), also referred to as genome instability or DNA damage response syndrome¹⁷³. This disease is characterized by genomic instability, sensitivity to DNA-damaging agents, and increased risk of developing cancer^{167,173}. The prevalence of A-T is estimated at 1:40,000 to 1:100,000 worldwide¹⁷³. In the appropriate clinical context, germline testing of ATM is recommended.

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

ATRX

ALTERATION

L764fs*2, Q193* - subclonal

TRANSCRIPT ID

NM_000489, NM_000489

CODING SEQUENCE EFFECT

2290_2291delTT, 577C>T

VARIANT ALLELE FREQUENCY (% VAF)

80.1%, 8.5%

strand break-induction with agents such as doxorubicin, irinotecan, and topotecan¹⁷⁸; however, these approaches have not been demonstrated clinically.

FREQUENCY & PROGNOSIS

Somatic mutation of ATRX has been reported in a number of solid tumor types, often associated with ALT¹⁷⁹. ATRX mutation correlating with ALT has been reported in 10–20% of pancreatic neuroendocrine tumors (PNETs)^{179–181}, 12.6% of pheochromocytomas and paragangliomas¹⁸², and 48% of adolescent and young adult (AYA) patients with glioblastoma (GBM) or neuroblastoma^{183–187}. ATRX loss in PNET^{180,188} and melanoma¹⁸⁹ and mutation in other neuroendocrine tumors¹⁸² is associated with poor prognosis. Pediatric patients with high-grade glioma and ATRX mutation were shown to have more aggressive disease but are more responsive to treatment with double-strand break therapy¹⁷⁸. ATRX mutation or loss of expression is more frequent in Grade 2/3 astrocytoma and secondary GBM than primary GBM, oligodendroglioma, and oligoastrocytoma^{190–193} and has been proposed as a distinguishing biomarker^{191–193}. ATRX mutation has not been detected in concurrence with MYCN amplification in glioma and neuroblastoma^{184–187}. Low-grade gliomas with both IDH1/2 mutation and ATRX mutation are associated with worse prognosis than those with IDH1/2 mutation but

no ATRX mutation¹⁹¹. Loss of ATRX protein expression has been reported in 33–39% of incidences of leiomyosarcoma (LMS) associating with ALT, a poor prognostic factor across all LMS subtypes, and with poor prognosis in extrauterine LMS but not in uterine LMS^{194–195}.

FINDING SUMMARY

ATRX encodes a SWI/SNF chromatin remodeling protein implicated in histone variant H3.3 deposition, transcriptional regulation, and telomere maintenance^{196–197}. ATRX inactivation or loss of expression is associated with alternative lengthening of telomeres (ALT)^{179,195,198–199}. Alterations that disrupt the ADD domain (aa 167–270) or helicase domain (aa 2010–2280) of ATRX are predicted to result in loss of ATRX function^{200–202}; however, the loss of ATRX function is not sufficient to induce ALT, which requires other undetermined factors^{196,203}. Germline mutations in ATRX give rise to alpha-thalassemia X-linked intellectual disability syndrome (ATR-X syndrome)²⁰⁴.

POTENTIAL DIAGNOSTIC IMPLICATIONS

ATRX mutations often co-occur with IDH1/2 mutations and may be indicative of Grade 2–3 astrocytoma or secondary glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v2.2021)^{80–81}.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

No targeted therapies are available to directly address ATRX inactivation. Based on preclinical^{174–175} and limited clinical data¹⁷⁶, ATRX alterations may confer sensitivity to combination strategies involving WEE1 inhibition. In a Phase 2 study evaluating the WEE1 inhibitor adavosertib plus irinotecan for the treatment of pediatric patients with neuroblastoma, prolonged SD was reported for 44% (4/9) of patients with ATRX-deficient tumors and responses were seen in two tumors that had evidence of ALT¹⁷⁶. Preclinical evidence also suggests that ATRX deficiency may impart sensitivity to synthetic lethal approaches involving PARP inhibition and irinotecan¹⁷⁷, combined PARP and ATR inhibition¹⁷⁵, or double-

GENE

CIC

ALTERATION

R215W, S146fs*1, R1214* - subclonal

TRANSCRIPT ID

NM_015125, NM_015125, NM_015125

CODING SEQUENCE EFFECT

643C>T, 436_437delAG, 3640C>T

VARIANT ALLELE FREQUENCY (% VAF)

17.4%, 35.8%, 7.7%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in CIC.

FREQUENCY & PROGNOSIS

CIC mutations have been described in various solid tumors, including 1–10% of sequenced gastric, endometrial, and colorectal carcinomas and melanoma tumors (cBioPortal, COSMIC, Jan 2022)^{95–97}, although the consequences of CIC mutations in these tumor types have not been studied. CIC mutations have been observed in 58–69% of oligodendrogliomas but are less

common in other gliomas, such as astrocytoma or oligoastrocytoma^{205–207}. Published data investigating the prognostic implications of CIC alterations are generally limited (PubMed, Jun 2022). Conflicting data have been reported regarding the prognostic significance of CIC mutation in oligodendroglioma^{206,208–209}.

FINDING SUMMARY

CIC encodes a transcriptional repressor that plays a role in central nervous system (CNS) development²¹⁰. CIC inactivation has been reported in various malignancies, and is highly recurrent in oligodendroglioma^{205–206}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

MLL2

ALTERATION

Q3910_Q3911del

TRANSCRIPT ID

NM_003482

CODING SEQUENCE EFFECT

11729_11734delAGCAAC

VARIANT ALLELE FREQUENCY (% VAF)

44.5%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in MLL2.

FREQUENCY & PROGNOSIS

One study detected MLL2 mutation in 19% of

IDH-wildtype glioblastoma (GBM) samples analyzed²¹¹. MLL2 alterations are observed in a number of solid tumor contexts (COSMIC, Jan 2022)⁹⁵, and are especially prevalent in lung squamous cell carcinoma (SCC)²¹² and small cell lung carcinoma (SCLC)²¹³. MLL2 mutation was found to be an independent prognostic factor of poor PFS and OS in non-small cell lung cancer, but not in SCLC²¹⁴. One study reported that MLL2 truncating mutations were more common in recurrent ovary granulosa cell tumors (GCT) compared with primary GCTs (24% [10/42] vs. 3.0% [1/32])²¹⁵. In a study of esophageal SCC, high MLL2 expression positively correlated with tumor stage, differentiation, and size, and negatively correlated with OS²¹⁶.

FINDING SUMMARY

MLL2 encodes an H3K4-specific histone methyltransferase that is involved in the transcriptional response to progesterone signaling²¹⁷. Germline de novo mutations of MLL2

are responsible for the majority of cases of Kabuki syndrome, a complex and phenotypically distinctive developmental disorder²¹⁸. A significant number of inactivating MLL2 alterations have been observed in multiple tumor types, suggesting a tumor suppressor role²¹⁹.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²²⁰⁻²²⁵. Comprehensive genomic profiling of solid tumors may detect nontumor alterations that are due to CH^{224,226-227}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

GENE

NOTCH2

ALTERATION

NOTCH2- ALG6 rearrangement

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Several approaches for inhibiting NOTCH2 signaling have been developed, including neutralizing NOTCH antibodies such as tarextumab (OMP-59R5)²²⁸, which targets NOTCH2 and NOTCH3, and pan-NOTCH inhibitors, such as gamma-secretase inhibitors (GSI). In a Phase 2 study, the GSI AL101 (BMS-906024) elicited PR in 15% (6/39) and SD in 54% (21/39) of patients with metastatic adenoid cystic carcinoma harboring NOTCH activating alterations²²⁹. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

FREQUENCY & PROGNOSIS

NOTCH2 mutations have been reported in <1% of glioblastomas⁶¹. NOTCH2 mutations have been observed in 3.3% (2/60) of astrocytoma samples analyzed in the COSMIC database (Aug 2021). Notch signaling has been shown in multiple studies to play oncogenic roles in glioblastoma pathogenesis²³⁰⁻²³³. However, while some studies have suggested that NOTCH2 promotes glioma cancer stem cell functions and tumorigenesis^{230,232,234-235}, others have suggested that NOTCH2, unlike NOTCH1, may act as a tumor suppressor in this context²³⁶. Published data investigating the prognostic implications of NOTCH2 alteration in glioma are limited (PubMed, Feb 2022).

FINDING SUMMARY

NOTCH2 encodes a member of the NOTCH family of receptors, which play a role in cell fate determination and various developmental processes. Upon binding of membrane-bound ligands, NOTCH signaling involves gamma-secretase (GS) cleavage of the NOTCH intracellular

domain (NICD), which subsequently forms part of a transcription factor complex that regulates downstream target genes²³⁷⁻²³⁸. Depending on cellular context, NOTCH2 can act as either a tumor suppressor or an oncogene²³⁹⁻²⁴³. NOTCH2 rearrangements and fusions that result in expression of the NOTCH2 intracellular domain (amino acids 1699-2471) without the transmembrane domain (amino acids 1678-1698) are predicted to result in GS-independent activation of NOTCH signaling²⁴⁴. Similar rearrangements leading to the fusion of SEC22B exon 1 to NOTCH2 exons 27-34 have previously been reported in breast cancer²⁴⁴ and in chronic lymphocytic leukemia²⁴⁵ and have been shown to be GS-inhibitor insensitive²⁴⁴. One or more alterations observed here have not been fully characterized, and the effect on NOTCH2 function is unclear; however, it has been reported in the context of cancer, which may indicate biological relevance.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS
GENE

SETD2

ALTERATION

K583fs*17 - subclonal

TRANSCRIPT ID

NM_014159

CODING SEQUENCE EFFECT

1748_1751delAACA

VARIANT ALLELE FREQUENCY (% VAF)

8.2%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in SETD2.

FREQUENCY & PROGNOSIS

Somatic inactivating alterations of SETD2 are documented to occur at low frequency in a number of solid tumors, most commonly in renal carcinoma²⁴⁶. SETD2 has been associated with favorable prognosis in gastric cancer²⁴⁷. SETD2 has also been associated with poor prognosis in RCC and MDS²⁴⁸, while data in other tumor types is limited (PubMed, Jun 2022).

FINDING SUMMARY

SETD2 encodes a histone lysine-36 methyltransferase²⁴⁹ that preferentially interacts with the expanded N-terminal polyglutamine tracts present in mutant huntingtin, implicating it in the pathogenesis of Huntington disease²⁵⁰. SETD2 mRNA expression has been observed to be consistently reduced in breast tumors relative to adjacent non-tumor tissue, suggesting a potential tumor suppressor role²⁵¹. SETD2 alterations such as observed here have been shown to be inactivating²⁵²⁻²⁵⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

M237I

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

711G>C

VARIANT ALLELE FREQUENCY (% VAF)

90.2%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁵⁸⁻²⁶¹, or p53 gene therapy and immunotherapeutics such as SGT-53²⁶²⁻²⁶⁶ and ALT-801²⁶⁷. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²⁶⁸. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁶⁹. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²⁷⁰. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁷¹. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel²⁷². A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations²⁷³. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal

cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²⁷⁴. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁶⁶. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutated p53 such as APR-246²⁷⁵⁻²⁷⁷. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR²⁷⁸. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies^{156,279}; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies²⁸⁰⁻²⁸¹. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

In the TCGA dataset, TP53 alterations have been reported in 35% of glioblastomas (GBMs), with a high incidence in pediatric and secondary GBMs and a low incidence in primary GBMs^{158,282}. One study detected TP53 alterations in 31% (73/232) of IDH-wildtype GBM samples analyzed, with most of the events being mutations²¹¹. TP53 mutations have been reported in 18-40% of astrocytoma samples, and preferentially in anaplastic astrocytoma; one study reported TP53 loss of function and partially/fully functional mutations in 15% and 25% of anaplastic astrocytomas, respectively²⁸³⁻²⁸⁸. Some studies suggest that the presence of a TP53 mutation is correlated with a favorable prognosis in patients with glioblastoma (GBM)²⁸⁹. One study reported that TP53 alterations were associated with poorer OS (12.9 months altered vs. 19.7 months wildtype, HR=1.58, p=0.0054) in IDH-wildtype GBM²¹¹. Mutation of TP53 is thought to be an early step in the tumorigenesis of astrocytomas, which can progress into anaplastic astrocytoma and then glioblastoma through gain of other genetic abnormalities such as loss of CDKN2A or RB1,

followed by loss of PTEN²⁹⁰.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²⁹¹. Alterations such as seen here may disrupt TP53 function or expression²⁹²⁻²⁹⁶.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Mar 2022)²⁹⁷. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²⁹⁸⁻³⁰⁰, including sarcomas³⁰¹⁻³⁰². Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000³⁰³ to 1:20,000³⁰². For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30³⁰⁴. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²²⁰⁻²²⁵. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²²⁰⁻²²¹. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease³⁰⁵. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{224,226-227}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Ivosidenib

Assay findings association
IDH1
R132H

AREAS OF THERAPEUTIC USE

Ivosidenib is an isocitrate dehydrogenase 1 (IDH1) inhibitor that is FDA approved to treat patients with a susceptible IDH1 mutation in relapsed or refractory acute myeloid leukemia (AML) or previously treated locally advanced or metastatic cholangiocarcinoma. It is also approved as a first-line treatment for patients with AML and a susceptible IDH1 mutation who are not eligible for intensive induction chemotherapy or who are ≥ 75 years old. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of extensive clinical evidence in AML³⁰⁶ and

cholangiocarcinoma³⁰⁷⁻³⁰⁸ and limited clinical data in myelodysplastic syndrome (MDS)³⁰⁶ and glioma^{47,309}, IDH1 R132 mutation may confer sensitivity to ivosidenib.

SUPPORTING DATA

In a Phase 1 study of ivosidenib for patients with IDH1-mutated advanced solid tumors, 1 patient achieved PR in the non-enhancing glioma population (ORR=2.9% [1/35]); for patients with non-enhancing glioma and enhancing glioma, SD rates were 85.7% (30/35) and 45.2% (14/31), respectively, and median PFS was 13.6 months and 1.4 months, respectively^{47,309}.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
ARID1A
RATIONALE
 ARID1A loss or inactivation may predict

sensitivity to ATR inhibitors.

ALTERATION
 splice site 3539+1G>A

NCT04768296
PHASE 2

Berzosertib + Topotecan in Relapsed Platinum-Resistant Small-Cell Lung Cancer (DDriver SCLC 250)

TARGETS
 TOP1, ATR

LOCATIONS: Zhejiang (China), Hangzhou (China), Nanjing (China), Wuhan (China), Xi'an (China), Osaka (Japan), Beijing (China), Hirakata-shi (Japan), Takatsuki-shi (Japan), Sichuan (China)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom), Villejuif (France)

NCT04802174
PHASE 1/2

Lurbinectedin With Berzosertib, an ATR Kinase Inhibitor in Small Cell Cancers and High-Grade Neuroendocrine Cancers

TARGETS
 ATR

LOCATIONS: Maryland

NCT04657068
PHASE 1/2

A Study of ART0380 for the Treatment of Advanced or Metastatic Solid Tumors

TARGETS
 ATR

LOCATIONS: London (United Kingdom), Colorado, Oklahoma, Pennsylvania, Tennessee, Florida

NCT02595931
PHASE 1

ATR Kinase Inhibitor VX-970 and Irinotecan Hydrochloride in Treating Patients With Solid Tumors That Are Metastatic or Cannot Be Removed by Surgery

TARGETS
 ATR

LOCATIONS: California, Missouri, Pennsylvania, Massachusetts, Connecticut, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

CLINICAL TRIALS
NCT04514497
PHASE 1

Testing the Addition of an Anti-cancer Drug, BAY 1895344, to Usual Chemotherapy for Advanced Stage Solid Tumors, With a Specific Focus on Patients With Small Cell Lung Cancer, Poorly Differentiated Neuroendocrine Cancer, and Pancreatic Cancer

TARGETS
ATR, TOP1

LOCATIONS: Minnesota, Oklahoma, Pennsylvania, Connecticut, Tennessee, Florida

NCT04266912
PHASE 1/2

Avelumab and M6620 for the Treatment of DDR Deficient Metastatic or Unresectable Solid Tumors

TARGETS
ATR, PD-L1

LOCATIONS: Texas

NCT03669601
PHASE 1

AZD6738 & Gemcitabine as Combination Therapy

TARGETS
ATR

LOCATIONS: Cambridge (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

CLINICAL TRIALS
GENE
ATM
RATIONALE
 Loss or inactivation of ATM may increase sensitivity to PARP inhibitors, ATR inhibitors, or

DNA-PKcs inhibitors.

ALTERATION
 rearrangement intron 58

NCT04768296
PHASE 2

Berzosertib + Topotecan in Relapsed Platinum-Resistant Small-Cell Lung Cancer (DDriver SCLC 250)

TARGETS
 TOP1, ATR

LOCATIONS: Zhejiang (China), Hangzhou (China), Nanjing (China), Wuhan (China), Xi'an (China), Osaka (Japan), Beijing (China), Hirakata-shi (Japan), Takatsuki-shi (Japan), Sichuan (China)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
 PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Nedlands (Australia), Port Macquarie (Australia), Darlinghurst (Australia), Adana (Turkey), Ankara (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom), Villejuif (France)

NCT04644068
PHASE 1/2

Study of AZD5305 as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Malignancies

TARGETS
 ERBB2, TROP2, PARP

LOCATIONS: Seoul (Korea, Republic of), Chuo-ku (Japan), Koto-ku (Japan), Melbourne (Australia), Warszawa (Poland), Gdynia (Poland), Grzegorz (Poland), Budapest (Hungary), Brno (Czechia), Padova (Italy)

NCT04740190
PHASE 2

Talazoparib - Carboplatin for Recurrent High-grade Glioma With DDRd

TARGETS
 PARP

LOCATIONS: Hong Kong (Hong Kong)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

NCT04715620
PHASE 2

Niraparib Combined With Radiotherapy in rGBM

TARGETS
PARP

LOCATIONS: Tianjin (China)

NCT05035745
PHASE 1/2

Selinexor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
XPO1, PARP

LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT05076513
PHASE NULL

Trial of Niraparib in Participants With Newly-diagnosed Glioblastoma and Recurrent Glioma

TARGETS
PARP

LOCATIONS: Arizona

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

CLINICAL TRIALS
GENE
IDH1
ALTERATION
R132H
RATIONALE

IDH1 mutations may predict sensitivity to IDH1 inhibitors. On the basis of preclinical data, IDH1 mutations may also confer sensitivity to PARP

inhibitors in solid tumors. Preclinical data indicate that IDH1 mutations may predict sensitivity to glutaminase inhibitors.

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1
LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom), Villejuif (France)

NCT04644068
PHASE 1/2

Study of AZD5305 as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Malignancies

TARGETS
ERBB2, TROP2, PARP
LOCATIONS: Seoul (Korea, Republic of), Chuo-ku (Japan), Koto-ku (Japan), Melbourne (Australia), Warszawa (Poland), Gdynia (Poland), Grzegorz (Poland), Budapest (Hungary), Brno (Czechia), Padova (Italy)

NCT04740190
PHASE 2

Talazoparib - Carboplatin for Recurrent High-grade Glioma With DDRd

TARGETS
PARP
LOCATIONS: Hong Kong (Hong Kong)

NCT04715620
PHASE 2

Niraparib Combined With Radiotherapy in rGBM

TARGETS
PARP
LOCATIONS: Tianjin (China)

NCT05035745
PHASE 1/2

Selineor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
XPO1, PARP
LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
PARP, AKTs, PD-L1
LOCATIONS: Singapore (Singapore)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

NCT05076513

PHASE NULL

Trial of Niraparib in Participants With Newly-diagnosed Glioblastoma and Recurrent Glioma

TARGETS
PARP

LOCATIONS: Arizona

NCT04614909

PHASE NULL

Phase 0/2 Study of Pamiparib in Newly Diagnosed and rGBM

TARGETS
PARP

LOCATIONS: Arizona

NCT04801966

PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
CDK4, CDK6, PI3K-alpha, PD-L1, MEK,
PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT04497116

PHASE 1/2

Study of RP-3500 in Advanced Solid Tumors

TARGETS
ATR, PARP

LOCATIONS: Copenhagen (Denmark), Newcastle Upon Tyne (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Illinois, Toronto (Canada), Massachusetts, Rhode Island, New York, Tennessee

ORDERED TEST # ORD-1384802-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

CIC

I155V

EPHB1

E605K

MTOR

T1834_T1837del

NOTCH3

P779S

PARP3

E476K

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMSB	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK1	NTRK2	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENTSC (FAM46C)	TET2	TGFBR2	TIPARP
TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL
WT1	XPO1	XRCC2	ZNF217	ZNF703				

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV1
ETV4	ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT
KMT2A (MLL)	MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA
RAF1	RARA	RET	ROS1	RSPO2	SDC4	SLC34A2	TERC*	TERT**
TMPRSS2								

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Homologous Recombination status
Loss of Heterozygosity (LOH) score
Microsatellite (MS) status
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX
About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials
Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score $\geq 16\%$ will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian,

peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
- Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant

patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX

About FoundationOne®CDx

CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version 6.3.0

The median exon coverage for this sample is 983x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX
References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Martinez R, et al. Oncology (2004) PMID: 15331927
7. Martinez R, et al. J. Cancer Res. Clin. Oncol. (2005) PMID: 15672285
8. Martinez R, et al. Cancer Genet. Cytogenet. (2007) PMID: 17498554
9. Szybka M, et al. Clin. Neuropathol. () PMID: 12908754
10. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
11. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
12. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
13. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
14. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
15. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
16. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
17. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
18. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
19. Cristescu R, et al. Science (2018) PMID: 30309915
20. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
21. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
22. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
24. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
25. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
26. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
27. Thout M, et al. Nature (2020) PMID: 32322066
28. Bouffett E, et al. J. Clin. Oncol. (2016) PMID: 27001570
29. Johanns TM, et al. Cancer Discov (2016) PMID: 27683556
30. Lukas RV, et al. J. Neurooncol. (2018) PMID: 30073642
31. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
32. Patel RR, et al. Pediatr Blood Cancer (2020) PMID: 32386112
33. Johnson A, et al. Oncologist (2017) PMID: 28912153
34. Draaisma K, et al. Acta Neuropathol Commun (2015) PMID: 26699864
35. Wang L, et al. BMC Cancer (2020) PMID: 32164609
36. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
37. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
38. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
39. Rizvi NA, et al. Science (2015) PMID: 25765070
40. Johnson BE, et al. Science (2014) PMID: 24336570
41. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
42. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
43. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
44. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
45. Nature (2012) PMID: 22810696
46. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
47. Fan B, et al. Invest New Drugs (2019) PMID: 31028664
48. De La Fuente et al., 2020; ASCO Abstract 2505
49. O'Carrigan et al., 2016; ASCO Abstract 2504
50. Philip B, et al. Cell Rep (2018) PMID: 29719265
51. Molenaar RJ, et al. Clin. Cancer Res. (2018) PMID: 29339439
52. Lu Y, et al. Cancer Res. (2017) PMID: 28202508
53. Sulkowski PL, et al. Sci Transl Med (2017) PMID: 28148839
54. McBrayer SK, et al. Cell (2018) PMID: 30220459
55. Chaumeil MM, et al. Nat Commun (2013) PMID: 24019001
56. Hartmann C, et al. Clin. Cancer Res. (2013) PMID: 23918605
57. Rossetto M, et al. Rev. Neurol. (Paris) (2011) PMID: 21885076
58. Shin JH, et al. J. Neurooncol. (2013) PMID: 24129546
59. Parsons DW, et al. Science (2008) PMID: 18772396
60. Cancer Genome Atlas Research Network, et al. N. Engl. J. Med. (2015) PMID: 26061751
61. Brennan CW, et al. Cell (2013) PMID: 24120142
62. Hartmann C, et al. Acta Neuropathol. (2010) PMID: 21088844
63. Sonoda Y, et al. Cancer Sci. (2009) PMID: 19765000
64. Ahmadi R, et al. J. Neurooncol. (2012) PMID: 22528790
65. Jiang H, et al. Neuro-oncology (2013) PMID: 23486687
66. Shibahara I, et al. Int. J. Clin. Oncol. (2012) PMID: 21971842
67. Juratli TA, et al. J. Neurooncol. (2012) PMID: 23015095
68. Weller M, et al. J. Clin. Oncol. (2009) PMID: 19805672
69. Reitman ZJ, et al. J. Natl. Cancer Inst. (2010) PMID: 20513808
70. Jin G, et al. PLoS ONE (2011) PMID: 21326614
71. Gross S, et al. J. Exp. Med. (2010) PMID: 20142433
72. Ward PS, et al. Cancer Cell (2010) PMID: 20171147
73. Leonardi R, et al. J. Biol. Chem. (2012) PMID: 22442146
74. Dang L, et al. Nature (2009) PMID: 19935646
75. Ward PS, et al. Oncogene (2012) PMID: 21996744
76. Figueroa ME, et al. Cancer Cell (2010) PMID: 21130701
77. Xu W, et al. Cancer Cell (2011) PMID: 21251613
78. Turcan S, et al. Nature (2012) PMID: 22343889
79. Duncan CG, et al. Genome Res. (2012) PMID: 22899282
80. Weller M, et al. Nat Rev Clin Oncol (2021) PMID: 33293629
81. Louis DN, et al. Acta Neuropathol (2016) PMID: 27157931
82. Williamson CT, et al. Nat Commun (2016) PMID: 27958275
83. Aggarwal et al., 2021; ESMO Abstract 5120
84. Thomas A, et al. J. Clin. Oncol. (2018) PMID: 29252124
85. Yap TA, et al. J Clin Oncol (2020) PMID: 32568634
86. Bitler BG, et al. Nat. Med. (2015) PMID: 25686104
87. Kim KH, et al. Nat. Med. (2015) PMID: 26552009
88. Wiegand KC, et al. BMC Cancer (2014) PMID: 24559118
89. Huang HN, et al. Mod. Pathol. (2014) PMID: 24336158
90. Samartzis EP, et al. Oncotarget (2014) PMID: 24979463
91. Okamura R, et al. J Immunother Cancer (2020) PMID: 32111729
92. Yokoyama Y, et al. J Gynecol Oncol (2014) PMID: 24459582
93. Katagiri A, et al. Mod. Pathol. (2012) PMID: 22101352
94. Xie C, et al. Tumour Biol. (2014) PMID: 24833095
95. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
96. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
97. Gao J, et al. Sci Signal (2013) PMID: 23550210
98. Wu RC, et al. Cancer Biol. Ther. (2014) PMID: 24618703
99. Jones S, et al. Hum. Mutat. (2012) PMID: 22009941
100. Dulak AM, et al. Nat. Genet. (2013) PMID: 23525077
101. Streppel MM, et al. Oncogene (2014) PMID: 23318448
102. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
103. Ross JS, et al. Oncologist (2014) PMID: 24563076
104. Huang HN, et al. Histopathology (2015) PMID: 25195947
105. Hussein YR, et al. Mod. Pathol. (2015) PMID: 25394778
106. Bosse T, et al. Mod. Pathol. (2013) PMID: 23702729
107. Allo G, et al. Mod. Pathol. (2014) PMID: 23887303
108. Chou A, et al. Hum. Pathol. (2014) PMID: 24925223
109. Ye J, et al. Hum. Pathol. (2014) PMID: 25311944
110. Wei XL, et al. World J. Gastroenterol. (2014) PMID: 25561809
111. Chen K, et al. Proc. Natl. Acad. Sci. U.S.A. (2015) PMID: 25583476
112. Wang K, et al. Nat. Genet. (2011) PMID: 22037554
113. Abe H, et al. Virchows Arch. (2012) PMID: 22915242
114. Wang DD, et al. PLoS ONE (2012) PMID: 22808142
115. Wiegand KC, et al. Hum. Pathol. (2014) PMID: 24767857
116. Katagiri A, et al. Int. J. Gynecol. Cancer (2012) PMID: 22274316
117. Cho H, et al. Hum. Pathol. (2013) PMID: 23427874
118. Gui Y, et al. Nat. Genet. (2011) PMID: 21822268
119. Balbás-Martínez C, et al. PLoS ONE (2013) PMID: 23650517
120. Faraj SF, et al. Hum. Pathol. (2014) PMID: 25175170
121. Rahman M, et al. Hum. Pathol. (2013) PMID: 22939958
122. Maeda D, et al. Int J Mol Sci (2010) PMID: 21614196
123. Lowery WJ, et al. Int. J. Gynecol. Cancer (2012) PMID: 22193641
124. Fadare O, et al. Mod. Pathol. (2013) PMID: 23524907
125. Mao TL, et al. Am. J. Surg. Pathol. (2013) PMID: 24076775
126. Zhang X, et al. Cancer Epidemiol (2012) PMID: 21889920
127. Mamo A, et al. Oncogene (2012) PMID: 21892209
128. Zhao J, et al. Tumour Biol. (2014) PMID: 24430365
129. Lichner Z, et al. Am. J. Pathol. (2013) PMID: 23416164
130. Feng F, et al. Int J Clin Oncol (2021) PMID: 33387086
131. Conci S, et al. Updates Surg (2020) PMID: 32020551
132. Simbolo M, et al. Sci Rep (2018) PMID: 29740198
133. Ruzzenante A, et al. Ann. Surg. Oncol. (2016) PMID: 26717940
134. Guan B, et al. Cancer Res. (2011) PMID: 21900401
135. Wiegand KC, et al. N. Engl. J. Med. (2010) PMID: 20942669
136. Jones S, et al. Science (2010) PMID: 20826764
137. Yan HB, et al. Carcinogenesis (2014) PMID: 24293408
138. Huang J, et al. Nat. Genet. (2012) PMID: 22922871
139. Chan-On W, et al. Nat. Genet. (2013) PMID: 24185513
140. Zang ZJ, et al. Nat. Genet. (2012) PMID: 22484628
141. Michels J, et al. Oncogene (2014) PMID: 24037533
142. Bryant HE, et al. Nucleic Acids Res. (2006) PMID: 16556909
143. Mateo J, et al. N. Engl. J. Med. (2015) PMID: 26510020
144. Mateo J, et al. Lancet Oncol. (2019) PMID: 31806540
145. Abida W, et al. Clin. Cancer Res. (2020) PMID: 32086346
146. Ma et al., 2022; ASCO GI Abstract 563
147. Dhawan et al., 2020; ASCO Abstract 3513
148. Papageorgiou GI, et al. Front Oncol (2021) PMID: 35004311
149. Olson D, et al. Clin Genitourin Cancer (2016) PMID: 27079472
150. Swisher EM, et al. Nat Commun (2021) PMID: 33941784
151. Zhang W, et al. Oncologist (2020) PMID: 32045060
152. Yap TA, et al. Cancer Discov (2021) PMID: 32988960
153. Menezes DL, et al. Mol. Cancer Res. (2015) PMID: 25232030

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX
References

154. Vendetti FP, et al. Oncotarget (2015) PMID: 26517239
155. Min A, et al. Mol. Cancer Ther. (2017) PMID: 28138034
156. Kwok M, et al. Blood (2016) PMID: 26563132
157. Riabinska A, et al. Sci Transl Med (2013) PMID: 23761041
158. Nature (2008) PMID: 18772890
159. Kheirollahi M, et al. Med. Oncol. (2011) PMID: 20077038
160. Seol HJ, et al. Oncol. Rep. (2011) PMID: 21617879
161. Tribius S, et al. Int. J. Radiat. Oncol. Biol. Phys. (2001) PMID: 11380241
162. Roy K, et al. Biochem. Biophys. Res. Commun. (2006) PMID: 16631604
163. Biddlestone-Thorpe L, et al. Clin. Cancer Res. (2013) PMID: 23620409
164. Carminati PO, et al. Mol. Biol. Rep. (2014) PMID: 24218165
165. Nadkarni A, et al. J. Neurooncol. (2012) PMID: 23054561
166. Eich M, et al. Mol. Cancer Ther. (2013) PMID: 23960094
167. Shiloh Y, et al. Nat. Rev. Mol. Cell Biol. (2013) PMID: 23847781
168. Cremona CA, et al. Oncogene (2014) PMID: 23851492
169. Jiang X, et al. J. Biol. Chem. (2006) PMID: 16603769
170. Fernandes N, et al. J. Biol. Chem. (2005) PMID: 15713674
171. Scott SP, et al. Proc. Natl. Acad. Sci. U.S.A. (2002) PMID: 11805335
172. van Os NJ, et al. Clin Genet (2016) PMID: 26662178
173. Rothblum-Oviatt C, et al. Orphanet J Rare Dis (2016) PMID: 27884168
174. Liang J, et al. Cancer Res (2020) PMID: 31551363
175. Garbarino J, et al. Transl Oncol (2021) PMID: 34118569
176. Cole et al., 2021; AACR Abstract CT059
177. George SL, et al. EBioMedicine (2020) PMID: 32846370
178. Koschmann C, et al. Sci Transl Med (2016) PMID: 26936505
179. Heaphy CM, et al. Science (2011) PMID: 21719641
180. Singhi et al., 2015; USCAP Abstract 1797
181. Jiao Y, et al. Science (2011) PMID: 21252315
182. Fishbein L, et al. Nat Commun (2015) PMID: 25608029
183. Morosini et al., 2014; ASCO Abstract 11008
184. Cheung NK, et al. JAMA (2012) PMID: 22416102
185. Molenaar JJ, et al. Nature (2012) PMID: 22367537
186. Pugh TJ, et al. Nat. Genet. (2013) PMID: 23334666
187. Cheung NK, et al. Nat. Rev. Cancer (2013) PMID: 23702928
188. Marinoni I, et al. Gastroenterology (2014) PMID: 24148618
189. Qadeer ZA, et al. J. Invest. Dermatol. (2014) PMID: 24468746
190. Kannan K, et al. Oncotarget (2012) PMID: 23104868
191. Haberler C, et al. Clin. Neuropathol. () PMID: 24559763
192. Reuss DE, et al. Acta Neuropathol. (2015) PMID: 25427834
193. Sahm F, et al. Acta Neuropathol. (2014) PMID: 25143301
194. Singhi et al., 2015; USCAP Abstract 93
195. Liao JY, et al. Am. J. Surg. Pathol. (2015) PMID: 25229770
196. Clynes D, et al. Trends Biochem. Sci. (2013) PMID: 23916100
197. Ratnakumar K, et al. Epigenetics (2013) PMID: 23249563
198. Lovejoy CA, et al. PLoS Genet. (2012) PMID: 22829774
199. Bower K, et al. PLoS ONE (2012) PMID: 23185534
200. Nan X, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17296936
201. Garrick D, et al. Gene (2004) PMID: 14729260
202. Eustermann S, et al. Nat. Struct. Mol. Biol. (2011) PMID: 21666677
203. Flynn RL, et al. Science (2015) PMID: 25593184
204. Gibbons RJ, et al. Cell (1995) PMID: 7697714
205. Bettgeowda C, et al. Science (2011) PMID: 21817013
206. Yip S, et al. J. Pathol. (2012) PMID: 22072542
207. Sahm F, et al. Acta Neuropathol. (2012) PMID: 22588899
208. Jiao Y, et al. Oncotarget (2012) PMID: 22869205
209. Chan AK, et al. Mod. Pathol. (2014) PMID: 24030748
210. Lee CJ, et al. Brain Res. Mol. Brain Res. (2002) PMID: 12393275
211. Yan et al. 2020; DOI:10.1200/PO.19.00385
212. Nature (2012) PMID: 22960745
213. Augert A, et al. J Thorac Oncol (2017) PMID: 28007623
214. Ardeshir-Larijani F, et al. Clin Lung Cancer (2018) PMID: 29627316
215. Hillman RT, et al. Nat Commun (2018) PMID: 29950560
216. Abudurehman A, et al. J. Cancer Res. Clin. Oncol. (2018) PMID: 29532228
217. Vicent GP, et al. Genes Dev. (2011) PMID: 21447625
218. Hannibal MC, et al. Am. J. Med. Genet. A (2011) PMID: 21671394
219. Fagan RJ, et al. Cancer Lett. (2019) PMID: 31128216
220. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
221. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
222. Xie M, et al. Nat. Med. (2014) PMID: 25326804
223. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
224. Severson EA, et al. Blood (2018) PMID: 29678827
225. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
226. Chabon JJ, et al. Nature (2020) PMID: 32269342
227. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
228. Yen WC, et al. Clin. Cancer Res. (2015) PMID: 25934888
229. Ferrarotto et al., 2020; ESMO Abstract 919MO
230. Chen J, et al. Genes Cancer (2010) PMID: 21127729
231. Wang J, et al. Stem Cells (2010) PMID: 19921751
232. Fan X, et al. Stem Cells (2010) PMID: 19904829
233. Dell'Albani P, et al. Neuro-oncology (2014) PMID: 24305720
234. Tchorz JS, et al. Cell Death Dis (2012) PMID: 22717580
235. Chudnovsky Y, et al. Cell Rep (2014) PMID: 24440720
236. Xu P, et al. PLoS ONE (2013) PMID: 23349727
237. Penton AL, et al. Semin. Cell Dev. Biol. (2012) PMID: 22306179
238. Kopan R, et al. Cell (2009) PMID: 19379690
239. Pancewicz J, et al. BMC Cancer (2011) PMID: 22128846
240. Sakata-Yanagimoto M, et al. Curr. Top. Microbiol. Immunol. (2012) PMID: 22695918
241. Kiel MJ, et al. J. Exp. Med. (2012) PMID: 22891276
242. Egloff AM, et al. Clin. Cancer Res. (2012) PMID: 22773520
243. Wang NJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2011) PMID: 22006338
244. Robinson DR, et al. Nat. Med. (2011) PMID: 22101766
245. Crotty et al., 2020; DOI: 10.1016/j.ehpc.2020.200408
246. Varela I, et al. Nature (2011) PMID: 21248752
247. Chen Z, et al. Biochem Biophys Res Commun (2018) PMID: 29522714
248. Chen BY, et al. Blood (2020) PMID: 32202636
249. Sun XJ, et al. J. Biol. Chem. (2005) PMID: 16118227
250. Faber PW, et al. Hum. Mol. Genet. (1998) PMID: 9700202
251. Al Sarakbi W, et al. BMC Cancer (2009) PMID: 19698110
252. Parker H, et al. Leukemia (2016) PMID: 27282254
253. Zhang J, et al. Nature (2012) PMID: 22237106
254. McKinney M, et al. Cancer Discov (2017) PMID: 28122867
255. Moffitt AB, et al. J. Exp. Med. (2017) PMID: 28424246
256. Zhu X, et al. Nat. Genet. (2014) PMID: 24509477
257. Lu C, et al. Science (2016) PMID: 27174990
258. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
259. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
260. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
261. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
262. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
263. Xu L, et al. Mol. Med. (2001) PMID: 11713371
264. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
265. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
266. Pirollo KF, et al. Mol. Ther. (2016) PMID: 27357628
267. Hajdenberg et al., 2012; ASCO Abstract e15010
268. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
269. Moore et al., 2019; ASCO Abstract 5513
270. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
271. Oza et al., 2015; ASCO Abstract 5506
272. Lee J, et al. Cancer Discov (2019) PMID: 31315834
273. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
274. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
275. Lehmann S, et al. J. Clin. Oncol. (2012) PMID: 22965953
276. Mohell N, et al. Cell Death Dis (2015) PMID: 26086967
277. Franssøn A, et al. J Ovarian Res (2016) PMID: 27179933
278. Gourley et al., 2016; ASCO Abstract 5571
279. Boudny M, et al. Haematologica (2019) PMID: 30975914
280. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
281. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
282. Jha P, et al. Diagn. Mol. Pathol. (2011) PMID: 22089350
283. Uno M, et al. Cancer Lett. (2005) PMID: 15914282
284. Uno M, et al. Int. J. Biol. Markers () PMID: 16711514
285. Lass U, et al. PLoS ONE (2012) PMID: 22844452
286. Faria MH, et al. APMIS (2012) PMID: 23009112
287. Milinkovic V, et al. PLoS ONE (2013) PMID: 24358143
288. Galatro TF, et al. PLoS ONE (2013) PMID: 23613880
289. Schmidt MC, et al. J. Neuropathol. Exp. Neurol. (2002) PMID: 11939587
290. Nozaki M, et al. Neuro-oncology (1999) PMID: 11550308
291. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
292. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
293. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
294. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
295. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
296. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
297. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
298. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
299. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
300. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
301. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
302. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
303. Lalloo F, et al. Lancet (2003) PMID: 12672316
304. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713
305. Hematology Am Soc Hematol Educ Program (2018)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1384802-01

APPENDIX

References

pmid: 30504320
 306. DiNardo CD, et al. N. Engl. J. Med. (2018) pmid: 29860938

307. Lowery MA, et al. Lancet Gastroenterol Hepatol (2019) pmid: 31300360
 308. Abou-Alfa GK, et al. Lancet Oncol. (2020) pmid:

32416072
 309. Mellingerhoff IK, et al. J. Clin. Oncol. (2020) pmid: 32530764

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 16 June 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531