



## Sample Information

**Patient Name:** 魏慧珍  
**Gender:** Female  
**ID No.:** J221117841  
**History No.:** 18365441  
**Age:** 53

**Ordering Doctor:** DOC1697J 蔡淳光  
**Ordering REQ.:** H44EEBK  
**Signing in Date:** 2022/10/06

**Path No.:** S111-97920  
**MP No.:** MY22029  
**Assay:** Oncomine Myeloid Assay  
**Sample Type:** Bone Marrow  
**Bone Marrow Aspirating Date:** 2022/09/26

**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)

**Note:**

## Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	3
Prognostic Details	4

**Report Highlights**  
1 Relevant Biomarkers  
1 Therapies Available  
0 Clinical Trials

## Relevant Myelodysplastic Syndrome Variants

Gene	Finding	Gene	Finding
ASXL1	None detected	NPM1	None detected
BCOR	None detected	NRAS	None detected
CBL	None detected	NUP214	None detected
CREBBP	None detected	RUNX1	None detected
ETV6	None detected	SF3B1	<b>SF3B1 p.(K700E) c.2098A&gt;G</b>
EZH2	None detected	SRSF2	None detected
FLT3	None detected	STAG2	None detected
GATA2	None detected	TP53	None detected
IDH2	None detected	U2AF1	None detected
KIT	None detected	WT1	None detected
KMT2A	None detected	ZRSR2	None detected
MECOM	None detected		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>SF3B1 p.(K700E) c.2098A&gt;G</b> splicing factor 3b subunit 1 Allele Frequency: 34.67% <b>Prognostic significance:</b> NCCN: Favorable	luspatercept	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
SF3B1	p.(K700E)	c.2098A>G	COSM84677	chr2:198266834	34.67%	NM_012433.4	missense	1647
TET2	p.(L1210P)	c.3629T>C	.	chr4:106164761	41.89%	NM_001127208.2	missense	1998

## Biomarker Descriptions

### SF3B1 (splicing factor 3b subunit 1)

**Background:** The SF3B1 gene encodes the splicing factor 3b subunit 1 protein, a core component of the U2 small nuclear ribonucleoprotein (snRNP) complex of the spliceosome responsible for RNA splicing. SF3B1 is involved in recognition of the branch point sequence during selection of the 3' splice site. Recurrent somatic mutations in SF3B1 and other components of the splicing machinery including SRSF2, U2AF1, and ZRSR2, are common in myelodysplasia. These components experience mutations in a mutually exclusive manner suggesting a common impact on RNA splicing and the pathogenesis of myelodysplasia<sup>1</sup>. SF3B1 mutations are believed to contribute to aberrant post-translational inactivation of the regulatory complex PPP2R5A of protein phosphatase 2A (PP2A), leading to the activation and stabilization of MYC activation and impairing apoptosis<sup>2</sup>.

**Alterations and prevalence:** SF3B1 mutations occur in the majority (70-80%) of myelodysplastic syndromes (MDS) with ring sideroblasts (RS) and at lower frequency in other myeloid neoplasms including MDS without RS (7%), chronic myelomonocytic leukemia (5-6%), therapy-related acute myeloid leukemia (AML) or AML with MDS features (5%), and de novo AML (3%)<sup>1,3,4</sup>. Recurrent somatic SF3B1 mutations are also common in certain solid cancers including uveal melanoma (20-30%) and breast cancer (2%) and at lower frequencies in diverse cancer types<sup>5,6,7,8,9,10,11,12</sup>. Cancer-associated recurrent missense mutations in SF3B1 occur within the HEAT repeat domains 5-9 at codon positions R625, K666, K700, G742, and D781<sup>13</sup>. The functional significance of recurrent SF3B1 mutations is to alter branch point selection thus inducing cryptic 3' splice site selection<sup>13,14,15</sup>.

**Potential relevance:** Currently, no therapies are approved for SF3B1 aberrations. SF3B1 mutations are associated with aggressive disease and shorter survival in patients diagnosed with chronic lymphocytic leukemia (CLL)<sup>16</sup>. Investigational inhibitors of the spliceosome are in early clinical development<sup>17,18</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

SF3B1 p.(K700E) c.2098A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
luspatercept	×	×	×	<input checked="" type="radio"/>	×

## Relevant Therapy Details

### Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2022-08-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

SF3B1 p.(K700E) c.2098A>G

### ☒ luspatercept

Cancer type: Myelodysplastic Syndrome

Variant class: SF3B1 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Myelodysplastic Syndromes [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.11.002>]

## Prognostic Details

### Current NCCN Information

NCCN information is current as of 2022-08-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### SF3B1 p.(K700E) c.2098A>G

### Prognostic significance: NCCN: Favorable

Cancer type: Myelodysplastic Syndrome

Variant class: SF3B1 K700E mutation

NCCN Recommendation category: 2A

#### Summary:

- NCCN Guidelines® independently associate the biomarker with favorable prognosis

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2022]

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Yoshida et al. Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature*. 2011 Sep 11;478(7367):64-9. PMID: 21909114
2. Liu et al. Mutations in the RNA Splicing Factor SF3B1 Promote Tumorigenesis through MYC Stabilization. *Cancer Discov*. 2020 Mar 18. PMID: 32188705
3. Patnaik et al. Refractory anemia with ring sideroblasts (RARS) and RARS with thrombocytosis (RARS-T): 2017 update on diagnosis, risk-stratification, and management. *Am. J. Hematol*. 2017 Mar;92(3):297-310. PMID: 28188970
4. Patnaik et al. Spliceosome mutations involving SRSF2, SF3B1, and U2AF35 in chronic myelomonocytic leukemia: prevalence, clinical correlates, and prognostic relevance. *Am. J. Hematol*. 2013 Mar;88(3):201-6. PMID: 23335386
5. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
7. Ellis et al. Whole-genome analysis informs breast cancer response to aromatase inhibition. *Nature*. 2012 Jun 10;486(7403):353-60. PMID: 22722193
8. Maguire et al. SF3B1 mutations constitute a novel therapeutic target in breast cancer. *J. Pathol*. 2015 Mar;235(4):571-80. PMID: 25424858
9. Harbour et al. Recurrent mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nat. Genet*. 2013 Feb;45(2):133-5. PMID: 23313955
10. Martin et al. Exome sequencing identifies recurrent somatic mutations in EIF1AX and SF3B1 in uveal melanoma with disomy 3. *Nat. Genet*. 2013 Aug;45(8):933-6. PMID: 23793026
11. Furney et al. SF3B1 mutations are associated with alternative splicing in uveal melanoma. *Cancer Discov*. 2013 Oct;3(10):1122-1129. PMID: 23861464
12. Dono et al. Mutation frequencies of GNAQ, GNA11, BAP1, SF3B1, EIF1AX and TERT in uveal melanoma: detection of an activating mutation in the TERT gene promoter in a single case of uveal melanoma. *Br. J. Cancer*. 2014 Feb 18;110(4):1058-65. PMID: 24423917
13. DeBoever et al. Transcriptome sequencing reveals potential mechanism of cryptic 3' splice site selection in SF3B1-mutated cancers. *PLoS Comput. Biol*. 2015 Mar;11(3):e1004105. PMID: 25768983
14. Darman et al. Cancer-Associated SF3B1 Hotspot Mutations Induce Cryptic 3' Splice Site Selection through Use of a Different Branch Point. *Cell Rep*. 2015 Nov 3;13(5):1033-45. PMID: 26565915
15. Alsafadi et al. Cancer-associated SF3B1 mutations affect alternative splicing by promoting alternative branchpoint usage. *Nat Commun*. 2016 Feb 4;7:10615. doi: 10.1038/ncomms10615. PMID: 26842708
16. Wan et al. SF3B1 mutations in chronic lymphocytic leukemia. *Blood*. 2013 Jun 6;121(23):4627-34. PMID: 23568491
17. Seiler et al. H3B-8800, an orally available small-molecule splicing modulator, induces lethality in spliceosome-mutant cancers. *Nat. Med*. 2018 May;24(4):497-504. PMID: 29457796
18. Lee et al. Therapeutic targeting of splicing in cancer. *Nat. Med*. 2016 Sep 7;22(9):976-86. PMID: 27603132