



## Sample Information

**Patient Name:** 姜靜宜  
**Gender:** Male  
**ID No.:** F103691438  
**History No.:** 28948110  
**Age:** 88

**Ordering Doctor:** DOC3049E 金彥承  
**Ordering REQ.:** 0BKYYKA  
**Signing in Date:** 2021/09/09

**Path No.:** S110-99501  
**MP No.:** F21076  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S110-92053A  
**Percentage of tumor cells:** 50%

**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)

**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Alert Details	3

**Report Highlights**  
1 Relevant Biomarkers  
0 Therapies Available  
0 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

## Relevant Biomarkers

No relevant biomarkers found in this sample.

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ALK	p.(R1436C)	c.4306C>T	.	chr2:29416647	48.75%	NM_004304.5	missense	2000

### Copy Number Variations

Gene	Locus	Copy Number
FGFR1	chr8:38271445	8.08

## Biomarker Descriptions

### FGFR1 (fibroblast growth factor receptor 1)

**Background:** The FGFR1 gene encodes fibroblast growth receptor 1, a member of the fibroblast growth factor receptor (FGFR) family that also includes FGFR2, 3, and 4. These proteins are single transmembrane receptors composed of three extracellular immunoglobulin (Ig)-type domains and an intracellular kinase domain. Upon FGF-mediated stimulation, FGFRs activate several oncogenic signaling pathways, including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, PLC/PKC, and JAK/STAT pathways influencing cell proliferation, migration, and survival<sup>1,2,3</sup>.

**Alterations and prevalence:** Recurrent somatic alterations common to the FGFR family include gene amplification, mutation, and chromosomal translocations leading to FGFR fusions<sup>4</sup>. Amplification of FGFR1 is observed in 15-20% of squamous lung cancer, 10-15% of breast cancer, 8% of bladder cancer, and 2-5% of uterine cancer cases<sup>5,6,7,8,9</sup>. The most common recurrent mutations, N546K and K656E, are relatively infrequent (<1%); they activate mutations in the kinase domain and are distributed in diverse cancer types<sup>10</sup>. FGFR1 translocations giving rise to expressed fusions are common in certain hematological cancers, but less common in solid tumors<sup>11,12,13</sup>.

**Potential relevance:** The FDA has granted fast-track designation (2018) to Debio 1347<sup>14</sup> for solid tumors harboring aberrations in FGFR1, FGFR2, or FGFR3. FDA has approved multi-kinase inhibitors, including regorafenib, ponatinib, lenvatinib, nintedanib, and pazopanib, that are known to inhibit FGFR family members. These inhibitors have demonstrated anti-tumor activity in select cancer types with FGFR alterations<sup>15,16,17,18,19,20,21</sup>. In a phase II clinical trial, dovitinib, a multi-tyrosine kinase inhibitor (TKI), exhibited an overall response rate (ORR) of 11.5% and a disease control rate (DCR) of 50% in patients with advanced squamous cell lung cancer possessing FGFR1 amplification. The patients had a median overall survival (OS) of 5 months and progression-free survival (PFS) of 2.9 months<sup>22</sup>. Likewise, in a phase Ib study testing the FGFR inhibitor AZD4547, median OS was 4.9 months in patients with FGFR1-amplified advanced squamous cell lung cancer. One of 13 (8%) patients achieved a partial response, 4 (31%) exhibited stable disease, and 2 (13.3%) demonstrated PFS at 12 weeks<sup>23</sup>.

## Clinical Trials in Taiwan region:

### Alerts Informed By Public Data Sources

#### Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2021-07-14. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### FGFR1 amplification

##### Debio 1347

**Cancer type:** Solid Tumor

**Variant class:** FGFR1 aberration

**Supporting Statement:**

The FDA has granted Fast Track Designation to the FGFR 1-3 inhibitor, debio 1347, for FGFR1/2/3 alterations in unresectable or metastatic solid tumors.

**Reference:**

<https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Babina et al. Advances and challenges in targeting FGFR signalling in cancer. *Nat. Rev. Cancer*. 2017 May;17(5):318-332. PMID: 28303906
2. Ahmad et al. Mechanisms of FGFR-mediated carcinogenesis. *Biochim. Biophys. Acta*. 2012 Apr;1823(4):850-60. PMID: 22273505
3. Sarabipour et al. Mechanism of FGF receptor dimerization and activation. *Nat Commun*. 2016 Jan 4;7:10262. doi: 10.1038/ncomms10262. PMID: 26725515
4. Helsten et al. The FGFR Landscape in Cancer: Analysis of 4,853 Tumors by Next-Generation Sequencing. *Clin. Cancer Res*. 2016 Jan 1;22(1):259-67. PMID: 26373574
5. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature*. 2012 Sep 27;489(7417):519-25. PMID: 22960745
6. Ciriello et al. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*. 2015 Oct 8;163(2):506-19. PMID: 26451490
7. Cancer et al. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013 May 2;497(7447):67-73. PMID: 23636398
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
10. Lew et al. The precise sequence of FGF receptor autophosphorylation is kinetically driven and is disrupted by oncogenic mutations. *Sci Signal*. 2009 Feb 17;2(58):ra6. PMID: 19224897
11. Jackson et al. 8p11 myeloproliferative syndrome: a review. *Hum. Pathol*. 2010 Apr;41(4):461-76. PMID: 20226962
12. Li et al. Identification of a novel partner gene, TPR, fused to FGFR1 in 8p11 myeloproliferative syndrome. *Genes Chromosomes Cancer*. 2012 Sep;51(9):890-7. PMID: 22619110
13. Wasag et al. The kinase inhibitor TKI258 is active against the novel CUX1-FGFR1 fusion detected in a patient with T-lymphoblastic leukemia/lymphoma and t(7;8)(q22;p11). *Haematologica*. 2011 Jun;96(6):922-6. PMID: 21330321
14. <https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>
15. Cha et al. FGFR2 amplification is predictive of sensitivity to regorafenib in gastric and colorectal cancers in vitro. *Mol Oncol*. 2018 Jun;12(7):993-1003. PMID: 29573334
16. Chae et al. Inhibition of the fibroblast growth factor receptor (FGFR) pathway: the current landscape and barriers to clinical application. *Oncotarget*. 2017 Feb 28;8(9):16052-16074. PMID: 28030802
17. Porta et al. FGFR a promising druggable target in cancer: Molecular biology and new drugs. *Crit. Rev. Oncol. Hematol*. 2017 May;113:256-267. PMID: 28427515
18. Gozgit et al. Ponatinib (AP24534), a multitargeted pan-FGFR inhibitor with activity in multiple FGFR-amplified or mutated cancer models. *Mol. Cancer Ther*. 2012 Mar;11(3):690-9. PMID: 22238366
19. Yamamoto et al. Lenvatinib, an angiogenesis inhibitor targeting VEGFR/FGFR, shows broad antitumor activity in human tumor xenograft models associated with microvessel density and pericyte coverage. *Vasc Cell*. 2014 Sep 6;6:18. doi: 10.1186/2045-824X-6-18. eCollection 2014. PMID: 25197551
20. Kim et al. Pazopanib, a novel multitargeted kinase inhibitor, shows potent in vitro antitumor activity in gastric cancer cell lines with FGFR2 amplification. *Mol. Cancer Ther*. 2014 Nov;13(11):2527-36. PMID: 25249557
21. Hibi et al. FGFR gene alterations in lung squamous cell carcinoma are potential targets for the multikinase inhibitor nintedanib. *Cancer Sci*. 2016 Nov;107(11):1667-1676. PMID: 27581340
22. Lim et al. Efficacy and safety of dovitinib in pretreated patients with advanced squamous non-small cell lung cancer with FGFR1 amplification: A single-arm, phase 2 study. *Cancer*. 2016 Oct;122(19):3024-31. PMID: 27315356
23. Paik et al. A Phase Ib Open-Label Multicenter Study of AZD4547 in Patients with Advanced Squamous Cell Lung Cancers. *Clin. Cancer Res*. 2017 Sep 15;23(18):5366-5373. PMID: 28615371