

**ABOUT THE TEST** FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

<b>PATIENT</b>	<b>DISEASE</b> Colon adenocarcinoma (CRC)	<b>PHYSICIAN</b>	<b>ORDERING PHYSICIAN</b> Yeh, Yi-Chen	<b>SPECIMEN</b>	<b>SPECIMEN SITE</b> Colon
	<b>NAME</b> Yang, Shih Chang		<b>MEDICAL FACILITY</b> Taipei Veterans General Hospital		<b>SPECIMEN ID</b> S110-34021 D (PF23035)
	<b>DATE OF BIRTH</b> 24 October 1969		<b>ADDITIONAL RECIPIENT</b> None		<b>SPECIMEN TYPE</b> Slide Deck
	<b>SEX</b> Male		<b>MEDICAL FACILITY ID</b> 205872		<b>DATE OF COLLECTION</b> 09 November 2021
	<b>MEDICAL RECORD #</b> 46204846		<b>PATHOLOGIST</b> Not Provided		<b>SPECIMEN RECEIVED</b> 10 April 2023

## Biomarker Findings

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 1 Muts/Mb

## Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

**KRAS** wildtype

**NRAS** wildtype

**NF1** loss exons 37-58

**APC** R232\*

**CCND2** amplification

**ERBB2** T798I

**FGF23** amplification

**FGF6** amplification

**KDM5A** amplification

**NOTCH3** splice site 119-156\_197+54>GGGGG

**SMAD4** loss

**TP53** P151R

3 Disease relevant genes with no reportable alterations: **BRAF**, **KRAS**, **NRAS**

## Report Highlights

- Targeted therapies with **NCCN categories of evidence** in this tumor type: **Cetuximab** (p. 11), **Panitumumab** (p. 11)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 14)

### BIOMARKER FINDINGS

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 1 Muts/Mb

### GENOMIC FINDINGS

**KRAS** - wildtype

0 Trials

**NRAS** - wildtype

0 Trials

### THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

No therapies or clinical trials. See Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)		THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
Cetuximab	2A	none
Panitumumab	2A	
Cetuximab	2A	none
Panitumumab	2A	

 NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

GENOMIC FINDINGS	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
<b>NF1</b> - loss exons 37-58	none	Cobimetinib
10 Trials see p. 19		Selumetinib
		Trametinib
<b>APC</b> - R232*	none	none
3 Trials see p. 14		
<b>CCND2</b> - amplification	none	none
10 Trials see p. 15		
<b>ERBB2</b> - T798I	none	none
9 Trials see p. 17		

☐ NCCN category

**GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS**

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

<b>FGF23</b> - amplification.....	p. 7	<b>NOTCH3</b> - splice site 119-156_197+54>GGGGG.....	p. 9
<b>FGF6</b> - amplification.....	p. 8	<b>SMAD4</b> - loss.....	p. 9
<b>KDM5A</b> - amplification.....	p. 8	<b>TP53</b> - P151R.....	p. 10

**NOTE** Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

## BIOMARKER FINDINGS

## BIOMARKER

# Microsatellite status

## RESULT

MS-Stable

## POTENTIAL TREATMENT STRATEGIES

### — Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors<sup>1-3</sup>, including approved therapies nivolumab and pembrolizumab<sup>4</sup>. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%,  $p=0.001$ )<sup>5</sup>. For patients with chemotherapy-refractory microsatellite-stable (MSS) metastatic colorectal cancer (CRC), a Phase 3 trial reported no OS advantage from the combination of the PD-L1 inhibitor atezolizumab plus cobimetinib relative to regorafenib (8.9 vs. 8.5 months, HR=1.00);

atezolizumab monotherapy similarly did not prolong OS (7.1 vs. 8.5 months, HR=1.19)<sup>6</sup>. For patients with MSS CRC, a Phase 2 study combining ipilimumab and nivolumab reported an overall DCR of 25% (10/40)<sup>7</sup>. Two Phase 1 studies for patients with MSS CRC treated with regorafenib and nivolumab reported PFSs of 7.9 months<sup>8</sup> and 5.7 months<sup>9</sup>, and a patient with MSS CRC refractory to chemotherapy treated with the PD-1 inhibitor sintilimab and regorafenib reported a CR<sup>10</sup>.

### — Nontargeted Approaches —

MSI has not been found to be a predictive biomarker for combination chemotherapy regimens, including FOLFOX<sup>11-12</sup> and FOLFIRI<sup>13-14</sup>. Patients with MSS CRC are more likely to benefit from postsurgical fluorouracil (FU)-based adjuvant therapy<sup>15-16</sup> but less likely to benefit from irinotecan chemotherapy<sup>17</sup>.

## FREQUENCY & PROGNOSIS

MSS colorectal cancers (CRCs) make up 70-85% of CRC cases<sup>3,18-22</sup>. MSS colorectal cancers are

molecularly heterogeneous, driven by diverse mechanisms such as extensive DNA methylation, oncogenic mutations in KRAS or BRAF, or chromosomal instability<sup>22</sup>. Multiple studies have shown that MSS CRCs have a worse prognosis than MSI-high tumors<sup>18,23-29</sup>.

## FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor<sup>20</sup>. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2<sup>20,30-31</sup>. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers<sup>19,32-33</sup>. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins<sup>19-20,31,33</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

## BIOMARKER FINDINGS

## BIOMARKER

# Tumor Mutational Burden

## RESULT

1 Muts/Mb

## POTENTIAL TREATMENT STRATEGIES

### — Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1<sup>34-36</sup>, anti-PD-1 therapies<sup>34-37</sup>, and combination nivolumab and ipilimumab<sup>38-43</sup>. In multiple pan-tumor studies, increased tissue tumor mutational burden (TMB) was associated with sensitivity to immune checkpoint inhibitors<sup>34-37,44-48</sup>. In the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors, significant improvement in ORR was observed for patients with TMB  $\geq 10$  Muts/Mb (as measured by this assay) compared with those with TMB  $< 10$  Muts/Mb in a large cohort that included multiple tumor types<sup>44</sup>; similar findings were observed in the KEYNOTE 028 and 012 trials<sup>37</sup>. At the same TMB cutpoint, retrospective analysis of patients with solid tumors treated with any checkpoint inhibitor identified that tissue TMB scores  $\geq 10$  Muts/Mb were associated with prolonged time to treatment failure compared with scores  $< 10$  Muts/Mb (HR=0.68)<sup>48</sup>. For patients with solid tumors treated with nivolumab plus ipilimumab in the CheckMate 848 trial, improved responses were observed in patients with a tissue TMB  $\geq 10$  Muts/Mb independent of blood TMB at any cutpoint in matched samples<sup>49</sup>. However, support for higher TMB thresholds and efficacy was observed in the prospective Phase 2 MyPathway trial of atezolizumab for patients with pan-solid tumors, where improved ORR and DCR

was seen in patients with TMB  $\geq 16$  Muts/Mb than those with TMB  $\geq 10$  and  $< 16$  Muts/Mb<sup>47</sup>. Similarly, analyses across several solid tumor types reported that patients with higher TMB (defined as  $\geq 16-20$  Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy compared with patients with higher TMB treated with chemotherapy<sup>50</sup> or those with lower TMB treated with PD-1 or PD-L1-targeting agents<sup>35</sup>. In CRC specifically, a retrospective analysis of immune checkpoint inhibitor efficacy reported significantly improved OS for patients with tumors harboring TMB  $\geq 9.8$  Muts/MB compared with those with tumors with TMB  $< 9.8$  Muts/Mb (~equivalency  $< 12$  Muts/Mb as measured by this assay)<sup>34</sup>. Another retrospective study reported that a TMB  $\geq 12$  Muts/Mb cutoff identifies  $> 99\%$  of MSI-High CRC cases but only 3% of MSS cases, indicating the utility of this cutoff for identification of patients with CRC likely to benefit from treatment with immune checkpoint inhibitors<sup>51</sup>.

## FREQUENCY & PROGNOSIS

Elevated tumor mutational burden (TMB) has been reported in 8-25% of colorectal cancer (CRC) samples<sup>21,52-53</sup>. Multiple studies have reported that up to 90% of hypermutated CRC cases exhibit high levels of microsatellite instability (MSI-H) and mismatch repair deficiency (MMR-D)<sup>21,52</sup>. Increased TMB is significantly associated with MSI-H and MMR-D, with studies reporting that 100% of MSI-H CRCs harbor elevated TMB and conversely that 100% of tumors with low TMB harbor intact MMR<sup>52</sup>. A subset of CRCs that harbor increased TMB but not MSI-H are driven by mutations in POLE, which leads to an "ultramutated" phenotype with especially high TMB<sup>21,52</sup>. Tumors with increased TMB harbor BRAF V600E mutations more frequently than those with low TMB<sup>21,52</sup>, whereas TMB-low tumors more frequently harbor mutations in TP53 and APC<sup>21</sup>. The prognostic value of tumor mutational burden (TMB) in colorectal cancer (CRC) is context- and therapy-dependent. A

study of tissue TMB (tTMB) in 145 CRC samples showed longer OS in TMB-high samples compared with TMB-low ones<sup>54</sup>. Similarly, for patients with metastatic CRC treated with first-line chemotherapy combined with bevacizumab or cetuximab, high tissue TMB (tTMB-H) was associated with longer OS<sup>55</sup>. For patients treated with adjuvant chemotherapy, tTMB-H was associated with better 5-year relapse-free survival<sup>56</sup>. However, for patients with EGFR/BRAF-inhibitor-treated, BRAF-mutated microsatellite stable (MSS) metastatic CRC, intermediate tTMB was associated with significantly poorer PFS and OS compared with TMB-low status; patients with primary resistance to EGFR/BRAF blockage had higher TMB than those sensitive to these therapies<sup>57</sup>. In a study for 61 patients with metastatic, MSS CRC treated with best standard of care, plasma TMB scores  $\geq 28$  Muts/Mb (approximately 14 Muts/Mb as measured by this assay) were associated with reduced OS compared with plasma TMB scores  $< 28$  Muts/Mb (3.0 vs. 5.3 months, HR=0.76,  $p=0.007$ ), whereas tTMB was not found to be prognostic in this population<sup>58</sup>.

## FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma<sup>59-60</sup> and cigarette smoke in lung cancer<sup>61-62</sup>, treatment with temozolomide-based chemotherapy in glioma<sup>63-64</sup>, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes<sup>21,65-68</sup>, and microsatellite instability (MSI)<sup>21,65,68</sup>. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents<sup>34,44,51</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

GENOMIC FINDINGS

GENE  
**KRAS**

ALTERATION  
wildtype

with clinical benefit of treatment with EGFR-targeting antibodies cetuximab<sup>69-72</sup> or panitumumab<sup>73-75</sup> for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

Numerous studies have reported that KRAS wild-type status is associated with decreased metastasis, better clinicopathological features, and longer survival of patients with CRC<sup>78-81,85-86</sup>.

**FINDING SUMMARY**

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor formation<sup>87-88</sup>. No alterations in KRAS were identified in this case.

**POTENTIAL TREATMENT STRATEGIES**

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated

**FREQUENCY & PROGNOSIS**

Approximately 50-65% of colorectal cancers (CRCs) have been reported to lack KRAS mutations<sup>76-84</sup>.

GENE  
**NRAS**

ALTERATION  
wildtype

targeting antibodies cetuximab<sup>69-72</sup> or panitumumab<sup>73-75</sup> for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

frequency of metastasis<sup>84</sup> and longer survival<sup>94-95</sup> of patients with CRC.

**FINDING SUMMARY**

NRAS encodes a member of the RAS family of small GTPases that mediate transduction of growth signals. Activation of RAS signaling causes cell growth, differentiation, and survival by activating the RAF-MAPK-ERK, PI3K, and other pathways<sup>87</sup>. No alterations in NRAS were identified in this case.

**POTENTIAL TREATMENT STRATEGIES**

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated with clinical benefit of treatment with EGFR-

**FREQUENCY & PROGNOSIS**

The majority of colorectal cancers (CRCs) (91-98%) have been reported to lack NRAS mutations<sup>21,84,89-94</sup>. NRAS wild-type status has been reported to be associated with decreased

GENE  
**NF1**

ALTERATION  
loss exons 37-58

nerve sheath tumors (MPNST)<sup>110</sup>.

— Potential Resistance —

Multiple clinical studies report that inhibitors of the PI3K-AKT-mTOR pathway have not produced significant clinical benefit as monotherapies to treat CRC, even for tumors that harbor alterations in PIK3CA or PTEN; data are more limited for alterations in other genes in this pathway<sup>111-113</sup>.

**FINDING SUMMARY**

NF1 encodes neurofibromin, a GTPase-activating protein (GAP) that is a key negative regulator of the RAS signaling pathway<sup>115</sup>. Neurofibromin acts as a tumor suppressor by repressing RAS signaling<sup>116</sup>. Alterations such as seen here may disrupt NF1 function or expression<sup>116-125</sup>.

**POTENTIAL TREATMENT STRATEGIES**

— Targeted Therapies —

On the basis of clinical evidence in neurofibromatosis Type 1-associated neurofibroma<sup>96-99</sup>, glioma or glioblastoma<sup>99-103</sup>, and non-small cell lung cancer<sup>104</sup>, NF1 inactivation may predict sensitivity to MEK inhibitors such as cobimetinib, trametinib, binimetinib, and selumetinib. On the basis of limited clinical data<sup>105-107</sup> and preclinical data<sup>108-109</sup>, loss or inactivation of NF1 may predict sensitivity to mTOR inhibitors, including everolimus and temsirolimus. A preclinical study suggests that combined mTOR and MEK inhibition is effective in a model of NF1-deficient malignant peripheral

**FREQUENCY & PROGNOSIS**

In the Colorectal Adenocarcinoma TCGA dataset, NF1 mutations have been found in 1.4% of sequenced tumors<sup>21</sup>. For patients with colorectal cancer (CRC), NF1 mutations have been reported to be associated with significantly worse OS following treatment with cetuximab or bevacizumab (p=0.04 and 0.007, HR=2.62 and 2.05); low NF1 expression was associated with significantly worse OS (p=0.003, HR=1.49) and PFS (p=0.02, HR=1.36) when compared with high NF1 expression<sup>114</sup>.

**POTENTIAL GERMLINE IMPLICATIONS**

Germline mutations in NF1 cause the autosomal dominant disorder neurofibromatosis type 1, which is characterized in part by increased risk of developing various tumors, including sarcoma, glioma, breast carcinoma, and neuroendocrine and hematological neoplasms<sup>126-128</sup>. Estimates for the prevalence of the disorder in the general population range from 1:2,500 to 1:3,000<sup>129-130</sup>, and in the appropriate clinical context, germline testing of NF1 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**GENOMIC FINDINGS**
**GENE**

# APC

**ALTERATION**

R232\*

**HGVS VARIANT**

NM\_000038.4: c.694C&gt;T (p.R232\*)

**VARIANT CHROMOSOMAL POSITION**

chr5:112128191

**VARIANT ALLELE FREQUENCY (% VAF)**

56.7%

**POTENTIAL TREATMENT STRATEGIES**
**— Targeted Therapies —**

There are no approved drugs targeting APC inactivation in cancer. Loss of APC function leads to accumulation of beta-catenin and upregulation of WNT pathway transcription programs<sup>131</sup>, and potential therapeutic approaches to target this pathway include CBP/beta-catenin antagonists, which interfere with the ability of beta-catenin to

interact with transcriptional co-activator CBP<sup>132-133</sup>. In a Phase 1 trial of the CBP/beta-catenin antagonist E7386, 1 patient with APC-mutated small bowel adenocarcinoma achieved a PR with tumor shrinkage of -69% and response duration of 165 days<sup>134</sup>; preclinical data support sensitivity of APC-deficient gastric or colorectal cancer models to E7386<sup>135-136</sup>.

**FREQUENCY & PROGNOSIS**

APC mutations have been found in 73% of tumors in the colorectal adenocarcinoma TCGA dataset<sup>21</sup>. In 1 study, loss of heterozygosity (LOH) of APC was observed in 32% of colorectal cancer (CRC) samples<sup>137</sup>. The prognostic significance of APC mutations in sporadic CRC remains unclear<sup>138</sup>. Solid tumors with WNT/beta-catenin pathway alterations, as seen here, were observed to have significantly less T-cell inflammation in one study<sup>139</sup>.

**FINDING SUMMARY**

APC (adenomatous polyposis coli) encodes a tumor

suppressor with critical roles in regulating cell division and adhesion. APC interacts with beta-catenin and controls signaling in the WNT pathway, which regulates embryonic development and cell differentiation<sup>140</sup>. Alterations such as seen here may disrupt APC function or expression<sup>141-145</sup>.

**POTENTIAL GERMLINE IMPLICATIONS**

One or more of the APC variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with familial adenomatous polyposis (ClinVar, Sep 2022)<sup>146</sup>. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in APC are found in more than 90% of patients with familial adenomatous polyposis (FAP)<sup>147-149</sup>. The prevalence for FAP in the general population is estimated to be 1:8,300 from birth<sup>150</sup>, and in the appropriate clinical context germline testing of APC is recommended.

**GENE**

# CCND2

**ALTERATION**

amplification

**POTENTIAL TREATMENT STRATEGIES**
**— Targeted Therapies —**

Although preclinical studies suggest that cyclin D2 activates CDK4/6<sup>151-152</sup>, it is unknown whether CCND2 amplification or activating mutation predicts response to CDK4/6 inhibitors such as

abemaciclib, palbociclib, and ribociclib. Clinical studies of CDK4/6 inhibitors have shown the most promise for estrogen receptor-positive breast cancer<sup>153-154</sup>.

**FREQUENCY & PROGNOSIS**

In the Colorectal Adenocarcinoma TCGA dataset, putative high-level amplification of CCND2 has been found in 1.4% of cases<sup>21</sup>. Cyclin D2 has also reportedly been overexpressed in 53% of colorectal adenocarcinomas<sup>155</sup>. Overexpression of CCND2 mRNA in CRC has been correlated with poor patient prognosis<sup>156</sup>. In addition, the expression of cyclin D2 at the invasive margin has been

associated with liver metastasis in colorectal cancer, and in patients with Stage 1 and 2 tumors, cyclin D2 expression has also been associated with reduced patient survival<sup>157</sup>.

**FINDING SUMMARY**

CCND2 encodes the protein cyclin D2, which binds and regulates the cyclin-dependent kinases that control cell cycle progression, and is a downstream target of cancer signaling pathways including hedgehog and PI3K<sup>158-159</sup>. CCND2 has been reported to be amplified in cancer<sup>160</sup>, and may be biologically relevant in this context<sup>161-162</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1606878-01

GENOMIC FINDINGS

GENE

**ERBB2**

ALTERATION

T798I

HGVS VARIANT

NM\_004448.2: c.2393C>T (p.T798I)

VARIANT CHROMOSOMAL POSITION

chr17:37881064

VARIANT ALLELE FREQUENCY (% VAF)

39.2%

tucatinib<sup>177-180</sup>, and dual EGFR/HER2 kinase inhibitors such as lapatinib<sup>181-189</sup>, afatinib<sup>168,190-199</sup>, neratinib<sup>200-203</sup>, dacomitinib<sup>204</sup>, and pyrotinib<sup>205-206</sup>. The Phase 1 trial of HER2-selective TKI BI-1810631 for patients with HER2-aberration-positive metastatic solid tumors reported a 3.7% ORR (7/19) and an 84% DCR; for patients with NSCLC, a 45% ORR (5/11) and a 91% DCR were reported<sup>207</sup>. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

4% and 2-6% of colorectal adenocarcinoma cases, respectively<sup>21,216-218</sup>. For patients with colorectal cancer, multiple studies have shown that ERBB2 overexpression does not correlate with survival<sup>218-220</sup> and is not considered prognostic (NCCN Colon Cancer Guidelines, v3.2022, NCCN Rectal Guidelines, v4.2022); however, studies have shown an association of ERBB2 amplification with reduced response and/or shorter survival for patients treated with anti-EGFR antibodies<sup>221-222</sup>.

FINDING SUMMARY

ERBB2 (also known as HER2) encodes a receptor tyrosine kinase which is in the same family as EGFR. The ERBB2 gatekeeper mutation T798I has been found to lack transforming ability<sup>212,215</sup>, but was reported to be an emergent mutation associated with acquired resistance to neratinib<sup>215</sup>. Preclinical studies have reported that this mutation also confers reduced sensitivity to lapatinib<sup>212-213</sup>. ERBB2 T798I retains sensitivity to other irreversible pan-HER inhibitors such as afatinib<sup>215</sup> and PD168393<sup>213</sup>.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of extensive clinical evidence, ERBB2 amplification or activating mutation may predict sensitivity to therapies targeting HER2, including antibodies such as trastuzumab<sup>163-168</sup>, pertuzumab in combination with trastuzumab<sup>165,169-171</sup>, and zanidatamab (ZW25)<sup>172</sup>, as well as antibody-directed conjugates such as ado-trastuzumab emtansine (T-DM1)<sup>173</sup> and fam-trastuzumab deruxtecan (T-DXd)<sup>174-176</sup>, HER2 kinase inhibitors such as

— Potential Resistance —

Tyrosine kinase inhibitors such as afatinib and lapatinib have yielded limited clinical efficacy as monotherapies in CRC, suggesting antibody therapeutics or combination therapies may be more beneficial in this tumor type<sup>208-211</sup>. Limited clinical and preclinical data indicate that ERBB2 T798I may confer resistance to lapatinib<sup>212-213</sup> and neratinib<sup>214</sup> and remain sensitive to afatinib<sup>215</sup> and PD168393<sup>213</sup>.

FREQUENCY & PROGNOSIS

ERBB2 mutation or amplification was observed in

GENE

**FGF23**

ALTERATION

amplification

that pan-FGFR inhibitors show activity in FGF amplified cancers; following treatment with a selective pan-FGFR inhibitor, a patient with head and neck squamous cell carcinoma (HNSCC) and amplification of 11q13 (FGF3, FGF4, FGF19) and 12p13 (FGF6 and FGF23) experienced a radiologic CR<sup>223</sup>.

sarcoma (4.3%), colorectal adenocarcinoma (4.2%), lung adenocarcinoma (3.7%), and head and neck squamous cell carcinoma (3.4%) (cBioPortal, 2023)<sup>160,224</sup>.

FINDING SUMMARY

FGF23 encodes a member of the fibroblast growth factor protein family that plays a central role in phosphate homeostasis<sup>225</sup>. Overexpression of FGF23 by tumor cells can cause hypophosphatemia through excessive renal phosphate clearance<sup>226</sup>, while germline gain-of-function (protein stabilizing) mutations in FGF23 cause autosomal dominant hypophosphatemic rickets<sup>227</sup>.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies that directly address genomic alterations in FGF23. Inhibitors of FGF receptors, however, are undergoing clinical trials in a number of different cancers. Limited data suggest

FREQUENCY & PROGNOSIS

FGF23 alterations have been reported with highest incidence in uterine carcinosarcoma (7.0%), ovarian carcinoma (6.5%), testicular germ cell cancer (5.4%), cutaneous melanoma (5.0%), low-grade glioma (4.9%), lung squamous cell carcinoma (4.5%),

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

GENOMIC FINDINGS

GENE  
**FGF6**

ALTERATION  
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies that directly address genomic alterations in FGF6. Inhibitors of FGF receptors, however, are undergoing clinical trials in a number of different cancers. Limited data suggest that pan-FGFR inhibitors show activity in FGF amplified cancers; following treatment with a selective pan-FGFR inhibitor, a patient with head and neck squamous cell carcinoma (HNSCC) and amplification of 11q13 (FGF3, FGF4, FGF19) and

12p13 (FGF6 and FGF23) experienced a radiologic CR<sup>223</sup>.

FREQUENCY & PROGNOSIS

Somatic alterations affecting FGF6 are infrequently documented, with the highest rates reported in penile cancer (4%), cutaneous melanoma (1-3%), stomach carcinoma (1-3%) and colorectal cancer (1%) (cBioPortal, COSMIC, 2023)<sup>160,224,228</sup>. Amplification of FGF6 has been frequently observed in testicular germ cell cancer (5%) and ovarian serous cystadenocarcinoma (5%), and in 2-6% of lower-grade gliomas, glioblastomas, sarcomas, breast invasive carcinomas, uterine carcinosarcomas, lung squamous cell carcinomas (SCC), head and neck SCC, pancreatic adenocarcinomas, and esophageal carcinomas (cBioPortal, 2023)<sup>160,224</sup>. FGF6 is co-localized with FGF23 and CCND2 at chromosomal locus 12p13

and has been reported to be co-amplified with these genes in 1.3% of patients with breast cancer<sup>229</sup>. FGF6 expression has been reported in 54% (14/26) of prostate cancer samples, which also frequently express FGFR4<sup>230</sup>. FGF6 expression has also been observed in 71% (12/17) of patients with childhood acute lymphoblastic leukemia<sup>231</sup>.

FINDING SUMMARY

FGF6 (also known as HST-2) encodes a member of the fibroblast growth factor protein family and is hypothesized to play a role in muscle tissue regeneration<sup>232</sup> by signaling through FGFR4, and to a lesser extent FGFR1 and FGFR2<sup>233</sup>. FGF6 expression has been observed in several cancers<sup>230-231,234</sup> and was shown to be oncogenic in preclinical models<sup>234-235</sup>. FGF6 has been reported as amplified in cancer<sup>160</sup> and may be biologically relevant in this context<sup>161-162</sup>.

GENE  
**KDM5A**

ALTERATION  
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to directly address genomic alterations in KDM5A. However, multiple preclinical studies have identified potential targets in KDM5A amplified or activated cells that may respond to therapy. KDM5A-mediated chromatin remodeling induces CCND1 expression and represses CDKI expression<sup>236-240</sup>; therefore, KDM5A activation or amplification may sensitize cells to CDK4/CDK6 inhibitors. Drug-resistant cell populations, characterized by elevated KDM5A expression, responded to histone deacetylase (HDAC) inhibition<sup>241</sup>, suggesting that HDAC inhibitors may be a potential therapeutic

option. KDM5A also induces expression of VEGF and promotes angiogenesis, oncogenic transformation, and tumorigenesis, which can be inhibited by KDM5A knockdown<sup>242-243</sup>, suggesting that tumors harboring KDM5A amplification may be sensitive to angiogenesis inhibitors, including kinase inhibitors that target the VEGF receptors, such as sunitinib, sorafenib, vandetanib, ponatinib, cabozantinib, regorafenib, pazopanib, and axitinib. However, these inhibitors have yet to be extensively tested in the context of KDM5A amplification or activation; therefore, it is not known if these therapeutic strategies are relevant.

FREQUENCY & PROGNOSIS

KDM5A amplification has been reported with the highest incidence in TCGA datasets in ovarian serous cystadenocarcinoma (7.2%), testicular germ cell cancer (5.8%), pancreatic adenocarcinoma (4.3%), and lung squamous cell carcinoma (3.9%) (cBioPortal, Jan 2023)<sup>160,224</sup>. Elevated levels of KDM5A expression have also been reported in a range of solid tumor types<sup>237-238,240,242,244-245</sup>, and

fusion of KDM5A to NUP98 has been documented in acute myeloid leukemia<sup>246-247</sup>. KDM5A expression is significantly correlated with HIF-1α and VEGF expression, as well as tumor size, angiogenesis, and poor patient prognosis in lung cancer<sup>243</sup>.

FINDING SUMMARY

KDM5A encodes a lysine-specific histone demethylase that potentiates the expression of genes involved in cellular proliferation, senescence, angiogenesis, and migration<sup>236-237,242,248-249</sup>. KDM5A overexpression alters the transcriptional regulation of cell cycle genes, including CCND1, and a variety of cyclin-dependent kinase inhibitors (CDKIs), including CDKN1A, CDKN1B, and CDKN2A, and results in cell cycle progression<sup>236-240,242</sup>. Additionally, elevated KDM5A expression and associated chromatin remodeling has been implicated in resistance to various tyrosine kinase inhibitors in vitro, including erlotinib and gefitinib<sup>241,244,250</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1606878-01

GENOMIC FINDINGS

GENE

# NOTCH3

ALTERATION

splice site 119-156\_197+54>GGGGG

HGVS VARIANT

NM\_000435.2: c.119-156\_197+54delinsGGGGG (p.?)

VARIANT CHROMOSOMAL POSITION

chr19:15308257-15308545

VARIANT ALLELE FREQUENCY (% VAF)

11.2%

## POTENTIAL TREATMENT STRATEGIES

### — Targeted Therapies —

Several approaches for inhibiting NOTCH3 signaling are being developed, including neutralizing NOTCH antibodies such as tarextumab (OMP-59R5)<sup>251</sup>, which targets NOTCH2 and NOTCH3, and pan-NOTCH inhibitors, such as gamma-secretase inhibitors (GSI)<sup>252-254</sup>. In a Phase 2 study, the GSI AL101 (BMS-906024) elicited PR in 15% (6/39) and SD in 54% (21/39) of patients

with metastatic adenoid cystic carcinoma harboring NOTCH activating alterations<sup>255</sup>. Phase 2 studies have evaluated the efficacy of tarextumab in combination with chemotherapy in metastatic pancreatic cancer or extensive-stage small cell lung cancer, though NOTCH3 expression was not found to be a predictor of OS or PFS in either study<sup>256</sup>. These approaches would not be relevant in the context of inactivating alterations, as seen here.

## FREQUENCY & PROGNOSIS

NOTCH3 mutations have been reported in 6% of colorectal adenocarcinomas (CRC), whereas NOTCH3 amplification was reported in 1% of cases<sup>21</sup>. NOTCH3 rearrangements were reported in <0.1% of colorectal cancers<sup>257</sup>. NOTCH3 activation has been reported to be frequent in CRC, with mRNA and/or protein overexpression reported in 19.7-38% of samples<sup>258-260</sup>. NOTCH3 expression is more frequent in aggressive colorectal micropapillary carcinoma (35/51, 69%) than in microsatellite-stable CRC (42/97, 43%) cases (P=0.005)<sup>261</sup>. Nuclear (activated) NOTCH3 expression has been associated with advanced

disease and tumor recurrence in patients with Stage 2 and 3 colorectal cancer<sup>259</sup>.

## FINDING SUMMARY

NOTCH3 encodes a member of the NOTCH family of receptors, which are involved in cell fate determination and various developmental processes. Upon binding of membrane-bound ligands, NOTCH signaling involves cleavage of the NOTCH intracellular domain (NICD), which subsequently forms part of a transcription factor complex that regulates downstream target genes<sup>262-263</sup>. Alterations that disrupt or remove the transmembrane domain (amino acids 1644-1664), RAM domain (amino acids 1665-1837), and/or ANK repeats region (amino acids 1838-2000), which are necessary for the transcriptional activity of NOTCH family proteins, as well as internal deletions that remove EGF repeats (7-10 and 21-22), have been shown in vitro to negatively affect ligand binding and reduce NOTCH3 transcriptional activity and are predicted to be inactivating<sup>263-267</sup>.

GENE

# SMAD4

ALTERATION

loss

## POTENTIAL TREATMENT STRATEGIES

### — Targeted Therapies —

There are no targeted therapies available to address genomic alterations in SMAD4. Preclinical studies in colorectal cancer have reported associations of SMAD4 inactivation or loss with sensitivity to inhibitors of Aurora kinase A<sup>268</sup> and the Wnt/beta-catenin pathway<sup>269</sup>.

### — Nontargeted Approaches —

Clinical studies have reported associations of SMAD4 loss or low SMAD4 expression with improved responses to chemotherapeutic agents in patients with pancreatic cancer<sup>270-272</sup> and non-small cell lung cancer (NSCLC)<sup>273</sup>. Other clinical studies in pancreatic cancer have reported an association of high SMAD4 expression with better

responses to neoadjuvant chemotherapy<sup>274</sup> and adjuvant chemoradiotherapy<sup>275</sup>.

## FREQUENCY & PROGNOSIS

SMAD4 mutation or homozygous deletion is most frequently observed in pancreatic adenocarcinoma (43%)<sup>276</sup>, pancreatic acinar cell carcinoma (26%)<sup>277</sup>, cholangiocarcinoma (25%)<sup>278</sup>, small intestine cancer (20%)<sup>279</sup>, appendiceal adenocarcinoma (14-20% mutation; 57% deletion)<sup>280-281</sup>, colorectal adenocarcinoma (CRC; 14%)<sup>21</sup>, esophageal adenocarcinoma (14%)<sup>282</sup>, and stomach adenocarcinoma (13%)<sup>283</sup>. In preclinical studies, SMAD4 loss of function has been implicated in the development of mucinous neoplasms of the pancreas, including mucinous cystic neoplasms (MCN)<sup>284</sup> and intraductal papillary mucinous neoplasms (IPMN)<sup>285</sup>; in clinical samples, SMAD4 homozygous deletion has been observed in 10% of IPMNs and 8% of MCNs, and mutation was also observed in 5% of IPMNs<sup>286</sup>. SMAD4 gene alterations have been associated with reduced OS for patients with pancreatic adenocarcinoma<sup>287</sup>. Reduced SMAD4 expression has been associated with worse prognosis in various cancer types, including CRC<sup>288-290</sup>, appendiceal mucinous

neoplasm<sup>291</sup>, gastric adenocarcinoma<sup>292-293</sup>, esophageal adenocarcinoma<sup>294</sup>, esophageal squamous cell carcinoma<sup>295</sup>, breast cancer<sup>296</sup>, and prostate cancer<sup>297</sup>.

## FINDING SUMMARY

SMAD4, also known as DPC4, encodes a tumor suppressor that regulates transcriptional activity downstream of TGF-beta receptor signaling<sup>298-299</sup>. SMAD4 alterations that result in loss or disruption of the MH1 domain (aa 18-142), MH2 domain (aa 323-552), or SAD domain (aa 275-320) are predicted to be inactivating<sup>300-313</sup>.

## POTENTIAL GERMLINE IMPLICATIONS

Germline SMAD4 mutations, including those at the R361 hotspot, have been observed in patients with juvenile polyposis syndrome<sup>314-316</sup>, which is associated with an increased risk of gastrointestinal cancers<sup>317</sup>. The penetrance of deleterious SMAD4 mutations in patients with colon cancer is estimated at 20% by age 35 and 70% by age 65<sup>318</sup>. In the appropriate clinical context, germline testing of SMAD4 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

## GENOMIC FINDINGS

## GENE

## TP53

## ALTERATION

P151R

## HGVS VARIANT

NM\_000546.4: c.452C&gt;G (p.P151R)

## VARIANT CHROMOSOMAL POSITION

chr17:7578478

## VARIANT ALLELE FREQUENCY (% VAF)

67.3%

## POTENTIAL TREATMENT STRATEGIES

## — Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib<sup>319-322</sup> or p53 gene therapy such as SGT53<sup>323-327</sup>. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype<sup>328</sup>. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer<sup>329</sup>. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer<sup>330</sup>. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone<sup>331</sup>. In the Phase 2

VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel<sup>332</sup>. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations<sup>333</sup>. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring<sup>334</sup>. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage<sup>327</sup>. Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR<sup>335</sup>. A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)<sup>336</sup>.

## FREQUENCY &amp; PROGNOSIS

TP53 mutations have been reported in up to 75% of colorectal cancer cases<sup>21,337-342</sup>. A study reported p53 expression in 49% of analyzed colorectal cancer cases<sup>343</sup>. TP53 mutation has not been consistently demonstrated to be a significant independent prognostic marker in the context of CRC<sup>344</sup>.

## FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers<sup>345</sup>. Alterations such as seen here may disrupt TP53 function or expression<sup>346-350</sup>.

## POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers<sup>351-353</sup>, including sarcomas<sup>354-355</sup>. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000<sup>356</sup> to 1:20,000<sup>355</sup>. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30<sup>357</sup>. In the appropriate clinical context, germline testing of TP53 is recommended.

## POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion<sup>358-363</sup>. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy<sup>358-359</sup>. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease<sup>364</sup>. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH<sup>362,365-366</sup>. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**THERAPIES WITH CLINICAL BENEFIT**
**IN PATIENT'S TUMOR TYPE**

## Cetuximab

*Assay findings association*
**KRAS**  
 wildtype

**NRAS**  
 wildtype

### AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Therapies targeting EGFR, including cetuximab, have been shown to have significant clinical activity for patients with CRC<sup>69-72,367-368</sup>; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of cetuximab in metastatic CRC (NCCN Colon Cancer Guidelines, v3.2022, NCCN Rectal Cancer Guidelines, v4.2022).

### SUPPORTING DATA

Cetuximab has been shown to improve OS, PFS, and response rate for patients with KRAS-wildtype colorectal cancer (CRC), both in combination with FOLFIRI, FOLFOX<sub>4</sub>, or irinotecan<sup>69-70,367-369</sup> and as monotherapy for chemotherapy-refractory patients<sup>72,370</sup>. The Phase 3 study STRATEGIC-1 reported a similar duration of disease control (DDC) for patients with unresectable metastatic CRC (mCRC) and KRAS-, NRAS-, and BRAF-wildtype status treated with mFOLFOX-bevacizumab alternated

with a cetuximab regimen in first or second line, respectively (overall DDC 22.5 vs. 23.5 months); in addition, the study reported similar OS (37.8 vs. 34.4 months) and higher numerical ORR for patients treated with cetuximab in the first line followed by mFOLFOX-bevacizumab compared with those receiving EGFR-directed antibodies in the second or third line<sup>371</sup>. A prospective study of cetuximab monotherapy for patients with KRAS-, NRAS-, and BRAF-wildtype mCRC reported 11% (2/19) PRs and 58% (11/19) SDs<sup>372</sup>. The Phase 2 AVETUX trial of cetuximab combined with avelumab and mFOLFOX<sub>6</sub> for patients with RAS- and BRAF-wildtype mCRC resulted in an ORR of 81% (4 CR and 27 PRs, n=37) and a DCR of 89%<sup>373</sup>. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)<sup>374</sup>. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)<sup>375</sup>.

## Panitumumab

*Assay findings association*
**KRAS**  
 wildtype

**NRAS**  
 wildtype

### AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

Therapies targeting EGFR, including panitumumab, have been shown to have significant clinical activity for patients with CRC<sup>73,374,376</sup>; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of panitumumab in metastatic CRC (NCCN Colon Cancer Guidelines v3.2022)(NCCN Rectal Cancers Guidelines, v4.2022).

### SUPPORTING DATA

Panitumumab has been shown to improve OS, PFS, and ORR for patients with KRAS-wildtype colorectal cancer (CRC), both in combination with FOLFOX<sub>4</sub>, FOLFIRI, irinotecan, or best supportive care<sup>73,377-380</sup>, and as monotherapy for chemotherapy-refractory patients<sup>340,374,376</sup>. The Phase 3 PARADIGM trial comparing panitumumab plus mFOLFOX<sub>6</sub> versus

bevacizumab plus mFOLFOX<sub>6</sub> as first-line treatment for patients with RAS-wildtype left-sided metastatic CRC demonstrated that treatment with panitumumab significantly improved median OS (mOS; 36.2 months vs. 31.3 months) compared with bevacizumab<sup>381</sup>. A Phase 2 trial reported that, for patients with unresectable RAS-wildtype colorectal adenocarcinoma treated with panitumumab plus FOLFOX<sub>4</sub>, maintenance with a combination of panitumumab plus fluorouracil and leucovorin was superior to panitumumab monotherapy (10-month PFS OF 59% vs. 49%)<sup>382</sup>. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)<sup>374</sup>. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)<sup>375</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**THERAPIES WITH CLINICAL BENEFIT**
**IN OTHER TUMOR TYPE**

## Cobimetinib

*Assay findings association*
**NF1**  
 loss exons 37-58

### AREAS OF THERAPEUTIC USE

Cobimetinib is a MEK inhibitor that is FDA approved to treat patients with histiocytic neoplasms. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma<sup>96-99,383-387</sup>, glioma<sup>99-103,388</sup>, and non-small cell lung cancer<sup>104</sup>, NF1 inactivation may predict sensitivity to MEK inhibitors.

### SUPPORTING DATA

The Phase 3 IMblaze370 study of cobimetinib plus atezolizumab for patients with chemotherapy-refractory metastatic colorectal cancer (CRC) reported similar median OS (8.9 vs. 7.1 vs. 8.5 months, HR=1.0–1.2), median PFS (1.9 vs. 1.9 vs. 2.0 months), and ORR (2.7% vs. 2.2% vs. 2.2%) to atezolizumab monotherapy or regorafenib, respectively<sup>389</sup>.

## Selumetinib

*Assay findings association*
**NF1**  
 loss exons 37-58

### AREAS OF THERAPEUTIC USE

Selumetinib is a MEK inhibitor that is FDA approved to treat pediatric patients with neurofibromatosis type 1 (NF1)-associated plexiform neurofibromas (PNs). Please see the drug label for full prescribing information.

### GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma<sup>96-99,383-387</sup>, glioma<sup>99-103,388</sup>, and non-small cell lung cancer<sup>104</sup>, NF1 inactivation may predict sensitivity to MEK inhibitors.

### SUPPORTING DATA

In Phase 2 studies for chemotherapy-refractory metastatic

colorectal cancer (CRC), single-agent selumetinib demonstrated similar outcomes compared with capecitabine in genomically unselected patients (ORR, 0% [0/34] vs. 2.8% [1/35]; median PFS [mPFS], 81 vs. 88 days)<sup>390</sup>, and combination of selumetinib with irinotecan elicited an ORR of 9.7% (3/31 PRs), mPFS of 105 days, and median OS of 267 days in patients with KRAS mutation<sup>391</sup>. Phase 1 and 2 studies have evaluated selumetinib in combination with the AKT inhibitor MK-2206<sup>392-393</sup>, the EGFR-targeting monoclonal antibody cetuximab<sup>394</sup>, the EGFR-targeting TKI afatinib<sup>395</sup>, or Cyclosporin A<sup>396</sup>, but have reported limited activity and no confirmed objective responses for patients with CRC.

## Trametinib

*Assay findings association*
**NF1**  
 loss exons 37-58

### AREAS OF THERAPEUTIC USE

Trametinib is a MEK inhibitor that is FDA approved as a monotherapy to treat patients with melanoma with BRAF V600E or V600K mutations. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

On the basis of clinical evidence in neurofibromatosis type 1 (NF1)-associated neurofibroma<sup>96-99,383-387</sup>, glioma<sup>99-103,388</sup>, and non-small cell lung cancer<sup>104</sup>, NF1 inactivation may predict sensitivity to MEK inhibitors.

### SUPPORTING DATA

Trametinib has shown limited activity as a single agent in advanced or metastatic colorectal cancer (CRC), with no objective responses in 28 patients (including 13 with KRAS and 3 with BRAF mutation) included in a Phase 1 study<sup>397</sup> nor in 7 patients with non-V600 BRAF mutation included in the Phase 2 NCI-MATCH trial<sup>398</sup>. In Phase 1/2 studies for metastatic CRC (mCRC), combination of trametinib with the EGFR-targeting antibody panitumumab demonstrated modest activity for BRAF/KRAS/NRAS-wildtype patients, with 38% (5/13) confirmed ORR, 92% (12/13) DCR, and 4.4 months

median PFS<sup>399</sup>, but limited activity for patients with BRAF V600 mutation, with 0% (0/31) ORR, 55% (17/31) DCR, 2.6 months mPFS, and 8.2 months median OS<sup>400</sup>. Combination of trametinib with the PD-1-targeting antibody pembrolizumab did not elicit any objective responses in 15 patients with genomically unselected CRC in the Phase 1/2 KEYNOTE-022 study<sup>401</sup>, and trametinib plus the PD-L1-targeting antibody durvalumab showed limited activity in microsatellite stable mCRC with 3.4% (1/29) ORR and 41% (12/29) DCR<sup>402</sup>. Early phase studies examining combination of trametinib with agents targeting the PI3K-AKT-mTOR pathway<sup>403-406</sup>, CDK4/6 inhibitors palbociclib or ribociclib<sup>407-408</sup>, the multi-TKI pazopanib<sup>409</sup>, or the BCL-2 inhibitor navitoclax<sup>410</sup> have shown minimal anti-tumor activity for included patients with CRC. Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors<sup>403</sup>, a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months<sup>411</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

**NOTE** Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | [www.rochefoundationmedicine.com](http://www.rochefoundationmedicine.com)

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**

**NOTE** Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually

updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that

may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://clinicaltrials.gov). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

**GENE**
**APC**
**ALTERATION**
**R232\***
**RATIONALE**

Based on preclinical and limited clinical data, APC inactivation may be associated with sensitivity to CBP/beta-catenin interaction inhibitors.

**NCT05091346**
**PHASE 1/2**

A Study of E7386 in Combination With Pembrolizumab in Previously Treated Participants With Selected Solid Tumors

**TARGETS**

CBP, Beta-catenin, PD-1

**LOCATIONS:** Fukuoka (Japan), Osaka (Japan), Shizouka (Japan), Tokyo (Japan), Chiba-shi (Japan), Kashiwa (Japan), Sapporo shi (Japan), Glasgow (United Kingdom), Manchester (United Kingdom), London (United Kingdom)

**NCT04008797**
**PHASE 1**

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

**TARGETS**

CBP, Beta-catenin, FGFRs, RET, PDGFRA, VEGFRs, KIT

**LOCATIONS:** Kurume (Japan), Matsuyama (Japan), Seodaemun (Korea, Republic of), Osakasayama (Japan), Nagoya (Japan), Kawasaki (Japan), Chuo-Ku (Japan), Koto-ku (Japan), Chiba (Japan), Kashiwa (Japan)

**NCT03264664**
**PHASE 1**

Study of E7386 in Participants With Selected Advanced Neoplasms

**TARGETS**

CBP, Beta-catenin

**LOCATIONS:** Glasgow (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Sutton (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | [www.rochefoundationmedicine.com](http://www.rochefoundationmedicine.com)

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1606878-01

CLINICAL TRIALS

## GENE

**CCND2**

## RATIONALE

CCND2 amplification or activation may predict sensitivity to CDK4/6 inhibitors.

## ALTERATION

**amplification**
**NCT04282031**
**PHASE 1/2**

A Study of BPI-1178 in Patients With Advanced Solid Tumor and HR+/HER2- Breast Cancer

**TARGETS**

CDK6, CDK4, ER, Aromatase

**LOCATIONS:** Shanghai (China)

**NCT05480280**
**PHASE 2**

mFOLFOX6 Combined With Dalpiciclib in Patients With Metastatic Colorectal Cancer

**TARGETS**

CDK6, CDK4

**LOCATIONS:** Guangzhou (China)

**NCT04801966**
**PHASE NULL**

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

**TARGETS**

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

**LOCATIONS:** Melbourne (Australia)

**NCT03994796**
**PHASE 2**

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

**TARGETS**

TRKB, ALK, TRKC, ROS1, TRKA, CDK4, CDK6, PI3K, mTOR

**LOCATIONS:** Washington, Oregon, Idaho, Montana

**NCT05252416**
**PHASE 1/2**

(VELA) Study of BLU-222 in Advanced Solid Tumors

**TARGETS**

ER, CDK4, CDK6, CDK2

**LOCATIONS:** Illinois, Massachusetts, Arkansas, New York, Virginia, Texas, Florida

**NCT02896335**
**PHASE 2**

Palbociclib In Progressive Brain Metastases

**TARGETS**

CDK4, CDK6

**LOCATIONS:** Massachusetts

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**
**NCT04616183**
**PHASE 1/2**

LY3214996 and Cetuximab Alone or in Combination With Abemaciclib for the Treatment of Unresectable or Metastatic Colorectal Cancer

**TARGETS**

ERK1, ERK2, EGFR, CDK4, CDK6

**LOCATIONS:** Texas

**NCT05159245**
**PHASE 2**

The Finnish National Study to Facilitate Patient Access to Targeted Anti-cancer Drugs

**TARGETS**

BRAF, VEGFRs, RET, KIT, ERBB2, TRKB, ALK, TRKC, ROS1, TRKA, SMO, PD-L1, MEK, CDK4, CDK6

**LOCATIONS:** Kuopio (Finland), Helsinki (Finland), Tampere (Finland), Turku (Finland)

**NCT03310879**
**PHASE 2**

Study of the CDK4/6 Inhibitor Abemaciclib in Solid Tumors Harboring Genetic Alterations in Genes Encoding D-type Cyclins or Amplification of CDK4 or CDK6

**TARGETS**

CDK4, CDK6

**LOCATIONS:** Massachusetts

**NCT03454035**
**PHASE 1**

Ulixertinib/Palbociclib in Patients With Advanced Pancreatic and Other Solid Tumors

**TARGETS**

MAPK3, MAPK1, CDK4, CDK6

**LOCATIONS:** North Carolina

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**
**GENE**
**ERBB2**
**ALTERATION**
**T798I**
**RATIONALE**

ERBB2 amplification or activating mutation may confer sensitivity to HER2-targeted and dual EGFR/HER2-directed therapies, and may enhance efficacy of HSP90 inhibitors. Based on limited clinical and preclinical evidence, ERBB2 T798I may confer reduced sensitivity to lapatinib and neratinib. Tyrosine kinase inhibitors such as erlotinib, gefitinib, afatinib, and lapatinib have

yielded limited clinical efficacy as monotherapies in CRC, suggesting that antibody therapeutics or combination therapies may be more beneficial in this tumor type. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

**NCT04589845**
**PHASE 2**

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

**TARGETS**

TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha, RAFs, NRAS

**LOCATIONS:** Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Shanghai City (China), Shanghai (China), Shatin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China)

**NCT04579380**
**PHASE 2**

Basket Study of Tucatinib and Trastuzumab in Solid Tumors With HER2 Alterations

**TARGETS**

ERBB2, ER

**LOCATIONS:** Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Osakasayama (Japan), Nagoya-shi (Japan), Kawasaki-shi (Japan), Chuo-Ku (Japan), Tokyo (Japan), Kashiwa-shi (Japan), Poznan (Poland), Berlin (Germany)

**NCT04999761**
**PHASE 1**

AB122 Platform Study

**TARGETS**

PD-1, HSP90, FGFRs

**LOCATIONS:** Ehime (Japan), Wakayama (Japan), Osaka (Japan), Aichi (Japan), Shizuoka (Japan), Kanagawa (Japan), Tokyo (Japan), Chiba (Japan), Hokkaido (Japan)

**NCT02693535**
**PHASE 2**

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

**TARGETS**

CDK4, CDK6, FLT3, VEGFRs, CSF1R, KIT, RET, mTOR, ERBB2, MEK, BRAF, PARP, PD-1, CTLA-4, PD-L1, TRKB, ALK, TRKC, ROS1, TRKA, FGFRs

**LOCATIONS:** Hawaii, Washington, Oregon, California

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**
**NCT03297606**
**PHASE 2**

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

**TARGETS**

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

**LOCATIONS:** Vancouver (Canada), Kelowna (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

**NCT05159245**
**PHASE 2**

The Finnish National Study to Facilitate Patient Access to Targeted Anti-cancer Drugs

**TARGETS**

BRAF, VEGFRs, RET, KIT, ERBB2, TRKB, ALK, TRKC, ROS1, TRKA, SMO, PD-L1, MEK, CDK4, CDK6

**LOCATIONS:** Kuopio (Finland), Helsinki (Finland), Tampere (Finland), Turku (Finland)

**NCT04817956**
**PHASE 2**

Improving Public Cancer Care by Implementing Precision Medicine in Norway

**TARGETS**

PD-L1, VEGFA, ERBB2, ALK, RET, PARP, SMO, TRKB, TRKC, ROS1, TRKA, MEK, BRAF, PI3K-alpha, FGFR1, FGFR2, FGFR3, MET, KIT, ABL

**LOCATIONS:** Tromsø (Norway), Bodø (Norway), Hamar (Norway), Oslo (Norway), Fredrikstad (Norway), Drammen (Norway), Trondheim (Norway), Skien (Norway), Førde (Norway), Bergen (Norway)

**NCT04294628**
**PHASE 1**

Testing the Biological Effects of DS-8201a on Patients With Advanced Cancer

**TARGETS**

ERBB2

**LOCATIONS:** Ohio, Pennsylvania, Massachusetts, New York, Maryland

**NCT05395052**
**PHASE 1**

FT536 Monotherapy and in Combination With Monoclonal Antibodies in Advanced Solid Tumors

**TARGETS**

MET, EGFR, PD-1, PD-L1, ERBB2

**LOCATIONS:** California, Arizona, New Jersey, Texas, North Carolina

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**
**GENE**
**NF1**
**ALTERATION**

loss exons 37-58

**RATIONALE**

On the basis of clinical evidence and strong preclinical evidence, NF1 inactivation may predict sensitivity to MEK inhibitors. Limited clinical data and strong preclinical data indicate that loss or inactivation of NF1 may also predict sensitivity to mTOR inhibitors. Several clinical studies have

shown that inhibitors of the PI3K-AKT-mTOR pathway have not produced significant clinical benefit when used as a monotherapy in patients with colorectal cancer; combination therapies may be required to overcome this lack of response.

**NCT04337463**
**PHASE NULL**

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

**TARGETS**

mTORC1, mTORC2, PD-1

**LOCATIONS:** Chongqing (China), Chengdu (China)

**NCT04803318**
**PHASE 2**

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

**TARGETS**

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

**LOCATIONS:** Guangzhou (China)

**NCT04985604**
**PHASE 1/2**

DAY101 Monotherapy or in Combination With Other Therapies for Patients With Solid Tumors

**TARGETS**

BRAF, MEK

**LOCATIONS:** Busan (Korea, Republic of), Seoul (Korea, Republic of), Clayton (Australia), Edegem (Belgium), Oregon, Barcelona (Spain), Madrid (Spain), California, Colorado

**NCT04801966**
**PHASE NULL**

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

**TARGETS**

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

**LOCATIONS:** Melbourne (Australia)

**NCT03905148**
**PHASE 1/2**

Study of the Safety and Pharmacokinetics of BGB-283 and PD-0325901 in Patients With Advanced or Refractory Solid Tumors

**TARGETS**

RAFs, EGFR, MEK

**LOCATIONS:** Nedlands (Australia), Blacktown (Australia), Randwick (Australia), Melbourne (Australia), California, Texas

**NCT04965818**
**PHASE 1/2**

Phase 1b/2 Study of Futibatinib in Combination With Binimetinib in Patients With Advanced KRAS Mutant Cancer

**TARGETS**

MEK, FGFRs

**LOCATIONS:** California, Indiana, Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**CLINICAL TRIALS**
**NCT05159245**
**PHASE 2**

The Finnish National Study to Facilitate Patient Access to Targeted Anti-cancer Drugs

**TARGETS**

 BRAF, VEGFRs, RET, KIT, ERBB2, TRKB,  
 ALK, TRKC, ROS1, TRKA, SMO, PD-L1,  
 MEK, CDK4, CDK6

**LOCATIONS:** Kuopio (Finland), Helsinki (Finland), Tampere (Finland), Turku (Finland)

**NCT04817956**
**PHASE 2**

Improving Public Cancer Care by Implementing Precision Medicine in Norway

**TARGETS**

 PD-L1, VEGFA, ERBB2, ALK, RET, PARP,  
 SMO, TRKB, TRKC, ROS1, TRKA, MEK,  
 BRAF, PI3K-alpha, FGFR1, FGFR2,  
 FGFR3, MET, KIT, ABL

**LOCATIONS:** Tromsø (Norway), Bodø (Norway), Hamar (Norway), Oslo (Norway), Fredrikstad (Norway), Drammen (Norway), Trondheim (Norway), Skien (Norway), Førde (Norway), Bergen (Norway)

**NCT04551521**
**PHASE 2**

CRAFT: The NCT-PMO-1602 Phase II Trial

**TARGETS**

 PD-L1, AKTs, MEK, BRAF, ALK, RET,  
 ERBB2

**LOCATIONS:** Würzburg (Germany), Mainz (Germany), Heidelberg (Germany), Tübingen (Germany)

**NCT04720976**
**PHASE 1/2**

JAB-3312 Activity in Adult Patients With Advanced Solid Tumors

**TARGETS**

MEK, SHP2, PD-1, EGFR, KRAS

**LOCATIONS:** Utah, California, Arizona, Minnesota, Illinois, Michigan, Oklahoma, Missouri, Indiana, Connecticut

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



ORDERED TEST # ORD-1606878-01

## APPENDIX

## Variants of Unknown Significance

**NOTE** One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

**ALK**

NM\_004304.4: c.1528C>T  
(p.R510W)  
chr2:29543635

**BRD4**

NM\_014299.2: c.682G>A  
(p.V228I)  
chr19:15376332

**ERBB4**

NM\_005235.2: c.2936G>A  
(p.R979Q)  
chr2:212286760

**FANCC**

NM\_000136.2: c.973G>A  
(p.A325T)  
chr9:97887391

**MTOR**

NM\_004958.3: c.836G>A  
(p.G279E)  
chr1:11313900

**POLE**

NM\_006231.2: c.4603G>A  
(p.G1535S)  
chr12:133219531 and  
NM\_006231.2: c.5035C>T  
(p.R1679C)  
chr12:133218901

**RAD52**

amplification

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**APPENDIX**

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

**DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS**

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAS	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFA1	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCC1 (PD-1)	PDCC1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFB2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

**DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS**

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

\*TERC is an NCRNA

\*\*Promoter region of TERT is interrogated

**ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS**


Homologous Recombination status  
 Loss of Heterozygosity (LOH) score  
 Microsatellite (MS) status  
 Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1606878-01

## APPENDIX

## About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplstraat 3, 2440 Geel, Belgium. 

## ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:  
[www.rochefoundationmedicine.com/f1cdxtech](http://www.rochefoundationmedicine.com/f1cdxtech).

## INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

## TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g. gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

## THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

### Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

### Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical

methodology has identified as being present in <10% of the assayed tumor DNA.

## Ranking of Therapies and Clinical Trials

### Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

### Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

## NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) ([www.nccn.org](http://www.nccn.org)). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to [NCCN.org](http://NCCN.org). NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

## Limitations

1. In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | [www.rochefoundationmedicine.com](http://www.rochefoundationmedicine.com)

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

## APPENDIX

## About FoundationOne®CDx

- analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/P170019B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf). The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
  - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score  $\geq 16\%$  will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
  - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and

extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
- Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

### REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal

hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

### VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

\*Interquartile Range = 1st Quartile to 3rd Quartile

### VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | [www.rochefoundationmedicine.com](http://www.rochefoundationmedicine.com)

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1606878-01

**APPENDIX**

About FoundationOne®CDx

tumor sequencing is germline or somatic.  
 Interpretation should be based on clinical context.

**VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS**

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

**LEVEL OF EVIDENCE NOT PROVIDED**

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

**NO GUARANTEE OF CLINICAL BENEFIT**

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

**NO GUARANTEE OF REIMBURSEMENT**

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

**TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN**

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

**SELECT ABBREVIATIONS**

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

**REFERENCE SEQUENCE INFORMATION**

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.7.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. The median exon coverage for this sample is 946x. The suitability of use.



ORDERED TEST # **ORD-1606878-01**
**APPENDIX**
**References**

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Ciardiello et al., 2018; ESMO Abstract LBA-004
7. Parikh et al., 2021; DOI: 10.1038/s43018-021-00269-7
8. Fukuoka S, et al. J. Clin. Oncol. (2020) PMID: 32343640
9. Kim et al., 2020; DOI: 10.1016/j.annonc.2020.04.073
10. Zhang Y, et al. BMC Gastroenterol (2021) PMID: 34688262
11. Sinicrope FA, et al. J. Clin. Oncol. (2013) PMID: 24019539
12. Gavin PG, et al. Clin. Cancer Res. (2012) PMID: 23045248
13. Bertagnolli MM, et al. J. Clin. Oncol. (2009) PMID: 19273709
14. Van Cutsem E, et al. J. Clin. Oncol. (2009) PMID: 19451425
15. Ribic CM, et al. N. Engl. J. Med. (2003) PMID: 12867608
16. Sargent DJ, et al. J. Clin. Oncol. (2010) PMID: 20498393
17. Fallik D, et al. Cancer Res. (2003) PMID: 14522894
18. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
19. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
20. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
21. Nature (2012) PMID: 22810696
22. Histopathology (2007) PMID: 17204026
23. Samowitz WS, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11535541
24. Elsaleh H, et al. Clin Colorectal Cancer (2001) PMID: 12445368
25. Brueckl WM, et al. Anticancer Res. ( ) PMID: 12820457
26. Guidoboni M, et al. Am. J. Pathol. (2001) PMID: 11438476
27. Gryfe R, et al. N. Engl. J. Med. (2000) PMID: 10631274
28. Sinicrope FA, et al. Gastroenterology (2006) PMID: 16952542
29. Laghi L, et al. Dig Dis (2012) PMID: 22722556
30. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
31. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
32. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
33. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
34. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
35. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
36. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
37. Cristescu R, et al. Science (2018) PMID: 30309915
38. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
39. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
40. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
41. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
42. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
43. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
44. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
45. Ott PA, et al. J. Clin. Oncol. (2019) PMID: 30557521
46. Cristescu R, et al. J Immunother Cancer (2022) PMID: 35101941
47. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
48. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
49. Schenker et al., 2022; AACR Abstract 7845
50. Legrand et al., 2018; ASCO Abstract 12000
51. Fabrizio DA, et al. J Gastrointest Oncol (2018) PMID: 30151257
52. Stadler ZK, et al. J. Clin. Oncol. (2016) PMID: 27022117
53. Shao C, et al. JAMA Netw Open (2020) PMID: 33119110
54. Schwartz et al., 2018; ASCO Abstract 572
55. Innocenti F, et al. J Clin Oncol (2019) PMID: 30865548
56. Lee DW, et al. Clin Cancer Res (2019) PMID: 31285374
57. Randon G, et al. Eur J Cancer (2022) PMID: 34933155
58. Chen EX, et al. JAMA Oncol (2020) PMID: 32379280
59. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
60. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
61. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
62. Rizvi NA, et al. Science (2015) PMID: 25765070
63. Johnson BE, et al. Science (2014) PMID: 24336570
64. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
65. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
66. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
67. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
68. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
69. Van Cutsem E, et al. J. Clin. Oncol. (2011) PMID: 21502544
70. Bokemeyer C, et al. Ann. Oncol. (2011) PMID: 21228335
71. Karapetis CS, et al. N. Engl. J. Med. (2008) PMID: 18946061
72. De Roock W, et al. Ann. Oncol. (2008) PMID: 17998284
73. Douillard JY, et al. Ann. Oncol. (2014) PMID: 24718886
74. Douillard JY, et al. N. Engl. J. Med. (2013) PMID: 24024839
75. Amado RG, et al. J. Clin. Oncol. (2008) PMID: 18316791
76. Lièvre A, et al. Cancer Res. (2006) PMID: 16618717
77. De Roock W, et al. Lancet Oncol. (2011) PMID: 21163703
78. Chen J, et al. BMC Cancer (2014) PMID: 25367198
79. Li W, et al. BMC Cancer (2015) PMID: 25929517
80. Hu J, et al. Medicine (Baltimore) (2016) PMID: 27977612
81. Zekri J, et al. Genet. Mol. Res. (2017) PMID: 28218784
82. Staudacher JJ, et al. Clin Transl Gastroenterol (2017) PMID: 29048416
83. Wang Y, et al. Virchows Arch. (2018) PMID: 29705968
84. Guo F, et al. Sci Rep (2018) PMID: 29666387
85. Mármol I, et al. Int J Mol Sci (2017) PMID: 28106826
86. Kwak MS, et al. Medicine (Baltimore) (2017) PMID: 28858102
87. Pylyayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) PMID: 21993244
88. Kahn S, et al. Anticancer Res. ( ) PMID: 3310850
89. Pentheroudakis G, et al. BMC Cancer (2013) PMID: 23374602
90. Vaughn CP, et al. Genes Chromosomes Cancer (2011) PMID: 21305640
91. Janku F, et al. Target Oncol (2013) PMID: 23400451
92. De Roock W, et al. Lancet Oncol. (2010) PMID: 20619739
93. Irahara N, et al. Diagn. Mol. Pathol. (2010) PMID: 20376745
94. Schirripa M, et al. Int. J. Cancer (2015) PMID: 24806288
95. Cercek A, et al. Clin. Cancer Res. (2017) PMID: 28446505
96. Dombi E, et al. N. Engl. J. Med. (2016) PMID: 28029918
97. Schalkwijk S, et al. Cancer Chemother Pharmacol (2021) PMID: 33903938
98. Toledano H, et al. Childs Nerv Syst (2021) PMID: 33751171
99. Ronsley R, et al. Cancer Med (2021) PMID: 33939292
100. Fangusaro J, et al. Lancet Oncol. (2019) PMID: 31151904
101. Manoharan N, et al. J Neurooncol (2020) PMID: 32780261
102. Kondyli M, et al. J Neurooncol (2018) PMID: 30097824
103. Awada G, et al. Case Rep Oncol ( ) PMID: 33082744
104. Middleton G, et al. Nature (2020) PMID: 32669708
105. Lim SM, et al. Oncotarget (2016) PMID: 26859683
106. Weiss B, et al. Neuro-oncology (2015) PMID: 25314964
107. Janku F, et al. Oncotarget (2014) PMID: 24931142
108. Johannessen CM, et al. Curr. Biol. (2008) PMID: 18164202
109. Johannessen CM, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15937108
110. Malone CF, et al. Cancer Discov (2014) PMID: 24913553
111. Ng K, et al. Clin. Cancer Res. (2013) PMID: 23743569
112. Ganesan P, et al. Mol. Cancer Ther. (2013) PMID: 24092809
113. Janku F, et al. Cell Rep (2014) PMID: 24440717
114. Arai H, et al. Oncogene (2022) PMID: 34728807
115. Hattori S, et al. Biochem. Biophys. Res. Commun. (1991) PMID: 1904223
116. Morcos P, et al. Mol. Cell. Biol. (1996) PMID: 8628317
117. Ballester R, et al. Cell (1990) PMID: 2121371
118. Xu GF, et al. Cell (1990) PMID: 2116237
119. Martin GA, et al. Cell (1990) PMID: 2121370
120. Thomas L, et al. Hum. Mutat. (2012) PMID: 22807134
121. Skuse GR, et al. Hum. Mol. Genet. (1997) PMID: 9300663
122. Messiaen LM, et al. Genet. Med. ( ) PMID: 11258625
123. Ars E, et al. Hum. Mol. Genet. (2000) PMID: 10607834
124. Messiaen LM, et al. J. Med. Genet. (2005) PMID: 15863657
125. Pouillet P, et al. Mol. Cell. Biol. (1994) PMID: 8264648
126. Jett K, et al. Genet. Med. (2010) PMID: 20027112
127. Patil S, et al. Oncologist (2012) PMID: 22240541
128. Evans DG, et al. Clin Sarcoma Res (2012) PMID: 23036231
129. Upadhyaya M, et al. J. Med. Genet. (1995) PMID: 8544190
130. Williams VC, et al. Pediatrics (2009) PMID: 19117870
131. Zhan T, et al. Oncogene (2017) PMID: 27617575
132. Jung YS, et al. Exp Mol Med (2020) PMID: 32037398
133. Krishnamurthy N, et al. Cancer Treat Rev (2018) PMID: 29169144
134. Kawazoe et al., 2021; ESMO Abstract 473P
135. Yamada K, et al. Cancer Res (2021) PMID: 33408116
136. Kanda Y, et al. Biochem Biophys Res Commun (2022) PMID: 34837838
137. Christie M, et al. Oncogene (2013) PMID: 23085758
138. Quyn AJ, et al. Surgeon (2008) PMID: 19110823
139. Luke JJ, et al. Clin Cancer Res (2019) PMID: 30635339
140. Logan CY, et al. Annu. Rev. Cell Dev. Biol. (2004) PMID: 15473860
141. Eklof Spink K, et al. EMBO J. (2001) PMID: 11707392
142. Liu J, et al. J. Mol. Biol. (2006) PMID: 16753179
143. Dikovskaya D, et al. J. Cell. Sci. (2010) PMID: 20144988
144. Murphy SJ, et al. Dig. Dis. Sci. (2007) PMID: 17410430
145. Aretz S, et al. Hum. Mutat. (2004) PMID: 15459959
146. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
147. Kerr SE, et al. J Mol Diagn (2013) PMID: 23159591
148. Annu Rev Pathol (2011) PMID: 21090969
149. Kastiris E, et al. Int. J. Cancer (2009) PMID: 18844223

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.



ORDERED TEST # **ORD-1606878-01**
**APPENDIX**
**References**

150. Half E, et al. Orphanet J Rare Dis (2009) PMID: 19822006
151. Busk PK, et al. Exp. Cell Res. (2005) PMID: 15707582
152. Busk PK, et al. Cell Cycle ( ) PMID: 12695654
153. Finn RS, et al. Lancet Oncol. (2015) PMID: 25524798
154. DeMichele A, et al. Clin. Cancer Res. (2015) PMID: 25501126
155. Mermelshtein A, et al. Br. J. Cancer (2005) PMID: 16012517
156. Liu Y, et al. Diagn. Mol. Pathol. (2010) PMID: 21052002
157. Sarkar R, et al. Colorectal Dis (2010) PMID: 19508551
158. Katoh Y, et al. Curr. Mol. Med. (2009) PMID: 19860666
159. White PC, et al. Oncogene (2006) PMID: 16301994
160. Gao J, et al. Sci Signal (2013) PMID: 23550210
161. Zack TI, et al. Nat. Genet. (2013) PMID: 24071852
162. Beroukhim R, et al. Nature (2010) PMID: 20164920
163. Slamon DJ, et al. N. Engl. J. Med. (2001) PMID: 11248153
164. Bang YJ, et al. Lancet (2010) PMID: 20728210
165. Chumsri S, et al. J Natl Compr Canc Netw (2015) PMID: 26358791
166. Cappuzzo F, et al. N. Engl. J. Med. (2006) PMID: 16775247
167. Falchook GS, et al. J Thorac Oncol (2013) PMID: 23328556
168. Mazieres J, et al. J. Clin. Oncol. (2013) PMID: 23610105
169. Baselga J, et al. N. Engl. J. Med. (2012) PMID: 22149875
170. Swain SM, et al. N. Engl. J. Med. (2015) PMID: 25693012
171. Meric-Bernstam F, et al. Lancet Oncol. (2019) PMID: 30857956
172. Meric-Bernstam F, et al. Lancet Oncol (2022) PMID: 36400106
173. Verma S, et al. N. Engl. J. Med. (2012) PMID: 23020162
174. Modi S, et al. N. Engl. J. Med. (2019) PMID: 31825192
175. Shitara K, et al. N. Engl. J. Med. (2020) PMID: 32469182
176. Li BT, et al. N Engl J Med (2021) PMID: 34534430
177. Murthy RK, et al. N. Engl. J. Med. (2020) PMID: 31825569
178. Borges VF, et al. JAMA Oncol (2018) PMID: 29955792
179. Murthy R, et al. Lancet Oncol. (2018) PMID: 29804905
180. Moulder SL, et al. Clin. Cancer Res. (2017) PMID: 28053022
181. Fan Y, et al. Mol Oncol (2020) PMID: 32478891
182. Cameron D, et al. Oncologist (2010) PMID: 20736298
183. Geyer CE, et al. N. Engl. J. Med. (2006) PMID: 17192538
184. Serra V, et al. Cancer Discov (2013) PMID: 23950206
185. Ali SM, et al. J. Clin. Oncol. (2014) PMID: 24516025
186. Grellety T, et al. Ann. Oncol. (2016) PMID: 26487584
187. Vornicova O, et al. Oncologist (2014) PMID: 25085898
188. Ronellenfisch MW, et al. J Clin Invest (2020) PMID: 32017710
189. Hou JY, et al. Gynecol Oncol Rep (2020) PMID: 32405522
190. Lin NU, et al. Breast Cancer Res. Treat. (2012) PMID: 22418700
191. Schwab CL, et al. Br. J. Cancer (2014) PMID: 25268372
192. De Grève J, et al. Lung Cancer (2015) PMID: 25682316
193. De Grève J, et al. Lung Cancer (2012) PMID: 22325357
194. Li BT, et al. Lung Cancer (2015) PMID: 26559459
195. Dziadziuszko R, et al. J Thorac Oncol (2019) PMID: 30825613
196. Lai WV, et al. Eur. J. Cancer (2019) PMID: 30685684
197. Liu Z, et al. Onco Targets Ther (2018) PMID: 30425522
198. Fang W, et al. Oncologist (2019) PMID: 31748336
199. Yuan B, et al. Front Oncol (2020) PMID: 32477948
200. Ben-Baruch NE, et al. J Natl Compr Canc Netw (2015) PMID: 26358790
201. Ma CX, et al. Clin. Cancer Res. (2017) PMID: 28679771
202. Hyman DM, et al. Nature (2018) PMID: 29420467
203. Smyth LM, et al. Cancer Discov (2019) PMID: 31806627
204. Kris MG, et al. Ann. Oncol. (2015) PMID: 25899785
205. Jiang et al., 2019; ASCO Abstract 1001
206. Xu et al., 2020; ASCO Abstract 1003
207. Opdam et al., 2022; ENA Abstract 1LBA
208. Johnsson A, et al. Ann. Oncol. (2013) PMID: 23788755
209. Ma BB, et al. Cancer (2013) PMID: 24114668
210. Frank D, et al. J Gastrointest Oncol (2012) PMID: 22811876
211. Bouche O, et al. Anticancer Res. (2011) PMID: 21737652
212. Trowe T, et al. Clin. Cancer Res. (2008) PMID: 18413839
213. Li G, et al. PLoS ONE (2014) PMID: 25238247
214. Yamashiroya HM, et al. Am. J. Pathol. (1988) PMID: 2827495
215. Harker AB, et al. Cancer Discov (2017) PMID: 28274957
216. Ross JS, et al. Cancer (2018) PMID: 29338072
217. Ingold Heppner B, et al. Br. J. Cancer (2014) PMID: 25211663
218. Seo AN, et al. PLoS ONE (2014) PMID: 24879338
219. Wu SW, et al. Diagn Pathol (2015) PMID: 26276145
220. Scialfani F, et al. Ann. Oncol. (2013) PMID: 24146218
221. Martin V, et al. Br. J. Cancer (2013) PMID: 23348520
222. Sartore-Bianchi A, et al. Oncologist (2019) PMID: 30952821
223. Dumbava et al., 2018; doi/full/10.1200/PO.18.00100
224. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
225. Jonsson KB, et al. N. Engl. J. Med. (2003) PMID: 12711740
226. Shimada T, et al. Proc. Natl. Acad. Sci. U.S.A. (2001) PMID: 11344269
227. Yu X, et al. Cytokine Growth Factor Rev. (2005) PMID: 15863037
228. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
229. Parish A, et al. Cell Cycle (2015) PMID: 25950492
230. Ropiquet F, et al. Cancer Res. (2000) PMID: 10945637
231. Niini T, et al. Leukemia (2002) PMID: 12399964
232. Neuhaus P, et al. Mol. Cell. Biol. (2003) PMID: 12917328
233. Ornitz DM, et al. J. Biol. Chem. (1996) PMID: 8663044
234. Iida S, et al. Oncogene (1992) PMID: 1549352
235. Marics I, et al. Oncogene (1989) PMID: 2649847
236. Zeng J, et al. Gastroenterology (2010) PMID: 19850045
237. Teng YC, et al. Cancer Res. (2013) PMID: 23722541
238. Liang X, et al. PLoS ONE (2013) PMID: 23922798
239. Jiping Z, et al. J. Cell. Biochem. (2013) PMID: 23794145
240. Vieira FQ, et al. Endocr. Relat. Cancer (2014) PMID: 24200674
241. Sharma SV, et al. Cell (2010) PMID: 20371346
242. Li L, et al. Mol. Cancer (2014) PMID: 24716659
243. Qi L, et al. PLoS ONE (2014) PMID: 25162518
244. Hou J, et al. Am J Transl Res (2012) PMID: 22937203
245. Paollicchi E, et al. Crit. Rev. Oncol. Hematol. (2013) PMID: 23266085
246. van Zutven LJ, et al. Genes Chromosomes Cancer (2006) PMID: 16419055
247. Wang GG, et al. Nature (2009) PMID: 19430464
248. Chicas A, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22615382
249. Lin W, et al. Mol. Cancer Res. (2015) PMID: 25537453
250. Dannenberg JH, et al. Cell (2010) PMID: 20371339
251. Yen WC, et al. Clin. Cancer Res. (2015) PMID: 25934888
252. Hu W, et al. Cancer Res. (2014) PMID: 24743243
253. Konishi J, et al. Cancer Res. (2007) PMID: 17804716
254. Xiao Y, et al. Oncogene (2011) PMID: 20838375
255. Ferrarotto et al., 2020; ESMO Abstract 919MO
256. Hu ZI, et al. Cancer Med (2019) PMID: 31347292
257. Nguyen B, et al. Cell (2022) PMID: 35120664
258. Serafin V, et al. J. Pathol. (2011) PMID: 21598247
259. Ozawa T, et al. Ann. Surg. Oncol. (2014) PMID: 24728738
260. Furukawa S, et al. PLoS ONE (2013) PMID: 24244701
261. Lee HJ, et al. Mod. Pathol. (2013) PMID: 23060121
262. Penton AL, et al. Semin. Cell Dev. Biol. (2012) PMID: 22306179
263. Kopan R, et al. Cell (2009) PMID: 19379690
264. Deregowski V, et al. J. Bone Miner. Res. (2006) PMID: 16869730
265. Beà S, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 24145436
266. McDaniel R, et al. Am. J. Hum. Genet. (2006) PMID: 16773578
267. Lin L, et al. Cancer Res. (2010) PMID: 20068176
268. Shi C, et al. Oncogene (2022) PMID: 35393542
269. Park JW, et al. Cancer Med (2022) PMID: 35274815
270. Ormanns S, et al. Int J Mol Sci (2017) PMID: 28534865
271. Fei N, et al. Clin Transl Sci (2021) PMID: 34002944
272. Bachet JB, et al. Ann. Oncol. (2012) PMID: 22377565
273. Ziemke M, et al. Lung Cancer (2017) PMID: 28577946
274. Kassardjian A, et al. Pancreas (2020) PMID: 32897998
275. Pen SL, et al. Radiother Oncol (2021) PMID: 33667587
276. Witkiewicz AK, et al. Nat Commun (2015) PMID: 25855536
277. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
278. Churi CR, et al. PLoS ONE (2014) PMID: 25536104
279. Takeda et al., 2022; ASCO GI Abstract 642
280. Liu X, et al. Clin. Chem. (2014) PMID: 24821835
281. Maru D, et al. Oncogene (2004) PMID: 14647445
282. Wang K, et al. Oncologist (2015) PMID: 26336083
283. Nature (2014) PMID: 25079317
284. Izeradjene K, et al. Cancer Cell (2007) PMID: 17349581
285. Bardeesy N, et al. Genes Dev. (2006) PMID: 17114584
286. Springer S, et al. Gastroenterology (2015) PMID: 26253305
287. Blackford A, et al. Clin. Cancer Res. (2009) PMID: 19584151
288. Yan P, et al. Clin. Cancer Res. (2016) PMID: 26861460
289. Kozak MM, et al. J. Clin. Pathol. (2015) PMID: 25681512
290. Roth AD, et al. J. Natl. Cancer Inst. (2012) PMID: 23104212
291. Davison JM, et al. Am. J. Surg. Pathol. (2014) PMID: 24618609
292. Kim YH, et al. Ann. Oncol. (2004) PMID: 15033661
293. Xiangming C, et al. Clin. Cancer Res. (2001) PMID: 11234879
294. Singhi AD, et al. Am. J. Surg. Pathol. (2015) PMID: 25634752
295. Natsugoe S, et al. Clin. Cancer Res. (2002) PMID: 12060625
296. de Kruif EM, et al. Ann. Oncol. (2013) PMID: 23022998
297. Shipitsin M, et al. Br. J. Cancer (2014) PMID: 25032733
298. Nat. Rev. Mol. Cell Biol. (2012) PMID: 22992590
299. Cell (2008) PMID: 18662538
300. Massagué J, et al. Genes Dev. (2005) PMID: 16322555
301. Morén A, et al. Oncogene (2000) PMID: 10980615
302. Xu J, et al. Proc. Natl. Acad. Sci. U.S.A. (2000) PMID: 10781087
303. Luo K, et al. Genes Dev. (1999) PMID: 10485843
304. Jones JB, et al. Nucleic Acids Res. (2000) PMID: 10871368
305. Fink SP, et al. Cancer Res. (2001) PMID: 11196171
306. De Bosscher K, et al. Biochem. J. (2004) PMID: 14715079
307. Shi Y, et al. Nature (1997) PMID: 9214508

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1606878-01

**APPENDIX**
**References**

308. Miyaki M, et al. *Oncogene* (1999) pmid: 10340381
309. Prokova V, et al. *Biochemistry* (2007) pmid: 17994767
310. Wu JW, et al. *J. Biol. Chem.* (2001) pmid: 11274206
311. Ding L, et al. *J. Clin. Invest.* (2009) pmid: 19139564
312. Kuang C, et al. *Oncogene* (2004) pmid: 14647410
313. Watanabe M, et al. *EMBO Rep.* (2000) pmid: 11265759
314. Houlston R, et al. *Hum. Mol. Genet.* (1998) pmid: 9811934
315. Woodford-Richens K, et al. *Gut* (2000) pmid: 10764709
316. Howe JR, et al. *J. Med. Genet.* (2004) pmid: 15235019
317. Brosens LA, et al. *World J. Gastroenterol.* (2011) pmid: 22171123
318. Kalia SS, et al. *Genet. Med.* (2017) pmid: 27854360
319. Hirai H, et al. *Cancer Biol. Ther.* (2010) pmid: 20107315
320. Bridges KA, et al. *Clin. Cancer Res.* (2011) pmid: 21799033
321. Rajeshkumar NV, et al. *Clin. Cancer Res.* (2011) pmid: 21389100
322. Osman AA, et al. *Mol. Cancer Ther.* (2015) pmid: 25504633
323. Xu L, et al. *Mol. Cancer Ther.* (2002) pmid: 12489850
324. Xu L, et al. *Mol. Med.* (2001) pmid: 11713371
325. Camp ER, et al. *Cancer Gene Ther.* (2013) pmid: 23470564
326. Kim SS, et al. *Nanomedicine* (2015) pmid: 25240597
327. Pirolo KF, et al. *Mol. Ther.* (2016) pmid: 27357628
328. Leijen S, et al. *J. Clin. Oncol.* (2016) pmid: 27601554
329. Moore et al., 2019; ASCO Abstract 5513
330. Leijen S, et al. *J. Clin. Oncol.* (2016) pmid: 27998224
331. Oza et al., 2015; ASCO Abstract 5506
332. Lee J, et al. *Cancer Discov* (2019) pmid: 31315834
333. Méndez E, et al. *Clin. Cancer Res.* (2018) pmid: 29535125
334. Seligmann JF, et al. *J Clin Oncol* (2021) pmid: 34538072
335. Gourley et al., 2016; ASCO Abstract 5571
336. Park H, et al. *ESMO Open* (2022) pmid: 36084396
337. Goh HS, et al. *Cancer Res.* (1995) pmid: 7585578
338. Berg M, et al. *PLoS ONE* (2010) pmid: 21103049
339. Han SW, et al. *PLoS ONE* (2013) pmid: 23700467
340. Peeters M, et al. *Clin. Cancer Res.* (2013) pmid: 23325582
341. Malhotra P, et al. *Tumour Biol.* (2013) pmid: 23526092
342. Di Bartolomeo M, et al. *Target Oncol* (2014) pmid: 23821376
343. Wangefjord S, et al. *Diagn Pathol* (2013) pmid: 23337059
344. Russo A, et al. *J. Clin. Oncol.* (2005) pmid: 16172461
345. Brown CJ, et al. *Nat. Rev. Cancer* (2009) pmid: 19935675
346. Joerger AC, et al. *Annu. Rev. Biochem.* (2008) pmid: 18410249
347. Kato S, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2003) pmid: 12826609
348. Kamada R, et al. *J. Biol. Chem.* (2011) pmid: 20978130
349. Zerdoumi Y, et al. *Hum. Mol. Genet.* (2017) pmid: 28472496
350. Yamada H, et al. *Carcinogenesis* (2007) pmid: 17690113
351. Bougeard G, et al. *J. Clin. Oncol.* (2015) pmid: 26014290
352. Sorrell AD, et al. *Mol Diagn Ther* (2013) pmid: 23355100
353. Nichols KE, et al. *Cancer Epidemiol. Biomarkers Prev.* (2001) pmid: 11219776
354. Kleihues P, et al. *Am. J. Pathol.* (1997) pmid: 9006316
355. Gonzalez KD, et al. *J. Clin. Oncol.* (2009) pmid: 19204208
356. Lalloo F, et al. *Lancet* (2003) pmid: 12672316
357. Mandelker D, et al. *Ann. Oncol.* (2019) pmid: 31050713
358. Jaiswal S, et al. *N. Engl. J. Med.* (2014) pmid: 25426837
359. Genovese G, et al. *N. Engl. J. Med.* (2014) pmid: 25426838
360. Xie M, et al. *Nat. Med.* (2014) pmid: 25326804
361. Acuna-Hidalgo R, et al. *Am. J. Hum. Genet.* (2017) pmid: 28669404
362. Severson EA, et al. *Blood* (2018) pmid: 29678827
363. Fuster JJ, et al. *Circ. Res.* (2018) pmid: 29420212
364. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
365. Chabon JJ, et al. *Nature* (2020) pmid: 32269342
366. Razavi P, et al. *Nat. Med.* (2019) pmid: 31768066
367. Cunningham D, et al. *N. Engl. J. Med.* (2004) pmid: 15269313
368. Jonker DJ, et al. *N. Engl. J. Med.* (2007) pmid: 18003960
369. Papamichael D, et al. *Eur J Cancer* (2022) pmid: 35033994
370. Karapetis CS, et al. *Clin. Cancer Res.* (2014) pmid: 24218517
371. Chibaudel et al., 2022; ASCO Abstract 3504
372. Moiseyenko VM, et al. *Clin Drug Investig* (2018) pmid: 29470838
373. Stein A, et al. *J Immunother Cancer* (2021) pmid: 34315821
374. Price TJ, et al. *Lancet Oncol.* (2014) pmid: 24739896
375. Sakai D, et al. *Eur J Cancer* (2020) pmid: 32526634
376. Van Cutsem E, et al. *J. Clin. Oncol.* (2007) pmid: 17470858
377. Peeters M, et al. *Clin. Cancer Res.* (2015) pmid: 26341920
378. Watanabe J, et al. *Int J Cancer* (2022) pmid: 35723084
379. Kim TW, et al. *Clin Colorectal Cancer* (2018) pmid: 29703606
380. Shitara K, et al. *Cancer Sci* (2016) pmid: 27712015
381. Yoshino et al., 2022; ASCO Abstract LBA1
382. Pietrantonio F, et al. *JAMA Oncol* (2019) pmid: 31268481
383. Glassberg et al., 2020; ASPHO Abstract 2015
384. Coyne et al., 2020; ASCO Abstract 3612
385. McCowage et al., 2018; ASCO Abstract 10504
386. Mueller et al., 2020; SNO Abstract NFB-17
387. Waldner et al., 2020; DOI: 10.1055/s-0040-1715638
388. Romo et al., 2019; SNO Abstract RARE-54
389. Eng C, et al. *Lancet Oncol.* (2019) pmid: 31003911
390. Bennaoui J, et al. *Invest New Drugs* (2011) pmid: 20127139
391. Hochster HS, et al. *Cancer Chemother. Pharmacol.* (2015) pmid: 25322874
392. Do K, et al. *Invest New Drugs* (2015) pmid: 25637165
393. Tolcher AW, et al. *Clin. Cancer Res.* (2015) pmid: 25516890
394. Deming DA, et al. *Invest New Drugs* (2016) pmid: 26666244
395. van Brummelen EMJ, et al. *Oncologist* (2021) pmid: 33296125
396. Krishnamurthy A, et al. *Cancer Res.* (2018) pmid: 30042150
397. Infante JR, et al. *Lancet Oncol.* (2012) pmid: 22805291
398. Johnson DB, et al. *Clin Cancer Res* (2020) pmid: 31924734
399. Alshammari K, et al. *Clin Colorectal Cancer* (2021) pmid: 34471744
400. Corcoran RB, et al. *Cancer Discov* (2018) pmid: 29431699
401. Maio M, et al. *Eur J Cancer* (2022) pmid: 34801354
402. Johnson et al., 2020; ASCO GI Abstract 152
403. Tolcher AW, et al. *Ann. Oncol.* (2015) pmid: 25344362
404. Tolcher AW, et al. *Cancer Chemother Pharmacol* (2015) pmid: 25417902
405. Grilley-Olson JE, et al. *Invest New Drugs* (2016) pmid: 27450049
406. Bedard PL, et al. *Clin. Cancer Res.* (2015) pmid: 25500057
407. LoRusso et al., 2020; ESMO Abstract 561P
408. Sullivan et al., 2015; AACR-NCI-EORTC Abstract PR06
409. Kurzrock R, et al. *Clin Cancer Res* (2019) pmid: 31186313
410. Corcoran et al., 2019; ESMO Abstract 447PD
411. Patterson et al., 2018; AACR Abstract 3891

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531