



Sample Information

Patient Name: 曹光昇
Gender: Male
ID No.: J121326884
History No.: 46840782
Age: 54

Ordering Doctor: DOC6300K 吳宜鴻
Ordering REQ.: 0BFFJBS
Signing in Date: 2021/04/28

Path No.: S110-98698
MP No.: TM21005
Assay: Oncomine Tumor Mutation Load Assay
Sample Type: FFPE
Block No.: S110-13078A+B
Percentage of tumor cells: 40%
Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	4
Relevant Therapy Summary	5
Relevant Therapy Details	8
Clinical Trials Summary	13

Report Highlights
 3 Relevant Biomarkers
 1 Therapies Available
 33 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	Not detected	MET	Not detected
BRAF	Not detected	NTRK1	Not detected
EGFR	Not detected	NTRK3	Not detected
ERBB2	Not detected	RET	Not detected
KRAS	Not detected	ROS1	Not detected

Genomic Alteration	Finding
Tumor Mutational Burden	14.23 Mut/Mb measured

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	<i>TP53 p.(R273L) c.818G>T</i> tumor protein p53 Allele Frequency: 29.75%	None	None	12
IIC	<i>ARID1A p.(T783fs)</i> <i>c.2348_2351delCAGGinsGA</i> AT-rich interaction domain 1A Allele Frequency: 33.10%	None	None	5
	<i>Tumor Mutational Burden</i> 14.23 Mut/Mb measured	pembrolizumab ¹	pembrolizumab	19

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ARID1A	p.(T783fs)	c.2348_2351delCAGGinsGA	.	chr1:27088739	33.10%	NM_006015.5	frameshift Block Substitution	1979
TP53	p.(R273L)	c.818G>T	COSM10779	chr17:7577120	29.75%	NM_000546.5	missense	2000
PAX7	p.(=)	c.609G>A	.	chr1:19018270	40.81%	NM_002584.2	synonymous	1529
PAX7	p.(=)	c.654A>C	.	chr1:19018315	36.47%	NM_002584.2	synonymous	1999
PAX7	p.(=)	c.777G>A	.	chr1:19018438	33.37%	NM_002584.2	synonymous	1999
DDR2	p.(D81Y)	c.241G>T	.	chr1:162724469	25.18%	NM_006182.2	missense	1998
PBX1	p.(=)	c.843A>G	.	chr1:164781232	26.69%	NM_002585.3	synonymous	1004
SOX11	p.(A174E)	c.521C>A	.	chr2:5833374	36.00%	NM_003108.3	missense	1814
PAX3	p.(I96L)	c.286A>T	.	chr2:223161732	28.19%	NM_181459.3	missense	1926
XPC	p.(R617S)	c.1851G>T	.	chr3:14199532	36.47%	NM_004628.4	missense	1999
LTF	p.(R23dup)	c.68_69insAAG	.	chr3:46501284	100.00%	NM_002343.5	nonframeshift Insertion	1844
MITF	p.(M212L)	c.634A>T	.	chr3:69988300	11.01%	NM_198159.2	missense	1999
EPHA3	p.(=)	c.666C>T	.	chr3:89259522	39.90%	NM_005233.5	synonymous	2000
PDGFRA	p.(=)	c.1701A>G	.	chr4:55141055	99.95%	NM_006206.5	synonymous	1985
AFF1	p.(P798L)	c.2393C>T	.	chr4:88036378	44.40%	NM_001166693.2	missense	2000
PKHD1	p.(=)	c.3846T>G	.	chr6:51890762	31.17%	NM_138694.3	synonymous	1986
RPS6KA2	p.(=)	c.1794G>A	.	chr6:166833420	34.90%	NM_001006932.2	synonymous	1954
EGFR	p.(P20R)	c.59C>G	.	chr7:55087029	8.11%	NM_005228.4	missense	1998
EPHB6	p.(E45G)	c.134A>G	.	chr7:142561422	31.70%	NM_004445.5	missense	2000

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.03(005).

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) (continued)

DNA Sequence Variants (continued)

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
EPHB6	p.(A266V)	c.797C>T	.	chr7:142562355	13.12%	NM_004445.5	missense	983
KMT2C	p.(A4748P)	c.14242G>C	.	chr7:151841899	28.79%	NM_170606.2	missense	1997
KMT2C	p.(=)	c.5958G>T	.	chr7:151878987	18.00%	NM_170606.2	synonymous	2000
ADGRA2	p.(T1315I)	c.3944C>T	.	chr8:37699800	56.38%	NM_032777.9	missense	1999
HOOK3	p.(=)	c.1716C>G	.	chr8:42863050	44.17%	NM_032410.3	synonymous	1997
PRKDC	p.(P473T)	c.1417C>A	.	chr8:48848322	10.45%	NM_006904.6	missense	1991
CSMD3	p.(G1104V)	c.3311G>T	.	chr8:113651140	8.59%	NM_198123.1	missense	1223
PTPRD	p.(P126L)	c.377C>T	.	chr9:8528755	29.26%	NM_002839.3	missense	1999
CYP2C19	p.(E318V)	c.953A>T	.	chr10:96580386	12.27%	NM_000769.2	missense	1996
NUP98	p.(=)	c.3774A>G	.	chr11:3720547	33.27%	NM_016320.4	synonymous	1999
NUMA1	p.(=)	c.3985T>C	.	chr11:71724564	3.53%	NM_006185.3	synonymous	170
ARID2	p.(S1137*)	c.3410C>A	.	chr12:46245316	41.48%	NM_152641.3	nonsense	1996
KMT2D	p.(G794E)	c.2381G>A	.	chr12:49445085	3.70%	NM_003482.3	missense	135
KMT2D	p.(A792T)	c.2374G>A	.	chr12:49445092	4.51%	NM_003482.3	missense	133
ERBB3	p.(?)	c.-10C>T	.	chr12:56474075	16.76%	NM_001982.3	unknown	1999
HNF1A	p.(I618M)	c.1854C>G	.	chr12:121438953	72.70%	NM_000545.6	missense	2000
EP400	p.(H1765P)	c.5294A>C	.	chr12:132512746	14.96%	NM_015409.4	missense	1999
FLT1	p.(F135*)	c.404_405delTCinsA A	.	chr13:29012466	21.82%	NM_002019.4	nonsense	1517
BCL11B	p.(=)	c.885G>C	.	chr14:99642288	70.27%	NM_138576.3	synonymous	713
TCF12	p.(P279L)	c.836C>T	.	chr15:57524920	20.38%	NM_207037.1	missense	947
TSC2	p.(=)	c.2055C>T	.	chr16:2121893	3.85%	NM_000548.4	synonymous	52
MYH11	p.(=)	c.1131G>A	.	chr16:15857672	67.33%	NM_001040114.1	synonymous	1999
CDH5	p.(I517T)	c.1550_1551delTCins CT	.	chr16:66432423	68.65%	NM_001795.4	missense	1968
PER1	p.(=)	c.738G>T	.	chr17:8052895	66.83%	NM_002616.2	synonymous	1999
CDH2	p.(=)	c.2085C>A	.	chr18:25565088	14.91%	NM_001792.4	synonymous	1998
CDH20	p.(=)	c.1662G>T	.	chr18:59217224	18.03%	NM_031891.3	synonymous	1048
CDH20	p.(V569I)	c.1705G>A	.	chr18:59217267	54.60%	NM_031891.3	missense	1066
BCL2	p.(P40fs)	c.119_127delCGGGG GCCGinsGGGGCCA	.	chr18:60985773	100.00%	NM_000633.2	frameshift Block Substitution	707
BCL2	p.(?)	c.-32C>A	.	chr18:60985931	16.77%	NM_000633.2	unknown	1998
TCF3	p.(=)	c.759C>T	.	chr19:1622116	59.85%	NM_001136139.3	synonymous	1995
KEAP1	p.(G558V)	c.1673G>T	.	chr19:10599903	38.99%	NM_203500.1	missense	913

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) (continued)

DNA Sequence Variants (continued)

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
PTPRT	p.(W146L)	c.437G>T	.	chr20:41419884	36.45%	NM_133170.3	missense	2000
MYH9	p.(=)	c.2061C>T	.	chr22:36702074	53.13%	NM_002473.5	synonymous	1999
MYH9	p.(=)	c.1455C>T	.	chr22:36710289	5.00%	NM_002473.5	synonymous	100

Biomarker Descriptions

ARID1A (AT-rich interaction domain 1A)

Background: The ARID1A gene encodes the AT-rich interaction domain 1A tumor suppressor protein¹. ARID1A (also known as BAF250A) belongs to the ARID1 subfamily that also includes ARID1B. ARID1A and ARID1B are mutually exclusive subunits of the BAF variant of the SWI/SNF chromatin-remodeling complex^{2,3}. The SWI/SNF core complex is a multisubunit protein that consists of SMARCB1/IN1, SMARCC1/BAF155, SMARCC2/BAF170, SMARCA4/BRG1, and SMARCA2/BRM³. The SWI/SNF complex remodels chromatin at promoter and enhancer elements to alter and regulate gene expression^{3,4}. ARID1A binds to transcription factors and coactivator/corepressor complexes to alter transcription². Recurrent inactivating mutations in SWI/SNF complex subunits, including ARID1A, lead to transcriptional dysfunction thereby, altering its tumor suppressor function².

Alterations and prevalence: Mutations in SWI/SNF complex subunits are the most commonly mutated chromatin modulators in cancer and have been observed in 20% of all tumors⁴. The majority of ARID1A inactivating mutations are nonsense or frameshift mutations². Somatic mutations in ARID1A have been identified in 50% of ovarian clear cell carcinoma, 30% of endometrioid carcinoma, and 24-43% of uterine corpus endometrial carcinoma, bladder urothelial carcinoma, and stomach adenocarcinoma^{3,5,6}.

Potential relevance: Currently, no therapies are approved for ARID1A aberrations.

TP53 (tumor protein p53)

Background: The TP53 gene encodes the p53 tumor suppressor protein that binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis. Alterations in TP53 is required for oncogenesis as they result in loss of protein function and gain of transforming potential⁷. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers^{8,9}.

Alterations and prevalence: TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)^{5,6,10,11,12,13}. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common including substitutions at codons R158, R175, Y220, R248, R273, and R282^{5,6}. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes^{14,15,16,17}.

Potential relevance: The small molecule p53 reactivator, PC14586, received a fast track designation (2020) by the FDA for advanced tumors harboring a TP53 Y220C mutation¹⁸. The FDA has granted fast track designation (2019) to the p53 reactivator, APR-246 alone,¹⁹ and breakthrough designation²⁰ (2020) in combination with azacitidine for myelodysplastic syndrome (MDS) and acute myeloid leukemia patients (AML) harboring a TP53 mutation. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation^{21,22}. TP53 mutations confer poor prognosis in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), chronic lymphocytic leukemia (CLL),^{23,24,25,26}. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant²⁷. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system²⁸.

Tumor Mutational Burden

Background: Tumor mutational burden (TMB), also known as tumor mutational load (TML), is the count of somatic mutations in the DNA of cancer cells. TMB is determined by next-generation sequencing and is expressed as the number of mutations per megabase (mut/Mb) of DNA coding sequence²⁹. Errors in DNA repair, including mutations in the POLE gene and in mismatch repair (MMR) genes, are associated with increased TMB^{30,31,32,33,34}. High TMB is associated with increased neo-antigen burden and has been linked to

Biomarker Descriptions (continued)

response to immune checkpoint inhibitors (ICIs) that target the cytotoxic T lymphocyte antigen-4 (CTLA4), programmed death protein 1 (PD1), and programmed death-ligand 1 (PD-L1) inhibitors^{35,36,37,38}.

Alterations and prevalence: In one study of over 100,000 tumor samples, the median TMB value was 3.6 mut/Mb although TMB values vary widely across cancers³⁹. Certain childhood cancers, leukemia, glioblastoma, and neuroblastoma typically have low mutation burden and median TMB values <1 mut/Mb^{36,39}. In comparison, cancers that experience genotoxic insults including skin cancer and lung cancer have higher median TMB values of approximately 10 mut/Mb^{36,39}. For example, within non-small cell lung cancer (NSCLC), higher TMB was observed in former/current smokers (10.5 mut/Mb) relative to never smokers (0.6 mut/Mb)^{36,39,40}. There is no consensus around the definition of high and low TMB that could be applied universally to all tumor types, instead multiple sources suggest that TMB status is a cancer type specific attribute^{39,41,42}. In NSCLC, several studies have suggested establishing a threshold between low and high TMB of 10 +/- 1 mut/Mb^{43,44,45,46}.

Potential relevance: ICIs stimulate a patient's own T-cells to kill tumors and have exhibited benefits in some patients. The first ICI to be approved by the FDA was ipilimumab (2011), an anti-CTLA4 antibody indicated for the treatment of metastatic melanoma. In 2014, anti-PD-1 antibodies, nivolumab (2014) and pembrolizumab (2014), were subsequently approved for the treatment of metastatic melanoma. Pembrolizumab was also approved (2014) for advanced esophageal squamous cell carcinoma. In 2020, the indication for pembrolizumab⁴⁷ was expanded to include TMB-H (>= 10 mut/Mb) solid tumors that have progressed on prior therapy. Indications have been expanded for these ICIs to include several other cancer types including NSCLC, advanced renal cell carcinoma, classical Hodgkin lymphoma, recurrent or metastatic squamous cell carcinoma of the head and neck, urothelial carcinoma, microsatellite instability (MSI)-High or mismatch repair deficient (dMMR) colorectal cancer, and hepatocellular carcinoma. Atezolizumab (2016), avelumab (2017), and durvalumab (2017), that target programmed death-ligand 1 (PD-L1), were subsequently approved by the FDA. However, the predictive biomarkers that underlie the clinical benefits of these approved immunotherapies, including TMB, are under active investigation. Several published studies including the CheckMate 586 and CheckMate 817 clinical trials have concluded that high TMB was associated with improved response to FDA approved checkpoint inhibitors^{44,48,49}. In contrast, several promising previous trials failed to show an improvement in survival outcomes between high and low TMB including CheckMate 227 (ipilimumab + nivolumab vs. chemotherapy), CheckMate 026 (nivolumab vs. chemotherapy), KEYNOTE 189 (pembrolizumab vs. chemotherapy), KEYNOTE 021 (pembrolizumab vs. pembrolizumab + chemotherapy), and Lung-MAP (nivolumab + ipilimumab vs. nivolumab). In response, suggestions to combine TMB score with PD-L1 expression as a way to increase the predictive power for patient stratification have been reported⁵⁰. Nivolumab alone or in combination with ipilimumab is recommended for use in NSCLC with evidence of high TMB⁵¹. Pembrolizumab is indicated for use in various cancer types with evidence of metastasis including Ewing sarcoma, salivary gland neoplasms, cervical cancer, uterine sarcoma, endometrial carcinoma, thyroid cancer, and germ cell tumors with high TMB^{52,53,54,55,56,57}. TMB score estimation is affected by the utilized assays, therefore efforts are underway to develop a standardized approach for score calculation with the aim to support consistent reporting of TMB values across laboratories^{58,59,60,61}.

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ① In this cancer type and other cancer types ✕ No evidence

TP53 p.(R273L) c.818G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
atezolizumab	✕	✕	✕	✕	● (II)
durvalumab + olaparib	✕	✕	✕	✕	● (II)
niraparib	✕	✕	✕	✕	● (II)
olaparib	✕	✕	✕	✕	● (II)
talazoparib	✕	✕	✕	✕	● (II)
eprenetapopt, pembrolizumab	✕	✕	✕	✕	● (I/II)
sintilimab	✕	✕	✕	✕	● (I/II)
BAY-1895344, niraparib	✕	✕	✕	✕	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ✕ No evidence

TP53 p.(R273L) c.818G>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
HWH-340	✕	✕	✕	✕	● (I)
talazoparib, palbociclib, axitinib, crizotinib	✕	✕	✕	✕	● (I)

ARID1A p.(T783fs) c.2348_2351delCAGGinsGA

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
niraparib	✕	✕	✕	✕	● (II)
talazoparib	✕	✕	✕	✕	● (II)
PLX-2853	✕	✕	✕	✕	● (I/II)
copanlisib, olaparib, durvalumab	✕	✕	✕	✕	● (I)
VX-803	✕	✕	✕	✕	● (I)

Tumor Mutational Burden

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
pembrolizumab	●	○	✕	✕	● (II)
pembrolizumab, chemotherapy	✕	✕	✕	✕	● (III)
atezolizumab	✕	✕	✕	✕	● (II)
atezolizumab + chemotherapy	✕	✕	✕	✕	● (II)
atezolizumab, bevacizumab	✕	✕	✕	✕	● (II)
atezolizumab, nivolumab, ipilimumab	✕	✕	✕	✕	● (II)
ipilimumab + nivolumab	✕	✕	✕	✕	● (II)
ipilimumab, nivolumab	✕	✕	✕	✕	● (II)
pembrolizumab, ipilimumab + nivolumab	✕	✕	✕	✕	● (II)
pembrolizumab, quavonlimab, MK-4280, lenvatinib	✕	✕	✕	✕	● (II)
sintilimab	✕	✕	✕	✕	● (II)
sintilimab, bevacizumab	✕	✕	✕	✕	● (II)
entinostat, nivolumab	✕	✕	✕	✕	● (I/II)
anti-PD-1	✕	✕	✕	✕	● (I)
BAY1905254	✕	✕	✕	✕	● (I)
KN046	✕	✕	✕	✕	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types ☒ No evidence

Tumor Mutational Burden (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
pembrolizumab, targinine	×	×	×	×	<input checked="" type="radio"/> (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2021-02-17. For the most up-to-date information, search www.fda.gov.

Tumor Mutational Burden

● pembrolizumab

Cancer type: Solid Tumor

Label as of: 2020-11-13

Variant class: Tumor Mutational Burden

Indications and usage:

KEYTRUDA® is a programmed death receptor-1 (PD-1)-blocking antibody indicated:

Melanoma

- for the treatment of patients with unresectable or metastatic melanoma.
- for the adjuvant treatment of patients with melanoma with involvement of lymph node(s) following complete resection.

Non-Small Cell Lung Cancer (NSCLC)

- in combination with pemetrexed and platinum chemotherapy, as first-line treatment of patients with metastatic nonsquamous NSCLC, with no EGFR or ALK genomic tumor aberrations.
- in combination with carboplatin and either paclitaxel or paclitaxel protein-bound, as first-line treatment of patients with metastatic squamous NSCLC.
- as a single agent for the first-line treatment of patients with NSCLC expressing PD-L1 [Tumor Proportion Score (TPS) \geq 1%] as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations, and is:
 - stage III where patients are not candidates for surgical resection or definitive chemoradiation, or
 - metastatic.
- as a single agent for the treatment of patients with metastatic NSCLC whose tumors express PD-L1 (TPS \geq 1%) as determined by an FDA-approved test, with disease progression on or after platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving KEYTRUDA®.

Small Cell Lung Cancer (SCLC)

- for the treatment of patients with metastatic SCLC with disease progression on or after platinum-based chemotherapy and at least one other prior line of therapy.¹

Head and Neck Squamous Cell Cancer (HNSCC)

- in combination with platinum and FU for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC.
- as a single agent for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC whose tumors express PD-L1 [Combined Positive Score (CPS) \geq 1] as determined by an FDA-approved test.
- as a single agent for the treatment of patients with recurrent or metastatic HNSCC with disease progression on or after platinum-containing chemotherapy.

Classical Hodgkin Lymphoma (cHL)

- for the treatment of adult patients with relapsed or refractory cHL.
- for the treatment of pediatric patients with refractory cHL, or cHL that has relapsed after 2 or more lines of therapy.

Primary Mediastinal Large B-Cell Lymphoma (PMBCL)

- for the treatment of adult and pediatric patients with refractory PMBCL, or who have relapsed after 2 or more prior lines of therapy.
- Limitations of Use: KEYTRUDA® is not recommended for treatment of patients with PMBCL who require urgent cytoreductive therapy.

Urothelial Carcinoma

Tumor Mutational Burden (continued)

- for the treatment of patients with locally advanced or metastatic urothelial carcinoma who are not eligible for cisplatin-containing chemotherapy and whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 10] as determined by an FDA-approved test, or in patients who are not eligible for any platinum-containing chemotherapy regardless of PD-L1 status.¹
- for the treatment of patients with locally advanced or metastatic urothelial carcinoma who have disease progression during or following platinum-containing chemotherapy or within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy.
- for the treatment of patients with Bacillus Calmette-Guerin (BCG)-unresponsive, high-risk, non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ (CIS) with or without papillary tumors who are ineligible for or have elected not to undergo cystectomy.

Microsatellite Instability-High or Mismatch Repair Deficient Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR)
 - solid tumors that have progressed following prior treatment and who have no satisfactory alternative treatment options,¹ or
 - colorectal cancer that has progressed following treatment with a fluoropyrimidine, oxaliplatin, and irinotecan.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with MSI-H central nervous system cancers have not been established.

Microsatellite Instability-High or Mismatch Repair Deficient Colorectal Cancer (CRC)

- for the first-line treatment of patients with unresectable or metastatic MSI-H or dMMR colorectal cancer (CRC).

Gastric Cancer

- for the treatment of patients with recurrent locally advanced or metastatic gastric or gastroesophageal junction adenocarcinoma whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by an FDA-approved test, with disease progression on or after 2 or more prior lines of therapy including fluoropyrimidine- and platinum-containing chemotherapy and if appropriate, HER2/neu-targeted therapy.¹

Esophageal Cancer

- for the treatment of patients with recurrent locally advanced or metastatic squamous cell carcinoma of the esophagus whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 10] as determined by an FDA-approved test, with disease progression after one or more prior lines of systemic therapy.

Cervical Cancer

- for the treatment of patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by an FDA-approved test.¹

Hepatocellular Carcinoma (HCC)

- for the treatment of patients with HCC who have been previously treated with sorafenib.¹

Merkel Cell Carcinoma (MCC)

- for the treatment of adult and pediatric patients with recurrent locally advanced or metastatic Merkel cell carcinoma.¹

Renal Cell Carcinoma (RCC)

- in combination with axitinib, for the first-line treatment of patients with advanced RCC.

Endometrial Carcinoma

- in combination with lenvatinib, for the treatment of patients with advanced endometrial carcinoma that is not MSI-H or dMMR, who have disease progression following prior systemic therapy and are not candidates for curative surgery or radiation.¹

Tumor Mutational Burden-High (TMB-H) Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] solid tumors, as determined by an FDA-approved test, that have progressed following prior treatment and who have no satisfactory alternative treatment options.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with TMB-H central nervous system cancers have not been established.

Cutaneous Squamous Cell Carcinoma (cSCC)

- for the treatment of patients with recurrent or metastatic cutaneous squamous cell carcinoma that is not curable by surgery or radiation.

Triple-Negative Breast Cancer (TNBC)

Tumor Mutational Burden (continued)

- in combination with chemotherapy, for the treatment of patients with locally recurrent unresectable or metastatic TNBC whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 10] as determined by an FDA approved test.²

Adult Indications: Additional Dosing Regimen of 400 mg Every 6 Weeks

- for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications.³

¹This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

²This indication is approved under accelerated approval based on progression-free survival. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

³This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125514s088lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2021-02-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

Tumor Mutational Burden

☐ pembrolizumab

Cancer type: Osteosarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2021]

☐ pembrolizumab

Cancer type: Breast Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IV; Recurrent, Invasive (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 1.2021]

☐ pembrolizumab

Cancer type: Cervical Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Metastatic, Progression (Second-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2021]

☐ pembrolizumab

Cancer type: Head and Neck Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Nasopharyngeal; Recurrent, Unresectable, Metastatic (Subsequent therapy); Useful in certain circumstances
- Salivary Gland Neoplasm; Recurrent, Unresectable, Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 1.2021]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Testicular Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Germ Cell Tumor; Metastatic (Third-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Testicular Cancer [Version 1.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Follicular Carcinoma, Thyroid Gland Hurthle Cell Carcinoma, Thyroid Gland Papillary Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Locally Recurrent, Advanced, Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2020]

○ pembrolizumab

Cancer type: Thyroid Gland Medullary Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Locally Recurrent, Unresectable, Symptomatic, Progression, Distant Metastases, Asymptomatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2020]

○ pembrolizumab

Cancer type: Thyroid Gland Anaplastic Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IVC; Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2020]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Endometrial Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Metastatic (Line of therapy not specified); Useful in certain circumstances
- Uterine Sarcoma; Recurrent, Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2021]

○ pembrolizumab

Cancer type: Ewing Sarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2021]

Clinical Trials Summary

TP53 p.(R273L) c.818G>T

NCT ID	Title	Phase
NCT04383938	Study of APR-246 in Combination With Pembrolizumab in Subjects With Solid Tumor Malignancies	I/II
No NCT ID	Single-arm Exploratory Study for Sintilimab Second-line or Later-line Treatment of Advanced Non-small Cell Lung Cancer with DDR Pathway Gene Mutation	I/II
NCT03334617	An Open-Label, Multi-Drug, Biomarker-Directed, Multi-Centre Phase II Umbrella Study in Patients with Non-Small Cell Lung Cancer, who Progressed on an anti-PD-1/PD-L1 Containing Therapy (HUDSON)	II
NCT03767075	Basket of Baskets: A Modular, Open-label, Phase II, Multicentre Study To Evaluate Targeted Agents in Molecularly Selected Populations With Advanced Solid Tumours	II
NCT03207347	A Phase II Trial of the PARP Inhibitor, Niraparib, in BAP1 and Other DNA Damage Response (DDR) Pathway Deficient Neoplasms (UF-STO-ETI-001).	II
NCT02029001	A Two-period, Multicenter, Randomized, Open-label, Phase II Study Evaluating the Clinical Benefit of a Maintenance Treatment Targeting Tumor Molecular Alterations in Patients With Progressive Locally-advanced or Metastatic Solid Tumors MOST: My own specific treatment	II
NCT03297606	Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR): A Phase II Basket Trial	II
NCT02401347	A Phase II Clinical Trial of the PARP Inhibitor Talazoparib in BRCA1 and BRCA2 Wild Type Patients With Advanced Triple Negative Breast Cancer and Homologous Recombination Deficiency or Advanced HER2 Negative Breast Cancer or Other Solid Tumors With a Mutation in Homologous Recombination Pathway Genes Talazoparib Beyond BRCA (TBB) Trial	II
NCT03415659	A Phase I, Open-label, Single-center, Single/Multiple-dose, Dose-escalation/Dose-expansion Clinical Study on Tolerance and Pharmacokinetics of HWH340 Tablet in Patients With Advanced Solid Tumors	I

Clinical Trials Summary (continued)

TP53 p.(R273L) c.818G>T (continued)

NCT ID	Title	Phase
NCT03233204	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice)- Phase 2 Subprotocol of Olaparib in Patients With Tumors Harboring Defects in DNA Damage Repair Genes	II
NCT04267939	An Open-label Phase Ib Study to Determine the Maximum Tolerated and/or Recommended Phase II Dose of the ATR Inhibitor BAY 1895344 in Combination With PARP Inhibitor Niraparib, in Patients With Recurrent Advanced Solid Tumors and Ovarian Cancer	I
NCT04693468	Modular Phase IB Hypothesis-Testing, Biomarker-Driven, Talazoparib Combination Trial (TalaCom)	I

ARID1A p.(T783fs) c.2348_2351delCAGGinsGA

NCT ID	Title	Phase
NCT03207347	A Phase II Trial of the PARP Inhibitor, Niraparib, in BAP1 and Other DNA Damage Response (DDR) Pathway Deficient Neoplasms (UF-STO-ETI-001).	II
NCT02286687	Phase II Study of the PARP Inhibitor Talazoparib in Advanced Cancer Patients With Somatic Alterations in BRCA1/2, Mutations/Deletions in Other BRCA Pathway Genes and Germline Mutation in BRCA1/2 (Not Breast or Ovarian Cancer)	II
NCT03297424	A Phase Ib/IIa Dose-escalation Study to Assess Safety, Pharmacokinetics, Pharmacodynamics, and Preliminary Efficacy of PLX2853 in Subjects With Advanced Malignancies	I/II
NCT03842228	A Phase Ib Biomarker-Driven Combination Trial of Copanlisib, Olaparib, and MEDI4736 (Durvalumab) in Patients With Advanced Solid Tumors	I
NCT02278250	An Open-Label Study of the Safety, Tolerability, and Pharmacokinetic/Pharmacodynamic Profile of M4344 (Formerly VX-803) as a Single Agent and in Combination With Cytotoxic Chemotherapy in Participants With Advanced Solid Tumors	I

Tumor Mutational Burden

NCT ID	Title	Phase
No NCT ID	Pembrolizumab Alone Versus Pembrolizumab-Chemotherapy in First line NSCLC	III
NCT03836066	A Phase II Open Label Study of Atezolizumab in Combination With Bevacizumab as First Line Treatment for Locally Advanced or Metastatic High-intermediate Tumor Mutation Burden Selected Non-squamous Non-small Cell Lung Cancer (NSCLC) Patients.	II
NCT04591431	The Rome Trial From Histology to Target: the Road to Personalize Target Therapy and Immunotherapy	II
NCT03516981	A Phase II Precision Oncology Study of Biomarker-Directed, Pembrolizumab-(MK-3475, SCH 900475) Based Combination Therapy for Advanced Non-Small Cell Lung Cancer (KEYNOTE-495; KeyImPaCT)	II
NCT03574402	An Open-label, Multi-center, Phase II Umbrella Study to Assess Efficacy of Targeted Therapy or Immunotherapy Directed by Next Generation Sequencing (NGS) in Chinese Patients With Advanced NSCLC (TRUMP)	II
NCT04213170	Prospective Phase II Clinical Study of Sintilimab Combined With Bevacizumab for Driving Gene-negative, Asymptomatic Brain Metastases From Non-small Cell Lung Cancer	II
No NCT ID	A Multicenter, Open and Prospective Study on the Effect of Immunotherapy on T Cell Surface Receptors and Cytokines in non-small-cell lung carcinoma	I

Clinical Trials Summary (continued)

Tumor Mutational Burden (continued)

NCT ID	Title	Phase
NCT03733951	An Open-Label, Multicenter, Dose-Escalation and Expansion Phase Ia/Ib Study to Evaluate Safety, Tolerability, Pharmacokinetics, and Antitumor Activity of KN046 Monotherapy in Subjects With Advanced Solid Tumors and Lymphoma	I
NCT03767075	Basket of Baskets: A Modular, Open-label, Phase II, Multicentre Study To Evaluate Targeted Agents in Molecularly Selected Populations With Advanced Solid Tumours	II
NCT04185831	MEGALiT - a Multicenter, Basket and Umbrella Explorative Trial on the Efficacy and Safety of Molecular Profile Selected Commercially Available Targeted Anti-cancer Drugs in Patients With Advanced Cancers Progressive on Standard Therapy	II
NCT04589845	Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial	II
NCT04632992	MyTACTIC: An Open-Label Phase II Study Evaluating Targeted Therapies in Patients Who Have Advanced Solid Tumors With Genomic Alterations or Protein Expression Patterns Predictive of Response	II
NCT03297606	Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR): A Phase II Basket Trial	II
NCT03668119	A Randomized, Open-Label, Phase II Study of Nivolumab in Combination With Ipilimumab or Nivolumab Monotherapy in Participants With Advanced or Metastatic Solid Tumors of High Tumor Mutational Burden (TMB-H)	II
NCT02628067	A Clinical Trial of Pembrolizumab (MK-3475) Evaluating Predictive Biomarkers in Subjects With Advanced Solid Tumors (KEYNOTE 158)	II
NCT02693535	Targeted Agent and Profiling Utilization Registry (TAPUR) Study	II
NCT03838042	INFORM2 Exploratory Multinational Phase I/II Combination Study of Nivolumab and Entinostat in Children and Adolescents with Refractory High-risk Malignancies	I/II
NCT03666273	An Open-label, Phase I, First-in-human, Dose Escalation and Expansion Study to Evaluate the Safety, Tolerability, Maximum Tolerated or Administered Dose, Pharmacokinetics, Pharmacodynamics and Tumor Response Profile of the ILDR2 Function-blocking Antibody BAY1905254 in Patients With Advanced Solid Tumors	I
NCT03236935	Phase Ib Trial of L-NMMA in Combination With Pembrolizumab in Patients With Melanoma, Non-Small Cell Lung Cancer, Head and Neck Squamous Cell Carcinoma, Classical Hodgkin Lymphoma, Urothelial Carcinoma, Cervical Cancer, Esophageal Cancer, Gastric Cancer, Hepatocellular Carcinoma, Merkel Cell Carcinoma, Primary Mediastinal Large B-cell Lymphoma, Renal Cell Carcinoma, Small Cell Lung Cancer, Microsatellite Instability-High/Mismatch Repair Deficient Cancer, or for the Treatment of Adult Patients With Unresectable or Metastatic Tumor Mutational Burden-High Solid Tumors	I

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Wu et al. ARID1A mutations in cancer: another epigenetic tumor suppressor?. *Cancer Discov.* 2013 Jan;3(1):35-43. PMID: 23208470
3. Wilson et al. SWI/SNF nucleosome remodellers and cancer. *Nat. Rev. Cancer.* 2011 Jun 9;11(7):481-92. PMID: 21654818
4. Alver et al. The SWI/SNF Chromatin Remodelling Complex Is Required for Maintenance of Lineage Specific Enhancers. *Nat Commun.* 8;14648. PMID: 28262751
5. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
6. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
7. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell.* 2014 Mar 17;25(3):304-17. PMID: 24651012
8. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. PMID: 20182602
9. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med.* 2017 Apr 3;7(4). PMID: 28270529
10. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature.* 2012 Sep 27;489(7417):519-25. PMID: 22960745
11. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
12. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet.* 2016 Jun;48(6):607-16. PMID: 27158780
13. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
14. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat.* 2002 Jun;19(6):607-14. PMID: 12007217
15. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011 Apr;2(4):466-74. PMID: 21779514
16. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene.* 2007 Apr 2;26(15):2157-65. PMID: 17401424
17. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat.* 2014 Jun;35(6):766-78. PMID: 24729566
18. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
19. <https://www.aprea.com/news/aprea-therapeutics-receives-fda-fast-track-designation-and-orphan-drug-designation-for-apr-246-for-the-treatment-of-myelodysplastic-syndromes-mds/>. <https://www.aprea.com/news/aprea-therapeutics-receives-fda-fast-track-designation-and-orphan-drug-designation-for-apr-246-for-the-treatment-of-myelodysplastic-syndromes-mds/>
20. <https://ir.aprea.com//news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>
21. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol.* 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
22. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci.* 2017 Nov;74(22):4171-4187. PMID: 28643165
23. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]
24. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2021]
25. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 1.2020]
26. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 2.2021]
27. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 1.2021]
28. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med.* 2020 Aug 3. PMID: 32747829
29. Yuan et al. Novel technologies and emerging biomarkers for personalized cancer immunotherapy. 4:3. PMID: 26788324

References (continued)

30. Temko et al. Somatic POLE exonuclease domain mutations are early events in sporadic endometrial and colorectal carcinogenesis, determining driver mutational landscape, clonal neoantigen burden and immune response. *J. Pathol.* 2018 Jul;245(3):283-296. PMID: 29604063
31. Mehnert et al. Immune activation and response to pembrolizumab in POLE-mutant endometrial cancer. *126(6):2334-40.* PMID: 27159395
32. Bouffet et al. Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. *J. Clin. Oncol.* 2016 Jul 1;34(19):2206-11. PMID: 27001570
33. Humphris et al. Hypermutation In Pancreatic Cancer. *Gastroenterology.* 2017 Jan;152(1):68-74.e2. PMID: 27856273
34. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
35. Snyder et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* 2014 Dec 4;371(23):2189-2199. PMID: 25409260
36. Alexandrov et al. Signatures of mutational processes in human cancer. *Nature.* 2013 Aug 22;500(7463):415-21. PMID: 23945592
37. Rizvi et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science.* 2015 Apr 3;348(6230):124-8. PMID: 25765070
38. Van et al. Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science.* 2015 Oct 9;350(6257):207-211. PMID: 26359337
39. Chalmers et al. Analysis of 100,000 human cancer genomes reveals the landscape of tumor mutational burden. *9:34.* PMID: 28420421
40. Govindan et al. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell.* 2012 Sep 14;150(6):1121-34. PMID:22980976
41. Endris et al. Measurement of tumor mutational burden (TMB) in routine molecular diagnostics: in silico and real-life analysis of three larger gene panels. *Int. J. Cancer.* 2019 May 1;144(9):2303-2312. PMID: 30446996
42. Meléndez et al. Methods of measurement for tumor mutational burden in tumor tissue. *Transl Lung Cancer Res.* 2018 Dec;7(6):661-667. PMID: 30505710
43. Hellmann et al. Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor Mutational Burden. *N. Engl. J. Med.* 2018 May 31;378(22):2093-2104. PMID: 29658845
44. Ready et al. First-Line Nivolumab Plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer (CheckMate 568): Outcomes by Programmed Death Ligand 1 and Tumor Mutational Burden as Biomarkers. *J. Clin. Oncol.* 2019 Apr 20;37(12):992-1000. PMID: 30785829
45. Alborelli et al. Tumor mutational burden assessed by targeted NGS predicts clinical benefit from immune checkpoint inhibitors in non-small cell lung cancer. *J. Pathol.* 2020 Jan;250(1):19-29. PMID: 31471895
46. Heeke et al. In-house Implementation of Tumor Mutational Burden Testing to Predict Durable Clinical Benefit in Non-small Cell Lung Cancer and Melanoma Patients. *Cancers (Basel).* 2019 Aug 29;11(9). PMID: 31470674
47. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125514s088lbl.pdf
48. F. et al. CheckMate 817: First-Line Nivolumab + Ipilimumab in Patients with ECOG PS 2 and Other Special Populations with Advanced NSCLC. Volume 14, Issue 10, Supplement, Pages S214–S215. Abstract OA04.02
49. Zhu et al. Association Between Tumor Mutation Burden (TMB) and Outcomes of Cancer Patients Treated With PD-1/PD-L1 Inhibitions: A Meta-Analysis. *Front Oncol*, 9:1161, 04 Nov 2019. PMID: 31258479
50. Castellanos et al. Tumor mutational burden (TMB) and PD-L1 expression as predictors of response to immunotherapy (IO) in NSCLC. *Journal of Clinical Oncology* 37, no. 15_suppl (May 20, 2019) 2630-2630. Abstract 2630
51. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 2.2021]
52. NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2021]
53. NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 1.2021]
54. NCCN Guidelines® - NCCN-Testicular Cancer [Version 1.2021]
55. NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2021]
56. NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2021]
57. NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2020]
58. <https://www.focr.org/tmb>
59. <http://www.iqnpath.org/category/tmb>

References (continued)

60. Stenzinger et al. Tumor mutational burden standardization initiatives: Recommendations for consistent tumor mutational burden assessment in clinical samples to guide immunotherapy treatment decisions. *Genes Chromosomes Cancer*. 2019 Aug;58(8):578-588. PMID: 30664300
61. Merino et al. Establishing guidelines to harmonize tumor mutational burden (TMB): in silico assessment of variation in TMB quantification across diagnostic platforms: phase I of the Friends of Cancer Research TMB Harmonization Project. *J Immunother Cancer*. 2020 Mar;8(1). PMID: 32217756