

**ABOUT THE TEST** FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

## PATIENT

**DISEASE** Brain glioblastoma (GBM)  
**NAME** Wen, Sheng-Ting  
**DATE OF BIRTH** 13 December 1956  
**SEX** Male  
**MEDICAL RECORD #** 47793715

## PHYSICIAN

**ORDERING PHYSICIAN** Yeh, Yi-Chen  
**MEDICAL FACILITY** Taipei Veterans General Hospital  
**ADDITIONAL RECIPIENT** None  
**MEDICAL FACILITY ID** 205872  
**PATHOLOGIST** Not Provided

## SPECIMEN

**SPECIMEN SITE** Brain  
**SPECIMEN ID** S110-32117H (PF21049)  
**SPECIMEN TYPE** Slide Deck  
**DATE OF COLLECTION** 26 October 2021  
**SPECIMEN RECEIVED** 24 November 2021

## Biomarker Findings

**Microsatellite status** - MS-Stable  
**Tumor Mutational Burden** - 3 Muts/Mb

## Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

**CDK4** amplification  
**MDM2** amplification  
**RB1** inversion exons 18-24  
**TERT** promoter -146C>T

**3 Disease relevant genes with no reportable alterations: EGFR, IDH1, PDGFRA**

0 Therapies with Clinical Benefit  
 0 Therapies with Resistance

15 Clinical Trials

## BIOMARKER FINDINGS

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 3 Muts/Mb

## GENOMIC FINDINGS

**CDK4** - amplification

10 Trials see p. 5

**MDM2** - amplification

6 Trials see p. 7

## THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

### THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

none

### THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

none

none

none

## GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

**RB1** - inversion exons 18-24 ..... p. 4    **TERT** - promoter -146C>T ..... p. 4

**NOTE** Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

ORDERED TEST # ORD-1243577-01

## BIOMARKER FINDINGS

## BIOMARKER

## Microsatellite status

## RESULT

MS-Stable

### POTENTIAL TREATMENT STRATEGIES

#### — Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors<sup>1-3</sup>, including approved therapies nivolumab and pembrolizumab<sup>4</sup>. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%,  $p=0.001$ )<sup>5</sup>.

### FREQUENCY & PROGNOSIS

Low-level MSI has been reported in 5-9% of glioblastoma (GBM) samples<sup>6-8</sup>. A large-scale study did not find high-level microsatellite instability (MSI-H) in any of 129 GBM samples<sup>6</sup>, although a small-scale study reported MSI-H in 4 of 15 pediatric GBMs and 1 of 12 adult GBMs<sup>9</sup>. The frequency of MSI has been reported to be increased in relapsed compared to primary GBM<sup>6</sup>, in GBMs with a previous lower grade astrocytoma<sup>7</sup>, and in giant cell GBM compared to classic GBM<sup>8</sup>.

### FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor<sup>10</sup>. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2<sup>10-12</sup>. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers<sup>13-15</sup>. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins<sup>10,12,14-15</sup>.

## BIOMARKER

## Tumor Mutational Burden

## RESULT

3 Muts/Mb

### POTENTIAL TREATMENT STRATEGIES

#### — Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1<sup>16-18</sup>, anti-PD-1 therapies<sup>16-19</sup>, and combination nivolumab and ipilimumab<sup>20-25</sup>. In glioma, a lack of association between TMB and clinical benefit from immune checkpoint inhibitors has been reported<sup>16,26-27</sup>. However, multiple case studies have reported that patients with ultramutated gliomas driven by POLE

mutations have benefited from treatment with anti-PD-1<sup>28-29</sup> or anti-PD-L1<sup>30</sup> therapies. Therefore, although increased TMB alone may not be a strong biomarker for PD-1 or PD-L1 inhibitors in this cancer type, these agents may have efficacy for patients with glioma harboring both high TMB and POLE mutation.

### FREQUENCY & PROGNOSIS

Glioblastoma (GBM) harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 4.2% of cases have high TMB (>20 muts/Mb)<sup>31</sup>. For pediatric patients, high TMB has been reported in a subset of high-grade gliomas, frequently in association with mutations in mismatch repair or proofreading genes and in TP53, whereas BRAF alterations or other oncogene fusions were observed more frequently in brain tumors harboring low TMB<sup>32-33</sup>. Increased TMB has been reported to correlate with higher tumor grade in glioma<sup>34</sup> and glioblastoma (GBM) tissue samples with biallelic mismatch repair deficiency

(bMMRD)<sup>28</sup>, as well as with shorter OS of patients with diffuse glioma<sup>35</sup>.

### FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma<sup>36-37</sup> and cigarette smoke in lung cancer<sup>38-39</sup>, treatment with temozolomide-based chemotherapy in glioma<sup>40-41</sup>, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes<sup>42-46</sup>, and microsatellite instability (MSI)<sup>42,45-46</sup>. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents<sup>16,26-30</sup>.

ORDERED TEST # ORD-1243577-01

## GENOMIC FINDINGS

## GENE

## CDK4

## ALTERATION

amplification

palbociclib<sup>47,52</sup>, and ribociclib<sup>53</sup>.

## — Potential Resistance —

Rb inactivation may predict resistance to CDK4/6 inhibitors such as palbociclib, abemaciclib, and ribociclib, which act upstream of Rb<sup>49,54-62</sup>.

has been reported to be associated with a poorer patient outcome in anaplastic astrocytoma and glioblastoma<sup>69-72</sup>.

## FINDING SUMMARY

CDK4 encodes the cyclin-dependent kinase 4, which regulates the cell cycle, senescence, and apoptosis<sup>73</sup>. CDK4 and its functional homolog CDK6 are activated by D-type cyclins and promote cell cycle progression by inactivating the tumor suppressor Rb<sup>74-75</sup>. Amplification of the chromosomal region that includes CDK4 has been reported in multiple cancer types, including lung cancer, glioblastoma, and liposarcoma, and has been associated with overexpression of CDK4 protein<sup>47,76-82</sup>.

## POTENTIAL TREATMENT STRATEGIES

## — Targeted Therapies —

CDK4 amplification or activation may predict sensitivity to CDK4/6 inhibitors such as abemaciclib, palbociclib, and ribociclib<sup>47-50</sup>. Clinical benefit has been reported for limited tumor types including patients with CDK4-amplified liposarcoma and sarcoma in response to treatment with abemaciclib<sup>51</sup>,

## FREQUENCY &amp; PROGNOSIS

Across TCGA and MKSCC studies, CDK4 amplification has been reported in 4.0-9.4% of glioma cases and 14% of glioblastoma multiforme cases (cBioPortal, Sep 2021)<sup>63-67</sup>. A study has reported amplification of the 12q14-15 region, where CDK4 and MDM2 reside, in 5% (2/42) of glioblastomas<sup>68</sup>. Amplification of CDK4 and corresponding increased CDK4 protein expression

## GENE

## MDM2

## ALTERATION

amplification

types of solid tumors that had failed immunotherapy, including 1 out of 14 patients with non-small cell lung cancer; 1 out of 5 patients with urothelial carcinoma; and 2 out of 5, 1 out of 5, and 1 out of 11 patients with mucosal, uveal, and cutaneous melanoma, respectively<sup>88</sup>. Phase 1b studies of the MDM2 inhibitor idasanutlin for refractory AML in combination with cytarabine or venetoclax reported anti-leukemic response rates of 33% (25/75) and 37% (11/30), respectively<sup>89-90</sup>; clinical benefit (58% ORR, 7/12) with idasanutlin monotherapy has been reported for patients with polycythemia vera<sup>91</sup>. The dual MDM2/MDM4 inhibitor ALRN-6924 led to an ORR of 27% (4/15) for patients with TP53 wildtype peripheral T-cell lymphoma in a Phase 2 study<sup>92</sup>; responses have also been observed in TP53 wildtype AML, MDS, Merkel cell carcinoma, colorectal cancer, and liposarcoma<sup>93-94</sup>.

MDM2 has been associated with poor survival in patients with glioblastoma<sup>68,95</sup>.

## FINDING SUMMARY

MDM2 encodes an E3 ubiquitin protein ligase, which mediates the ubiquitination and subsequent degradation of p53, Rb1, and other proteins<sup>96-98</sup>. MDM2 acts to prevent the activity of the tumor suppressor p53; therefore, overexpression or amplification of MDM2 may be oncogenic<sup>99-100</sup>. Overexpression or amplification of MDM2 is frequent in cancer<sup>101</sup>. Although two retrospective clinical studies suggest that MDM2 amplification may predict a short time-to-treatment failure on anti-PD-1/PD-L1 immune checkpoint inhibitors, with 4/5 patients with MDM2 amplification<sup>102</sup> and 2/3 patients with MDM2 or MDM4 amplification<sup>103</sup> experiencing tumor hyperprogression, amplification of MDM2 or MDM4 was not associated with shorter progression-free survival (PFS) in a retrospective analysis of non-small cell lung cancer (NSCLC) outcomes with immune checkpoint inhibitors (hazard ratio of 1.4, p=0.44)<sup>104</sup>. The latter study reported PFS of >2 months for 5/8 patients with MDM2/MDM4 amplification<sup>104</sup>.

## POTENTIAL TREATMENT STRATEGIES

## — Targeted Therapies —

MDM2 antagonists disrupt the MDM2-p53 interaction, thereby stabilizing p53<sup>83</sup>. Preclinical studies have suggested that the amplification of MDM2, in the absence of concurrent TP53 mutations, may increase sensitivity to these agents<sup>84-85</sup>. Preliminary Phase 1 studies of the MDM2-p53 antagonist alrizomadlin (APG-115) reported a PR in a patient with liposarcoma harboring an MDM2 amplification and wildtype for TP53 and SD in 21%-38% (6/28 and 5/13, respectively) of patients in genomically unselected solid tumors<sup>86-87</sup>. A Phase 2 trial of alrizomadlin in combination with pembrolizumab reported a PR in 1 of 3 patients with malignant peripheral nerve sheath tumor that had failed standard therapy, as well as PRs in patients with multiple

## FREQUENCY &amp; PROGNOSIS

In the Glioblastoma Multiforme (GBM) TCGA dataset, amplification of MDM2 has been found in 8% of cases<sup>65</sup>. A study has reported amplification of the 12q14-15 region, where MDM2 and CDK4 reside, in 5% (2/42) of GBMs<sup>68</sup>. Amplification of

ORDERED TEST # ORD-1243577-01

GENOMIC FINDINGS

GENE

**RB1**

ALTERATION

inversion exons 18-24

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of limited clinical data<sup>105</sup> and strong preclinical data<sup>106-108</sup>, RB1 inactivation may be associated with sensitivity to inhibitors of Aurora kinase A, particularly in small cell lung cancer. It should be noted that a trial of the Aurora kinase A inhibitor alisertib in advanced prostate cancer did not find an association between RB1 deletion and clinical benefit<sup>109</sup>. Other approaches to target RB1 inactivation under investigation in preclinical studies include inhibitors of BCL-2 family members<sup>110</sup> and activation of the NOTCH pathway<sup>111</sup>.

— Potential Resistance —

Rb inactivation may predict resistance to CDK4/6 inhibitors such as palbociclib, abemaciclib, and ribociclib, which act upstream of Rb<sup>49,54-62</sup>.

— Nontargeted Approaches —

Loss of Rb function has been associated with increased sensitivity to cytotoxic agents and chemotherapeutics in both preclinical studies and in patients with bladder or breast cancer<sup>112-113</sup>.

FREQUENCY & PROGNOSIS

In the TCGA datasets, RB1 mutation or homozygous deletion was observed in 9% of glioblastomas<sup>65</sup> and 2.5% of lower grade glioma cases<sup>114</sup>. In one study, loss of RB1 transcript expression was observed in 10.6% of glioblastomas and occurred more frequently in the proneural subtype<sup>115</sup>. One study reports that mutation of RB1 is correlated with shorter survival in glioblastoma patients<sup>116</sup>. Several studies suggest that RB1, PTEN, and/or TP53 mutations

are early events in the development of glioblastoma<sup>117-119</sup>.

FINDING SUMMARY

RB1 encodes the retinoblastoma protein (Rb), a tumor suppressor and negative regulator of the cell cycle<sup>113,120</sup>. Alterations such as seen here may disrupt RB1 function or expression<sup>121-127</sup>.

POTENTIAL GERMLINE IMPLICATIONS

Mutations in RB1 underlie the development of retinoblastoma (RB), a rare tumor that arises at a rate of approximately 1:20,000 live births, with nearly 5,000 new cases worldwide per year<sup>128</sup>. Germline mutations in RB1 account for approximately 40% of RB tumors<sup>129</sup> and are associated with an increased risk of developing secondary malignancies that include soft tissue and bone sarcoma and malignant melanoma<sup>130-131</sup>. In the appropriate clinical context, germline testing of RB1 is recommended.

GENE

**TERT**

ALTERATION

promoter -146C>T

TRANSCRIPT ID

NM\_198253

CODING SEQUENCE EFFECT

-146C>T

VARIANT ALLELE FREQUENCY (% VAF)

43.0%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapeutic options for targeting tumors with TERT mutations are limited, although a variety of approaches have been investigated, including immunotherapies using TERT as a tumor-associated antigen and antisense oligonucleotide- or peptide-based therapies. TERT peptide vaccines showed limited anticancer efficacy in clinical trials<sup>132</sup>; however, in one preclinical study, the combination of a TERT peptide vaccine and anti-CTLA-4 therapy suppressed tumor growth<sup>133</sup>. A Phase 2 study of the TERT inhibitor imetelstat for patients with advanced non-small cell lung cancer reported no improvement in PFS or OS<sup>134</sup>.

FREQUENCY & PROGNOSIS

TERT promoter mutations have been reported in 51-59% of gliomas<sup>135-136</sup>, most frequently in glioblastoma (GBM, 54-84%), gliosarcoma (81%), oligodendroglioma (78%), and historically in oligoastrocytomas (25-31%) but less frequently in lower grade astrocytomas (10-18%) and in only 1% of ependymomas<sup>135-139</sup>. In patients with glioblastoma (GBM), the prevalence of TERT promoter mutation is lower in pediatric primary GBM (11%) and adult secondary GBM (28%) compared with adult primary GBM (58-83%)<sup>135,137</sup>. One study detected TERT promoter mutations in 78% (181/232) of IDH-wildtype GBM samples analyzed<sup>140</sup>. The significance of the TERT promoter mutation as an independent prognostic indicator in patients with glioma is not clear. While TERT promoter mutations significantly associate with poor prognosis in patients with GBM, this correlation may be due to the association with primary GBM as opposed to IDH-positive secondary GBM<sup>135,137,141-142</sup>. In the context of IDH-wildtype glioma, TERT mutations are associated with reduced OS, whereas in IDH-mutated, 1p/19q co-deleted oligodendroglioma, TERT mutations are associated with improved OS (NCCN CNS Cancers Guidelines, v5.2020). TERT promoter mutation has been shown to be significantly associated with increased TERT gene expression in astrocytoma, oligodendroglioma,

and GBM<sup>142</sup>.

FINDING SUMMARY

Telomerase reverse transcriptase (TERT, or hTERT) is a catalytic subunit of the telomerase complex, which is required to maintain appropriate chromosomal length<sup>143</sup>. Activation of TERT is a hallmark of cancer, being detected in up to 80-90% of malignancies and absent in quiescent cells<sup>144-146</sup>. Mutations within the promoter region of TERT that confer enhanced TERT promoter activity have been reported in two hotspots, located at -124 bp and -146 bp upstream of the transcriptional start site (also termed C228T and C250T, respectively)<sup>147-149</sup>, as well as tandem mutations at positions -124/-125 bp and -138/-139 bp<sup>147</sup>.

POTENTIAL DIAGNOSTIC IMPLICATIONS

TERT mutations are associated with 1p/19q co-deletion in oligodendrogliomas, and are highly recurrent in IDH/ATRX-wildtype glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v5.2020). Co-occurring TERT mutation, IDH mutation, and 1p/19q co-deletion is indicative of oligodendroglioma, whereas IDH mutation in the absence of TERT mutation is suggestive of astrocytoma (NCCN CNS Cancers Guidelines, v5.2020).

ORDERED TEST # ORD-1243577-01

**CLINICAL TRIALS**

**NOTE** Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.foundationmedicine.com/genomic-testing#support-services). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

**GENE**  
**CDK4**
**ALTERATION**  
amplification

**RATIONALE**  
CDK4 amplification may predict sensitivity to CDK4/6 inhibitors. As clinical and preclinical studies have suggested that loss of RB may predict

resistance to CDK4/6 inhibitors, it is not known whether these therapeutic approaches would be effective.

**NCT03239015**
**PHASE 2**

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

**TARGETS**  
EGFR, ERBB2, ERBB4, PARP, mTOR, MET, RET, ROS1, VEGFRs, BRAF, CDK4, CDK6

**LOCATIONS:** Shanghai (China)

**NCT04594005**
**PHASE 1/2**

CDK4/6 Tumor, Abemaciclib, Paclitaxel

**TARGETS**  
CDK4, CDK6

**LOCATIONS:** Seoul (Korea, Republic of)

**NCT03099174**
**PHASE 1**

This Study in Patients With Different Types of Cancer (Solid Tumours) Aims to Find a Safe Dose of Xentuzumab in Combination With Abemaciclib With or Without Hormonal Therapies. The Study Also Tests How Effective These Medicines Are in Patients With Lung and Breast Cancer.

**TARGETS**  
CDK4, CDK6, IGF-1, IGF-2, Aromatase, ER

**LOCATIONS:** Seoul (Korea, Republic of), Goyang (Korea, Republic of), Aichi, Nagoya (Japan), Kanagawa, Isehara (Japan), Tokyo, Chuo-ku (Japan), Tokyo, Koto-ku (Japan), Chiba, Kashiwa (Japan), Helsinki (Finland), Tampere (Finland), Turku (Finland)

**NCT04801966**
**PHASE NULL**

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

**TARGETS**  
CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

**LOCATIONS:** Melbourne (Australia)

ORDERED TEST # ORD-1243577-01

**CLINICAL TRIALS**
**NCT03994796**
**PHASE 2**

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

**TARGETS**  
ALK, ROS1, TRKA, TRKB, TRKC, CDK4,  
CDK6, PI3K, mTOR

**LOCATIONS:** Alaska, Washington

**NCT03158389**
**PHASE 1/2**

NCT Neuro Master Match - N<sup>2</sup>M<sup>2</sup> (NOA-20)

**TARGETS**  
ALK, RET, CDK4, CDK6, mTOR,  
MDM2, PD-L1, SMO

**LOCATIONS:** Berlin (Germany), Dresden (Germany), Regensburg (Germany), Bochum (Germany), Frankfurt am Main (Germany), Essen (Germany), Mainz (Germany), Heidelberg (Germany), Cologne (Germany), Mannheim (Germany)

**NCT04116541**
**PHASE 2**

A Study Evaluating the Activity of Anti-cancer Treatments Targeting Tumor Molecular Alterations/  
Characteristics in Advanced / Metastatic Tumors.

**TARGETS**  
CDK6, CDK4, MDM2, MET, RET, ROS1,  
VEGFRs

**LOCATIONS:** Nice (France), Lyon (France), Marseille (France), Toulouse (France), Bordeaux (France)

**NCT02981940**
**PHASE 2**

A Study of Abemaciclib in Recurrent Glioblastoma

**TARGETS**  
CDK4, CDK6

**LOCATIONS:** Utah, California, Massachusetts

**NCT02896335**
**PHASE 2**

Palbociclib In Progressive Brain Metastases

**TARGETS**  
CDK4, CDK6

**LOCATIONS:** Massachusetts

**NCT03310879**
**PHASE 2**

Study of the CDK4/6 Inhibitor Abemaciclib in Solid Tumors Harboring Genetic Alterations in Genes  
Encoding D-type Cyclins or Amplification of CDK4 or CDK6

**TARGETS**  
CDK4, CDK6

**LOCATIONS:** Massachusetts



ORDERED TEST # ORD-1243577-01

CLINICAL TRIALS

# GENE MDM2

ALTERATION  
amplification

## RATIONALE

Inhibitors of the MDM2-p53 interaction are being tested in clinical trials. Overexpression or

amplification of MDM2 may increase sensitivity to these agents, but more data are required.

### NCT04589845

#### PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

#### TARGETS

ALK, ROS1, TRKA, TRKB, TRKC, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

**LOCATIONS:** Zhongzheng Dist. (Taiwan), Taipei City (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Beijing (China), Woolloongabba (Australia), Darlinghurst (Australia), Randwick (Australia), Melbourne (Australia), Haifa (Israel)

### NCT03449381

#### PHASE 1

This Study Aims to Find the Best Dose of BI 907828 in Patients With Different Types of Advanced Cancer (Solid Tumors)

#### TARGETS

MDM2

**LOCATIONS:** Tokyo, Chuo-ku (Japan), Ottawa (Canada), Connecticut, New York, Tennessee, Florida

### NCT03611868

#### PHASE 1/2

A Study of APG-115 in Combination With Pembrolizumab in Patients With Metastatic Melanomas or Advanced Solid Tumors

#### TARGETS

MDM2, PD-1

**LOCATIONS:** Brisbane (Australia), California, Arizona, Missouri, Arkansas, Pennsylvania, New York, Tennessee, Texas

### NCT03158389

#### PHASE 1/2

NCT Neuro Master Match - N<sup>2</sup>M<sup>2</sup> (NOA-20)

#### TARGETS

ALK, RET, CDK4, CDK6, mTOR, MDM2, PD-L1, SMO

**LOCATIONS:** Berlin (Germany), Dresden (Germany), Regensburg (Germany), Bochum (Germany), Frankfurt am Main (Germany), Essen (Germany), Mainz (Germany), Heidelberg (Germany), Cologne (Germany), Mannheim (Germany)

### NCT03107780

#### PHASE 1

MDM2 Inhibitor AMG-232 in Treating Patients With Recurrent or Newly Diagnosed Glioblastoma

#### TARGETS

MDM2

**LOCATIONS:** California, Michigan, Pennsylvania, Massachusetts, New York, Maryland, North Carolina, Alabama

### NCT03725436

#### PHASE 1

ALRN-6924 and Paclitaxel in Treating Patients With Advanced, Metastatic, or Unresectable Solid Tumors

#### TARGETS

MDM2, MDM4

**LOCATIONS:** Texas

ORDERED TEST # ORD-1243577-01

**APPENDIX**
**Variants of Unknown Significance**

**NOTE** One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

**ATR**  
M1617V

**BRCA2**  
D974N

**CBL**  
L857F

**ERCC4**  
S334L

**MSH3**  
A62\_P63insSAA

**SF3B1**  
R524C

**SPEN**  
S2306del

**TBX3**  
A192T



ORDERED TEST # ORD-1243577-01

**APPENDIX**
**Genes Assayed in FoundationOne®CDx**

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

**DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS**

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTB	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXO2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDMS5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

**DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS**

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

\*TERC is an NCRNA

\*\*Promoter region of TERT is interrogated

**ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS**

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

ORDERED TEST # ORD-1243577-01

**APPENDIX**
**About FoundationOne®CDx**

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


**ABOUT FOUNDATIONONE CDx**

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:  
[www.rochefoundationmedicine.com/f1cdxtech](http://www.rochefoundationmedicine.com/f1cdxtech).

**INTENDED USE**

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

**TEST PRINCIPLES**

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

**THE REPORT**

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

**Diagnostic Significance**

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

**Qualified Alteration Calls (Equivocal and Subclonal)**

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

**Ranking of Therapies and Clinical Trials**
**Ranking of Therapies in Summary Table**

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

**Ranking of Clinical Trials**

Pediatric trial qualification → Geographical proximity → Later trial phase.

**NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION**

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) ([www.nccn.org](http://www.nccn.org)). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2021. All rights reserved. To view the most recent and complete version of the guidelines, go online to [NCCN.org](http://NCCN.org). NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

**Limitations**

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 06 December 2021  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1243577-01

## APPENDIX

## About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
2. TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation [https://www.accessdata.fda.gov/cdrh\\_docs/pdf17/P170019B.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf). The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
  3. The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
  4. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
  5. Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

6. Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

### VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

#### Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS		%CV*
Repeatability		5.11 - 10.40
Reproducibility		5.95 - 12.31
INDELS		%CV*
Repeatability		6.29 - 10.00
Reproducibility		7.33 - 11.71

\*Interquartile Range = 1<sup>st</sup> Quartile to 3<sup>rd</sup> Quartile

### VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert

panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

### VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

### LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

### NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

### NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or

ORDERED TEST # ORD-1243577-01

APPENDIX

About FoundationOne®CDx

governmental, will reimburse a patient for the cost of FoundationOne CDx.

#### TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

#### SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 5.1.1

The median exon coverage for this sample is 906x

© 2021 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 06 December 2021  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531



**ORDERED TEST #** ORD-1243577-01

**APPENDIX**
**References**

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Martinez R, et al. Oncology (2004) PMID: 15331927
7. Martinez R, et al. J. Cancer Res. Clin. Oncol. (2005) PMID: 15672285
8. Martinez R, et al. Cancer Genet. Cytogenet. (2007) PMID: 17498554
9. Szybka M, et al. Clin. Neuropathol. ( ) PMID: 12908754
10. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
11. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
12. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
13. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
14. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
15. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
16. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
17. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
18. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
19. Cristescu R, et al. Science (2018) PMID: 30309915
20. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
21. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
22. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
24. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
25. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
26. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
27. Touat M, et al. Nature (2020) PMID: 32322066
28. Bouffett E, et al. J. Clin. Oncol. (2016) PMID: 27001570
29. Johanns TM, et al. Cancer Discov (2016) PMID: 27683556
30. Lukas RV, et al. J. Neurooncol. (2018) PMID: 30073642
31. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
32. Patel RR, et al. Pediatr Blood Cancer (2020) PMID: 32386112
33. Johnson A, et al. Oncologist (2017) PMID: 28912153
34. Draaisma K, et al. Acta Neuropathol Commun (2015) PMID: 26699864
35. Wang L, et al. BMC Cancer (2020) PMID: 32164609
36. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
37. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
38. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
39. Rizvi NA, et al. Science (2015) PMID: 25765070
40. Johnson BE, et al. Science (2014) PMID: 24336570
41. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
42. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
43. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
44. Heitz E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
45. Nature (2012) PMID: 22810696
46. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
47. Dickson MA, et al. J. Clin. Oncol. (2013) PMID: 23569312
48. Flaherty KT, et al. Clin. Cancer Res. (2012) PMID: 22090362
49. Patnaik A, et al. Cancer Discov (2016) PMID: 27217383
50. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
51. Dickson et al., 2019; ASCO Abstract 11004
52. Dickson MA, et al. JAMA Oncol (2016) PMID: 27124835
53. Peguero et al., 2016; ASCO Abstract 2528
54. Condorelli R, et al. Ann. Oncol. (2018) PMID: 29236940
55. Fry DW, et al. Mol. Cancer Ther. (2004) PMID: 15542782
56. Dean JL, et al. Oncogene (2010) PMID: 20473330
57. Dean JL, et al. Cell Cycle (2012) PMID: 22767154
58. Garnett MJ, et al. Nature (2012) PMID: 22460902
59. Roberts PJ, et al. J. Natl. Cancer Inst. (2012) PMID: 22302033
60. O'Leary B, et al. Cancer Discov (2018) PMID: 30206110
61. Costa C, et al. Cancer Discov (2019) PMID: 31594766
62. Chen SH, et al. Oncogene (2018) PMID: 29059158
63. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
64. Gao J, et al. Sci Signal (2013) PMID: 23550210
65. Brennan CW, et al. Cell (2013) PMID: 24120142
66. Jonsson P, et al. Clin. Cancer Res. (2019) PMID: 31263031
67. Ceccarelli M, et al. Cell (2016) PMID: 26824661
68. Zheng S, et al. Genes Dev. (2013) PMID: 23796897
69. Kim H, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20080666
70. Ruano Y, et al. Am. J. Clin. Pathol. (2009) PMID: 19141386
71. Fischer U, et al. Mol. Cancer Res. (2008) PMID: 18403636
72. Bäcklund LM, et al. Br. J. Cancer (2005) PMID: 15970925
73. Choi YJ, et al. Oncogene (2014) PMID: 23644662
74. Cell (1995) PMID: 7736585
75. Musgrove EA, et al. Nat. Rev. Cancer (2011) PMID: 21734724
76. Wikman H, et al. Genes Chromosomes Cancer (2005) PMID: 15543620
77. Rao SK, et al. J. Neurooncol. (2010) PMID: 19609742
78. Chung L, et al. Am. J. Surg. Pathol. (2009) PMID: 19574885
79. Ragazzini P, et al. Histol. Histopathol. (2004) PMID: 15024701
80. Dujardin F, et al. Mod. Pathol. (2011) PMID: 21336260
81. Zhang K, et al. Cancer Res. (2013) PMID: 23393200
82. Horvai AE, et al. Mod. Pathol. (2009) PMID: 19734852
83. Cheok CF, et al. Nat Rev Clin Oncol (2011) PMID: 20975744
84. Ohnstad HO, et al. Cancer (2013) PMID: 23165797
85. Gamble LD, et al. Oncogene (2012) PMID: 21725357
86. Zhang et al., 2019; ASCO Abstract 3124
87. Rasco et al., 2019; ASCO Abstract 3126
88. Tolcher et al., 2021; ASCO Abstract 2506
89. Martinelli et al., 2016; EHA21 Abstract S504
90. Daver et al., 2018; ASH Abstract 767
91. Mascarenhas et al., 2019; ASH Abstract 134
92. Shustov et al., 2018; ASH Abstract 1623
93. Sallman et al., 2018; ASH Abstract 4066
94. Meric-Bernstam et al., 2017; ASCO Abstract 2505
95. Fischer U, et al. Int. J. Cancer (2010) PMID: 19839052
96. Sdek P, et al. Mol. Cell (2005) PMID: 16337594
97. Brady M, et al. Mol. Cell. Biol. (2005) PMID: 15632057
98. Li M, et al. Mol. Cell (2004) PMID: 15053880
99. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
100. Cordon-Cardo C, et al. Cancer Res. (1994) PMID: 8306343
101. Beroukhim R, et al. Nature (2010) PMID: 20164920
102. Kato S, et al. Clin. Cancer Res. (2017) PMID: 28351930
103. Singavi et al., 2017; ESMO Abstract 1140PD
104. Rizvi H, et al. J. Clin. Oncol. (2018) PMID: 29337640
105. Owonikoko et al., 2016; ESMO Abstract 14230
106. Hook KE, et al. Mol. Cancer Ther. (2012) PMID: 22222631
107. Gong X, et al. Cancer Discov (2019) PMID: 30373917
108. Oser MG, et al. Cancer Discov (2019) PMID: 30373918
109. Beltran H, et al. Clin. Cancer Res. (2019) PMID: 30232224
110. Allaman-Pillet N, et al. Ophthalmic Genet. ( ) PMID: 21955141
111. Viatour P, et al. J. Exp. Med. (2011) PMID: 21875955
112. Derenzini M, et al. Clin. Cancer Res. (2008) PMID: 18381962
113. Knudsen ES, et al. Nat. Rev. Cancer (2008) PMID: 19143056
114. Cancer Genome Atlas Research Network, et al. N. Engl. J. Med. (2015) PMID: 26061751
115. Goldhoff P, et al. J. Neuropathol. Exp. Neurol. (2012) PMID: 22157621
116. Bäcklund LM, et al. Clin. Cancer Res. (2003) PMID: 14519639
117. Chow LM, et al. Cancer Cell (2011) PMID: 21397855
118. Nobusawa S, et al. Brain Pathol. (2010) PMID: 20406234
119. Deighton RF, et al. Brain Pathol. (2010) PMID: 20175778
120. Burkhardt DL, et al. Nat. Rev. Cancer (2008) PMID: 18650841
121. Berge EO, et al. Mol. Cancer (2010) PMID: 20594292
122. Giacinti C, et al. Oncogene (2006) PMID: 16936740
123. Otterson GA, et al. Proc. Natl. Acad. Sci. U.S.A. (1997) PMID: 9342358
124. Otterson GA, et al. Am. J. Hum. Genet. (1999) PMID: 10486322
125. Qin XQ, et al. Genes Dev. (1992) PMID: 1534305
126. Rubin SM, et al. Cell (2005) PMID: 16360038
127. Sun H, et al. Mol. Cell. Biol. (2006) PMID: 16449662
128. Chen Z, et al. Hum. Mutat. (2014) PMID: 24282159
129. Yun J, et al. Int J Ophthalmol (2011) PMID: 22553621
130. Houston SK, et al. Int Ophthalmol Clin (2011) PMID: 21139478
131. Ng AK, et al. Semin Radiat Oncol (2010) PMID: 19959033
132. Nat Rev Clin Oncol (2017) PMID: 27245281
133. Duperret EK, et al. Mol Ther (2018) PMID: 29249395
134. Chiappori AA, et al. Ann Oncol (2015) PMID: 25467017
135. Killela PJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23530248
136. Killela PJ, et al. Oncotarget (2014) PMID: 24722048
137. Nonoguchi N, et al. Acta Neuropathol. (2013) PMID: 23955565
138. Liu X, et al. Cell Cycle (2013) PMID: 23603989
139. Koelsche C, et al. Acta Neuropathol. (2013) PMID: 24154961
140. Yan et al. 2020; DOI:10.1200/P0.19.00385
141. Reitman ZJ, et al. Acta Neuropathol. (2013) PMID: 24217890
142. Arita H, et al. Acta Neuropathol. (2013) PMID: 23764841
143. Shay JW, et al. Semin. Cancer Biol. (2011) PMID: 22015685
144. Shay JW, et al. Eur. J. Cancer (1997) PMID: 9282118
145. Kim NW, et al. Science (1994) PMID: 7605428
146. Hanahan D, et al. Cell (2000) PMID: 10647931
147. Horn S, et al. Science (2013) PMID: 23348503
148. Huang FW, et al. Science (2013) PMID: 23348506
149. Vinagre J, et al. Nat Commun (2013) PMID: 23887589

© 2021 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 06 December 2021  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D0207531  
 Shakti Ramkissoon, M.D., Ph.D., M.M. Sc, Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D0207531  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D0207531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D0207531