

ACT Onco[®] + Report

| PATIENT | | |
|------------------------------|-----------------------------|----------------------|
| Identifier: 李鈺棋 | | Patient ID: 47556672 |
| Date of Birth: Feb 28, 1966 | | Gender: Male |
| Diagnosis: Esophageal cancer | | |
| ORDERING PHYSICIAN | | |
| Name: 陳明晃醫師 | | Tel: 886-228712121 |
| Facility: 臺北榮總 | | |
| Address: 臺北市北投區石牌路二段 201 號 | | |
| SPECIMEN | | |
| Specimen ID: S11107471I | Collection site: Lymph node | Type: FFPE tissue |
| Date received: May 15, 2023 | Lab ID: AA-23-03052 | D/ID: NA |

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

| Genomic Alterations/Biomarkers | Probable Effects in Patient's Cancer Type | | Probable Sensitive in Other Cancer Types |
|--------------------------------|---|-----------|--|
| | Sensitive | Resistant | |
| Not detected | | | |

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

| Genomic Alterations/Biomarkers | Possibly Sensitive | Possibly Resistant |
|--------------------------------|--------------------------------------|--------------------|
| CCND1 Amplification | Abemaciclib, Palbociclib, Ribociclib | - |

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACT Onco[®] + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

| Gene | Amino Acid Change | Allele Frequency |
|-------------|-------------------|------------------|
| <i>TP53</i> | E285K | 41.8% |

- Copy Number Alterations

| Chromosome | Gene | Variation | Copy Number |
|------------|-------------------|-----------------------|-------------|
| Chr13 | <i>RB1</i> | Heterozygous deletion | 1 |
| Chr17 | <i>FLCN, TP53</i> | Heterozygous deletion | 1 |
| Chr22 | <i>CHEK2, NF2</i> | Heterozygous deletion | 1 |
| Chr9 | <i>CDKN2A</i> | Heterozygous deletion | 1 |
| Chr11 | <i>CCND1</i> | Amplification | 7 |
| Chr1 | <i>NRAS</i> | Amplification | 16 |
| Chr18 | <i>TYMS</i> | Amplification | 20 |

- Fusions

| Fusion Gene & Exon | Transcript ID |
|--|---------------|
| No fusion gene detected in this sample | |

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

| Biomarker | Results |
|----------------------------------|-----------------------------|
| Tumor Mutational Burden (TMB) | 3.2 muts/Mb |
| Microsatellite Instability (MSI) | Microsatellite stable (MSS) |

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 48% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

| Genomic Alterations | Therapies | Effect |
|----------------------------|--------------------------------------|------------------|
| Level 3B | | |
| CCND1 Amplification | Abemaciclib, Palbociclib, Ribociclib | sensitive |

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

| Level | Description |
|-----------|--|
| 1 | FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication |
| 2 | Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication |
| 3A | Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type |
| 3B | Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required) |
| 4 | Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies |

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

| Genomic Alterations | Potential Clinical Effects |
|---------------------|----------------------------|
| Not detected | |

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

| Genomic Alterations | Therapies | Effect | Level of Evidence | Cancer Type |
|------------------------------|--------------|-----------------------|-------------------|-------------------|
| TYMS Amplification | Fluorouracil | Less sensitive | Clinical | Colorectal cancer |
| | Pemetrexed | Less sensitive | Clinical | Lung cancer |

HORMONAL THERAPIES

| Genomic Alterations | Therapies | Effect | Level of Evidence | Cancer Type |
|-------------------------------|-------------|-----------------------|-------------------|---------------|
| CCND1 Amplification | Anastrozole | Less sensitive | Clinical | Breast cancer |
| | Tamoxifen | Less sensitive | Clinical | Breast cancer |

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

TP53 E285K, Heterozygous deletion

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[1]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[2].

E285K mutation is located in the DNA-binding domain (DBD) of the p53 protein^[3]. This mutation resulted in a temperature-dependent decrease in transcriptional activation by p53 in vitro^[4].

Loss of the second wild-type allele resulted in the biallelic inactivation of the gene.

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[5].

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[6]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[7].

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[8][9][10]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[11]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[12][13]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[14].

CCND1 Amplification

Biological Impact

The cyclin D1 (CCND1) gene encodes a protein involved in the control of cell growth, proliferation, transcription, and DNA repair^[15]. CCND1 forms a complex with CDK4 and CDK6, leading to G1-S cell-cycle transition by inhibiting the retinoblastoma (RB) protein^[15]. Amplification or overexpression of CCND1 could be oncogenic and is associated with carcinogenesis of various cancer types^[16].

Therapeutic and prognostic relevance

Several CDK4 inhibitors, including palbociclib (PD0332991), LEE011, and LY2835219 have entered clinical trials for tumors with CCND1 amplification^{[17][18]}. In the Phase II study of palbociclib and letrozole in patients with ER-positive HER2-negative metastatic breast cancer, patient selection based on CCND1 amplification or p16 loss did not further improve patient outcome^[19]. Preclinical studies also demonstrated conflicting results regarding the correlation between high-level CCND1 and palbociclib sensitivity^{[20][21][22]}.

CCND1 amplification has been implicated in predicting poor clinical outcomes in postmenopausal breast cancer patients treated with either anastrozole or tamoxifen^[23]. In lung cancer patients, the increased CCND1 copy number is

ACT Onco[®] + Report

associated with poorer overall survival^[24]. A retrospective study showed that melanoma patients whose tumor harboring CCND1, cRAF or KRAS gene copy number gain had better treatment response with CPS (carboplatin, paclitaxel, and sorafenib)^[25]. Of note, 3 of 4 patients treated with ribociclib for the longest duration had CCND1 amplification in a phase I trial^[26].

Amplification of CCND1 are frequent and contributes to dedifferentiation and cellular proliferative activity of intrahepatic cholangiocarcinoma (ICC), and also indicates a poor prognosis for ICC patients^[27]. Of note, CCND1 amplification has been selected as an inclusion criterion for the trial examining CDK4/6 inhibitors in different types of malignant solid tumors (NCT02187783, NCT02896335, NCT03526250, NCT02693535, NCT01037790, NCT03454919, NCT03310879, and NCT03356223).

CDKN2A Heterozygous deletion

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^{[28][29][30]}. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[31]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[32][33]}.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[34][19]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[35][36][37]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[38][26][39]}. CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

The phase II TAPUR trial demonstrated clinical benefits to palbociclib monotherapy in advanced NSCLC or head and neck cancer harboring a CDKN2A mutation or copy number loss. However, pancreatic and biliary cancer patients harboring a CDKN2A mutation or copy number loss did not demonstrate an objective response or stable disease when treated with palbociclib monotherapy for 16 weeks (DOI: 10.1200/JCO.2021.39.15_suppl.6043)^{[40][41]}.

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[19][42][43]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[36]. Administration of combined palbociclib and MEK inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[44].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[45].

ACT Onco[®] + Report

CHEK2 Heterozygous deletion

Biological Impact

The checkpoint kinase 2 (CHEK2 or CHK2) gene encodes a serine/threonine protein kinase involved in transducing DNA damage signals that are required for both the intra-S phase and G2/M checkpoints^[46]. CHEK2 heterozygosity has been shown to cause haploinsufficient phenotypes that can contribute to tumorigenesis through inappropriate S phase entry, accumulation of DNA damage during replication, and failure to restrain mitotic entry^{[47][48]}. CHEK2 aberrations are associated with glioblastoma, breast, ovarian, prostate, colorectal, gastric, thyroid, and lung cancers^{[49][50][51][52][53]}.

Therapeutic and prognostic relevance

Olaparib is FDA-approved for treating mCRPC patients harboring mutations in homologous recombination repair (HRR) genes, including CHEK2.

CHEK2 mutation has been determined as an inclusion criterion for the trials evaluating olaparib, rucaparib, niraparib, and talazoparib efficacies in various types of solid tumors (NCT03297606, NCT01968213, NCT03840967, NCT02401347, NCT03148795).

In TBCRC 048 trial, olaparib treatment did not show response in 7 metastatic breast cancer patients with germline mutations in CHEK2 (SD: n=3, PD: n=4)^[54]. In TRITON2 trial, rucaparib treatment had limited response in 12 mCRPC patients with CHEK2 alterations^[55].

FLCN Heterozygous deletion

Biological Impact

The FLCN gene encodes the tumor suppressor, Folliculin, a GTPase activating protein (GAP) for RagC/D GTPase proteins involved in amino acid sensing and signaling to mTORC1^[56]. FLCN has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[57][58]}. Inactivation of the FLCN gene by mutation or deletion results in the activation of the mTOR pathway and AKT signaling^{[59][60]}. Germline mutation of the FLCN gene causes the Birt-Hogg-Dubé syndrome, a rare disorder that is characterized by benign hamartomatous skin lesions and an increased risk of pneumothorax and renal tumors^[61].

Therapeutic and prognostic relevance

In a prospective Phase 2 study, four anaplastic thyroid cancer (ATC)/ poorly differentiated thyroid cancer (PDTC) patients who had PI3K/mTOR/AKT alterations, including TSC2, FLCN or NF1, showed impressive progression-free survival (PFS) of 15.2 months after receiving everolimus^[62]. mTOR inhibition via rapamycin also demonstrated potential in inhibiting the growth of renal cells deficient in FLCN in the preclinical setting^[63].

NF2 Heterozygous deletion

Biological Impact

The neurofibromin (NF2) gene encodes the protein Merlin, a tumor suppressor that functions as a negative regulator of the PI3K/AKT/mTOR pathway^{[64][65][66]}. NF2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[67]. Inactivation germline mutations in the NF2 are associated with the hereditary neurofibromatosis type 2, a disorder characterized by the growth of noncancerous tumors in the nervous system^{[64][68]}. Somatic mutations or deletion of NF2 are frequently observed in human cancers, including 20-50% of pleural mesotheliomas^[69], 6% papillary renal cell carcinoma, 5% pancreas cancer, and 4% melanoma (cbioPortal; June 2015), and less frequently in other cancers^[70].

ACT Onco[®] + Report

Therapeutic and prognostic relevance

Genomic alterations with activating effects on the mTOR signaling pathway have been identified to confer sensitivity to everolimus across multiple cancer types^{[71][72][73][74]}. There are at least two case studies indicating the clinical efficacy of everolimus in bladder cancer and urothelial carcinoma^{[75][76]}, both harboring NF2 truncating mutations. Preclinical evidence has shown the efficacy of MEK1/2 inhibitor selumetinib in KRAS-mutant thyroid cancer model with NF2 loss^[77].

Analysis of afatinib-plus-cetuximab-resistant biopsy specimens revealed a loss-of-function alteration in genes that modulate mTOR signaling pathway, including NF2 and TSC1^[78].

NRAS Amplification

Biological Impact

The neuroblastoma RAS viral oncogene homolog (NRAS) gene encodes a membrane-associated RAS protein which belongs to the large family of small GTPases. RAS proteins cycle between an active (GTP-bound) and an inactive (GDP-bound) state, to regulate intracellular oncogenic MAPK and PI3K signaling pathways^[79]. Activated RAS proteins mediate the regulation of cellular proliferation and other cellular functions through the activation of distinct intracellular signaling pathways, including the RAF/MEK/ERK and PI3K/AKT/mTOR pathways. Mutations in NRAS are present in thyroid cancer, ovarian cancers, melanoma and hematological cancers^{[80][81][82]}.

Overexpression of wild-type NRAS has been shown to enhance the chemical-induced tumorigenesis in mouse model^[83].

Therapeutic and prognostic relevance

Cetuximab and panitumumab are FDA-approved for treating RAS wild-type metastatic colorectal cancer. The NCCN for CRC recommends cetuximab and panitumumab use only if both KRAS and NRAS genes are normal.

NRAS mutation has been determined as an inclusion criterion for the trials evaluating cobimetinib or selumetinib efficacies in various types of solid tumors (NCT04109456, NCT03732703, NCT03181100, NCT02639546, NCT02664935).

Studies have shown that MEK1/2 inhibitors trametinib and binimetinib are effective in treating melanoma and biliary tract cancer patients with an NRAS mutation^{[84][85][86]}. However, a phase II trial with 47 patients with a refractory solid tumor harboring a codon 12, 13, or 61 NRAS mutation showed limited response to binimetinib monotherapy, with an ORR of only 2.1%^[87]. Combining MEK and mTOR inhibitors is being evaluated as a potential strategy in CRC^{[88][89]}.

Preclinical studies have demonstrated the efficacy of MEK inhibitors, such as selumetinib and trametinib, against various NRAS-mutant cancer cell lines^{[90][91][92][93]}. In a mouse model of melanoma with inducible NRAS Q61K mutation, tumor regression occurred with the combination of MEK and CDK4/6 inhibitors, leading to the extinction of the mutant NRAS^[94].

A retrospective study showed that relatively high expression of NRAS associates with longer progression-free survival and overall survival in patients with pancreatic adenocarcinoma after initial surgery^[95]. Of note, elevated expression of NRAS isoform 2 was found to be associated with acquired resistance to BRAF inhibitor, vemurafenib in vitro^[96].

RB1 Heterozygous deletion

Biological Impact

The Retinoblastoma (RB1) gene encodes a tumor suppressor that negatively regulates the cell cycle, cell division, and DNA replication^[97]. Loss-of-function RB1 could lead to unregulated cell division and growth, abrogation of multiple mechanisms that safeguard against cellular transformation, and tumorigenesis^[98]. RB1 has also been implicated as a haploinsufficient tumor suppressor with one copy loss may lead to weak protein expression and is insufficient to execute

ACT Onco[®] + Report

its original physiological functions^{[99][100][101]}. Deletion or inactivating mutation of RB1 is found in a number of tumors, including lung, prostate, bladder, breast cancers and sarcomas. RB1 mutations are found in approximately half of all retinoblastoma cases^[102].

Therapeutic and prognostic relevance

A deleterious mutation in one or more of the three DNA repair genes ATM, RB1, and FANCC predicted pathologic response and better overall survival to cisplatin-based chemotherapy for muscle-invasive bladder cancer patients^[103]. High RB loss was found to be associated with improved pathologic clinical response in breast cancer patients treated with 5-fluorouracil/adriamycin/cytosine (FAC), T/FAC, and Taxane/Adriamycin neoadjuvant therapy^[104].

Clinical and experimental data suggested that a non-functional retinoblastoma pathway is associated with resistance to tamoxifen in breast cancer^{[105][106]}.

Acquired RB1 mutations were found in hormone receptor positive breast cancer patients who developed resistance to palbociclib or ribociclib treatment^[107]. Preclinical data also showed that knockdown of RB1 would impair antitumor activity of CDK4/6 inhibitor, abemaciclib^[108].

Two large-scale genome-sequencing projects have identified a high prevalence of mutations in TP53 and RB1 in small cell lung cancer (SCLC)^{[109][110]}. Analyses of repeat biopsy samples from patients with EGFR-mutant adenocarcinoma that had transformed to the SCLC subtype have revealed that 100% of these patients have loss of RB1 and may be the alteration that induces this non-small-cell to small-cell transformation^{[106][111]}.

TYMS Amplification

Biological Impact

TYMS (Thymidylate Synthetase) gene encodes the thymidylate synthase that catalyzes the methylation of deoxyuridylate to deoxythymidylate. The enzyme is critical for DNA replication and repair^{[112][113][114]}. TYMS polymorphisms may be associated with etiology of neoplasia, including acute lymphoblastic leukemia^[115], breast cancer, and response to chemotherapy^[116].

Therapeutic and prognostic relevance

Thymidylate synthase gene amplification was associated with pemetrexed resistance in patients with advanced non-small cell lung cancer^{[117][118][119][120]}, and 5-FU resistance in CRC patients^[121].

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Abemaciclib (VERZENIO)

Abemaciclib is a cyclin-dependent kinase 4/6 (CDK4/6) inhibitor. Abemaciclib is developed and marketed by Eli Lilly under the trade name VERZENIO.

- FDA Approval Summary of Abemaciclib (VERZENIO)

| | |
|--|---|
| MONARCH E NCT03155997 | Breast cancer (Approved on 2021/10/12) |
| | HR+/HER2- Abemaciclib + tamoxifen/aromatase inhibitor vs. Tamoxifen/aromatase inhibitor [IDFS at 36 months(%): 86.1 vs. 79.0] |
| MONARCH 3 ^[122] NCT02246621 | Breast cancer (Approved on 2018/02/26) |
| | HR+/HER2- Abemaciclib + anastrozole/letrozole vs. Placebo + anastrozole/letrozole [PFS(M): 28.2 vs. 14.8] |
| MONARCH 2 ^[43] NCT02107703 | Breast cancer (Approved on 2017/09/28) |
| | HR+/HER2- Abemaciclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 16.4 vs. 9.3] |
| MONARCH 1 ^[123] NCT02102490 | Breast cancer (Approved on 2017/09/28) |
| | HR+/HER2- Abemaciclib [ORR(%): 19.7 vs. 17.4] |

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

| | |
|--|--|
| RADIANT-4 ^[124] NCT01524783 | Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26) |
| | - Everolimus vs. Placebo [PFS(M): 11 vs. 3.9] |
| BOLERO-2 ^[125] NCT00863655 | Breast cancer (Approved on 2012/07/20) |
| | ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2] |
| EXIST-2 NCT00790400 | Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26) |
| | - Everolimus vs. Placebo [ORR(%): 41.8 vs. 0] |
| RADIANT-3 ^[126] NCT00510068 | Pancreatic neuroendocrine tumor (Approved on 2011/05/05) |
| | - Everolimus vs. Placebo [PFS(M): 11 vs. 4.6] |
| EXIST-1 ^[127] NCT00789828 | Subependymal giant cell astrocytoma (Approved on 2010/10/29) |
| | - Everolimus vs. Placebo [ORR(%): 35.0] |
| RECORD-1 ^[128] NCT00410124 | Renal cell carcinoma (Approved on 2009/05/30) |
| | - Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9] |

ACT Onco[®] + Report

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (ZEJULA)

| | |
|--|---|
| PRIMA NCT02655016 | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29) |
| | - |
| NOVA^[129] NCT01847274 | Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2] |
| | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27) |
| | - |
| | Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7] |

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (LYNPARZA)

| | |
|---|---|
| OlympiA NCT02032823 | Her2-negative high-risk early breast cancer (Approved on 2022/03/11) |
| | HER2-/gBRCA mutation |
| PROfound^[130] NCT02987543 | Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):] |
| | Prostate cancer (Approved on 2020/05/19) |
| PAOLA-1^[131] NCT02477644 | HRR genes mutation |
| | Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5] |
| POLO^[132] NCT02184195 | Ovarian cancer (Approved on 2020/05/08) |
| | HRD+ |
| SOLO-1^[133] NCT01844986 | Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7] |
| | Pancreatic adenocarcinoma (Approved on 2019/12/27) |
| SOLO-2/ENGOT-Ov21^[135] NCT01874353 | gBRCA mutation |
| | Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8] |
| OlympiAD^[134] NCT02000622 | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19) |
| | gBRCA mutation or sBRCA mutation |
| Study19^[136] NCT00753545 | Olaparib vs. Placebo [PFS(M): NR vs. 13.8] |
| | Breast cancer (Approved on 2018/02/06) |
| | HER2-/gBRCA mutation |
| | Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2] |
| | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17) |
| | gBRCA mutation |
| | Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5] |
| | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17) |
| | - |
| | Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8] |

ACT Onco® + Report

Palbociclib (IBRANCE)

Palbociclib is an oral, cyclin-dependent kinase (CDK) inhibitor specifically targeting CDK4 and CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Palbociclib is developed and marketed by Pfizer under the trade name IBRANCE.

- FDA Approval Summary of Palbociclib (IBRANCE)

| | |
|---|---|
| PALOMA-2 ^[137] NCT01740427 | Breast cancer (Approved on 2017/03/31) |
| | ER+/HER2- Palbociclib + letrozole vs. Placebo + letrozole [PFS(M): 24.8 vs. 14.5] |
| PALOMA-3 ^[138] NCT01942135 | Breast cancer (Approved on 2016/02/19) |
| | ER+/HER2- Palbociclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 9.5 vs. 4.6] |

Ribociclib (KISQALI)

Ribociclib is a cyclin-dependent kinase (CDK) inhibitor specifically targeting cyclin D1/CDK4 and cyclin D3/CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Ribociclib is developed by Novartis and Astex Pharmaceuticals and marketed by Novartis under the trade name KISQALI.

- FDA Approval Summary of Ribociclib (KISQALI)

| | |
|---|--|
| MONALEESA-2 ^[42] NCT01958021 | Breast cancer (Approved on 2017/03/13) |
| | HR+/HER2- Ribociclib vs. Letrozole [PFS(M): NR vs. 14.7] |

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

| | |
|---|--|
| TRITON2 NCT02952534 | Prostate cancer (Approved on 2020/05/15) |
| | gBRCA mutation or sBRCA mutation Rucaparib [ORR(%): 44.0, DOR(M): NE] |
| ARIEL3 ^[139] NCT01968213 | Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06) |
| | - Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4] |

ACT Onco[®] + Report

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

| | |
|---|--|
| EMBRACA ^[140] NCT01945775 | Breast cancer (Approved on 2018/10/16) |
| | HER2-/gBRCA mutation |
| | Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6] |

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

| | |
|----------------------|--|
| [141] NCT00065468 | Renal cell carcinoma (Approved on 2007/05/30) |
| | - |
| | Temsirolimus vs. Ifn- α [OS(M): 10.9 vs. 7.3] |

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

| Gene | Amino Acid Change | Exon | cDNA Change | Accession Number | COSMIC ID | Allele Frequency | Coverage |
|------|-------------------|------|-------------|------------------|-----------|------------------|----------|
| TP53 | E285K | 8 | c.853G>A | NM_000546 | COSM10722 | 41.8% | 595 |

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

| Gene | Amino Acid Change | Exon | cDNA Change | Accession Number | COSMIC ID | Allele Frequency | Coverage |
|---------|----------------------|------|----------------------|------------------|-------------|------------------|----------|
| ALK | A371T | 4 | c.1111G>A | NM_004304 | COSM5949450 | 20.0% | 894 |
| ARID1B | Q124_Q131del | 1 | c.369_392del | NM_017519 | COSM7345407 | 40.0% | 936 |
| BRIP1 | M473L | 10 | c.1417A>C | NM_032043 | - | 18.5% | 395 |
| ERBB2 | R143Q | 3 | c.428G>A | NM_004448 | COSM1382867 | 57.6% | 819 |
| ERCC3 | S764L | 15 | c.2291C>T | NM_000122 | COSM3042186 | 26.3% | 612 |
| ESR1 | G145S | 1 | c.433G>A | NM_000125 | - | 31.2% | 564 |
| ETV4 | V247I | 8 | c.739G>A | NM_001079675 | - | 46.1% | 1248 |
| FAM46C | S203C | 2 | c.608C>G | NM_017709 | - | 7.7% | 2046 |
| INSR | R410Q | 5 | c.1229G>A | NM_000208 | COSM4766585 | 40.5% | 1764 |
| KMT2C | A3692T | 43 | c.11074G>A | NM_170606 | - | 58.7% | 1113 |
| PALB2 | I374L | 4 | c.1120A>C | NM_024675 | - | 13.3% | 732 |
| PIK3C2G | H524Q | 11 | c.1572C>A | NM_004570 | - | 15.8% | 1692 |
| PTPRT | G1111E | 24 | c.3332G>A | NM_007050 | COSM3546438 | 16.8% | 888 |
| SYNE1 | Splice region | - | c.24642+3A>G | NM_182961 | - | 39.9% | 343 |
| SYNE1 | Splice region | - | c.9807+6G>A | NM_182961 | - | 62.5% | 1463 |
| TERT | L1019_P1020 delinsPS | 14 | c.3056_3058delinsCCT | NM_198253 | - | 25.0% | 416 |


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Feb 24, 2023
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11107471I
- Collection site: Lymph node
- Examined by: Dr. Yun-An Chen
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 20%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 40%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 3%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 10%
 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 882x
- Target Base Coverage at 100x: 95%

RNA test

- Average unique RNA Start Sites per control GSP2: 118

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

ACT Onco[®] + Report

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

ACT Onco[®] + Report

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

解剖病理專科醫師朱盈霞
Ying-Hsia Chu, M.D.
病解字第 000653 號



Sign Off

解剖病理專科醫師朱盈霞
Ying-Hsia Chu, M.D.
病解字第 000653 號



ACT Onco[®] + Report

GENE LIST SNV & CNV

| | | | | | | | | | | | |
|----------|---------|---------|----------|----------|---------|-----------|-----------|----------|----------|----------|----------|
| ABCB1* | ABCC2* | ABCG2* | ABL1 | ABL2 | ADAMTS1 | ADAMTS13 | ADAMTS15 | ADAMTS16 | ADAMTS18 | ADAMTS6 | ADAMTS9 |
| ADAMTS11 | ADGRA2 | ADH1C* | AKT1 | AKT2 | AKT3 | ALDH1A1* | ALK | AMER1 | APC | AR | ARAF |
| ARID1A | ARID1B | ARID2 | ASXL1 | ATM | ATR | ATRX | AURKA | AURKB | AXIN1 | AXIN2 | AXL |
| B2M | BAP1 | BARD1 | BCL10 | BCL2* | BCL2L1 | BCL2L2* | BCL6 | BCL9 | BCOR | BIRC2 | BIRC3 |
| BLM | BMPR1A | BRAF | BRCA1 | BRCA2 | BRD4 | BRIP1 | BTG1 | BTG2* | BTB | BUB1B | CALR |
| CANX | CARD11 | CASP8 | CBFB | CBL | CCNA1 | CCNA | CCNB1 | CCNB2 | CCNB3 | CCND1 | CCND2 |
| CCND3 | CCNE1 | CCNE2 | CCNH | CD19 | CD274 | CD58 | CD70* | CD79A | CD79B | CDC73 | CDH1 |
| CDK1 | CDK12 | CDK2 | CDK4 | CDK5 | CDK6 | CDK7 | CDK8 | CDK9 | CDKN1A | CDKN1B | CDKN2A |
| CDKN2B | CDKN2C | CEBPA* | CHEK1 | CHEK2 | CIC | CREBBP | CRKL | CRLF2 | CSF1R | CTCF | CTLA4 |
| CTNNA1 | CTNNB1 | CUL3 | CYLD | CYP1A1* | CYP2B6* | CYP2C19* | CYP2C8* | CYP2D6 | CYP2E1* | CYP3A4* | CYP3A5* |
| DAXX | DCUN1D1 | DDR2 | DICER1 | DNMT3A | DOT1L | DPYD | DTX1 | E2F3 | EGFR | EP300 | EPCAM |
| EPHA2 | EPHA3 | EPHA5 | EPHA7 | EPHB1 | ERBB2 | ERBB3 | ERBB4 | ERCC1 | ERCC2 | ERCC3 | ERCC4 |
| ERCC5 | ERG | ESR1 | ESR2 | ETV1 | ETV4 | EZH2 | FAM46C | FANCA | FANCC | FANCD2 | FANCE |
| FANCF | FANCG | FANCL | FAS | FAT1 | FBXW7 | FCGR2B | FGF1* | FGF10 | FGF14 | FGF19* | FGF23 |
| FGF3 | FGF4* | FGF6 | FGFR1 | FGFR2 | FGFR3 | FGFR4 | FH | FLCN | FLT1 | FLT3 | FLT4 |
| FOXL2* | FOXP1 | FRG1 | FUBP1 | GATA1 | GATA2 | GATA3 | GNA11 | GNA13 | GNAQ | GNAS | GREM1 |
| GRIN2A | GSK3B | GSTP1* | GSTT1* | HGF | HIF1A | HIST1H1C* | HIST1H1E* | HNF1A | HR | HRAS* | HSP90AA1 |
| HSP90AB1 | HSPA4 | HSPA5 | IDH1 | IDH2 | IFNL3* | IGF1 | IGF1R | IGF2 | IKBK | IKBKE | IKZF1 |
| IL6 | IL7R | INPP4B | INSR | IRF4 | IRS1 | IRS2* | JAK1 | JAK2 | JAK3 | JUN* | KAT6A |
| KDM5A | KDM5C | KDM6A | KDR | KEAP1 | KIT | KMT2A | KMT2C | KMT2D | KRAS | LCK | LIG1 |
| LIG3 | LMO1 | LRP1B | LYN | MALT1 | MAP2K1 | MAP2K2 | MAP2K4 | MAP3K1 | MAP3K7 | MAPK1 | MAPK3 |
| MAX | MCL1 | MDM2 | MDM4 | MED12 | MEF2B | MEN1 | MET | MITF | MLH1 | MPL | MRE11 |
| MSH2 | MSH6 | MTHFR* | MTOR | MUC16 | MUC4 | MUC6 | MUTYH | MYC | MYCL | MYCN | MYD88 |
| NAT2* | NBN | NEFH | NF1 | NF2 | NFE2L2 | NFKB1 | NFKBIA | NKX2-1* | NOTCH1 | NOTCH2 | NOTCH3 |
| NOTCH4 | NPM1 | NQO1* | NRAS | NSD1 | NTRK1 | NTRK2 | NTRK3 | PAK3 | PALB2 | PARP1 | PAX5 |
| PAX8 | PBRM1 | PDCD1 | PDCD1LG2 | PDGFRA | PDGFRB | PDIA3 | PGF | PHOX2B* | PIK3C2B | PIK3C2G | PIK3C3 |
| PIK3CA | PIK3CB | PIK3CD | PIK3CG | PIK3R1 | PIK3R2 | PIK3R3 | PIM1 | PMS1 | PMS2 | POLB | POLD1 |
| POLE | PPARG | PPP2R1A | PRDM1 | PRKAR1A | PRKCA | PRKCB | PRKCG | PRKCI | PRKCQ | PRKDC | PRKN |
| PSMB8 | PSMB9 | PSME1 | PSME2 | PSME3 | PTCH1 | PTEN | PTGS2 | PTPN11 | PTPRD | PTPRT | RAC1 |
| RAD50 | RAD51 | RAD51B | RAD51C | RAD51D | RAD52 | RAD54L | RAF1 | RARA | RB1 | RBM10 | RECQL4 |
| REL | RET | RHOA | RICTOR | RNF43 | ROS1 | RPPH1 | RPTOR | RUNX1 | RUNX1T1 | RXRA | SDHA |
| SDHB | SDHC | SDHD | SERPINB3 | SERPINB4 | SETD2 | SF3B1 | SGK1 | SH2D1A* | SLC19A1* | SLC22A2* | SLC01B1* |
| SLC01B3* | SMAD2 | SMAD3 | SMAD4 | SMARCA4 | SMARCB1 | SMO | SOC1* | SOX2* | SOX9 | SPEN | SPOP |
| SRC | STAG2 | STAT3 | STK11 | SUFU | SYK | SYNE1 | TAF1 | TAP1 | TAP2 | TAPBP | TBX3 |
| TEK | TERT | TET1 | TET2 | TGFBR2 | TMSB4X* | TNF | TNFAIP3 | TNFRSF14 | TNFSF11 | TOP1 | TP53 |
| TPMT* | TSC1 | TSC2 | TSHR | TYMS | U2AF1 | UBE2A* | UBE2K | UBR5 | UGT1A1* | USH2A | VDR* |
| VEGFA | VEGFB | VHL | WT1 | XIAP | XPO1 | XRCC2 | ZNF217 | | | | |

*Analysis of copy number alterations NOT available.

FUSION

| | | | | | | | | | | | | |
|-----|------|------|-------|-------|-------|-----|------|-------|-------|-------|-----|------|
| ALK | BRAF | EGFR | FGFR1 | FGFR2 | FGFR3 | MET | NRG1 | NTRK1 | NTRK2 | NTRK3 | RET | ROS1 |
|-----|------|------|-------|-------|-------|-----|------|-------|-------|-------|-----|------|


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION


| Gene | Therapies | Possible effect |
|---------------|---|------------------|
| <i>CDKN2A</i> | Abemaciclib, Palbociclib, Ribociclib | sensitive |
| <i>FLCN</i> | Everolimus, Temsirolimus | sensitive |
| <i>NF2</i> | Everolimus, Temsirolimus | sensitive |
| <i>CHEK2</i> | Niraparib, Olaparib, Rucaparib, Talazoparib | sensitive |
| <i>RB1</i> | Abemaciclib, Palbociclib, Ribociclib | resistant |

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Palbociclib, Ribociclib, Abemaciclib

ACT Onco[®] + Report



1: Olaparib, Niraparib, Rucaparib, Talazoparib



1: Everolimus, Temsirolimus

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
2. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
3. PMID: 22713868; 2012, Genes Dev;26(12):1268-86
Mutant p53: one name, many proteins.
4. PMID: 15613472; 2005, J Biol Chem;280(8):6682-91
Hsp90 is essential for restoring cellular functions of temperature-sensitive p53 mutant protein but not for stabilization and activation of wild-type p53: implications for cancer therapy.
5. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months.
6. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
7. PMID: 25669829; 2015, Ann Oncol;26(5):1012-1018
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
8. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
9. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
10. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
11. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
12. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
13. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
14. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
15. PMID: 8114739; 1994, Mol Cell Biol;14(3):2077-86
Identification of G1 kinase activity for cdk6, a novel cyclin D partner.
16. PMID: 12432268; 2002, Cancer Biol Ther;1(3):226-31
Cycling to cancer with cyclin D1.
17. PMID: 24795392; 2014, Clin Cancer Res;20(13):3379-83
Molecular pathways: CDK4 inhibitors for cancer therapy.
18. PMID: 27030077; 2016, Nat Rev Clin Oncol;13(7):417-30
Treating cancer with selective CDK4/6 inhibitors.

ACT Onco[®] + Report

19. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
20. PMID: 27334220; 2016, Ann Surg Oncol;23(9):2745-52
Targeting Cyclin-Dependent Kinases in Synovial Sarcoma: Palbociclib as a Potential Treatment for Synovial Sarcoma Patients.
21. PMID: 19874578; 2009, Breast Cancer Res;11(5):R77
PD 0332991, a selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-positive human breast cancer cell lines in vitro.
22. PMID: 23898052; 2013, Anticancer Res;33(8):2997-3004
PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity.
23. PMID: 22475046; 2012, Breast Cancer Res;14(2):R57
Effects of cyclin D1 gene amplification and protein expression on time to recurrence in postmenopausal breast cancer patients treated with anastrozole or tamoxifen: a TransATAC study.
24. PMID: 17070615; 2007, Lung Cancer;55(1):1-14
Cyclin D1 in non-small cell lung cancer: a key driver of malignant transformation.
25. PMID: 26307133; 2016, Clin Cancer Res;22(2):374-82
Copy Number Changes Are Associated with Response to Treatment with Carboplatin, Paclitaxel, and Sorafenib in Melanoma.
26. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
27. PMID: 11495045; 2001, J Hepatol;35(1):74-9
The role of overexpression and gene amplification of cyclin D1 in intrahepatic cholangiocarcinoma.
28. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
29. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
30. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
31. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.
32. PMID: 7550353; 1995, Nat Genet;11(2):210-2
Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.
33. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
34. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
35. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
36. PMID: 27217383; 2016, Cancer Discov;6(7):740-53
Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.

ACT Onco[®] + Report

37. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
38. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
39. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.
40. PMID: 35050752; 2020, JCO Precis Oncol;4():757-766
Palbociclib in Patients With Non-Small-Cell Lung Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
41. PMID: 35100714; 2019, JCO Precis Oncol;3():1-8
Palbociclib in Patients With Pancreatic and Biliary Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
42. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
43. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
44. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
45. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
46. PMID: 21088254; 2011, Clin Cancer Res;17(3):401-5
Tumor suppressor CHK2: regulator of DNA damage response and mediator of chromosomal stability.
47. PMID: 15261141; 2004, Cancer Cell;6(1):45-59
Chk1 is haploinsufficient for multiple functions critical to tumor suppression.
48. PMID: 15539958; 2005, Cell Cycle;4(1):131-9
Chk1 is essential for tumor cell viability following activation of the replication checkpoint.
49. PMID: 23296741; 2013, Fam Cancer;12(3):473-8
The risk of gastric cancer in carriers of CHEK2 mutations.
50. PMID: 24713400; 2014, Hered Cancer Clin Pract;12(1):10
A risk of breast cancer in women - carriers of constitutional CHEK2 gene mutations, originating from the North - Central Poland.
51. PMID: 25583358; 2015, Int J Cancer;137(3):548-52
CHEK2 mutations and the risk of papillary thyroid cancer.
52. PMID: 12052256; 2002, Breast Cancer Res;4(3):R4
Mutation analysis of the CHK2 gene in breast carcinoma and other cancers.
53. PMID: 15125777; 2004, Mol Cancer;3():14
CHK2 kinase expression is down-regulated due to promoter methylation in non-small cell lung cancer.
54. PMID: 33119476; 2020, J Clin Oncol;38(36):4274-4282
TBCRC 048: Phase II Study of Olaparib for Metastatic Breast Cancer and Mutations in Homologous Recombination-Related Genes.

ACT Onco[®] + Report

55. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
56. PMID: 24095279; 2013, Mol Cell;52(4):495-505
The folliculin tumor suppressor is a GAP for the RagC/D GTPases that signal amino acid levels to mTORC1.
57. PMID: 26342594; 2016, Fam Cancer;15(1):127-32
Birt-Hogg-Dubé syndrome: novel FLCN frameshift deletion in daughter and father with renal cell carcinomas.
58. PMID: 23223565; 2013, J Clin Pathol;66(3):178-86
Birt-Hogg-Dube syndrome: clinicopathological features of the lung.
59. PMID: 19850877; 2009, Proc Natl Acad Sci U S A;106(44):18722-7
Homozygous loss of BHD causes early embryonic lethality and kidney tumor development with activation of mTORC1 and mTORC2.
60. PMID: 24908670; 2014, Hum Mol Genet;23(21):5706-19
Folliculin (Flcn) inactivation leads to murine cardiac hypertrophy through mTORC1 deregulation.
61. PMID: 15956655; 2005, J Natl Cancer Inst;97(12):931-5
High frequency of somatic frameshift BHD gene mutations in Birt-Hogg-Dubé-associated renal tumors.
62. PMID: 29301825; 2018, Clin Cancer Res;24(7):1546-1553
Genomic Correlates of Response to Everolimus in Aggressive Radioiodine-refractory Thyroid Cancer: A Phase II Study.
63. PMID: 26418749; 2015, Oncotarget;6(32):32761-73
Flcn-deficient renal cells are tumorigenic and sensitive to mTOR suppression.
64. PMID: 25893302; 2016, Oncogene;35(5):537-48
Role of Merlin/NF2 inactivation in tumor biology.
65. PMID: 19451229; 2009, Mol Cell Biol;29(15):4235-49
Loss of the tumor suppressor gene NF2, encoding merlin, constitutively activates integrin-dependent mTORC1 signaling.
66. PMID: 19451225; 2009, Mol Cell Biol;29(15):4250-61
NF2/merlin is a novel negative regulator of mTOR complex 1, and activation of mTORC1 is associated with meningioma and schwannoma growth.
67. PMID: 17655741; 2007, Brain Pathol;17(4):371-6
Role of NF2 haploinsufficiency in NF2-associated polyneuropathy.
68. PMID: 19545378; 2009, Orphanet J Rare Dis;4():16
Neurofibromatosis type 2 (NF2): a clinical and molecular review.
69. PMID: 21642991; 2011, Nat Genet;43(7):668-72
The nuclear deubiquitinase BAP1 is commonly inactivated by somatic mutations and 3p21.1 losses in malignant pleural mesothelioma.
70. PMID: 24393766; 2014, Oncotarget;5(1):67-77
NF2/merlin in hereditary neurofibromatosis 2 versus cancer: biologic mechanisms and clinical associations.
71. PMID: 27091708; 2016, J Clin Oncol;34(18):2115-24
Molecular Alterations and Everolimus Efficacy in Human Epidermal Growth Factor Receptor 2-Overexpressing Metastatic Breast Cancers: Combined Exploratory Biomarker Analysis From BOLERO-1 and BOLERO-3.
72. PMID: 26503204; 2016, J Clin Oncol;34(5):419-26
Correlative Analysis of Genetic Alterations and Everolimus Benefit in Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: Results From BOLERO-2.
73. PMID: 24833916; 2014, Breast Cancer (Dove Med Press);6():43-57
Use of mTOR inhibitors in the treatment of breast cancer: an evaluation of factors that influence patient outcomes.

ACT Onco[®] + Report

74. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
75. PMID: 22923433; 2012, Science;338(6104):221
Genome sequencing identifies a basis for everolimus sensitivity.
76. PMID: 25630452; 2015, Eur Urol;67(6):1195-1196
Exceptional Response on Addition of Everolimus to Taxane in Urothelial Carcinoma Bearing an NF2 Mutation.
77. PMID: 26359368; 2015, Cancer Discov;5(11):1178-93
NF2 Loss Promotes Oncogenic RAS-Induced Thyroid Cancers via YAP-Dependent Transactivation of RAS Proteins and Sensitizes Them to MEK Inhibition.
78. PMID: 24813888; 2014, Cell Rep;7(4):999-1008
Acquired resistance of EGFR-mutant lung adenocarcinomas to afatinib plus cetuximab is associated with activation of mTORC1.
79. PMID: 22189424; 2011, Nat Rev Mol Cell Biol;13(1):39-51
Regulating the regulator: post-translational modification of RAS.
80. PMID: 21993244; 2011, Nat Rev Cancer;11(11):761-74
RAS oncogenes: weaving a tumorigenic web.
81. PMID: 22589270; 2012, Cancer Res;72(10):2457-67
A comprehensive survey of Ras mutations in cancer.
82. PMID: 24651010; 2014, Cancer Cell;25(3):272-81
Dragging ras back in the ring.
83. PMID: 14633661; 2004, Carcinogenesis;25(4):535-9
Complex effects of Ras proto-oncogenes in tumorigenesis.
84. PMID: 22805292; 2012, Lancet Oncol;13(8):782-9
Activity of the oral MEK inhibitor trametinib in patients with advanced melanoma: a phase 1 dose-escalation trial.
85. PMID: 28152546; 2017, Br J Cancer;116(5):575-583
A phase 1 dose-escalation and expansion study of binimetinib (MEK162), a potent and selective oral MEK1/2 inhibitor.
86. PMID: 28514312; 2017, Medicine (Baltimore);96(20):e6931
Treatment inferred from mutations identified using massive parallel sequencing leads to clinical benefit in some heavily pretreated cancer patients.
87. PMID: 33637626; 2021, Clin Cancer Res;27(11):2996-3004
Differential Outcomes in Codon 12/13 and Codon 61 NRAS-Mutated Cancers in the Phase II NCI-MATCH Trial of Binimetinib in Patients with NRAS-Mutated Tumors.
88. PMID: 2734037; 1989, Pediatr Nurs;15(3):226
The darker side of competition.
89. PMID: 22392911; 2012, Clin Cancer Res;18(9):2515-25
Inhibition of MEK and PI3K/mTOR suppresses tumor growth but does not cause tumor regression in patient-derived xenografts of RAS-mutant colorectal carcinomas.
90. PMID: 23515407; 2013, Clin Cancer Res;19(9):2584-91
Characteristics of lung cancers harboring NRAS mutations.
91. PMID: 26343583; 2015, Cancer Cell;28(3):384-98
Inhibition of RAF Isoforms and Active Dimers by LY3009120 Leads to Anti-tumor Activities in RAS or BRAF Mutant Cancers.
92. PMID: 25110411; 2014, World J Gastroenterol;20(29):9732-43

ACT Onco[®] + Report

Beyond KRAS: Predictive factors of the efficacy of anti-EGFR monoclonal antibodies in the treatment of metastatic colorectal cancer.

93. PMID: 24024839; 2013, N Engl J Med;369(11):1023-34
Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer.
94. PMID: 22983396; 2012, Nat Med;18(10):1503-10
Oncogenic NRAS signaling differentially regulates survival and proliferation in melanoma.
95. PMID: 29101736; 2019, Pathol Oncol Res;25(1):269-278
Clinical Implications of NRAS Overexpression in Resectable Pancreatic Adenocarcinoma Patients.
96. PMID: 28827320; 2017, Proc Natl Acad Sci U S A;114(36):9629-9634
Identification of NRAS isoform 2 overexpression as a mechanism facilitating BRAF inhibitor resistance in malignant melanoma.
97. PMID: 22293180; 2012, J Clin Invest;122(2):425-34
Understanding pRb: toward the necessary development of targeted treatments for retinoblastoma.
98. PMID: 6320372; 1984, Science;223(4640):1028-33
Retinoblastoma: clues to human oncogenesis.
99. PMID: 27308386; 2015, Mol Cell Oncol;2(1):e968069
Conditional haploinsufficiency of the retinoblastoma tumor suppressor gene.
100. PMID: 23687339; 2013, Cancer Res;73(14):4247-55
Rb1 haploinsufficiency promotes telomere attrition and radiation-induced genomic instability.
101. PMID: 28169375; 2017, Sci Rep;7():42056
The Rb1 tumour suppressor gene modifies telomeric chromatin architecture by regulating TERRA expression.
102. PMID: 15884040; 2005, Hum Mutat;25(6):566-74
Sensitive multistep clinical molecular screening of 180 unrelated individuals with retinoblastoma detects 36 novel mutations in the RB1 gene.
103. PMID: 26238431; 2015, Eur Urol;68(6):959-67
Defects in DNA Repair Genes Predict Response to Neoadjuvant Cisplatin-based Chemotherapy in Muscle-invasive Bladder Cancer.
104. PMID: 22811582; 2012, Clin Cancer Res;18(18):5110-22
RB-pathway disruption is associated with improved response to neoadjuvant chemotherapy in breast cancer.
105. PMID: 21358261; 2011, Cell Cycle;10(6):956-62
A non-functional retinoblastoma tumor suppressor (RB) pathway in premenopausal breast cancer is associated with resistance to tamoxifen.
106. PMID: 17160137; 2007, J Clin Invest;117(1):218-28
The retinoblastoma tumor suppressor modifies the therapeutic response of breast cancer.
107. PMID: 29236940; 2018, Ann Oncol;29(3):640-645
Polyclonal RB1 mutations and acquired resistance to CDK 4/6 inhibitors in patients with metastatic breast cancer.
108. PMID: 29483214; 2018, Mol Cancer Ther;17(5):897-907
Preclinical Activity of Abemaciclib Alone or in Combination with Antimitotic and Targeted Therapies in Breast Cancer.
109. PMID: 22941188; 2012, Nat Genet;44(10):1104-10
Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer.
110. PMID: 22941189; 2012, Nat Genet;44(10):1111-6
Comprehensive genomic analysis identifies SOX2 as a frequently amplified gene in small-cell lung cancer.
111. PMID: 25846096; 2015, Lancet Oncol;16(4):e165-72
Transformation from non-small-cell lung cancer to small-cell lung cancer: molecular drivers and cells of origin.
112. PMID: 11502877; 2001, Mol Pharmacol;60(3):474-9

ACT Onco[®] + Report

Inhibition of thymidylate synthase activity by antisense oligodeoxynucleotide and possible role in thymineless treatment.

113. PMID: 10482907; 1999, Br J Pharmacol;127(8):1777-86
Antisense down-regulation of thymidylate synthase to suppress growth and enhance cytotoxicity of 5-FUdR, 5-FU and Tomudex in HeLa cells.
114. PMID: 16818500; 2006, Mol Cancer Ther;5(6):1423-33
Therapeutic potential of antisense oligodeoxynucleotides to down-regulate thymidylate synthase in mesothelioma.
115. PMID: 29500934; 2018, J Clin Pharm Ther;43(4):507-512
Genotype and allele frequencies of TYMS rs2790 A > G polymorphism in a Chinese paediatric population with acute lymphoblastic leukaemia.
116. PMID: 28899623; 2018, Clin Breast Cancer;18(3):e301-e304
TYMS Gene Polymorphisms in Breast Cancer Patients Receiving 5-Fluorouracil-Based Chemotherapy.
117. PMID: 26220094; 2016, Clin Transl Oncol;18(1):107-12
Thymidylate synthase gene amplification predicts pemetrexed resistance in patients with advanced non-small cell lung cancer.
118. PMID: 21487406; 2011, Br J Cancer;104(10):1594-601
Thymidylate synthase as a determinant of pemetrexed sensitivity in non-small cell lung cancer.
119. PMID: 26502926; 2015, BMC Pulm Med;15():132
Thymidylate synthase expression as a predictive biomarker of pemetrexed sensitivity in advanced non-small cell lung cancer.
120. PMID: 23242435; 2013, J Thorac Oncol;8(1):19-30
Significance of folate receptor alpha and thymidylate synthase protein expression in patients with non-small-cell lung cancer treated with pemetrexed.
121. PMID: 20727737; 2010, Eur J Cancer;46(18):3358-64
Amplification of thymidylate synthetase in metastatic colorectal cancer patients pretreated with 5-fluorouracil-based chemotherapy.
122. PMID: 28968163; 2017, J Clin Oncol;35(32):3638-3646
MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer.
123. PMID: 28533223; 2017, Clin Cancer Res;23(17):5218-5224
MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR(+)/HER2(-) Metastatic Breast Cancer.
124. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
125. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
126. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
127. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
128. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
129. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
130. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.

ACT Onco[®] + Report

131. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
132. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
133. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
134. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
135. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
136. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
137. PMID: 27959613; 2016, N Engl J Med;375(20):1925-1936
Palbociclib and Letrozole in Advanced Breast Cancer.
138. PMID: 26030518; 2015, N Engl J Med;373(3):209-19
Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer.
139. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
140. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
141. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsilolimus, interferon alfa, or both for advanced renal-cell carcinoma.