



## Sample Information

**Patient Name:** 謝桂香**Gender:** Female**ID No.:** F222751013**History No.:** 22558674**Age:** 57**Ordering Doctor:** DOC3064F 陳育民**Ordering REQ.:** D5EMP73**Signing in Date:** 2020/10/14**Path No.:** S109-89735**MP No.:** F20086**Assay:** Oncomine Focus Assay**Sample Type:** FFPE**Block No.:** S109-78125A**Percentage of tumor cells:** 50%**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

### Table of Contents

### Page

Variants (Exclude variant in Taiwan  
BioBank with >1% allele frequency)

2

Biomarker Descriptions

3

Relevant Therapy Summary

3

Relevant Therapy Details

11

### Report Highlights

2 Relevant Biomarkers

39 Therapies Available

50 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Findings

| Gene  | Finding   | Gene  | Finding      |
|-------|---|-------|--------------|
| ALK   | Not detected  | NTRK1 | Not detected |
| BRAF  | Not detected  | NTRK2 | Not detected |
| EGFR  | Not detected  | NTRK3 | Not detected |
| ERBB2 | <b>ERBB2 amplification, ERBB2 exon 20 insertion</b> | RET   | Not detected |
| KRAS  | Not detected  | ROS1  | Not detected |
| MET   | Not detected  |       |              |



## Relevant Biomarkers

| Tier | Genomic Alteration  | Relevant Therapies<br>(In this cancer type) | Relevant Therapies<br>(In other cancer type)  | Clinical Trials |
|------|---|---|---|-----------------|
| IIC  | <b>ERBB2 amplification</b><br>erb-b2 receptor tyrosine kinase 2                                 | None  | ado-trastuzumab emtansine <sup>1,2</sup><br>irbinitinib + trastuzumab + chemotherapy <sup>1</sup><br>lapatinib + hormone therapy <sup>1,2</sup><br>lapatinib + chemotherapy <sup>1,2</sup><br>neratinib <sup>1,2</sup><br>neratinib + chemotherapy <sup>1</sup><br>pertuzumab + trastuzumab + chemotherapy <sup>1,2</sup><br>trastuzumab* + chemotherapy <sup>1,2</sup><br>trastuzumab* <sup>1,2</sup><br>trastuzumab deruxtecan <sup>1</sup><br>abemaciclib + hormone therapy<br>hormone therapy<br>palbociclib + hormone therapy<br>ribociclib + hormone therapy<br>lapatinib + trastuzumab <sup>2</sup><br>lapatinib + trastuzumab + hormone therapy<br>pertuzumab + trastuzumab<br>pertuzumab + trastuzumab + hormone therapy + chemotherapy<br>trastuzumab* + hormone therapy <sup>2</sup><br>trastuzumab + hormone therapy + chemotherapy<br>trastuzumab containing regimen<br>pertuzumab + trastuzumab + hormone therapy | 33              |
|      | <b>ERBB2 exon 20 insertion</b><br>erb-b2 receptor tyrosine kinase 2<br>Allele Frequency: 71.56% | ado-trastuzumab emtansine                   | None  | 26              |

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

**Tier Reference:** Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

\* Includes biosimilars

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

| Gene  | Amino Acid Change | Coding            | Variant ID | Locus          | Allele Frequency | Transcript  | Variant Effect          | Coverage |
|-------|-------------------|-------------------|------------|----------------|------------------|-------------|-------------------------|----------|
| ERBB2 | p.(G776delinsVC)  | c.2326_2327insTGT | COSM12553  | chr17:37880997 | 71.56%           | NM_004448.3 | nonframeshift Insertion | 1955     |

### Copy Number Variations

| Gene  | Locus          | Copy Number |
|-------|----------------|-------------|
| ERBB2 | chr17:37868126 | 7.18        |



## Biomarker Descriptions

### ERBB2 (erb-b2 receptor tyrosine kinase 2)

**Background:** The ERBB2 gene encodes the erb-b2 receptor tyrosine kinase 2, a member of the human epidermal growth factor receptor (HER) family. Along with ERBB2/HER2, EGFR/ERBB1/HER1, ERBB3/HER3, and ERBB4/HER4 make up the HER protein family<sup>1</sup>. All ERBB/HER proteins encode transmembrane receptor tyrosine kinases. However, ERBB2/HER2 is an orphan receptor with no known ligand. ERBB2 preferentially binds other ligand bound ERBB/HER family members to form hetero-dimers resulting in the activation of ERBB2 tyrosine kinase activity and subsequent activation of the PI3K/AKT/MTOR and RAS/RAF/MAPK/ERK signaling pathways which promote cell proliferation, differentiation, and survival<sup>2</sup>. Recurrent focal amplification of the ERBB2 gene leads to increased expression in several cancer types. ERBB2 overexpression in immortalized cell lines is oncogenic and leads to ERBB2 homo-dimerization and activation without ligand binding<sup>3,4,5</sup>.

**Alterations and prevalence:** ERBB2 gene amplification occurs in 10-20% of breast, esophageal, and gastric cancers, 5-10% of bladder, cervical, pancreas, and uterine cancers, and 1-5% of colorectal, lung, and ovarian cancers<sup>6,7,8,9,10,11,12,13</sup>. Recurrent somatic activating mutations in ERBB2/HER2 occur at low frequencies (<1%) in diverse cancer types<sup>13,14,15</sup>. In breast, bladder, and colorectal cancers, the most common recurrent ERBB2 activating mutations include kinase domain mutations L755S and V777L and the extracellular domain mutation S310F. In lung cancer, the most common recurrent ERBB2 activating mutations include in-frame exon 20 insertions, particularly Y772\_A775dup.

**Potential relevance:** The discovery of ERBB2/HER2 as an important driver of breast cancer in 1987 led to the development of trastuzumab, a humanized monoclonal antibody with specificity to the extracellular domain of HER2<sup>16,17</sup>. Trastuzumab<sup>18</sup> was FDA approved for the treatment of HER2 positive breast cancer in 1998, and subsequently in HER2 positive metastatic gastric and gastroesophageal junction adenocarcinoma in 2010. Additional monoclonal antibody therapies have been approved by the FDA for HER2-positive breast cancer including pertuzumab<sup>19</sup> (2012), a humanized monoclonal antibody that inhibits HER2 dimerization, and ado-trastuzumab emtansine<sup>20</sup> (2013), a conjugate of trastuzumab and a potent antimicrotubule agent. The combination of pertuzumab, trastuzumab, and a taxane is the preferred front-line regimen for HER2-positive metastatic breast cancer<sup>21</sup>. In addition to monoclonal antibodies, the small molecule inhibitor lapatinib<sup>22</sup>, with specificity for both EGFR and ERBB2, was FDA approved (2007) for the treatment of patients with advanced HER2-positive breast cancer who have received prior therapy including trastuzumab. In 2017, the FDA approved the use of neratinib<sup>23</sup>, an irreversible kinase inhibitor of EGFR, ERBB2/HER2, and ERBB4, for the extended adjuvant treatment of adult patients with early stage HER2-positive breast cancer. In 2020, the FDA approved neratinib<sup>23</sup> in combination with capecitabine for HER2-positive advanced or metastatic patients after two or more prior HER2-directed therapies. The vaccine, nelipepimut-S<sup>24</sup>, was granted fast-track designation by the FDA (2016) in patients with low to intermediate HER2 expressing (IHC score 1+ or 2+) breast cancer. Additionally, fast-track designation was granted (2018) to the monoclonal antibody margetuximab<sup>25</sup> in patients with ERBB2 positive breast cancer previously treated with an anti-HER2 therapy as well as the novel bispecific antibody ZW25<sup>26</sup> (2019) in combination with standard chemotherapy for patients with HER2-overexpressing gastroesophageal adenocarcinoma (GEA). Certain activating mutations have been observed to impart sensitivity to neratinib, afatinib, lapatinib, and trastuzumab, or dacomitinib in early and ongoing clinical studies<sup>27,28,29,30,31</sup>. Additionally, acquired HER2 mutations in estrogen receptor-positive (ER+) breast cancer have been shown to confer resistance to hormone therapy<sup>32</sup>. However, this was shown to be overcome by neratinib in combination with therapies targeting ER<sup>32</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ Contraindicated
 ☒ Both for use and contraindicated
 ☒ No evidence

### ERBB2 amplification

| Relevant Therapy          | FDA                   | NCCN                  | EMA                   | ESMO                  | Clinical Trials*                      |
|---------------------------|-----------------------|-----------------------|-----------------------|-----------------------|---------------------------------------|
| ado-trastuzumab emtansine | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input checked="" type="radio"/> (II) |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 amplification (continued)

| Relevant Therapy                                   | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| pertuzumab + trastuzumab + chemotherapy            | ○   | ○    | ○   | ○    | ✕                |
| pertuzumab + trastuzumab + docetaxel               | ○   | ○    | ○   | ○    | ✕                |
| trastuzumab + capecitabine + cisplatin             | ○   | ○    | ○   | ○    | ✕                |
| trastuzumab + cisplatin + fluorouracil             | ○   | ○    | ○   | ○    | ✕                |
| lapatinib + capecitabine                           | ○   | ○    | ○   | ✕    | ✕                |
| trastuzumab  | ○   | ○    | ○   | ✕    | ✕                |
| trastuzumab + carboplatin + docetaxel              | ○   | ○    | ○   | ✕    | ✕                |
| trastuzumab + docetaxel                            | ○   | ○    | ○   | ✕    | ✕                |
| trastuzumab + paclitaxel                           | ○   | ○    | ○   | ✕    | ✕                |
| irbinitinib + trastuzumab + capecitabine           | ○   | ○    | ✕   | ✕    | ✕                |
| neratinib + capecitabine                           | ○   | ○    | ✕   | ✕    | ✕                |
| trastuzumab deruxtecan                             | ○   | ○    | ✕   | ✕    | ✕                |
| lapatinib + letrozole                              | ○   | ✕    | ○   | ✕    | ✕                |
| neratinib  | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon)                               | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + capecitabine + cisplatin    | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + carboplatin + docetaxel     | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + cisplatin + fluorouracil    | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + docetaxel                   | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + paclitaxel                  | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion)                            | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + capecitabine + cisplatin | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + carboplatin + docetaxel  | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + cisplatin + fluorouracil | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + docetaxel                | ○   | ✕    | ○   | ✕    | ✕                |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 amplification (continued)

| Relevant Therapy   | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| trastuzumab (Celltrion) + paclitaxel                     | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer)                                     | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + capecitabine + cisplatin          | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + carboplatin + docetaxel           | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + cisplatin + fluorouracil          | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + docetaxel                         | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + paclitaxel                        | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis)                            | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + capecitabine + cisplatin | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + carboplatin + docetaxel  | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + cisplatin + fluorouracil | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + docetaxel                | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + paclitaxel               | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon)                                    | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + capecitabine + cisplatin         | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + carboplatin + docetaxel          | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + cisplatin + fluorouracil         | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + docetaxel                        | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + paclitaxel                       | ○   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Enhance)                                    | ○   | ✕    | ✕   | ✕    | ✕                |
| trastuzumab (Enhance) + carboplatin + docetaxel          | ○   | ✕    | ✕   | ✕    | ✕                |
| trastuzumab (Enhance) + docetaxel                        | ○   | ✕    | ✕   | ✕    | ✕                |
| trastuzumab (Enhance) + paclitaxel                       | ○   | ✕    | ✕   | ✕    | ✕                |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

In this cancer type  
  In other cancer type  
  In this cancer type and other cancer types  
  Contraindicated  
  Both for use and contraindicated  
  No evidence

### ERBB2 amplification (continued)

| Relevant Therapy  | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|---|-----|------|-----|------|------------------|
| lapatinib + trastuzumab                                   | ×   | ○    | ○   | ○    | ×                |
| pertuzumab + trastuzumab                                  | ×   | ○    | ×   | ○    | ● (II)           |
| pertuzumab + trastuzumab + hormone therapy + chemotherapy | ×   | ○    | ×   | ○    | ×                |
| pertuzumab + trastuzumab + paclitaxel                     | ×   | ○    | ×   | ○    | ×                |
| tamoxifen   | ×   | ○    | ×   | ○    | ×                |
| trastuzumab + chemotherapy                                | ×   | ○    | ×   | ○    | ×                |
| trastuzumab + hormone therapy + chemotherapy              | ×   | ○    | ×   | ○    | ×                |
| trastuzumab + vinorelbine                                 | ×   | ○    | ×   | ○    | ×                |
| aromatase inhibitor                                       | ×   | ○    | ×   | ⊘    | ×                |
| abemaciclib + fulvestrant                                 | ×   | ○    | ×   | ×    | ×                |
| anastrozole   | ×   | ○    | ×   | ×    | ×                |
| anastrozole + fulvestrant                                 | ×   | ○    | ×   | ×    | ×                |
| exemestane  | ×   | ○    | ×   | ×    | ×                |
| fulvestrant   | ×   | ○    | ×   | ×    | ×                |
| fulvestrant + letrozole                                   | ×   | ○    | ×   | ×    | ×                |
| hormone therapy   | ×   | ○    | ×   | ×    | ×                |
| lapatinib + aromatase inhibitor                           | ×   | ○    | ×   | ×    | ×                |
| lapatinib + trastuzumab + aromatase inhibitor             | ×   | ○    | ×   | ×    | ×                |
| letrozole   | ×   | ○    | ×   | ×    | ×                |
| neratinib + paclitaxel                                    | ×   | ○    | ×   | ×    | ×                |
| palbociclib + fulvestrant                                 | ×   | ○    | ×   | ×    | ×                |
| pertuzumab + trastuzumab + carboplatin + docetaxel        | ×   | ○    | ×   | ×    | ×                |
| ribociclib + fulvestrant                                  | ×   | ○    | ×   | ×    | ×                |
| toremifene citrate  | ×   | ○    | ×   | ×    | ×                |
| trastuzumab + aromatase inhibitor                         | ×   | ○    | ×   | ×    | ×                |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ⓘ In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 amplification (continued)

| Relevant Therapy   | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| trastuzumab + capecitabine                               | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + capecitabine + oxaliplatin                 | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + carboplatin + docetaxel + fluorouracil     | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + carboplatin + paclitaxel                   | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + chemotherapy (other)                       | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + cisplatin + docetaxel                      | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + cisplatin + docetaxel + fluorouracil       | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + cisplatin + paclitaxel                     | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + cyclophosphamide + docetaxel               | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + docetaxel + fluorouracil + oxaliplatin     | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + fluorouracil                               | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + fluorouracil + irinotecan                  | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + fluorouracil + oxaliplatin                 | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + fulvestrant                                | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab + tamoxifen                                  | ✕   | ○    | ✕   | ✕    | ✕                |
| trastuzumab (Biocon) + anastrozole                       | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Biocon) + CMF + doxorubicin + paclitaxel    | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + anastrozole                    | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Celltrion) + CMF + doxorubicin + paclitaxel | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + anastrozole                       | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Pfizer) + CMF + doxorubicin + paclitaxel    | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Samsung Bioepis) + anastrozole              | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab (Synthon) + anastrozole                      | ✕   | ✕    | ○   | ✕    | ✕                |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 amplification (continued)

| Relevant Therapy   | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| trastuzumab (Synthon) + CMF + doxorubicin + paclitaxel     | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab + anastrozole                                  | ✕   | ✕    | ○   | ✕    | ✕                |
| trastuzumab + CMF + doxorubicin + paclitaxel               | ✕   | ✕    | ○   | ✕    | ✕                |
| aromatase inhibitor + luteinizing hormone-releasing factor | ✕   | ✕    | ✕   | ○    | ✕                |
| pertuzumab + trastuzumab + capecitabine                    | ✕   | ✕    | ✕   | ○    | ✕                |
| pertuzumab + trastuzumab + hormone therapy                 | ✕   | ✕    | ✕   | ○    | ✕                |
| pertuzumab + trastuzumab + nab-paclitaxel                  | ✕   | ✕    | ✕   | ○    | ✕                |
| pertuzumab + trastuzumab + vinorelbine                     | ✕   | ✕    | ✕   | ○    | ✕                |
| trastuzumab + hormone therapy                              | ✕   | ✕    | ✕   | ○    | ✕                |
| trastuzumab + taxane                                       | ✕   | ✕    | ✕   | ○    | ✕                |
| trastuzumab containing regimen                             | ✕   | ✕    | ✕   | ○    | ✕                |
| lapatinib + trastuzumab + chemotherapy                     | ✕   | ✕    | ✕   | ⛔    | ✕                |
| pertuzumab, trastuzumab                                    | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| targeted therapy, chemotherapy                             | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| trastuzumab, pertuzumab                                    | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| A-166  | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| BAT 1306, BAT-8001   | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| BDTX-189   | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| CART-PD1   | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| zotatifin  | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| AC-101 (AbClon)  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| ado-trastuzumab (Shanghai Fosun Pharma)                    | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| ARX-788  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| BDC-1001, pembrolizumab                                    | ✕   | ✕    | ✕   | ✕    | ● (I)            |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.





## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ⓘ In this cancer type and other cancer types    ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 amplification (continued)

| Relevant Therapy                               | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| BTRC-4017A, trastuzumab                        | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| CART-HER2                                      | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| disitamab vedotin                              | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| KN026, KN046                                   | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| M802   | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| MBS301   | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| MGD-013, margetuximab                          | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| MP-0274  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| MT-5111  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| neratinib, palbociclib, everolimus, trametinib | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| NJH-395  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| pirotinib                                      | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| PRS-343  | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| PRS-343, atezolizumab                          | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| trastuzumab deruxtecan, pembrolizumab          | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| ZW-25  | ✕   | ✕    | ✕   | ✕    | ● (I)            |

### ERBB2 exon 20 insertion

| Relevant Therapy               | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--------------------------------|-----|------|-----|------|------------------|
| ado-trastuzumab emtansine      | ✕   | ●    | ✕   | ✕    | ● (II)           |
| afatinib                       | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| anti-PD-L1 antibody, pyrotinib | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| neratinib                      | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| pertuzumab + trastuzumab       | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| pertuzumab, trastuzumab        | ✕   | ✕    | ✕   | ✕    | ● (II)           |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

● In this cancer type    ○ In other cancer type    ● In this cancer type and other cancer types    
 ⛔ Contraindicated    ⚠ Both for use and contraindicated    ✕ No evidence

### ERBB2 exon 20 insertion (continued)

| Relevant Therapy                               | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|------------------|
| poziotinib                                     | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| pyrotinib                                      | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| sintilimab                                     | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| targeted therapy, chemotherapy                 | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| tarloxotinib                                   | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| trastuzumab deruxtecan                         | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| trastuzumab, pertuzumab, chemotherapy          | ✕   | ✕    | ✕   | ✕    | ● (II)           |
| BDTX-189                                       | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| CBT-502, anlotinib hydrochloride               | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| DZD-9008                                       | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| zotatifin                                      | ✕   | ✕    | ✕   | ✕    | ● (I/II)         |
| disitamab vedotin                              | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| neratinib, palbociclib, everolimus, trametinib | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| pirotenib                                      | ✕   | ✕    | ✕   | ✕    | ● (I)            |
| trastuzumab deruxtecan, pembrolizumab          | ✕   | ✕    | ✕   | ✕    | ● (I)            |

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

FDA information is current as of 2020-05-26. For the most up-to-date information, search [www.fda.gov](http://www.fda.gov).

### ERBB2 amplification

#### ☐ ado-trastuzumab emtansine

Cancer type: Breast Cancer

Label as of: 2019-05-03

Variant class: ERBB2 overexpression or ERBB2 amplification

##### Indications and usage:

KADCYLA® is a HER2-targeted antibody and microtubule inhibitor conjugate indicated, as a single agent, for:

- the treatment of patients with HER2-positive, metastatic breast cancer who previously received trastuzumab and a taxane, separately or in combination. Patients should have either:
  - received prior therapy for metastatic disease, or
  - developed disease recurrence during or within six months of completing adjuvant therapy.
- the adjuvant treatment of patients with HER2-positive early breast cancer who have residual invasive disease after neoadjuvant taxane and trastuzumab-based treatment.

Select patients for therapy based on an FDA-approved companion diagnostic for KADCYLA®

##### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/125427s105lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/125427s105lbl.pdf)

#### ☐ irbinitinib + trastuzumab + capecitabine

Cancer type: Breast Cancer

Label as of: 2020-04-17

Variant class: ERBB2 amplification or ERBB2 overexpression

##### Indications and usage:

TUKYSA™ is a kinase inhibitor indicated in combination with trastuzumab and capecitabine for treatment of adult patients with advanced unresectable or metastatic HER2-positive breast cancer, including patients with brain metastases, who have received one or more prior anti-HER2-based regimens in the metastatic setting.

##### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/213411s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/213411s000lbl.pdf)



## ERBB2 amplification (continued)

### ○ lapatinib + capecitabine, lapatinib + letrozole

**Cancer type:** Breast Cancer

**Label as of:** 2018-12-06

**Variant class:** ERBB2 overexpression

**Other criteria:** ER positive, PR positive

#### Indications and usage:

TYKERB® is a kinase inhibitor indicated in combination with:

- capecitabine for the treatment of patients with advanced or metastatic breast cancer whose tumors overexpress human epidermal growth factor receptor 2 (HER2) and who have received prior therapy including an anthracycline, a taxane, and trastuzumab.
- Limitations of Use: Patients should have disease progression on trastuzumab prior to initiation of treatment with TYKERB® in combination with capecitabine.
- letrozole for the treatment of postmenopausal women with hormone receptor-positive metastatic breast cancer that overexpresses the HER2 receptor for whom hormonal therapy is indicated.

TYKERB® in combination with an aromatase inhibitor has not been compared to a trastuzumab-containing chemotherapy regimen for the treatment of metastatic breast cancer.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/022059s024lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/022059s024lbl.pdf)

### ○ neratinib, neratinib + capecitabine

**Cancer type:** Breast Cancer

**Label as of:** 2020-02-25

**Variant class:** ERBB2 overexpression

#### Indications and usage:

NERLYNX® is a kinase inhibitor indicated:

- As a single agent, for the extended adjuvant treatment of adult patients with early stage HER2-positive breast cancer, to follow adjuvant trastuzumab-based therapy.
- In combination with capecitabine, for the treatment of adult patients with advanced or metastatic HER2-positive breast cancer who have received two or more prior anti-HER2 based regimens in the metastatic setting.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/208051s005s006lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208051s005s006lbl.pdf)



## ERBB2 amplification (continued)

### ○ pertuzumab + trastuzumab + chemotherapy, pertuzumab + trastuzumab + docetaxel

Cancer type: Breast Cancer

Label as of: 2020-01-16

Variant class: ERBB2 amplification or ERBB2 overexpression

#### Indications and usage:

PERJETA® is a HER2/neu receptor antagonist indicated for:

- Use in combination with trastuzumab and docetaxel for treatment of patients with HER2-positive metastatic breast cancer (MBC) who have not received prior anti-HER2 therapy or chemotherapy for metastatic disease.
- Use in combination with trastuzumab and chemotherapy as
  - neoadjuvant treatment of patients with HER2-positive, locally advanced, inflammatory, or early stage breast cancer (either greater than 2 cm in diameter or node positive) as part of a complete treatment regimen for early breast cancer.
  - adjuvant treatment of patients with HER2-positive early breast cancer at high risk of recurrence

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/125409s124lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125409s124lbl.pdf)

### ○ trastuzumab (Biocon), trastuzumab (Biocon) + docetaxel, trastuzumab (Biocon) + paclitaxel, trastuzumab (Biocon) + capecitabine + cisplatin, trastuzumab (Biocon) + carboplatin + docetaxel, trastuzumab (Biocon) + cisplatin + fluorouracil

Cancer type: Breast Cancer, Gastric Cancer, Gastroesophageal Junction Adenocarcinoma

Label as of: 2019-04-17

Variant class: ERBB2 overexpression or ERBB2 amplification

#### Indications and usage:

OGIVRI™ is a HER2/neu receptor antagonist indicated for:

- The treatment of HER2-overexpressing breast cancer.
- The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761074s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761074s004lbl.pdf)



## ERBB2 amplification (continued)

- **trastuzumab (Celltrion), trastuzumab (Celltrion) + docetaxel, trastuzumab (Celltrion) + paclitaxel, trastuzumab (Celltrion) + capecitabine + cisplatin, trastuzumab (Celltrion) + carboplatin + docetaxel, trastuzumab (Celltrion) + cisplatin + fluorouracil**

**Cancer type:** Breast Cancer, Gastric Cancer, Gastroesophageal Junction Adenocarcinoma  
**Label as of:** 2019-05-16

**Variant class:** ERBB2 overexpression or ERBB2 amplification

### Indications and usage:

HERZUMA® is a HER2/neu receptor antagonist indicated for:

- the treatment of HER2-overexpressing breast cancer.
- the treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761091s001s002lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761091s001s002lbl.pdf)

- **trastuzumab (Enhance), trastuzumab (Enhance) + docetaxel, trastuzumab (Enhance) + paclitaxel, trastuzumab (Enhance) + carboplatin + docetaxel**

**Cancer type:** Breast Cancer

**Label as of:** 2019-02-28

**Variant class:** ERBB2 overexpression or ERBB2 amplification

### Indications and usage:

HERCEPTIN HYLECTA™ is a combination of trastuzumab, a HER2/neu receptor antagonist, and hyaluronidase, an endoglycosidase, indicated in adults for:

- The treatment of HER2-overexpressing breast cancer.

Select patients for therapy based on an FDA-approved companion diagnostic for trastuzumab.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761106Orig1s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761106Orig1s000lbl.pdf)



## ERBB2 amplification (continued)

- **trastuzumab (Pfizer), trastuzumab (Pfizer) + docetaxel, trastuzumab (Pfizer) + paclitaxel, trastuzumab (Pfizer) + capecitabine + cisplatin, trastuzumab (Pfizer) + carboplatin + docetaxel, trastuzumab (Pfizer) + cisplatin + fluorouracil**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-03-11  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 amplification or ERBB2 overexpression

### Indications and usage:

TRAZIMERA™ is a HER2/neu receptor antagonist indicated for:

- The treatment of HER2-overexpressing breast cancer.
- The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761081s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761081s000lbl.pdf)

- **trastuzumab (Samsung Bioepis), trastuzumab (Samsung Bioepis) + docetaxel, trastuzumab (Samsung Bioepis) + paclitaxel, trastuzumab (Samsung Bioepis) + capecitabine + cisplatin, trastuzumab (Samsung Bioepis) + carboplatin + docetaxel, trastuzumab (Samsung Bioepis) + cisplatin + fluorouracil**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-01-18  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 amplification or ERBB2 overexpression

### Indications and usage:

Ontruzant® is a HER2/neu receptor antagonist indicated for:

- The treatment of HER2-overexpressing breast cancer.
- The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761100s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761100s000lbl.pdf)



## ERBB2 amplification (continued)

- **trastuzumab (Synthon), trastuzumab (Synthon) + docetaxel, trastuzumab (Synthon) + paclitaxel, trastuzumab (Synthon) + capecitabine + cisplatin, trastuzumab (Synthon) + carboplatin + docetaxel, trastuzumab (Synthon) + cisplatin + fluorouracil**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-06-13  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression or ERBB2 amplification

### Indications and usage:

KANJINTI™ is a HER2/neu receptor antagonist indicated for:

- the treatment of HER2 overexpressing breast cancer.
- the treatment of HER2 overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761073Orig1s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761073Orig1s000lbl.pdf)

- **trastuzumab (Synthon), trastuzumab (Synthon) + paclitaxel, trastuzumab (Synthon) + capecitabine + cisplatin, trastuzumab (Synthon) + carboplatin + docetaxel, trastuzumab (Synthon) + cisplatin + fluorouracil**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-06-13  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression or ERBB2 amplification

### Indications and usage:

KANJINTI™ is a HER2/neu receptor antagonist indicated for:

- the treatment of HER2 overexpressing breast cancer.
- the treatment of HER2 overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for a trastuzumab product.

### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761073Orig1s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761073Orig1s000lbl.pdf)



**ERBB2 amplification (continued)****○ trastuzumab deruxtecan****Cancer type:** Breast Cancer**Label as of:** 2019-12-20**Variant class:** ERBB2 amplification or ERBB2 overexpression**Indications and usage:**

ENHERTU® is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received two or more prior anti-HER2-based regimens in the metastatic setting.

This indication is approved under accelerated approval based on tumor response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/761139s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/761139s000lbl.pdf)

**○ trastuzumab, trastuzumab + docetaxel, trastuzumab + paclitaxel, trastuzumab + capecitabine + cisplatin, trastuzumab + carboplatin + docetaxel, trastuzumab + cisplatin + fluorouracil****Cancer type:** Breast Cancer, Gastric Cancer, Gastroesophageal Junction Adenocarcinoma**Label as of:** 2018-11-29**Variant class:** ERBB2 overexpression or ERBB2 amplification**Indications and usage:**

HERCEPTIN® is a HER2/neu receptor antagonist indicated for:

- The treatment of HER2-overexpressing breast cancer.
- The treatment of HER2-overexpressing metastatic gastric or gastroesophageal junction adenocarcinoma.

Select patients for therapy based on an FDA-approved companion diagnostic for HERCEPTIN®.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/103792s5345lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/103792s5345lbl.pdf)



## Current NCCN Information

☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

NCCN information is current as of 2020-05-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### ERBB2 amplification

#### ☐ abemaciclib + fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

#### ☐ anastrozole + fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

#### ☐ fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ fulvestrant + letrozole

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ irbinitinib + trastuzumab + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; One or more lines of prior HER2-targeted therapy (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ irbinitinib + trastuzumab + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; One or more lines of prior HER2-targeted therapy (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ palbociclib + fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ pertuzumab + trastuzumab + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ ribociclib + fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + capecitabine + cisplatin

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ☐ trastuzumab + capecitabine + cisplatin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + chemotherapy

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER negative, PR negative

**NCCN Recommendation category:** 1

**Population segment (Line of therapy):**

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $>1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases  $>2$  mm) (Adjuvant therapy)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + chemotherapy

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 1

**Population segment (Line of therapy):**

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $>1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases  $>2$  mm) (Adjuvant therapy)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + cisplatin + fluorouracil

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 1

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + cisplatin + fluorouracil

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ ado-trastuzumab emtansine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ anastrozole

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ aromatase inhibitor

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ exemestane

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ fulvestrant

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ hormone therapy

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm or Tumor  $> 1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node Positive (1 or more ipsilateral metastases  $> 2$  mm) (Adjuvant Therapy)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ○ hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm or Tumor  $>1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node Positive (1 or more ipsilateral metastases  $>2$  mm) (Adjuvant Therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ lapatinib + aromatase inhibitor

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ lapatinib + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]





## ERBB2 amplification (continued)

### ○ lapatinib + trastuzumab

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ lapatinib + trastuzumab

Cancer type: Colon Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: BRAF wild type, RAS wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Advanced or Metastatic Colon Cancer; Not appropriate for intensive therapy (Initial Therapy)
- Advanced or Metastatic Colon Cancer; Progression after initial therapy; If no previous treatment with HER2 inhibitor (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2020]

### ○ lapatinib + trastuzumab

Cancer type: Rectal Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: BRAF wild type, RAS wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Advanced or Metastatic Rectal Cancer; Not appropriate for intensive therapy (Initial therapy)
- Advanced or Metastatic Rectal Cancer; Progression after initial therapy; If no previous treatment with HER2 inhibitor (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ lapatinib + trastuzumab + aromatase inhibitor

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ letrozole

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ neratinib + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ○ pertuzumab + trastuzumab

Cancer type: Colon Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: BRAF wild type, RAS wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Advanced or Metastatic Colon Cancer; Not appropriate for intensive therapy (Initial Therapy)
- Advanced or Metastatic Colon Cancer; Progression after initial therapy; If no previous treatment with HER2 inhibitor (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2020]

### ○ pertuzumab + trastuzumab

Cancer type: Rectal Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: BRAF wild type, RAS wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Advanced or Metastatic Rectal Cancer; Not appropriate for intensive therapy (Initial therapy)
- Advanced or Metastatic Rectal Cancer; Progression after initial therapy; If no previous treatment with HER2 inhibitor (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 4.2020]

### ○ pertuzumab + trastuzumab + carboplatin + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ pertuzumab + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases >2 mm) (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ pertuzumab + trastuzumab + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) given after doxorubicin/cyclophosphamide (Other Recommended)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ pertuzumab + trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases >2 mm) (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ pertuzumab + trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ pertuzumab + trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) given after doxorubicin/cyclophosphamide (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ tamoxifen

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ tamoxifen

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ toremifene citrate

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (First Line Therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + aromatase inhibitor

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + capecitabine

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER negative, ER positive, PR negative, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + capecitabine + oxaliplatin

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + capecitabine + oxaliplatin

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + carboplatin + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + carboplatin + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + carboplatin + paclitaxel

Cancer type: Endometrial Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage III/IV or Recurrent Uterine Serous Carcinoma (Adjuvant therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 1.2020]

### ○ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + chemotherapy (other)

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + cyclophosphamide + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) (Useful in Certain Circumstances)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]





## ERBB2 amplification (continued)

### ○ trastuzumab + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) given after doxorubicin/cyclophosphamide (Other Recommended)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + fluorouracil + oxaliplatin

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + fluorouracil + oxaliplatin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Preferred)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + fulvestrant

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm or Tumor  $> 1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases  $> 2$  mm) (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; pN0 or pN1mi ( $\leq 2$  mm axillary node metastasis); Tumor  $\leq 0.5$  cm and pN1mi or Tumor 0.6-1.0 cm or Tumor  $> 1$  cm (Adjuvant therapy)
- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; Node positive (1 or more ipsilateral metastases  $> 2$  mm) (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1 and pN0; Tumor  $\leq 0.5$  cm or Tumor 0.6-1.0 cm (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Preoperative (Adjuvant therapy) given after doxorubicin/cyclophosphamide (Preferred)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + tamoxifen

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer; Postmenopausal or Premenopausal receiving ovarian ablation or suppression (Not specified)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + vinorelbine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, ER positive, PR negative, PR positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent or Stage IV Invasive Breast Cancer (Not specified) (Other recommended regimens)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ☐ trastuzumab deruxtecan

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER negative, ER positive, PR negative, PR positive

**NCCN Recommendation category:** 2A

**Population segment (Line of therapy):**

- Recurrent or Stage IV Invasive Breast Cancer; Two or more lines of prior HER2-targeted therapy (Not specified) (Other recommended regimens)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ hormone therapy

**Cancer type:** Breast Cancer

**Variant class:** ERBB2 amplification or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; Tumor ≤0.5 cm and pN0 (Adjuvant therapy)

**Reference:** NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + capecitabine

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + capecitabine

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + carboplatin + docetaxel + fluorouracil

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + carboplatin + docetaxel + fluorouracil

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + carboplatin + paclitaxel

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + carboplatin + paclitaxel

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; Tumor ≤0.5 cm and pN0 (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + cisplatin + docetaxel

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + cisplatin + docetaxel

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + cisplatin + docetaxel + fluorouracil

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + cisplatin + docetaxel + fluorouracil

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + cisplatin + paclitaxel

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + cisplatin + paclitaxel

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + docetaxel

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy)  
(Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + docetaxel

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy)  
(Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + docetaxel + fluorouracil + oxaliplatin

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy)  
(Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + docetaxel + fluorouracil + oxaliplatin

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy)  
(Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]





## ERBB2 amplification (continued)

### ○ trastuzumab + fluorouracil

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + fluorouracil

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ trastuzumab + fluorouracil + irinotecan

**Cancer type:** Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + fluorouracil + irinotecan

**Cancer type:** Gastric Cancer

**Variant class:** ERBB2 overexpression

**NCCN Recommendation category:** 2B

**Population segment (Line of therapy):**

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

**Reference:** NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]



## ERBB2 amplification (continued)

### ☐ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; Tumor ≤0.5 cm and pN0 (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1, pT2, or pT3; Tumor ≤0.5 cm and pN0 (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ☐ trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1 and pN0; Tumor ≤0.5 cm (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER negative, PR negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Ductal/No Special Type, Lobular, Mixed, Micropapillary Invasive Breast Cancer; pT1 and pN0; Tumor  $\leq 0.5$  cm (Adjuvant therapy)

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ○ trastuzumab + paclitaxel

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### ○ trastuzumab + paclitaxel

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Unresectable Locally Advanced, Recurrent or Metastatic Adenocarcinoma; Local therapy is not indicated (First-line therapy) (Other recommended regimen)

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### ○ irbinitinib + trastuzumab + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Breast Cancer; Brain metastases; Use agents active against primary tumor (if previously treated with 1 or more anti-HER2-based regimen) (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### ☐ lapatinib + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Breast Cancer; Brain metastases; Use agents active against primary tumor (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2020]

### ☐ neratinib + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Breast Cancer; Brain metastases; Use agents active against primary tumor (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2020]

### ☐ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Breast Cancer; Leptomeningeal and spine metastases; Intra-CSF chemotherapy (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2020]

### ☐ neratinib + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Breast Cancer; Brain metastases; Use agents active against primary tumor (Not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### ○ trastuzumab

Cancer type: Head and Neck Cancer

Variant class: ERBB2 positive

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Recurrent, Unresectable, or Metastatic Salivary Gland Tumors; PS 0-3 (Useful in Certain Circumstances)

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 1.2020]

### ⊖ pertuzumab + trastuzumab + cyclophosphamide + docetaxel + doxorubicin

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided."

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ⊖ pertuzumab + trastuzumab + cyclophosphamide + doxorubicin + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided."

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### ⊖ trastuzumab + capecitabine + cisplatin + epirubicin

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"
- "The use of trastuzumab in combination with anthracyclines is not recommended"

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### – trastuzumab + capecitabine + cisplatin + epirubicin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### – trastuzumab + capecitabine + epirubicin + oxaliplatin

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"
- "The use of trastuzumab in combination with anthracyclines is not recommended"

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]

### – trastuzumab + capecitabine + epirubicin + oxaliplatin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### – trastuzumab + cisplatin + epirubicin + fluorouracil

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"
- "The use of trastuzumab in combination with anthracyclines is not recommended"

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### – trastuzumab + cisplatin + epirubicin + fluorouracil

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

### – trastuzumab + cyclophosphamide + docetaxel + doxorubicin

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided."

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### – trastuzumab + cyclophosphamide + doxorubicin + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab given in combination with an anthracycline is associated with significant cardiac toxicity. Concurrent use of trastuzumab and pertuzumab with an anthracycline should be avoided."

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]

### – trastuzumab + epirubicin + fluorouracil + oxaliplatin

Cancer type: Esophageal Cancer,  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"
- "The use of trastuzumab in combination with anthracyclines is not recommended"

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 2.2020]



## ERBB2 amplification (continued)

### — trastuzumab + epirubicin + fluorouracil + oxaliplatin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

#### Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Trastuzumab is not recommended for use with anthracyclines"

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 2.2020]

## ERBB2 exon 20 insertion

### ● ado-trastuzumab emtansine

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 mutation

NCCN Recommendation category: 2A

#### Population segment (Line of therapy):

- Non-Small Cell Lung Cancer; Emerging biomarker in metastatic disease (Not specified)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2020]

### — afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 mutation

#### Summary:

NCCN Guidelines® include the following supporting statement(s):

- "The NCCN NSCLC Panel does not recommend single-agent therapy with trastuzumab or afatinib (both for ERBB2 mutations), because response rates are lower and treatment is less effective when these agents are used for patients with ERBB2 mutations."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2020]

### — trastuzumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 mutation

#### Summary:

NCCN Guidelines® include the following supporting statement(s):

- "The NCCN NSCLC Panel does not recommend single-agent therapy with trastuzumab or afatinib (both for ERBB2 mutations), because response rates are lower and treatment is less effective when these agents are used for patients with ERBB2 mutations."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2020]





## Current EMA Information

☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

EMA information is current as of 2020-05-26. For the most up-to-date information, search [www.ema.europa.eu/ema](http://www.ema.europa.eu/ema).

### ERBB2 amplification

#### ☐ ado-trastuzumab emtansine

Cancer type: Breast Cancer

Label as of: 2020-01-20

Variant class: ERBB2 overexpression or ERBB2 amplification

Reference:

[https://www.ema.europa.eu/en/documents/product-information/kadcyla-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/kadcyla-epar-product-information_en.pdf)

#### ☐ lapatinib + capecitabine, lapatinib + letrozole, lapatinib + trastuzumab

Cancer type: Breast Cancer

Label as of: 2019-10-15

Variant class: ERBB2 overexpression or ERBB2 amplification

Other criteria: ER positive, PR positive or Hormone receptor negative

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tyverb-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tyverb-epar-product-information_en.pdf)

#### ☐ neratinib

Cancer type: Breast Cancer

Label as of: 2020-02-11

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/nerlynx-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/nerlynx-epar-product-information_en.pdf)

#### ☐ pertuzumab + trastuzumab + chemotherapy, pertuzumab + trastuzumab + docetaxel

Cancer type: Breast Cancer

Label as of: 2019-04-15

Variant class: ERBB2 amplification or ERBB2 overexpression

Reference:

[https://www.ema.europa.eu/documents/product-information/perjeta-epar-product-information\\_en.pdf](https://www.ema.europa.eu/documents/product-information/perjeta-epar-product-information_en.pdf)



## ERBB2 amplification (continued)

- **trastuzumab (Biocon), trastuzumab (Biocon) + anastrozole, trastuzumab (Biocon) + docetaxel, trastuzumab (Biocon) + paclitaxel, trastuzumab (Biocon) + capecitabine + cisplatin, trastuzumab (Biocon) + carboplatin + docetaxel, trastuzumab (Biocon) + cisplatin + fluorouracil, trastuzumab (Biocon) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2020-03-18  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/ogivri-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/ogivri-epar-product-information_en.pdf)

- **trastuzumab (Biocon), trastuzumab (Biocon) + anastrozole, trastuzumab (Biocon) + docetaxel, trastuzumab (Biocon) + paclitaxel, trastuzumab (Biocon) + carboplatin + docetaxel, trastuzumab (Biocon) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer **Label as of:** 2020-03-18

**Variant class:** ERBB2 amplification

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/ogivri-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/ogivri-epar-product-information_en.pdf)

- **trastuzumab (Celltrion), trastuzumab (Celltrion) + anastrozole, trastuzumab (Celltrion) + docetaxel, trastuzumab (Celltrion) + paclitaxel, trastuzumab (Celltrion) + capecitabine + cisplatin, trastuzumab (Celltrion) + carboplatin + docetaxel, trastuzumab (Celltrion) + cisplatin + fluorouracil, trastuzumab (Celltrion) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2020-02-14  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/herzuma-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/herzuma-epar-product-information_en.pdf)

- **trastuzumab (Celltrion), trastuzumab (Celltrion) + anastrozole, trastuzumab (Celltrion) + docetaxel, trastuzumab (Celltrion) + paclitaxel, trastuzumab (Celltrion) + carboplatin + docetaxel, trastuzumab (Celltrion) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer **Label as of:** 2020-02-14

**Variant class:** ERBB2 amplification

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/herzuma-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/herzuma-epar-product-information_en.pdf)



## ERBB2 amplification (continued)

- trastuzumab (Pfizer), trastuzumab (Pfizer) + anastrozole, trastuzumab (Pfizer) + docetaxel, trastuzumab (Pfizer) + paclitaxel, trastuzumab (Pfizer) + capecitabine + cisplatin, trastuzumab (Pfizer) + carboplatin + docetaxel, trastuzumab (Pfizer) + cisplatin + fluorouracil, trastuzumab (Pfizer) + CMF + doxorubicin + paclitaxel

Cancer type: Breast Cancer, Gastric Cancer, Label as of: 2020-03-24  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/trazimera-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/trazimera-epar-product-information_en.pdf)

- trastuzumab (Pfizer), trastuzumab (Pfizer) + anastrozole, trastuzumab (Pfizer) + docetaxel, trastuzumab (Pfizer) + paclitaxel, trastuzumab (Pfizer) + carboplatin + docetaxel, trastuzumab (Pfizer) + CMF + doxorubicin + paclitaxel

Cancer type: Breast Cancer Label as of: 2020-03-24

Variant class: ERBB2 amplification

Other criteria: ER positive, PR positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/trazimera-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/trazimera-epar-product-information_en.pdf)

- trastuzumab (Samsung Bioepis), trastuzumab (Samsung Bioepis) + anastrozole, trastuzumab (Samsung Bioepis) + docetaxel, trastuzumab (Samsung Bioepis) + paclitaxel, trastuzumab (Samsung Bioepis) + capecitabine + cisplatin, trastuzumab (Samsung Bioepis) + carboplatin + docetaxel, trastuzumab (Samsung Bioepis) + cisplatin + fluorouracil

Cancer type: Breast Cancer, Gastric Cancer, Label as of: 2020-05-19  
Gastroesophageal Junction Adenocarcinoma

Variant class: ERBB2 overexpression

Other criteria: ER positive, PR positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/ontruzant-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/ontruzant-epar-product-information_en.pdf)

- trastuzumab (Samsung Bioepis), trastuzumab (Samsung Bioepis) + anastrozole, trastuzumab (Samsung Bioepis) + docetaxel, trastuzumab (Samsung Bioepis) + paclitaxel, trastuzumab (Samsung Bioepis) + carboplatin + docetaxel

Cancer type: Breast Cancer Label as of: 2020-05-19

Variant class: ERBB2 amplification

Other criteria: ER positive, PR positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/ontruzant-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/ontruzant-epar-product-information_en.pdf)



## ERBB2 amplification (continued)

- **trastuzumab (Synthon), trastuzumab (Synthon) + anastrozole, trastuzumab (Synthon) + docetaxel, trastuzumab (Synthon) + paclitaxel, trastuzumab (Synthon) + capecitabine + cisplatin, trastuzumab (Synthon) + carboplatin + docetaxel, trastuzumab (Synthon) + cisplatin + fluorouracil, trastuzumab (Synthon) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-11-29  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression or ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/kanjinti-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/kanjinti-epar-product-information_en.pdf)

- **trastuzumab (Synthon), trastuzumab (Synthon) + anastrozole, trastuzumab (Synthon) + docetaxel, trastuzumab (Synthon) + paclitaxel, trastuzumab (Synthon) + carboplatin + docetaxel, trastuzumab (Synthon) + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer **Label as of:** 2019-11-29

**Variant class:** ERBB2 amplification or ERBB2 amplification

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/kanjinti-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/kanjinti-epar-product-information_en.pdf)

- **trastuzumab, trastuzumab + anastrozole, trastuzumab + docetaxel, trastuzumab + paclitaxel, trastuzumab + capecitabine + cisplatin, trastuzumab + carboplatin + docetaxel, trastuzumab + cisplatin + fluorouracil, trastuzumab + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer, Gastric Cancer, **Label as of:** 2019-10-14  
Gastroesophageal Junction Adenocarcinoma

**Variant class:** ERBB2 overexpression

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information_en.pdf)

- **trastuzumab, trastuzumab + anastrozole, trastuzumab + docetaxel, trastuzumab + paclitaxel, trastuzumab + carboplatin + docetaxel, trastuzumab + CMF + doxorubicin + paclitaxel**

**Cancer type:** Breast Cancer **Label as of:** 2019-10-14

**Variant class:** ERBB2 amplification

**Other criteria:** ER positive, PR positive

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/herceptin-epar-product-information_en.pdf)



## Current ESMO Information

- ☒ In this cancer type  
 ☐ In other cancer type  
 ☒ In this cancer type and other cancer types  
 ☒ Contraindicated  
 ☒ Not recommended  
 ☒ Resistance

ESMO information is current as of 2020-05-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### ERBB2 amplification

#### ☐ ado-trastuzumab emtansine

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Residual Invasive Early Breast Cancer; No pathological complete response after completion of neoadjuvant chemotherapy combined with anti-HER2 therapy; substitute for adjuvant trastuzumab (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

#### ☐ pertuzumab + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Non-luminal Early Breast Cancer (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

#### ☐ pertuzumab + trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]



## ERBB2 amplification (continued)

### ☐ trastuzumab + capecitabine + cisplatin

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Inoperable or Metastatic Gastric Adenocarcinoma (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Gastric Cancer [Ann Oncol (2016) 27 (suppl 5): v38-v49. (eUpdate: 6 May 2019, 4 November 2019)]

### ☐ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Non-luminal Early Breast Cancer (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ☐ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER negative, PR negative

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- ERBB2(+) Non-Luminal Cancer; Except very low risk, such as T1aN0 (Neoadjuvant therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Primary Breast Cancer [Ann Oncol (2015) 26 (suppl 5): v8-v30.]

### ☐ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Primary Breast Cancer (Neoadjuvant therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Primary Breast Cancer [Ann Oncol (2015) 26 (suppl 5): v8-v30.]



## ERBB2 amplification (continued)

### ○ trastuzumab + cisplatin + fluorouracil

Cancer type: Gastric Cancer

Variant class: ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Inoperable or Metastatic Gastric Adenocarcinoma (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Gastric Cancer [Ann Oncol (2016) 27 (suppl 5): v38-v49. (eUpdate: 6 May 2019, 4 November 2019)]

### ○ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ trastuzumab + hormone therapy + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Luminal B ERBB2-positive Breast Cancer; Except low-risk T1a (Neoadjuvant therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Primary Breast Cancer [Ann Oncol (2015) 26 (suppl 5): v8-v30.]

### ○ pertuzumab + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Early Breast Cancer; Higher-risk cases (Primary therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]



## ERBB2 amplification (continued)

### ○ trastuzumab containing regimen

Cancer type: Esophageal Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Metastatic Esophageal Adenocarcinoma (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Oesophageal Cancer [Ann Oncol (2016) 27 (suppl 5): v50-v57.]

### ○ pertuzumab + trastuzumab + hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer; Low-risk; T1abN0 (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ trastuzumab + hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer; Low-risk; T1abN0 (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ tamoxifen

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: IV / A

Population segment (Line of therapy):

- Luminal-like Ductal Invasive Early Male Breast Cancer (Adjuvant therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]





## ERBB2 amplification (continued)

### ○ aromatase inhibitor + luteinizing hormone-releasing factor

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Luminal-like Ductal Invasive Early Breast Cancer; If a strong contraindication exists for the use of tamoxifen (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ trastuzumab + hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer; Small, node-negative tumours; If contraindication for chemotherapy (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ trastuzumab + hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 overexpression

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Luminal B-like Early Breast Cancer; Small, node-negative tumours; If contraindication to chemotherapy (Not Specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ○ trastuzumab + hormone therapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Luminal B ERBB2-positive; If contraindication or refusal of chemotherapy (Neoadjuvant therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Primary Breast Cancer [Ann Oncol (2015) 26 (suppl 5): v8-v30.]



## ERBB2 amplification (continued)

### ○ ado-trastuzumab emtansine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer; Progression after one line of trastuzumab-based therapy (Second-line therapy) (Preferred)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ○ pertuzumab + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer; Previously untreated with anti-HER2 therapy (First-line therapy)
- Advanced Breast Cancer; Previously treated (in the (neo)adjuvant setting) with anti-HER2 therapy (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ○ pertuzumab + trastuzumab + docetaxel

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ○ trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer; Previously treated in the adjuvant setting (First-line therapy)
- Advanced Breast Cancer; Untreated with trastuzumab (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]



## ERBB2 amplification (continued)

### ☐ trastuzumab + taxane

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer; Pertuzumab is not given (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ trastuzumab + vinorelbine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced Breast Cancer; Pertuzumab is not given (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ lapatinib + trastuzumab

Cancer type: Breast Cancer

Variant class: ERBB2 positive

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced Breast Cancer; First-line therapy was endocrine therapy and anti-HER2 therapy (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ lapatinib + trastuzumab

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced Breast Cancer; Progression on trastuzumab-based therapy (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]



## ERBB2 amplification (continued)

### ☐ pertuzumab + trastuzumab

Cancer type: Breast Cancer

Variant class: ERBB2 positive

Other criteria: ER positive

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced Breast Cancer; First-line therapy was endocrine therapy and anti-HER2 therapy (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ pertuzumab + trastuzumab + paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced Breast Cancer (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ pertuzumab + trastuzumab + capecitabine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- Advanced Breast Cancer (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ☐ pertuzumab + trastuzumab + vinorelbine

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- Advanced Breast Cancer (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]



## ERBB2 amplification (continued)

### ○ pertuzumab + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Advanced Breast Cancer; Previously untreated with the combination of chemotherapy + trastuzumab + pertuzumab (After first-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ○ pertuzumab + trastuzumab + nab-paclitaxel

Cancer type: Breast Cancer

Variant class: ERBB2 positive

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Advanced Breast Cancer (Not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-ESO-ESMO Advanced Breast Cancer [Ann Oncol (2018); 29: 1634–1657.]

### ⊗ lapatinib + trastuzumab + chemotherapy

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / E

Population segment (Line of therapy):

- Male Breast Cancer

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### ⊗ aromatase inhibitor

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

Other criteria: ER positive, PR negative, PR positive

ESMO Level of Evidence/Grade of Recommendation: IV / E

Population segment (Line of therapy):

- Male Breast Cancer

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]



## ERBB2 amplification (continued)

### — trastuzumab + anthracycline

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

ESMO Level of Evidence/Grade of Recommendation: I / D

#### Summary:

ESMO Clinical Practice Guidelines include the following supporting statement(s):

- "Trastuzumab should usually not be given concomitantly with anthracycline-based ChT [I, D]".

Reference: ESMO Clinical Practice Guidelines - ESMO-Early Breast Cancer [Ann Oncol (2019); 30: 1194-1220.]

### — pertuzumab + trastuzumab

Cancer type: Breast Cancer

Variant class: ERBB2 amplification or ERBB2 overexpression

#### Summary:

ESMO Clinical Practice Guidelines include the following supporting statement:

- "The role of dual HER2 blockade (including a combination of trastuzumab and pertuzumab) is not well proven and such treatment is not recommended for routine use, although it may be discussed on a case-by-case basis."

Reference: ESMO Clinical Practice Guidelines - ESMO-Primary Breast Cancer [Ann Oncol (2015) 26 (suppl 5): v8-v30.]



## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:



## References

1. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985 Sep 6;229(4717):974-6. PMID: 2992089
2. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
3. Di et al. erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science*. 1987 Jul 10;237(4811):178-82. PMID: 2885917
4. Hudziak et al. Increased expression of the putative growth factor receptor p185HER2 causes transformation and tumorigenesis of NIH 3T3 cells. *Proc. Natl. Acad. Sci. U.S.A.* 1987 Oct;84(20):7159-63. PMID: 2890160
5. Lonardo et al. The normal erbB-2 product is an atypical receptor-like tyrosine kinase with constitutive activity in the absence of ligand. *New Biol.* 1990 Nov;2(11):992-1003. PMID: 1983208
6. Ciriello et al. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell*. 2015 Oct 8;163(2):506-19. PMID: 26451490
7. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature*. 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
8. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*. 2014 Mar 20;507(7492):315-22. doi: 10.1038/nature12965. Epub 2014 Jan 29. PMID: 24476821
9. Cancer Genome Atlas Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature*. 2012 Jul 18;487(7407):330-7. PMID: 22810696
10. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
11. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011 Jun 29;474(7353):609-15. PMID: 21720365
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
13. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
14. Petrelli et al. Clinical and pathological characterization of HER2 mutations in human breast cancer: a systematic review of the literature. *Breast Cancer Res. Treat.* 2017 Nov;166(2):339-349. PMID: 28762010
15. Bose et al. Activating HER2 mutations in HER2 gene amplification negative breast cancer. *Cancer Discov.* 2013 Feb;3(2):224-37. doi: 10.1158/2159-8290.CD-12-0349. Epub 2012 Dec 7. PMID: 23220880
16. Hudis. Trastuzumab--mechanism of action and use in clinical practice. *N. Engl. J. Med.* 2007 Jul 5;357(1):39-51. PMID: 17611206
17. Slamon et al. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science*. 1987 Jan 9;235(4785):177-82. PMID: 3798106
18. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/103792s5345lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/103792s5345lbl.pdf)
19. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/125409s124lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125409s124lbl.pdf)
20. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/125427s105lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/125427s105lbl.pdf)
21. NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2020]
22. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/022059s024lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/022059s024lbl.pdf)
23. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/208051s005s006lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208051s005s006lbl.pdf)
24. <https://www.globenewswire.com/news-release/2016/06/01/845166/0/en/Galena-Biopharma-Receives-Fast-Track-Designation-for-NeuVax-nelipepimut-S-PRESENT-Clinical-Trial.html>
25. <http://ir.macrogenics.com/news-releases/news-release-details/macrogenics-announces-continuation-sophia-study-margetuximab?ReleaseID=1055055>
26. <https://ir.zymeworks.com/News-Releases/news-details/2019/Zymeworks-Lead-Asset-ZW25-Granted-Fast-Track-Designation-from-the-FDA/default.aspx>





## References (continued)

27. Ma et al. Neratinib Efficacy and Circulating Tumor DNA Detection of HER2 Mutations in HER2 Nonamplified Metastatic Breast Cancer. *Clin. Cancer Res.* 2017 Oct 1;23(19):5687-5695. PMID: 28679771
28. De et al. Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu. *Lung Cancer.* 2012 Apr;76(1):123-7. PMID: 22325357
29. Kris et al. Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors. *Ann. Oncol.* 2015 Jul;26(7):1421-7. PMID: 25899785
30. Falchook et al. Non-small-cell lung cancer with HER2 exon 20 mutation: regression with dual HER2 inhibition and anti-VEGF combination treatment. *J Thorac Oncol.* 2013 Feb;8(2):e19-20. PMID: 23328556
31. David et al. Neratinib in HER2- or HER3-mutant solid tumors: SUMMIT, a global, multi-histology, open-label, phase 2 'basket' study. *AACR 2017. Abstract CT001*
32. Nayar et al. Acquired HER2 mutations in ER+ metastatic breast cancer confer resistance to estrogen receptor-directed therapies. *Nat. Genet.* 2019 Feb;51(2):207-216. PMID: 30531871