



Sample Information

Patient Name: 黃明娟  
Gender: Female  
ID No.: A222663308  
History No.: 49467756  
Age: 58

Ordering Doctor: DOC4205A 柯博仲  
Ordering REQ.: 0CKHGHU  
Signing in Date: 2023/05/05

Path No.: M112-00095  
MP No.: MY23025  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/05/01

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	13 Therapies Available
Relevant Therapy Summary	3	0 Clinical Trials
Relevant Therapy Details	4	
Prognostic Details	15	
Alert Details	16	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	None detected	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	FLT3 ITD mutation	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>FLT3 ITD mutation</b> fms related receptor tyrosine kinase 3 Allele Frequency: 5.08%	<b>gilteritinib</b> <sup>1,2</sup> <b>midostaurin + chemotherapy</b> <sup>1,2</sup> azacitidine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine sorafenib sorafenib + chemotherapy venetoclax + chemotherapy	None	0
<b>Prognostic significance:</b> ELN 2017: Adverse				

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

### Prevalent cancer biomarkers without relevant evidence based on included data sources

DNMT3A p.(R882S) c.2644C>A

### Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

#### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
DNMT3A	p.(R882S)	c.2644C>A	COSM87001	chr2:25457243	24.56%	NM_022552.4	missense	1999
FLT3	p.(D600_L601insPSD NEYFYVDFREYEYD)	c.1800_1801insCCCT . CAGATAATGAGTAC TTCTACGTTGATTT CAGAGAATATGAAT ATGAT		chr13:28608255	5.08%	NM_004119.3	nonframeshift Insertion	1987

## Biomarker Descriptions

### DNMT3A (DNA methyltransferase 3 alpha)

**Background:** The DNMT3A gene encodes the DNA methyltransferase 3 alpha which functions as a de novo methyltransferase (DNMT) with equal methylation efficiency for unmethylated and hemimethylated DNA<sup>1</sup>. Methylation of DNA occurs at CpG islands, a region of DNA consisting of sequential cytosine/guanine dinucleotide pairs. CpG island methylation plays an important role in development as well as stem cell regulation. Alterations to global DNA methylation patterns are dependent on DNMTs, which are associated with cancer initiation and progression<sup>2,3</sup>.

**Alterations and prevalence:** DNMT3A mutations are observed in approximately 25% of all acute myeloid leukemia (AML) including 29-34% of AML with normal karyotype (NK-AML)<sup>4,5,6,7,8,9,10</sup>. Mutations in DNMT3A are also reported in 12-18% of myelodysplastic syndromes (MDS) as well as 4-6% of melanoma, lung adenocarcinoma, and uterine cancer<sup>9,11</sup>. The majority of mutations in DNMT3A are missense however, frameshift, nonsense, and splice site mutations have also been reported<sup>4,9</sup>. Missense mutations at R882 are most prevalent and are observed to coexist with NPM1 and FLT3 mutations<sup>12,13</sup>. The R882 mutations occur at the dimer/tetramer interface within the catalytic domain, which leads to disruption of DNMT3A tetramerization and loss of CpG methylation<sup>14,15</sup>. However,

## Biomarker Descriptions (continued)

DNMT3A mutations observed in AML at positions other than R882 also contribute to pathogenesis by mechanisms that do not involve methyltransferase activity<sup>16</sup>.

Potential relevance: DNMT3A mutations confer shorter overall survival (OS) in patients with AML including those with NK-AML<sup>4,7,8,13</sup>. DNMT3A mutations are a useful in the diagnosis of angioimmunoblastic T-cell lymphoma (AITCL) when trying to differentiate from other peripheral T-cell lymphomas (PTCL)<sup>17</sup>.

### FLT3 (fms related receptor tyrosine kinase 3)

Background: The FLT3 gene encodes the fms related tyrosine kinase 3, a tyrosine kinase receptor that is a member of the class III receptor tyrosine kinase family that also includes PDGFR, FMS, and KIT<sup>18</sup>. FLT3 is highly expressed in hematopoietic progenitor cells<sup>19</sup>. Genomic alterations in FLT3 activate downstream oncogenic pathways including PI3K/AKT/mTOR and RAS/RAF/MEK/ERK pathways which promote cellular proliferation, survival, and inhibition of differentiation<sup>18</sup>.

Alterations and prevalence: Somatic mutations occur in approximately 30% of acute myeloid leukemia (AML), 7-10% of melanoma, and up to 8% of uterine cancer<sup>9,20,21,22</sup>. The most common activating FLT3 mutations are internal tandem duplications (ITD) that range from 3 to 400 base pairs in length within exons 14 and 15 in the juxtamembrane (JM) domain<sup>23</sup>. The second most frequent mutations are point mutations in exon 20 within the tyrosine kinase domain (TKD)<sup>24</sup>. FLT3 is amplified in up to 8% of colorectal cancer, 3% of stomach cancer, and is commonly overexpressed in AML<sup>9,22,25</sup>.

Potential relevance: The presence of FLT3-ITD confers poor prognosis in myelodysplastic syndrome (MDS)<sup>11</sup>. Concurrent expression of FLT-ITD with mutant or wild-type NPM1 (when lacking adverse risk genetic lesions) confers intermediate risk in AML<sup>10,26</sup>. FLT3 TKD mutation at D835 confers poor prognosis in MDS<sup>11</sup>. Midostaurin<sup>27</sup> (2017) and gilteritinib<sup>28</sup> (2018) are kinase inhibitors approved for AML patients with FLT3-ITD and TKD mutations including D835 and I836 mutations. The FDA granted fast track designations in 2017 to crenolanib<sup>29</sup> and in 2022 to tuspentinib (HM43239)<sup>30</sup> for FLT3 mutation-positive relapsed or refractory AML. In 2018 the FDA granted breakthrough therapy designation to quizartinib<sup>31</sup> for AML with FLT3-ITD. A phase II trial testing crenolanib in 34 patients with FLT3-ITD and TKD mutated relapsed/refractory AML, reported that FLT3 inhibitor naïve patients demonstrated a longer overall survival (OS) and event free survival (EFS) in comparison to previously treated patients (median OS: 55 weeks vs 13 weeks; median EFS: 13 weeks vs 7 weeks)<sup>32</sup>. Another phase II trial of crenolanib with chemotherapy in newly diagnosed FLT3 mutated AML reported complete remission in 24/29 (83%) patients<sup>33</sup>. Several multi-targeted tyrosine kinase inhibitors such as sorafenib (2005), sunitinib (2006), cabozantinib (2012), and ponatinib (2012) are FDA approved and include FLT3 as a target. Sorafenib is recommended in combination with chemotherapy in FLT3-ITD mutated AML<sup>10</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### FLT3 ITD mutation

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
gilteritinib	●	●	●	●	×
midostaurin + cytarabine + daunorubicin	●	●	●	●	×
azacitidine	×	●	×	×	×
cytarabine + daunorubicin	×	●	×	×	×
cytarabine + daunorubicin + etoposide	×	●	×	×	×
cytarabine + etoposide + idarubicin	×	●	×	×	×
cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×

## Relevant Therapy Summary (continued)

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types    ☒ No evidence

### FLT3 ITD mutation (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
decitabine	×	●	×	×	×
midostaurin + cytarabine	×	●	×	×	×
sorafenib	×	●	×	×	×
sorafenib + azacitidine	×	●	×	×	×
sorafenib + decitabine	×	●	×	×	×
venetoclax + azacitidine	×	●	×	×	×
venetoclax + cytarabine	×	●	×	×	×
venetoclax + decitabine	×	●	×	×	×

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

FDA information is current as of 2023-03-15. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### FLT3 ITD mutation

#### ● gilteritinib

Cancer type: Acute Myeloid Leukemia

Label as of: 2022-01-12

Variant class: FLT3 ITD mutation

#### Indications and usage:

XOSPATA® is a kinase inhibitor indicated for the treatment of adult patients who have relapsed or refractory acute myeloid leukemia (AML) with a FLT3 mutation as detected by an FDA-approved test.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/211349s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/211349s003lbl.pdf)

## FLT3 ITD mutation (continued)

### ● midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Label as of: 2021-11-15

Variant class: FLT3 ITD mutation

#### Indications and usage:

RYDAPT® is a kinase inhibitor indicated for the treatment of adult patients with:

- Newly diagnosed acute myeloid leukemia (AML) that is FLT3 mutation-positive as detected by an FDA-approved test, in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation.  
Limitations of Use: RYDAPT® is not indicated as a single-agent induction therapy for the treatment of patients with AML.
- Aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/207997s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/207997s008lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-03-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### FLT3 ITD mutation

#### ☒ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

#### ☒ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

#### ☒ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## FLT3 ITD mutation (continued)

### ● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● gilteritinib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## FLT3 ITD mutation (continued)

### ● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● midostaurin + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]



## FLT3 ITD mutation (continued)

### ● midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● sorafenib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● sorafenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## FLT3 ITD mutation (continued)

### ● sorafenib + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## FLT3 ITD mutation (continued)

### ● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## FLT3 ITD mutation (continued)

### ● cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### ● venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

## Current EMA Information

- ☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

EMA information is current as of 2023-03-15. For the most up-to-date information, search [www.ema.europa.eu/ema](https://www.ema.europa.eu/ema).

### FLT3 ITD mutation

#### ☒ gilteritinib

Cancer type: Acute Myeloid Leukemia

Label as of: 2021-09-08

Variant class: FLT3 ITD mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/xospata-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/xospata-epar-product-information_en.pdf)

#### ☒ midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Label as of: 2022-09-23

Variant class: FLT3 ITD mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/rydapt-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/rydapt-epar-product-information_en.pdf)

## Current ESMO Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

ESMO information is current as of 2023-03-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### FLT3 ITD mutation

#### ☒ gilteritinib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

#### ☒ midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Prognostic Details

### Current NCCN Information

NCCN information is current as of 2023-03-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### FLT3 ITD mutation

#### Prognostic significance: ELN 2017: Adverse

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

#### Summary:

- FLT3-ITD<sup>high</sup>; High defined as allelic ratio ( $\geq 0.5$ )

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

### Current ESMO Information

ESMO information is current as of 2023-03-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

#### FLT3 ITD mutation

#### Prognostic significance: ELN 2017: Adverse

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

#### Summary:

- FLT3-ITD<sup>High</sup>; High defined as allelic ratio ( $\geq 0.5$ )

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated    Not recommended    Resistance    Breakthrough    Fast Track

FDA information is current as of 2023-03-15. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### FLT3 ITD mutation

#### crenolanib

**Cancer type:** Acute Myeloid Leukemia

**Variant class:** FLT3 mutation

##### Supporting Statement:

The FDA has granted Fast Track Designation to the benzimidazole type I kinase inhibitor, crenolanib, for:

- FLT3 mutation-positive relapsed or refractory acute myeloid leukemia (AML)
- PDGFRA D842V mutated unresectable or metastatic gastrointestinal stromal tumors (GIST)

##### Reference:

<https://www.globenewswire.com/news-release/2017/12/01/1216122/0/en/Arog-Pharmaceuticals-Receives-FDA-Fast-Track-Designation-for-Crenolanib-in-Relapsed-or-Refractory-FLT3-Positive-AML.html>

#### tuspetinib

**Cancer type:** Acute Myeloid Leukemia

**Variant class:** FLT3 mutation

##### Supporting Statement:

The FDA has granted Fast Track Designation to tuspetinib (HM43239), a myeloid kinase inhibitor, for relapsed or refractory (R/R) acute myeloid leukemia (AML) with FLT3 mutation.

##### Reference:

<https://www.aptose.com/news-media/press-releases/detail/230/aptose-receives-fast-track-designation-for-hm43239-in>



## References

1. Okano et al. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet.* 1998 Jul;19(3):219-20. PMID: 9662389
2. Fernandez et al. A DNA methylation fingerprint of 1628 human samples. *Genome Res.* 2012 Feb;22(2):407-19. PMID: 21613409
3. Jones et al. The epigenomics of cancer. *Cell.* 2007 Feb 23;128(4):683-92. PMID: 17320506
4. Ley et al. DNMT3A mutations in acute myeloid leukemia. *N. Engl. J. Med.* 2010 Dec 16;363(25):2424-33. PMID: 21067377
5. Marková et al. Prognostic impact of DNMT3A mutations in patients with intermediate cytogenetic risk profile acute myeloid leukemia. *Eur. J. Haematol.* 2012 Feb;88(2):128-35. PMID: 21967546
6. Yang et al. DNMT3A in haematological malignancies. *Nat. Rev. Cancer.* 2015 Mar;15(3):152-65. PMID: 25693834
7. Renneville et al. Prognostic significance of DNA methyltransferase 3A mutations in cytogenetically normal acute myeloid leukemia: a study by the Acute Leukemia French Association. *Leukemia.* 2012 Jun;26(6):1247-54. PMID: 22289988
8. Marcucci et al. Age-related prognostic impact of different types of DNMT3A mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J. Clin. Oncol.* 2012 Mar 1;30(7):742-50. PMID: 22291079
9. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
10. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]
11. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
12. Kumar et al. DNMT3A (R882) mutation features and prognostic effect in acute myeloid leukemia in Coexistent with NPM1 and FLT3 mutations. *Hematol Oncol Stem Cell Ther.* 2018 Jun;11(2):82-89. PMID: 29079128
13. Thol et al. Incidence and prognostic influence of DNMT3A mutations in acute myeloid leukemia. *J. Clin. Oncol.* 2011 Jul 20;29(21):2889-96. PMID: 21670448
14. Sandoval et al. Mutations in the DNMT3A DNA methyltransferase in acute myeloid leukemia patients cause both loss and gain of function and differential regulation by protein partners. *J. Biol. Chem.* 2019 Mar 29;294(13):4898-4910. PMID: 30705090
15. Holz-Schietinger et al. Mutations in DNA methyltransferase (DNMT3A) observed in acute myeloid leukemia patients disrupt processive methylation. *J. Biol. Chem.* 2012 Sep 7;287(37):30941-51. PMID: 22722925
16. Russler-Germain et al. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell.* 2014 Apr 14;25(4):442-54. PMID: 24656771
17. NCCN Guidelines® - NCCN-T-Cell Lymphomas [Version 1.2023]
18. Grafone et al. An overview on the role of FLT3-tyrosine kinase receptor in acute myeloid leukemia: biology and treatment. *Oncol Rev.* 2012 Mar 5;6(1):e8. PMID: 25992210
19. Short et al. Emerging treatment paradigms with FLT3 inhibitors in acute myeloid leukemia. *Ther Adv Hematol.* 2019; 10: 2040620719827310. PMID: 30800259
20. Small. FLT3 mutations: biology and treatment. *Hematology Am Soc Hematol Educ Program.* 2006:178-84. PMID: 17124058
21. Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med.* 2013 May 30;368(22):2059-74. doi: 10.1056/NEJMoa1301689. Epub 2013 May 1. PMID: 23634996
22. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
23. Nakao et al. Internal tandem duplication of the *flt3* gene found in acute myeloid leukemia. *Leukemia.* 1996 Dec;10(12):1911-8. PMID: 8946930
24. Yamamoto et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood.* 2001 Apr 15;97(8):2434-9. PMID: 11290608
25. Carow et al. Expression of the hematopoietic growth factor receptor FLT3 (STK-1/Flk2) in human leukemias. *Blood.* 1996 Feb 1;87(3):1089-96. PMID: 8562934
26. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
27. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/207997s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/207997s008lbl.pdf)
28. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/211349s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/211349s003lbl.pdf)
29. <https://www.globenewswire.com/news-release/2017/12/01/1216122/0/en/Arog-Pharmaceuticals-Receives-FDA-Fast-Track-Designation-for-Crenolanib-in-Relapsed-or-Refractory-FLT3-Positive-AML.html>
30. <https://www.aptose.com/news-media/press-releases/detail/230/aptose-receives-fast-track-designation-for-hm43239-in>
31. [https://www.daiichisankyo.com/media\\_investors/media\\_relations/press\\_releases/detail/006896.html](https://www.daiichisankyo.com/media_investors/media_relations/press_releases/detail/006896.html)

---

## References (continued)

32. Jasleen et al. Results of a Phase II Study of Crenolanib in Relapsed/Refractory Acute Myeloid Leukemia Patients (Pts) with Activating FLT3 Mutations. *Blood*. 124:389
33. Bartus et al. Crenolanib in combination with standard chemotherapy in newly diagnosed FLT3-mutated AML patients. ASH 2017. Abstract #566