



## Sample Information

**Patient Name:** 韓鐘磊  
**Gender:** Male  
**ID No.:** C120260498  
**History No.:** 46869232  
**Age:** 50

**Ordering Doctor:** DOC1878G 沈佳儀  
**Ordering REQ.:** 0BXNMHK  
**Signing in Date:** 2022/07/21

**Path No.:** S111-99876  
**MP No.:** F22072  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S111-26016A  
**Percentage of tumor cells:** 60%

**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)

**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	4
Clinical Trials Summary	16
Alert Details	16

**Report Highlights**  
2 Relevant Biomarkers  
8 Therapies Available  
5 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	<b>EML4-ALK fusion</b>	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	<b>MET amplification</b>		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>EML4-ALK fusion</b> EMAP like 4 - ALK receptor tyrosine kinase	<b>alectinib</b> <sup>1,2</sup> <b>brigatinib</b> <sup>1,2</sup> <b>ceritinib</b> <sup>1,2</sup> <b>crizotinib</b> <sup>1,2</sup> <b>lorlatinib</b> <sup>1,2</sup> next-generation ALK inhibitor	brigatinib ceritinib crizotinib lorlatinib	4
IA	<b>MET amplification</b> MET proto-oncogene, receptor tyrosine kinase	capmatinib crizotinib tepotinib	None	1

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

**Tier Reference:** Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
EML4-ALK	EML4-ALK.E18A20.COSF487.1	chr2:42543190 - chr2:29446394	25783

### Copy Number Variations

Gene	Locus	Copy Number
MET	chr7:116313480	5.1

## Biomarker Descriptions

### ALK (ALK receptor tyrosine kinase)

**Background:** The ALK gene encodes the ALK receptor tyrosine kinase (RTK) with sequence similarity to the insulin receptor subfamily of kinases<sup>1</sup>. ALK is the target of recurrent alterations in cancer, the most common being chromosomal rearrangements that generate fusion genes containing the intact ALK tyrosine kinase domain combined with multiple partner genes<sup>2</sup>. ALK fusion kinases are constitutively activated and drive oncogenic transformation via activation of downstream STAT3, PI3K/AKT/MTOR, and RAS/RAF/MEK/ERK pathways<sup>2,3,4,5</sup>.

**Alterations and prevalence:** ALK was discovered by positional cloning of translocations involving nucleophosmin (NPM) on 5q35 with a previously unidentified RTK on 2p23 (ALK), which occur in over 50% of anaplastic large cell lymphoma cases<sup>1,6</sup>. In contrast, about 5% of non-small cell lung cancer (NSCLC) cases generate recurrent ALK fusions with EML4, KIF5B, and HIP1<sup>7,8,9</sup>.

**Potential relevance:** The first generation small molecule tyrosine kinase inhibitor (TKI), crizotinib<sup>10</sup>, was FDA approved (2011) for the treatment of ALK positive advanced NSCLC. Kinase domain mutations including L1196M, G1269A, F1174L, G1202R, as well as other variants have been shown to confer acquired resistance to crizotinib in ALK positive NSCLC<sup>11,12,13,14</sup>. Other mechanisms of acquired resistance involve amplification of the ALK fusion gene and activation of alternate or bypass signaling pathways involving EGFR, KIT, MET, and IGF1R<sup>15</sup>. In order to overcome acquired resistance, second and third-generation ALK inhibitors including ceritinib<sup>16</sup> (2014), alectinib<sup>17</sup> (2015), brigatinib<sup>18</sup> (2017), and lorlatinib<sup>19</sup> (2018) were developed and approved by the FDA. Two phase III trials evaluating crizotinib and alectinib as first line therapy in NSCLC, including patients with asymptomatic central nervous system (CNS) disease, were conducted and both studies showed consistent higher objective response rates (ORR) with alectinib relative to crizotinib<sup>20,21</sup>. For this reason, alectinib is a preferred first-line treatment of ALK positive NSCLC<sup>22</sup>.

### MET (MET proto-oncogene, receptor tyrosine kinase)

**Background:** The MET proto-oncogene encodes a receptor tyrosine kinase for the hepatocyte growth factor (HGF) protein, which is expressed by mesenchymal cells. MET is expressed as multiple isoforms with transcript variant 1 (NM\_001127500.3) encoding a

## Biomarker Descriptions (continued)

1408 amino acid protein and transcript variant 2 (NM\_000245.4) encoding a 1390 amino acid protein, both of which possess an intact protein kinase domain<sup>23</sup>. Ubiquitin-dependent proteolysis is responsible for regulating the steady state level of the MET protein via recognition of the tyrosine phosphorylation site Y1003(NM\_000245.4), sometimes referred to as Y1021 (NM\_001127500.3), in the MET Cbl-binding domain within the juxtamembrane region<sup>24,25,26</sup>. Growth factor signaling leads to MET dimerization and subsequent initiation of downstream effectors including those involved in the RAS/RAF/MEK/ERK and PI3K/AKT signaling pathways, which regulate cell migration, proliferation, and survival<sup>27,28</sup>.

**Alterations and prevalence:** Somatic mutations in MET are observed in 10% of uterine corpus endometrial carcinoma, 9% of skin cutaneous melanoma, 8% of papillary renal cell carcinoma (PRCC), and 4% of lung adenocarcinoma, colorectal adenocarcinoma, bladder urothelial carcinoma, and uterine carcinosarcoma<sup>29,30</sup>. Recurrent somatic MET alterations include activating mutations, gene amplification, and translocations generating MET gene fusions. Recurrent somatic mutations fall into two classes, mutations in the MET kinase domain, which are uncommon, and splice-site mutations affecting exon 14. Recurrent kinase domain mutations are observed in PRCC and include M1250T, H1094Y, and V1070E (NM\_000245.4)<sup>29,30</sup>. Mutation of the Y1003 phosphorylation site is reported in approximately 2% of MET altered lung cancer<sup>31</sup>. In contrast, splice-site mutations flanking exon 14 are observed in 3-4% of all non-small cell lung cancer (NSCLC)<sup>32</sup>. These mutations include canonical splice site mutations affecting exon 14 and deletions that extend into the splicing motifs within intron 13<sup>31,33</sup>. Such mutations disrupt splicing leading to the formation of an alternative transcript that joins exon 13 directly to exon 15 and skips exon 14 entirely. The MET exon 14 skipping transcript lacks the juxtamembrane domain that contains the recognition motif for ubiquitin-dependent proteolysis and thus leads to a marked increase in the steady-state level of the MET protein<sup>34</sup>. MET exon 14 skipping mutations act as oncogenic drivers in lung cancer mutually exclusive to activating mutations in EGFR and KRAS and other oncogenic fusions such as ALK and ROS1<sup>33,35,36</sup>. MET is amplified in 2-5% of ovarian cancer, esophageal adenocarcinoma, stomach adenocarcinoma, glioblastoma, and lung adenocarcinoma<sup>29,37,38</sup>. Recurrent MET fusions, although infrequent, are observed in adult and pediatric glioblastoma, papillary renal cell carcinoma, lung cancer, liver cancer, thyroid cancer, and melanoma<sup>39,40,41</sup>. MET alterations are believed to be enriched in late-stage cancers where they drive tumor progression and metastasis<sup>42,43,44</sup>.

**Potential relevance:** In 2020, the FDA granted accelerated approval to capmatinib<sup>45</sup> for NSCLC harboring MET exon 14 skipping positive as detected by an FDA-approved test. The kinase inhibitor, tepotinib<sup>46</sup>, is also approved (2021) for MET exon 14 skipping mutations in NSCLC. MET exon 14 skipping mutations confer sensitivity to approved kinase inhibitors including crizotinib (2011), which is recommended for MET amplifications and exon 14 skipping mutations<sup>22,33,35,36</sup>. The FDA also granted breakthrough therapy designation (2018) to crizotinib for metastatic non-small cell lung cancer (NSCLC) with MET exon 14 alterations with disease progression on or after platinum-based chemotherapy<sup>47</sup>. Conversely, amplification of MET has been observed to mediate resistance to EGFR tyrosine kinase inhibitors (TKIs)<sup>48,49,50,51,52</sup>. However, the FDA has granted Fast Track designation (2021) to the MET/CSF1R/SRC small molecule inhibitor, TPX-0022<sup>53</sup>, for MET amplified advanced or metastatic gastric cancer, including gastroesophageal junction adenocarcinoma (GEJ) after prior chemotherapy. Tepotinib has also been recommended for treatment of NSCLC with high-level MET amplification<sup>22</sup>. In a phase II trial testing the MET inhibitor savolitinib, patients with advanced PRCC exhibited median progression free survival (PFS) of 6.2 and 1.4 months for MET-driven and MET-independent PRCC, respectively<sup>54</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### EML4-ALK fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
ceritinib	●	ⓘ	●	●	● (IV)
brigatinib	●	ⓘ	●	●	×
crizotinib	●	ⓘ	●	●	×
lorlatinib	●	ⓘ	●	●	×
allectinib	●	●	●	●	×
next-generation ALK inhibitor	×	×	×	●	×

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### EML4-ALK fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
brigatinib, alectinib	×	×	×	×	● (III)
repotrectinib	×	×	×	×	● (I/II)

### MET amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
capmatinib	×	●	×	×	×
crizotinib	×	●	×	×	×
tepotinib	×	●	×	×	×
HLX55	×	×	×	×	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2022-06-15. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### EML4-ALK fusion

#### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-09-03

Variant class: ALK fusion or ALK overexpression

#### Indications and usage:

ALECENSA® is a kinase inhibitor indicated for the treatment of patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/208434s012lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208434s012lbl.pdf)

## EML4-ALK fusion (continued)

### ● brigatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2022-02-28

**Variant class:** ALK fusion

**Indications and usage:**

ALUNBRIG® is a kinase inhibitor indicated for the treatment of adult patients with anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/208772s013lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/208772s013lbl.pdf)

### ● ceritinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2021-10-07

**Variant class:** ALK fusion or ALK overexpression

**Indications and usage:**

ZYKADIA® is a kinase inhibitor indicated for the treatment of adults with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)-positive as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/211225s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/211225s004lbl.pdf)

### ● crizotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2021-09-22

**Variant class:** ALK fusion

**Indications and usage:**

XALKORI® is a kinase inhibitor indicated for the treatment of

- patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK) or ROS1-positive as detected by an FDA-approved test.
- pediatric patients 1 year of age and older and young adults with relapsed or refractory, systemic anaplastic large cell lymphoma (ALCL) that is ALK-positive.
  - Limitations of Use: The safety and efficacy of XALKORI® have not been established in older adults with relapsed or refractory, systemic ALK-positive ALCL.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/202570s031lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202570s031lbl.pdf)

### ● lorlatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2021-03-03

**Variant class:** ALK fusion or ALK overexpression

**Indications and usage:**

LORBRENA® is a kinase inhibitor indicated for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors are anaplastic lymphoma kinase (ALK)-positive as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/210868s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/210868s004lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-06-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### EML4-ALK fusion

#### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

#### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

#### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

#### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

## EML4-ALK fusion (continued)

### ● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion or ALK overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]

### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Metastatic, Advanced (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]

## EML4-ALK fusion (continued)

### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Metastatic, Advanced (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]

### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion or ALK overexpression

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]



## EML4-ALK fusion (continued)

### ● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Metastatic, Advanced (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 2.2021]

### ○ brigatinib

Cancer type: Inflammatory Myofibroblastic Tumor Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2022]

### ○ ceritinib

Cancer type: Inflammatory Myofibroblastic Tumor Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2022]

### ○ crizotinib

Cancer type: Inflammatory Myofibroblastic Tumor Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2022]

## EML4-ALK fusion (continued)

### ○ lorlatinib

Cancer type: Inflammatory Myofibroblastic Tumor    Variant class: ALK fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2022]

## MET amplification

### ● capmatinib

Cancer type: Non-Small Cell Lung Cancer    Variant class: MET amplification

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer    Variant class: MET amplification

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### ● tepotinib

Cancer type: Non-Small Cell Lung Cancer    Variant class: MET amplification

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

## Current EMA Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

EMA information is current as of 2022-06-15. For the most up-to-date information, search [www.ema.europa.eu/ema](https://www.ema.europa.eu/ema).

### EML4-ALK fusion

#### ☒ alectinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-10-11

Variant class: ALK fusion or ALK overexpression

Reference:

[https://www.ema.europa.eu/en/documents/product-information/alecensa-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/alecensa-epar-product-information_en.pdf)

#### ☒ brigatinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-05-18

Variant class: ALK fusion

Reference:

[https://www.ema.europa.eu/en/documents/product-information/alunbrig-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/alunbrig-epar-product-information_en.pdf)

#### ☒ crizotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-02-21

Variant class: ALK fusion

Reference:

[https://www.ema.europa.eu/en/documents/product-information/xalkori-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/xalkori-epar-product-information_en.pdf)

#### ☒ ceritinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-02-25

Variant class: ALK positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/zykadia-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/zykadia-epar-product-information_en.pdf)

#### ☒ lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-04-07

Variant class: ALK positive

Reference:

[https://www.ema.europa.eu/en/documents/product-information/lorviqua-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/lorviqua-epar-product-information_en.pdf)

## Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2022-06-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### EML4-ALK fusion

#### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / A

**Population segment (Line of therapy):**

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 4
- Advanced, Progression (Second-line therapy, Subsequent therapy); ESMO-MCBS v1.1 score: 4

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / A

**Population segment (Line of therapy):**

- Stage IV (First-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / A

**Population segment (Line of therapy):**

- Advanced, Progression (Second-line therapy, Subsequent therapy); ESMO-MCBS v1.1 score: 4

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EML4-ALK fusion (continued)

### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 4
- (Second-line therapy, Subsequent therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● next-generation ALK inhibitor

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (Second-line therapy, Subsequent therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 4

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 4

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EML4-ALK fusion (continued)

### ● next-generation ALK inhibitor

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- Advanced, Progression (Second-line therapy, Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● alectinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Advanced (Second-line therapy, Subsequent therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● lorlatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV; Advanced, Progression (Second-line therapy, Subsequent therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EML4-ALK fusion (continued)

### ● brigatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Advanced (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● ceritinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## Clinical Trials in Taiwan region:

### Clinical Trials Summary

#### EML4-ALK fusion

NCT ID	Title	Phase
NCT03596866	A Phase III Randomized Open-label Study of Brigatinib (Alunbrig) Versus Alectinib (Alecensa) in Advanced Anaplastic Lymphoma Kinase-Positive Non Small-Cell Lung Cancer Patients Who Have Progressed on Crizotinib (Xalkori)	III
NCT02584933	An Open-label, Multi-center, Phase IV Roll-over Study in Patients With ALK Positive Malignancies Who Have Completed a Novartis-sponsored Ceritinib (LDK378) Study and Are Judged by the Investigator to Benefit From Continued Treatment With Ceritinib	IV
NCT03093116	A Phase I/II, Open-Label, Multi-Center, First-in-Human Study of the Safety, Tolerability, Pharmacokinetics, and Anti-Tumor Activity of TPX-0005 in Patients With Advanced Solid Tumors Harboring ALK, ROS1, or NTRK1-3 Rearrangements (TRIDENT-1)	I/II
NCT04094610	A Phase I/II, Open-Label, Safety, Tolerability, Pharmacokinetics, and Anti-Tumor Activity Study of Repotrectinib in Pediatric and Young Adult Subjects With Advanced or Metastatic Malignancies Harboring ALK, ROS1, NTRK1-3 Alterations	I/II

#### MET amplification

NCT ID	Title	Phase
NCT04169178	A Phase I Dose Finding/Expansion Study of HLX55, A Monoclonal Antibody Targeting Tyrosine-Protein Kinase MET (C-MET) in Patients With Advanced Solide Tumors Refractory to Standard Therapy	I

## Alerts Informed By Public Data Sources

### Current NCCN Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

NCCN information is current as of 2022-06-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### EML4-ALK fusion

##### atezolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** ALK fusion

##### Summary:

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK rearrangements."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]



## EML4-ALK fusion (continued)

### – nivolumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK rearrangements."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

### – pembrolizumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: ALK fusion

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK rearrangements."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Webb et al. Anaplastic lymphoma kinase: role in cancer pathogenesis and small-molecule inhibitor development for therapy. *Expert Rev Anticancer Ther.* 2009 Mar;9(3):331-56. PMID: 19275511
2. Shaw et al. Tyrosine kinase gene rearrangements in epithelial malignancies. *Nat. Rev. Cancer.* 2013 Nov;13(11):772-87. PMID: 24132104
3. Chiarle et al. Stat3 is required for ALK-mediated lymphomagenesis and provides a possible therapeutic target. *Nat. Med.* 2005 Jun;11(6):623-9. PMID: 15895073
4. Bai et al. Nucleophosmin-anaplastic lymphoma kinase associated with anaplastic large-cell lymphoma activates the phosphatidylinositol 3-kinase/Akt antiapoptotic signaling pathway. *Blood.* 2000 Dec 15;96(13):4319-27. PMID: 11110708
5. Hrustanovic et al. RAS signaling in ALK fusion lung cancer. *Small GTPases.* 2016;7(1):32-3. PMID: 26901483
6. Morris et al. Fusion of a kinase gene, ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. *Science.* 1994 Mar 4;263(5151):1281-4. PMID: 8122112
7. Kwak et al. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N. Engl. J. Med.* 2010 Oct 28;363(18):1693-703. PMID: 20979469
8. Yu et al. Frequencies of ALK rearrangements in lung adenocarcinoma subtypes: a study of 2299 Chinese cases. *Springerplus.* 2016 Jun 27;5(1):894. doi: 10.1186/s40064-016-2607-5. eCollection 2016. PMID: 27386342
9. Dai et al. Incidence and patterns of ALK FISH abnormalities seen in a large unselected series of lung carcinomas. *Send to Mol Cytogenet.* 2012 Dec 3;5(1):44. doi: 10.1186/1755-8166-5-44. PMID: 23198868
10. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/202570s031lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/202570s031lbl.pdf)
11. Choi et al. EML4-ALK mutations in lung cancer that confer resistance to ALK inhibitors. *N. Engl. J. Med.* 2010 Oct 28;363(18):1734-9. PMID: 20979473
12. Awad et al. ALK inhibitors in non-small cell lung cancer: crizotinib and beyond. *Clin Adv Hematol Oncol.* 2014 Jul;12(7):429-39. PMID: 25322323
13. Kim et al. Heterogeneity of genetic changes associated with acquired crizotinib resistance in ALK-rearranged lung cancer. *J Thorac Oncol.* 2013 Apr;8(4):415-22. PMID: 23344087
14. Katayama et al. Mechanisms of acquired crizotinib resistance in ALK-rearranged lung Cancers. *Sci Transl Med.* 2012 Feb 8;4(120):120ra17. doi: 10.1126/scitranslmed.3003316. Epub 2012 Jan 25. PMID: 22277784
15. Katayama. Drug resistance in anaplastic lymphoma kinase-rearranged lung cancer. *Cancer Sci.* 2018 Mar;109(3):572-580. PMID: 29336091
16. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/211225s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/211225s004lbl.pdf)
17. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/208434s012lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208434s012lbl.pdf)
18. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/208772s013lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/208772s013lbl.pdf)
19. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/210868s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/210868s004lbl.pdf)
20. Peters et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N. Engl. J. Med.* 2017 Aug 31;377(9):829-838. PMID: 28586279
21. Hida et al. Alectinib versus crizotinib in patients with ALK-positive non-small-cell lung cancer (J-ALEX): an open-label, randomised phase 3 trial. *Lancet.* 2017 Jul 1;390(10089):29-39. PMID: 28501140
22. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2022]
23. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
24. Peschard et al. A conserved DpYR motif in the juxtamembrane domain of the Met receptor family forms an atypical c-Cbl/Cbl-b tyrosine kinase binding domain binding site required for suppression of oncogenic activation. *J. Biol. Chem.* 2004 Jul 9;279(28):29565-71. PMID: 15123609
25. Peschard et al. Mutation of the c-Cbl TKB domain binding site on the Met receptor tyrosine kinase converts it into a transforming protein. *Mol. Cell.* 2001 Nov;8(5):995-1004. PMID: 11741535
26. Abella et al. Met/Hepatocyte growth factor receptor ubiquitination suppresses transformation and is required for Hrs phosphorylation. *Mol. Cell. Biol.* 2005 Nov;25(21):9632-45. PMID: 16227611
27. Sierra et al. c-MET as a potential therapeutic target and biomarker in cancer. *Ther Adv Med Oncol.* 2011 Nov;3(1 Suppl):S21-35. PMID: 22128285
28. Mo et al. Targeting MET in cancer therapy. *Chronic Dis Transl Med.* 2017 Sep;3(3):148-153. PMID: 29063069

## References (continued)

29. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
30. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
31. Schrock et al. Characterization of 298 Patients with Lung Cancer Harboring MET Exon 14 Skipping Alterations. *J Thorac Oncol.* 2016 Sep;11(9):1493-502. PMID: 27343443
32. Socinski et al. MET Exon 14 Skipping Mutations in Non-Small-Cell Lung Cancer: An Overview of Biology, Clinical Outcomes, and Testing Considerations. *JCO Precis Oncol.* 2021;5. PMID: 34036238
33. Frampton et al. Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov.* 2015 Aug;5(8):850-9. PMID: 25971938
34. Pilotto et al. MET exon 14 juxtamembrane splicing mutations: clinical and therapeutical perspectives for cancer therapy. *Ann Transl Med.* 2017 Jan;5(1):2. doi: 10.21037/atm.2016.12.33. PMID: 28164087
35. Reungwetwattana et al. The race to target MET exon 14 skipping alterations in non-small cell lung cancer: The Why, the How, the Who, the Unknown, and the Inevitable. *Lung Cancer.* 2017 Jan;103:27-37. PMID: 28024693
36. Saffroy et al. MET exon 14 mutations as targets in routine molecular analysis of primary sarcomatoid carcinoma of the lung. *Oncotarget.* 2017 Jun 27;8(26):42428-42437. PMID: 28418914
37. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
38. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
39. Yeh et al. Activating MET kinase rearrangements in melanoma and Spitz tumours. *Nat Commun.* 2015 May 27;6:7174. doi: 10.1038/ncomms8174. PMID: 26013381
40. Bao et al. RNA-seq of 272 gliomas revealed a novel, recurrent PTPRZ1-MET fusion transcript in secondary glioblastomas. *Genome Res.* 2014 Nov;24(11):1765-73. PMID: 25135958
41. Sebastian et al. Recurrent MET fusion genes represent a drug target in pediatric glioblastoma. *Nat. Med.* 2016 Nov;22(11):1314-1320. PMID: 27748748
42. Zeng et al. c-Met gene amplification is associated with advanced stage colorectal cancer and liver metastases. *Cancer Lett.* 2008 Jul 8;265(2):258-69. PMID: 18395971
43. Tsugawa et al. Amplification of the c-met, c-erbB-2 and epidermal growth factor receptor gene in human gastric cancers: correlation to clinical features. *Oncology.* 1998 Sep-Oct;55(5):475-81. PMID: 9732228
44. Di et al. Overexpression and amplification of the met/HGF receptor gene during the progression of colorectal cancer. *Clin. Cancer Res.* 1995 Feb;1(2):147-54. PMID: 9815967
45. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/213591s002lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/213591s002lbl.pdf)
46. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/214096s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214096s000lbl.pdf)
47. [https://www.pfizer.com/news/press-release/press-release-detail/pfizer\\_s\\_xalkori\\_crizotinib\\_receives\\_fda\\_breakthrough\\_therapy\\_designation\\_in\\_two\\_new\\_indications-0](https://www.pfizer.com/news/press-release/press-release-detail/pfizer_s_xalkori_crizotinib_receives_fda_breakthrough_therapy_designation_in_two_new_indications-0)
48. Bean et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc. Natl. Acad. Sci. U.S.A.* 2007 Dec 26;104(52):20932-7. PMID: 18093943
49. Chen et al. Clinicopathologic and molecular features of epidermal growth factor receptor T790M mutation and c-MET amplification in tyrosine kinase inhibitor-resistant Chinese non-small cell lung cancer. *Pathol Oncol Res.* 2009 Dec;15(4):651-8. doi: 10.1007/s12253-009-9167-8. Epub 2009 Apr 21. PMID: 19381876
50. Suda et al. Reciprocal and complementary role of MET amplification and EGFR T790M mutation in acquired resistance to kinase inhibitors in lung cancer. *Clin. Cancer Res.* 2010 Nov 15;16(22):5489-98. PMID: 21062933
51. Zhang et al. Current mechanism of acquired resistance to epidermal growth factor receptor-tyrosine kinase inhibitors and updated therapy strategies in human nonsmall cell lung cancer. *J Cancer Res Ther.* 2016 Dec;12(Supplement):C131-C137. PMID: 28230005
52. Nguyen et al. Acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancers dependent on the epidermal growth factor receptor pathway. *Clin Lung Cancer.* 2009 Jul;10(4):281-9. PMID: 19632948
53. [https://www.sec.gov/Archives/edgar/data/1595893/000156459021042621/tptx-ex991\\_20.htm](https://www.sec.gov/Archives/edgar/data/1595893/000156459021042621/tptx-ex991_20.htm)
54. Choueiri et al. Biomarker-Based Phase II Trial of Savolitinib in Patients With Advanced Papillary Renal Cell Cancer. *J. Clin. Oncol.* 2017 Sep 10;35(26):2993-3001. PMID: 28644771