



Sample Information

Patient Name: 張馨勻
Gender: Female
ID No.: A228948640
History No.: 22747086
Age: 28

Ordering Doctor: DOC1901H 高志平
Ordering REQ.: D61374E
Signing in Date: 2021/02/18

Path No.: S110-98251
MP No.: MY21001
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2021/02/09
Note:

Sample Cancer Type: Acute Promyelocytic Leukemia

Table of Contents	Page	Report Highlights
Variant Details	2	1 Relevant Biomarkers
Biomarker Descriptions	3	9 Therapies Available
Relevant Therapy Summary	4	7 Clinical Trials
Relevant Therapy Details	5	
Diagnostic Details	11	

Relevant Acute Promyelocytic Leukemia Variants

Gene	Finding
RARA	<i>PML-RARA fusion</i>

Relevant Biomarkers

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
<i>PML-RARA fusion</i> promyelocytic leukemia - retinoic acid receptor alpha	arsenic trioxide ^{1,2} arsenic trioxide + tretinoin ^{1,2} anthracycline + arsenic trioxide anthracycline + arsenic trioxide + tretinoin arsenic trioxide + idarubicin + tretinoin cytarabine + daunorubicin + tretinoin gemtuzumab ozogamicin	None	7

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Relevant Biomarkers (continued)

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
	gemtuzumab ozogamicin + chemotherapy idarubicin + tretinoin		
Prognostic significance: None Diagnostic significance: Acute Promyelocytic Leukemia, Acute Myeloid Leukemia			

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

ETV6 p.(Q413fs) c.1238_1239insCA

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ETV6	p.(Q413fs)	c.1238_1239insCA	.	chr12:12038945	4.06%	NM_001987.4	frameshift Insertion	1994
SF3B1	p.(=)	c.2631T>C	.	chr2:198265526	99.95%	NM_012433.3	synonymous	1996
TET2	p.(V218M)	c.652G>A	.	chr4:106155751	49.75%	NM_001127208.2	missense	1998
TET2	p.(=)	c.4140T>C	.	chr4:106190862	47.67%	NM_001127208.2	synonymous	1999
HRAS	p.(=)	c.81T>C	.	chr11:534242	99.50%	NM_001130442.2	synonymous	1999
WT1	p.(=)	c.1107A>G	.	chr11:32417945	99.95%	NM_024426.4	synonymous	2000
ETV6	p.(=)	c.258G>A	.	chr12:11992168	51.35%	NM_001987.4	synonymous	1996
ETV6	p.(Y391N)	c.1171T>A	.	chr12:12038878	14.81%	NM_001987.4	missense	1999
KRAS	p.(=)	c.483G>A	.	chr12:25368462	99.90%	NM_033360.3	synonymous	2000
SH2B3	p.(F135V)	c.403T>G	.	chr12:111856352	48.61%	NM_005475.2	missense	1329
SH2B3	p.(W262R)	c.784T>C	.	chr12:111884608	99.90%	NM_005475.2	missense	1999
TP53	p.(P72R)	c.215C>G	.	chr17:7579472	81.23%	NM_000546.5	missense	1998
NF1	p.(=)	c.702G>A	.	chr17:29508775	44.79%	NM_001042492.2	synonymous	1085
NF1	p.(=)	c.2034G>A	.	chr17:29553485	51.33%	NM_001042492.2	synonymous	1999
SRSF2	p.(=)	c.144C>T	.	chr17:74733099	100.00%	NM_003016.4	synonymous	2000
ASXL1	p.(L815P)	c.2444T>C	.	chr20:31022959	99.95%	NM_015338.5	missense	1997
ASXL1	p.(=)	c.3759T>C	.	chr20:31024274	99.95%	NM_015338.5	synonymous	1999
ZRSR2	p.(=)	c.864C>T	.	chrX:15838366	100.00%	NM_005089.3	synonymous	1996
BCOR	p.(=)	c.1692A>G	.	chrX:39932907	47.42%	NM_001123385.1	synonymous	1976
BCOR	p.(=)	c.1260T>C	.	chrX:39933339	99.95%	NM_001123385.1	synonymous	1999

Variant Details (continued)

Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
PML-RARA	PML-RARA.P3R3	chr15:74315749 - chr17:38504568	70849

Biomarker Descriptions

ETV6 (ETS variant 6)

Background: The ETV6 gene encodes the E twenty-six (ETS) variant 1 transcription factor. ETV6 contains an N-terminal pointed (PNT) domain responsible for protein-protein interactions and a C-terminal ETS domain involved in DNA binding¹. ETV6 plays a critical role in embryonic development as well as hematopoiesis and is the target of chromosomal rearrangement and missense mutations in hematological malignancies as well as solid tumors^{2,3}. Hereditary mutations in ETV6 are associated with a predisposition to hematological cancers, including acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), and myelodysplastic syndromes (MDS)^{4,5,6}.

Alterations and prevalence: ETV6 translocations are prevalent in hematological malignancies and have been observed with numerous fusion partners⁷. The most recurrent translocation is t(12;21)(q34;q11) which results in ETV6-RUNX1 fusion and is observed in 20-25% childhood acute lymphoblastic leukemia (ALL)^{7,8,9}. ETV6-RUNX1 fusions are also observed in adult ALL (2%)^{8,9}. The t(5;12)(q33;p13) translocation which results in the ETV6-PDGFRB fusion is recurrent in chronic myelomonocytic leukemia (CMML)^{7,10}. Other ETV6 fusions including ETV6-PDGFRB, ETV6-NTRK2, ETV6-NTRK3, and ETV6-ABL1 are reported in hematological malignancies as well as solid tumors^{3,7,11}. ETV6 fusions involving a receptor tyrosine kinase (RTK) fusion partner retains the ETV6 PNT domain and the tyrosine kinase domain of the RTK, leading to constitutive kinase activation^{7,11}. Mutations in ETV6 are primarily missense, nonsense, or frameshift and are observed in about 1-5% of select myeloid malignancies and solid tumors, including chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), diffuse large B-cell lymphoma (DLBCL), MDS, AML, ALL, melanoma, lung, bladder, stomach, colorectal, and uterine cancers^{1,12,13}. ETV6 mutations occur in the PNT and ETS domain of ETV6 and may impair ETV6 oligomerization or DNA-binding, respectively¹.

Potential relevance: ETV6-NTRK3 fusions are used as an ancillary diagnostic marker in congenital/infantile fibrosarcoma¹⁴. Nonsense or frameshift mutations in ETV6 are independently associated with poor prognosis in MDS⁶. However, ETV6-RUNX1 fusions are associated with favorable outcomes in ALL and good risk in B-cell ALL (B-ALL)⁹. ETV6 fusions that partner with a RTKs demonstrate response to various tyrosine kinase inhibitors such as imatinib, nilotinib, and entrectinib. Specifically, individual case reports of an ETV6-PDGFRB fusion chronic eosinophilic leukemia patient and an ETV6-PDGFRB fusion CMML patient treated with imatinib demonstrated complete cytogenetic response (CCyR) and complete hematological responses, respectively^{15,16}. Additionally, an ETV6-ABL1 fusion Ph-negative CML patient treated with nilotinib demonstrated CCyR and major molecular response (MMR) at 22 months from diagnosis¹⁷. In another case report, an ETV6-NTRK3 fusion mammary analogue secretory carcinoma (MASC) patient demonstrated partial response to entrectinib with 89% reduction in tumor burden¹⁸.

RARA (retinoic acid receptor alpha)

Background: The RARA gene encodes the retinoic acid receptor alpha, a transcription factor and a member of the retinoic acid (RA) nuclear receptor family. RARA binds DNA as a heterodimer with its cofactor the retinoid X receptor alpha (RXRA)¹⁹. Binding of the RARA/RXRA complex to specific RA response elements (RAREs) causes activation of transcription¹⁹. RARA is also involved in white blood cell (WBC) differentiation and hematopoietic stem cell specification^{20,21}. RARA translocations are the genetic driver of acute promyelocytic leukemia (APL) where the 3' region of the RARA gene is translocated to the 5' region of partner genes such as the promyelocytic leukemia (PML) gene²². The PML-RARA fusion protein contributes to the pathogenesis of APL by blocking differentiation and promoting aberrant self-renewal of APL cells, leading to a buildup of immature white blood cells in the blood and bone marrow²³.

Alterations and prevalence: More than 95% of APL patients harbor the t(15;17)(q22;q21) translocation that results in PML-RARA fusion^{19,24}. Other RARA fusion partners, including PLZF, NPM, NUMA, STAT5b, PRKAR1A, FIP1L1, TBLXR1, FNDC3B, GTF2I, IRF2BP2, account for the rest^{24,25,26,27,28,29}.

Potential relevance: The presence of PML-RARA fusion characterized by the presence of t(15;17)(q22;q21) translocation is diagnostic of APL³⁰. Arsenic trioxide³¹ is approved (2000) alone or in combination with tretinoin (ATRA) for the treatment of APL harboring PML-RARA fusions. Somatic missense mutations in PML-RARA fusion including A216V, S214L, A216T, L217F, and S220G are associated with acquired resistance to treatment with arsenic trioxide³².

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

PML-RARA fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
arsenic trioxide + tretinoin	●	●	●	●	×
arsenic trioxide	●	●	●	×	×
anthracycline + arsenic trioxide + tretinoin	×	●	×	●	×
idarubicin + tretinoin	×	●	×	●	×
anthracycline + arsenic trioxide	×	●	×	×	×
arsenic trioxide + idarubicin + tretinoin	×	●	×	×	×
cytarabine + daunorubicin + tretinoin	×	●	×	×	×
gemtuzumab ozogamicin + arsenic trioxide	×	●	×	×	×
gemtuzumab ozogamicin + arsenic trioxide + tretinoin	×	●	×	×	×
gemtuzumab ozogamicin + tretinoin	×	●	×	×	×
chemotherapy	×	×	×	×	● (IV)
chemotherapy, supplement	×	×	×	×	● (IV)
gemtuzumab ozogamicin, ganetespib, chemotherapy, allogeneic stem cells, supplement	×	×	×	×	● (III)
supplement, chemotherapy	×	×	×	×	● (III)
chemotherapy, valproic acid	×	×	×	×	● (II)
T-cell therapy	×	×	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

FDA information is current as of 2020-12-16. For the most up-to-date information, search www.fda.gov.

PML-RARA fusion

☒ arsenic trioxide, arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia **Label as of:** 2020-10-20

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

Indications and usage:

TRISENOX® is an arsenical indicated:

- In combination with tretinoin for treatment of adults with newly-diagnosed low-risk acute promyelocytic leukemia (APL) whose APL is characterized by the presence of the t(15;17) translocation or PML/RAR-alpha gene expression.
- For induction of remission and consolidation in patients with APL who are refractory to, or have relapsed from, retinoid and anthracycline chemotherapy, and whose APL is characterized by the presence of the t(15;17) translocation or PML/RAR-alpha gene expression.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021248s019lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2020-12-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

PML-RARA fusion

● arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● idarubicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● anthracycline + arsenic trioxide

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● anthracycline + arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

PML-RARA fusion (continued)

● arsenic trioxide

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● arsenic trioxide + idarubicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● cytarabine + daunorubicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● gemtuzumab ozogamicin + arsenic trioxide

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

PML-RARA fusion (continued)

● gemtuzumab ozogamicin + arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention
- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● gemtuzumab ozogamicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention, Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

● idarubicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

Current EMA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

EMA information is current as of 2020-12-16. For the most up-to-date information, search www.ema.europa.eu/ema.

PML-RARA fusion

☒ arsenic trioxide, arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia **Label as of:** 2019-11-26

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/trisenox-epar-product-information_en.pdf

Current ESMO Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2020-12-01. For the most up-to-date information, search www.esmo.org.

PML-RARA fusion

● arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia **Variant class:** PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

● anthracycline + arsenic trioxide + tretinoin

Cancer type: Acute Promyelocytic Leukemia **Variant class:** PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

● idarubicin + tretinoin

Cancer type: Acute Promyelocytic Leukemia **Variant class:** PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Diagnostic Details

Current NCCN Information

NCCN information is current as of 2020-12-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

PML-RARA fusion

Diagnostic significance: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

NCCN Recommendation category: 2A

Diagnostic notes:

- WHO 2016 classification

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2021]

Diagnostic significance: Acute Myeloid Leukemia

Variant class: t(15;17)

NCCN Recommendation category: 2A

Diagnostic notes:

- Karyotypes t(8;21), inv16, and t(15;17) are considered to be Acute Myeloid Leukemia even if the marrow blast count is less than 20%

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2021]

Current ESMO Information

ESMO information is current as of 2020-12-01. For the most up-to-date information, search www.esmo.org.

PML-RARA fusion

Diagnostic significance: Acute Promyelocytic Leukemia

Variant class: PML-RARA fusion [t(15;17)(q22;q21) or t(15;17)(q24;q21)]

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Wang et al. ETV6 mutation in a cohort of 970 patients with hematologic malignancies. *Haematologica*. 2014 Oct;99(10):e176-8. PMID: 24997145
2. Wang et al. The TEL/ETV6 gene is required specifically for hematopoiesis in the bone marrow. *Genes Dev*. 1998 Aug 1;12(15):2392-402. PMID: 9694803
3. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res*. 2013 Jan;41(Database issue):D920-4. PMID: 23161685
4. Feurstein et al. Germline ETV6 mutations and predisposition to hematological malignancies. *Int. J. Hematol*. 2017 Aug;106(2):189-195. PMID: 28555414
5. Melazzini et al. Clinical and pathogenic features of ETV6-related thrombocytopenia with predisposition to acute lymphoblastic leukemia. *Haematologica*. 2016 Nov;101(11):1333-1342. PMID: 27365488
6. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2021]
7. De et al. ETV6 fusion genes in hematological malignancies: a review. *Leuk. Res*. 2012 Aug;36(8):945-61. PMID: 22578774
8. Pui et al. Acute lymphoblastic leukemia. *N. Engl. J. Med*. 2004 Apr 8;350(15):1535-48. PMID: 15071128
9. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
10. Golub et al. Fusion of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell*. 1994 Apr 22;77(2):307-16. PMID: 8168137
11. Taylor et al. Oncogenic TRK fusions are amenable to inhibition in hematologic malignancies. *J. Clin. Invest*. 2018 Aug 31;128(9):3819-3825. PMID: 29920189
12. Bejar et al. Clinical effect of point mutations in myelodysplastic syndromes. *N. Engl. J. Med*. 2011 Jun 30;364(26):2496-506. PMID: 21714648
13. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
14. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 1.2021]
15. Curtis et al. Two novel imatinib-responsive PDGFRA fusion genes in chronic eosinophilic leukaemia. *Br. J. Haematol*. 2007 Jul;138(1):77-81. PMID: 17555450
16. Curtis et al. A novel ETV6-PDGFRB fusion transcript missed by standard screening in a patient with an imatinib responsive chronic myeloproliferative disease. *Leukemia*. 2007 Aug;21(8):1839-41. Epub 2007 May 17. PMID: 17508004
17. Gancheva et al. Myeloproliferative neoplasm with ETV6-ABL1 fusion: a case report and literature review. *Mol Cytogenet*. 2013 Sep 20;6(1):39. doi: 10.1186/1755-8166-6-39. PMID: 24053143
18. Drilon et al. What hides behind the MASC: clinical response and acquired resistance to entrectinib after ETV6-NTRK3 identification in a mammary analogue secretory carcinoma (MASC). *Ann Oncol*. 2016 May;27(5):920-6. doi: 10.1093/annonc/mdw042. Epub 2016 Feb 15. PMID: 26884591
19. Nasr et al. Therapy-induced PML/RARA proteolysis and acute promyelocytic leukemia cure. *Clin. Cancer Res*. 2009 Oct 15;15(20):6321-6. PMID: 19808868
20. Laursen et al. Epigenetic regulation by RARα maintains ligand-independent transcriptional activity. *Nucleic Acids Res*. 2012 Jan;40(1):102-15. PMID: 21911359
21. Chanda et al. Retinoic acid signaling is essential for embryonic hematopoietic stem cell development. *Cell*. 2013 Sep 26;155(1):215-27. PMID: 24074870
22. Borrow et al. Molecular analysis of acute promyelocytic leukemia breakpoint cluster region on chromosome 17. *Science*. 1990 Sep 28;249(4976):1577-80. PMID: 2218500
23. Fasan et al. Molecular landscape of acute promyelocytic leukemia at diagnosis and relapse. *Haematologica*. 2017 Jun;102(6):e222-e224. PMID: 28341736
24. Subramaniam et al. Do RARA/PML fusion gene deletions confer resistance to ATRA-based therapy in patients with acute promyelocytic leukemia?. 2006 Dec;20(12):2193-5. PMID: 17008891
25. Juliane et al. FIP1L1/RARA with breakpoint at FIP1L1 intron 13: a variant translocation in acute promyelocytic leukemia. *Haematologica*. 2011 Oct; 96(10): 1565–1566. PMID: 21750086
26. Osumi et al. Acute promyelocytic leukemia with a cryptic insertion of RARA into TBL1XR1. *Genes Chromosomes Cancer*. 2019 Jul 27. PMID: 31350930
27. Cheng et al. FNDC3B is another novel partner fused to RARA in the t(3;17)(q26;q21) variant of acute promyelocytic leukemia. *Blood*. 2017 May 11;129(19):2705-2709. PMID: 28314734

References (continued)

28. Li et al. GTF2I-RARA is a novel fusion transcript in a t(7;17) variant of acute promyelocytic leukaemia with clinical resistance to retinoic acid. *Br. J. Haematol.* 2015 Mar;168(6):904-8. PMID: 25284716
29. Yin et al. Identification of a novel fusion gene, IRF2BP2-RARA, in acute promyelocytic leukemia. *J Natl Compr Canc Netw.* 2015 Jan;13(1):19-22. PMID: 25583766
30. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
31. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021248s019lbl.pdf
32. Liu et al. Varying responses of PML-RARA with different genetic mutations to arsenic trioxide. *Blood.* 2016 Jan 14;127(2):243-50. PMID: 26537301