



Sample Information

Patient Name: 吳德卿
Gender: Male
ID No.: F101709768
History No.: 47876255
Age: 68

Ordering Doctor: DOC5390F 高冠鈞
Ordering REQ.: 0BPRKQX
Signing in Date: 2021/12/17

Path No.: S110-94848
MP No.: TM21014
Assay: Oncomine Tumor Mutation Load Assay
Sample Type: FFPE
Block No.: S110-36096A
Percentage of tumor cells: 20%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Other Solid Tumor

| Table of Contents | Page | Report Highlights |
|--|------|-----------------------|
| Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) | 2 | 3 Relevant Biomarkers |
| Biomarker Descriptions | 4 | 1 Therapies Available |
| Relevant Therapy Summary | 6 | 5 Clinical Trials |
| Relevant Therapy Details | 8 | |
| Clinical Trials Summary | 15 | |
| Alert Details | 15 | |

Relevant Biomarkers

| Tier | Genomic Alteration | Relevant Therapies (In this cancer type) | Relevant Therapies (In other cancer type) | Clinical Trials |
|------|--|---|--|-----------------|
| IIC | <i>Tumor Mutational Burden</i> 8.35 Mut/Mb measured Prognostic significance: None Diagnostic significance: None | None | pembrolizumab ¹ | 3 |

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Relevant Biomarkers (continued)

| Tier | Genomic Alteration | Relevant Therapies (In this cancer type) | Relevant Therapies (In other cancer type) | Clinical Trials |
|------|---|---|--|-----------------|
| IIC | <i>NF1</i> p.(R816*) c.2446C>T neurofibromin 1 Allele Frequency: 28.85% Prognostic significance: None Diagnostic significance: None | None | None | 1 |
| IIC | <i>TP53</i> p.(R273H) c.818G>A tumor protein p53 Allele Frequency: 32.60% Prognostic significance: None Diagnostic significance: None | None | None | 1 |

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources

TSC2 p.(R1032*) c.3094C>T, *MSH2* p.(R929*) c.2785C>T, *TET2* p.(F868L) c.2604T>G, *MSH6* p.(L1081Afs*11) c.3240_3241delGT

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

| Gene | Amino Acid Change | Coding | Variant ID | Locus | Allele Frequency | Transcript | Variant Effect | Coverage |
|--------|-------------------|-------------------|------------|----------------|------------------|----------------|----------------------------|----------|
| MSH2 | p.(R929*) | c.2785C>T | . | chr2:47710068 | 35.14% | NM_000251.3 | nonsense | 1998 |
| MSH6 | p.(L1081Afs*11) | c.3240_3241delGT | . | chr2:48030625 | 26.49% | NM_000179.3 | frameshift Deletion | 1763 |
| TET2 | p.(F868L) | c.2604T>G | COSM87107 | chr4:106157703 | 65.13% | NM_001127208.2 | missense | 1606 |
| TSC2 | p.(R1032*) | c.3094C>T | . | chr16:2129160 | 30.18% | NM_000548.5 | nonsense | 1173 |
| TP53 | p.(R273H) | c.818G>A | COSM10660 | chr17:7577120 | 32.60% | NM_000546.5 | missense | 2000 |
| NF1 | p.(R816*) | c.2446C>T | . | chr17:29556079 | 28.85% | NM_001042492.3 | nonsense | 2000 |
| ARID1A | p.(Q1334dup) | c.4001_4002insGCA | . | chr1:27100181 | 24.16% | NM_006015.6 | nonframeshift Insertion | 327 |
| TAL1 | p.(P276Rfs*4) | c.826_827insG | . | chr1:47685561 | 4.00% | NM_003189.5 | frameshift Insertion | 50 |
| MSH2 | p.(T564A) | c.1690A>G | . | chr2:47698132 | 39.19% | NM_000251.3 | missense | 1998 |
| LRP1B | p.(R1464=) | c.4392A>G | . | chr2:141625346 | 65.85% | NM_018557.3 | synonymous | 1909 |
| STK36 | p.(A134T) | c.400G>A | . | chr2:219540162 | 63.90% | NM_015690.5 | missense | 1967 |
| LTF | p.(R23dup) | c.68_69insAAG | . | chr3:46501284 | 100.00% | NM_002343.6 | nonframeshift Insertion | 1966 |
| PHOX2B | p.(S286P) | c.856T>C | . | chr4:41747913 | 14.93% | NM_003924.4 | missense | 1822 |
| PDGFRA | p.(P567=) | c.1701A>G | . | chr4:55141055 | 99.90% | NM_006206.6 | synonymous | 1977 |
| NOTCH4 | p.(G534S) | c.1600G>A | . | chr6:32185796 | 14.21% | NM_004557.4 | missense | 1189 |

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.11(003).

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) (continued)

DNA Sequence Variants (continued)

| Gene | Amino Acid Change | Coding | Variant ID | Locus | Allele Frequency | Transcript | Variant Effect | Coverage |
|----------|-------------------|--------------------------|------------|-----------------|------------------|----------------|------------------------|----------|
| NOTCH4 | p.(L13_L16del) | c.36_47delIGCTGCTGCTGCT | . | chr6:32191658 | 46.77% | NM_004557.4 | nonframeshift Deletion | |
| ROS1 | p.(Y1353S) | c.4058A>C | . | chr6:117677875 | 36.18% | NM_002944.2 | missense | 1114 |
| SYNE1 | p.(D4661=) | c.13983C>T | . | chr6:152651837 | 36.37% | NM_182961.4 | synonymous | 1911 |
| SYNE1 | p.(K4121S) | c.12362_12363delAG insGT | . | chr6:152658141 | 33.17% | NM_182961.4 | missense | 1993 |
| SYNE1 | p.(K4121R) | c.12362A>G | . | chr6:152658142 | 65.43% | NM_182961.4 | missense | 1993 |
| SYNE1 | p.(M3369V) | c.10105A>G | . | chr6:152686022 | 63.93% | NM_182961.4 | missense | 1999 |
| IGF2R | p.(S1137=) | c.3411C>T | . | chr6:160482789 | 13.30% | NM_000876.3 | synonymous | 2000 |
| POT1 | p.(?) | c.-15T>G | . | chr7:124537242 | 24.16% | NM_015450.3 | unknown | 1668 |
| ADGRA2 | p.(A927T) | c.2779G>A | . | chr8:37698635 | 28.14% | NM_032777.10 | missense | 1997 |
| ADGRA2 | p.(R1277H) | c.3830G>A | . | chr8:37699686 | 5.56% | NM_032777.10 | missense | 126 |
| NBN | p.(V346M) | c.1036G>A | . | chr8:90971041 | 37.75% | NM_002485.5 | missense | 2000 |
| PTPRD | p.(A545V) | c.1634C>T | . | chr9:8507344 | 35.58% | NM_002839.4 | missense | 163 |
| PTPRD | p.(G35=) | c.105G>T | . | chr9:8636804 | 33.42% | NM_002839.4 | synonymous | 1885 |
| RET | p.(R189C) | c.565C>T | . | chr10:43598017 | 29.57% | NM_020975.6 | missense | 1197 |
| KAT6B | p.(E1101Rfs*35) | c.3301_3302delGA | . | chr10:76781916 | 99.58% | NM_012330.4 | frameshift Deletion | 1443 |
| FGFR2 | p.(I422T) | c.1265T>C | . | chr10:123274653 | 29.54% | NM_000141.5 | missense | 1997 |
| GUCY1A2 | p.(G723=) | c.2169C>T | . | chr11:106558305 | 64.96% | NM_000855.3 | synonymous | 1618 |
| ADAMTS20 | p.(Y459=) | c.1377C>T | . | chr12:43858526 | 13.28% | NM_025003.5 | synonymous | 1958 |
| HCAR1 | p.(P283=) | c.849C>T | . | chr12:123214038 | 45.58% | NM_032554.4 | synonymous | 1744 |
| EP400 | p.(E1853D) | c.5559G>T | . | chr12:132514423 | 8.35% | NM_015409.5 | missense | 1832 |
| IRS2 | p.(Q1269P) | c.3806A>C | . | chr13:110434595 | 44.80% | NM_003749.3 | missense | 433 |
| NKX2-1 | p.(A302=) | c.906G>C | . | chr14:36986783 | 16.52% | NM_001079668.3 | synonymous | 666 |
| MAP2K1 | p.(A257=) | c.771G>A | . | chr15:66777405 | 28.75% | NM_002755.4 | synonymous | 2000 |
| COL1A1 | p.(S1199G) | c.3595A>G | . | chr17:48264220 | 68.82% | NM_000088.4 | missense | 1966 |
| RNF213 | p.(R4963C) | c.14887C>T | . | chr17:78360656 | 40.65% | NM_001256071.3 | missense | 2000 |
| TCF3 | p.(S253=) | c.759C>T | . | chr19:1622116 | 99.90% | NM_001136139.4 | synonymous | 959 |
| AXL | p.(I708=) | c.2124C>T | . | chr19:41762444 | 64.20% | NM_021913.5 | synonymous | 2000 |
| ITGB2 | p.(V761=) | c.2283C>T | . | chr21:46306310 | 99.95% | NM_000211.5 | synonymous | 1999 |
| MYH9 | p.(T807=) | c.2421C>T | . | chr22:36698692 | 62.14% | NM_002473.6 | synonymous | 1474 |
| KDM6A | p.(S418I) | c.1253G>T | . | chrX:44919325 | 41.60% | NM_021140.3 | missense | 1077 |
| TFE3 | p.(A474T) | c.1420G>A | . | chrX:48887977 | 26.74% | NM_006521.6 | missense | 258 |

Biomarker Descriptions

MSH2 (mutS homolog 2)

Background: The MSH2 gene encodes the mutS homolog 2 protein¹. MSH2 is a tumor suppressor gene that heterodimerizes with MSH6 to form the MutSa complex or with MSH3 to form the MutSβ complex². Both MutS complexes function in DNA damage recognition of base-base mismatches or insertion/deletion (indels) mispairs². Specifically, the MutSa complex recognizes 1-2 nucleotide indels while MutSβ recognizes longer indel mispairs². DNA damage recognition initiates the mismatch repair (MMR) process that repairs mismatch errors which typically occur during DNA replication. Mutations in MSH2 result in the degradation of MSH6³. Loss of MSH2 protein expression correlate to mutations in the genes and are used to pre-screen colorectal cancer or endometrial hyperplasia⁴. MSH2, along with MLH1, MSH6, and PMS2 form the core components of the MMR pathway⁵. Deficiency in MMR (dMMR) is characterized by mutations and loss of expression in these genes². dMMR is associated with microsatellite instability (MSI), which is defined as a change in the length of a microsatellite in a tumor as compared to normal tissue^{5,6,7}. MSI-high (MSI-H) is a hallmark of Lynch Syndrome (LS), also known as hereditary non-polyposis colorectal cancer, which is caused by germline mutations in MMR genes^{5,8}. LS is associated with an increased risk of developing colorectal cancer, as well as other cancers, including endometrial and stomach cancer.^{6,8,9,10}. Specifically, MSH2 mutations are associated with increased risk of ovarian and pancreatic cancer¹¹.

Alterations and prevalence: Somatic mutations in MSH2 are observed in 8% of uterine corpus endometrial carcinoma, as well as 2-3% of bladder urothelial carcinoma, melanoma, and colorectal adenocarcinoma^{12,13}.

Potential relevance: Pembrolizumab (2014) is an anti-PD-1 immune checkpoint inhibitor that is approved for patients with dMMR solid tumors that have progressed on prior therapies¹⁴. Nivolumab (2015), an anti-PD-1 immune checkpoint inhibitor, is approved alone or in combination with the cytotoxic T-lymphocyte antigen 4 (CTLA-4) blocking antibody, ipilimumab (2011), for patients with dMMR colorectal cancer that have progressed on prior treatment^{15,16}.

MSH6 (mutS homolog 6)

Background: The MSH6 gene encodes the mutS homolog 6 protein¹. MSH6 is a tumor suppressor gene that heterodimerizes with MSH2 to form the MutSa complex². The MutSa complex functions in the DNA damage recognition of base-base mismatches or insertion/deletion (indels) of 1-2 nucleotides². DNA damage recognition initiates the mismatch repair (MMR) process that repairs mismatch errors which typically occur during DNA replication. Mutations in MSH2 result in the degradation of MSH6³. MSH6, along with MLH1, MSH2, and PMS2 form the core components of the MMR pathway². The MMR pathway is critical to the repair of mismatch errors which typically occur during DNA replication. Deficiency in MMR (dMMR) is characterized by mutations and loss of expression in these genes. dMMR is associated with microsatellite instability (MSI), which is defined as a change in the length of a microsatellite in a tumor as compared to normal tissue^{5,6,7}. MSI-high (MSI-H) is a hallmark of Lynch Syndrome (LS), also known as hereditary non-polyposis colorectal cancer, which is caused by germline mutations in MMR genes^{5,8}. LS is associated with an increased risk of developing colorectal cancer, as well as other cancers, including endometrial and stomach cancer^{6,8,9,10}. Specifically, MSH6 mutations are associated with increased risk of ovarian and pancreatic cancer¹¹.

Alterations and prevalence: Somatic mutations in MSH6 are observed in 11% of uterine corpus endometrial carcinoma, 4% colorectal adenocarcinoma, and 3% skin cutaneous melanoma^{12,13}.

Potential relevance: Pembrolizumab (2014) is an anti-PD-1 immune checkpoint inhibitor that is approved for patients with dMMR solid tumors that have progressed on prior therapies¹⁴. Nivolumab (2015), an anti-PD-1 immune checkpoint inhibitor, is approved alone or in combination with the cytotoxic T-lymphocyte antigen 4 (CTLA-4) blocking antibody, ipilimumab (2011), for patients with dMMR colorectal cancer that have progressed on prior treatment^{15,16}.

NF1 (neurofibromin 1)

Background: The NF1 gene encodes the neurofibromin protein, a tumor suppressor within the Ras-GTPase-activating protein (GAP) family¹⁷. NF1 regulates cellular levels of activated RAS proteins including KRAS, NRAS, and HRAS, by down regulating the active GTP-bound state to an inactive GDP-bound state^{17,18}. Inactivation of NF1 due to missense mutations results in sustained intracellular levels of RAS-GTP and prolonged activation of the RAS/RAF/MAPK and PI3K/AKT/mTOR signaling pathways leading to increased proliferation and survival¹⁷. Constitutional mutations in NF1 are associated with neurofibromatosis type 1, a RASopathy autosomal dominant tumor syndrome with predisposition to myeloid malignancies such as juvenile myelomonocytic leukemia (JMML) and myeloproliferative neoplasms (MPN)^{17,19,20}.

Alterations and prevalence: NF1 aberrations include missense mutations, insertions, indels, aberrant splicing, microdeletions, and rearrangements¹⁷. The majority of NF1 mutated tumors exhibit biallelic inactivation of NF1, supporting the 'two-hit' hypothesis of carcinogenesis^{17,21}. Somatic mutations in NF1 have been identified in over 30% of ovarian serous carcinoma, 12-30% of melanoma, 10-20% of chronic myelomonocytic leukemia (CMML), and 7% of acute myeloid leukemia (AML)^{17,20}.

Biomarker Descriptions (continued)

Potential relevance: Somatic NF1 mutations are an ancillary diagnostic criteria for malignant peripheral nerve sheath tumor (MPNST) in soft tissue sarcoma^{22,23}.

TET2 (tet methylcytosine dioxygenase 2)

Background: TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3²⁴. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to 5-hydroxymethylcytosine^{25,26}. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded β -helix domain (DSBH)²⁷. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies^{24,25,26}.

Alterations and prevalence: Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)²⁰. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies^{25,28}. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

Potential relevance: The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations^{22,29}. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia^{29,30}.

TP53 (tumor protein p53)

Background: The TP53 gene encodes the p53 tumor suppressor protein that binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis. Alterations in TP53 is required for oncogenesis as they result in loss of protein function and gain of transforming potential³¹. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers^{32,33}.

Alterations and prevalence: TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)^{12,13,34,35,36,37}. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common including substitutions at codons R158, R175, Y220, R248, R273, and R282^{12,13}. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes^{38,39,40,41}.

Potential relevance: The small molecule p53 reactivator, PC14586, received a fast track designation (2020) by the FDA for advanced tumors harboring a TP53 Y220C mutation⁴². The FDA has granted fast track designation (2019) to the p53 reactivator, eprenetapopt⁴³ and breakthrough designation⁴⁴ (2020) in combination with azacitidine or azacitidine and venetoclax for acute myeloid leukemia patients (AML) and myelodysplastic syndrome (MDS) harboring a TP53 mutation, respectively. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation^{45,46}. TP53 mutations confer poor prognosis in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), chronic lymphocytic leukemia (CLL)^{20,29,47,48}. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant⁴⁹. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system⁵⁰.

TSC2 (TSC complex subunit 2)

Background: The TSC2 gene encodes the tuberlin protein. TSC2 and TSC1 (also known as hamartin) form a complex through their respective coiled-coil domains⁵¹. The TSC1-TSC2 complex is a negative regulator of the mTOR signaling pathway that regulates cell growth, cell proliferation, and protein and lipid synthesis⁵². Specifically, the TSC1-TSC2 complex acts as a GTPase activating (GAP) protein that inhibits the G-protein RHEB and keeps it in an inactivated state (RHEB-GDP). GTP bound RHEB (RHEB-GTP) is required to activate the mTOR complex 1 (mTORC1). TSC1 and TSC2 are tumor suppressor genes. Loss of function mutations in TSC1 and TSC2 lead to dysregulation of the mTOR pathway^{51,53}. Inactivating germline mutations in TSC1 and TSC2 are associated with tuberous sclerosis complex (TSC), an autosomal dominant neurocutaneous and progressive disorder that presents with multiple benign tumors in different organs⁵¹.

Alterations and prevalence: Somatic mutations are observed in up to 8% of skin cutaneous melanoma, 7% of uterine corpus endometrial carcinoma, and 4% of cervical squamous cell carcinoma^{12,13}.

Potential relevance: Currently, no therapies are approved for TSC2 aberrations.

Biomarker Descriptions (continued)

Tumor Mutational Burden

Background: Tumor mutational burden (TMB), also known as tumor mutational load (TML), is the count of somatic mutations in the DNA of cancer cells. TMB is determined by next-generation sequencing and is expressed as the number of mutations per megabase (mut/Mb) of DNA coding sequence⁵⁴. Errors in DNA repair, including mutations in the POLE gene and in mismatch repair (MMR) genes, are associated with increased TMB^{55,56,57,58,59}. High TMB is associated with increased neo-antigen burden and has been linked to response to immune checkpoint inhibitors (ICIs) that target the cytotoxic T lymphocyte antigen-4 (CTLA4), programmed death protein 1 (PD1), and programmed death-ligand 1 (PD-L1) inhibitors^{60,61,62,63}.

Alterations and prevalence: In one study of over 100,000 tumor samples, the median TMB value was 3.6 mut/Mb although TMB values vary widely across cancers⁶⁴. Certain childhood cancers, leukemia, glioblastoma, and neuroblastoma typically have low mutation burden and median TMB values <1 mut/Mb^{61,64}. In comparison, cancers that experience genotoxic insults including skin cancer and lung cancer have higher median TMB values of approximately 10 mut/Mb^{61,64}. For example, within non-small cell lung cancer (NSCLC), higher TMB was observed in former/current smokers (10.5 mut/Mb) relative to never smokers (0.6 mut/Mb)^{61,64,65}. There is no consensus around the definition of high and low TMB that could be applied universally to all tumor types, instead multiple sources suggest that TMB status is a cancer type specific attribute^{64,66,67}. In NSCLC, several studies have suggested establishing a threshold between low and high TMB of 10 +/- 1 mut/Mb^{68,69,70,71}.

Potential relevance: ICIs stimulate a patient's own T-cells to kill tumors and have exhibited benefits in some patients. The first ICI to be approved by the FDA was ipilimumab (2011), an anti-CTLA4 antibody indicated for the treatment of metastatic melanoma. In 2014, anti-PD-1 antibodies, nivolumab (2014) and pembrolizumab (2014), were subsequently approved for the treatment of metastatic melanoma. Pembrolizumab was also approved (2014) for advanced esophageal squamous cell carcinoma. In 2020, the indication for pembrolizumab¹⁴ was expanded to include TMB-H (>= 10 mut/Mb) solid tumors that have progressed on prior therapy. Indications have been expanded for these ICIs to include several other cancer types including NSCLC, advanced renal cell carcinoma, classical Hodgkin lymphoma, recurrent or metastatic squamous cell carcinoma of the head and neck, urothelial carcinoma, microsatellite instability (MSI)-High or mismatch repair deficient (dMMR) colorectal cancer, and hepatocellular carcinoma. Atezolizumab (2016), avelumab (2017), and durvalumab (2017), that target programmed death-ligand 1 (PD-L1), were subsequently approved by the FDA. However, the predictive biomarkers that underlie the clinical benefits of these approved immunotherapies, including TMB, are under active investigation. Several published studies including the CheckMate 586 and CheckMate 817 clinical trials have concluded that high TMB was associated with improved response to FDA approved checkpoint inhibitors^{69,72,73}. In contrast, several promising previous trials failed to show an improvement in survival outcomes between high and low TMB including CheckMate 227 (ipilimumab + nivolumab vs. chemotherapy), CheckMate 026 (nivolumab vs. chemotherapy), KEYNOTE 189 (pembrolizumab vs. chemotherapy), KEYNOTE 021 (pembrolizumab vs. pembrolizumab + chemotherapy), and Lung-MAP (nivolumab + ipilimumab vs. nivolumab). In response, suggestions to combine TMB score with PD-L1 expression as a way to increase the predictive power for patient stratification have been reported⁷⁴. Nivolumab alone or in combination with ipilimumab is recommended for use in NSCLC with evidence of high TMB⁷⁵. Pembrolizumab is indicated for use in various cancer types with evidence of metastasis including Ewing sarcoma, salivary gland neoplasms, cervical cancer, uterine sarcoma, endometrial carcinoma, thyroid cancer, ovarian cancer, esophageal cancer, esophagogastric junction cancer, breast cancer, and germ cell tumors with high TMB^{76,77,78,79,80,81,82,83,84}. TMB score estimation is affected by the utilized assays, therefore efforts are underway to develop a standardized approach for score calculation with the aim to support consistent reporting of TMB values across laboratories^{85,86,87,88}.

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ① In this cancer type and other cancer types ✕ No evidence

Tumor Mutational Burden

| Relevant Therapy | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|---|-----|------|-----|------|------------------|
| pembrolizumab | ○ | ○ | ✕ | ✕ | ● (II) |
| atezolizumab | ✕ | ✕ | ✕ | ✕ | ● (II) |
| pembrolizumab, quavonlimab, favezelimab, lenvatinib | ✕ | ✕ | ✕ | ✕ | ○ (II) |

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

NF1 p.(R816*) c.2446C>T

| Relevant Therapy | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|-------------------------|-----|------|-----|------|---|
| RMC-4630, pembrolizumab | × | × | × | × | <input checked="" type="radio"/> (I/II) |

TP53 p.(R273H) c.818G>A

| Relevant Therapy | FDA | NCCN | EMA | ESMO | Clinical Trials* |
|--|-----|------|-----|------|-----------------------------|
| pembrolizumab, chemotherapy, radiation therapy | × | × | × | × | <input type="radio"/> (III) |

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2021-10-13. For the most up-to-date information, search www.fda.gov.

Tumor Mutational Burden

☐ pembrolizumab

Cancer type: Solid Tumor

Label as of: 2021-08-31

Variant class: Tumor Mutational Burden

Indications and usage:

KEYTRUDA® is a programmed death receptor-1 (PD-1)-blocking antibody indicated:

Melanoma

- for the treatment of patients with unresectable or metastatic melanoma.
- for the adjuvant treatment of patients with melanoma with involvement of lymph node(s) following complete resection.

Non-Small Cell Lung Cancer (NSCLC)

- in combination with pemetrexed and platinum chemotherapy, as first-line treatment of patients with metastatic nonsquamous NSCLC, with no EGFR or ALK genomic tumor aberrations.
- in combination with carboplatin and either paclitaxel or paclitaxel protein-bound, as first-line treatment of patients with metastatic squamous NSCLC.
- as a single agent for the first-line treatment of patients with NSCLC expressing PD-L1 [Tumor Proportion Score (TPS) \geq 1%] as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations, and is:
 - stage III where patients are not candidates for surgical resection or definitive chemoradiation, or
 - metastatic.
- as a single agent for the treatment of patients with metastatic NSCLC whose tumors express PD-L1 (TPS \geq 1%) as determined by an FDA-approved test, with disease progression on or after platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving KEYTRUDA®.

Head and Neck Squamous Cell Cancer (HNSCC)

- in combination with platinum and FU for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC.
- as a single agent for the first-line treatment of patients with metastatic or with unresectable, recurrent HNSCC whose tumors express PD-L1 [Combined Positive Score (CPS) \geq 1] as determined by an FDA-approved test.
- as a single agent for the treatment of patients with recurrent or metastatic HNSCC with disease progression on or after platinum-containing chemotherapy.

Classical Hodgkin Lymphoma (cHL)

- for the treatment of adult patients with relapsed or refractory cHL.
- for the treatment of pediatric patients with refractory cHL, or cHL that has relapsed after 2 or more lines of therapy.

Primary Mediastinal Large B-Cell Lymphoma (PMBCL)

- for the treatment of adult and pediatric patients with refractory PMBCL, or who have relapsed after 2 or more prior lines of therapy.
- Limitations of Use: KEYTRUDA® is not recommended for treatment of patients with PMBCL who require urgent cytoreductive therapy.

Urothelial Carcinoma

- for the treatment of patients with locally advanced or metastatic urothelial carcinoma who:
 - are not eligible for any platinum-containing chemotherapy, or
 - who have disease progression during or following platinum-containing chemotherapy or within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy

Tumor Mutational Burden (continued)

- for the treatment of patients with Bacillus Calmette-Guerin (BCG)-unresponsive, high-risk, non-muscle invasive bladder cancer (NMIBC) with carcinoma in situ (CIS) with or without papillary tumors who are ineligible for or have elected not to undergo cystectomy.

Microsatellite Instability-High or Mismatch Repair Deficient Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) solid tumors that have progressed following prior treatment and who have no satisfactory alternative treatment options.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with MSI-H central nervous system cancers have not been established.

Microsatellite Instability-High or Mismatch Repair Deficient Colorectal Cancer (CRC)

- for the treatment of patients with unresectable or metastatic MSI-H or dMMR colorectal cancer (CRC).

Gastric Cancer

- in combination with trastuzumab, fluoropyrimidine- and platinum-containing chemotherapy, for the first-line treatment of patients with locally advanced unresectable or metastatic HER2-positive gastric or gastroesophageal junction (GEJ) adenocarcinoma.¹
- as a single agent for the treatment of patients with recurrent locally advanced or metastatic gastric or GEJ adenocarcinoma whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by an FDA-approved test, with disease progression on or after 2 or more prior lines of therapy including fluoropyrimidine- and platinum-containing chemotherapy and if appropriate, HER2/neu-targeted therapy.¹

Esophageal Cancer

- for the treatment of patients with locally advanced or metastatic esophageal or gastroesophageal junction (GEJ) (tumors with epicenter 1 to 5 centimeters above the GEJ) carcinoma that is not amenable to surgical resection or definitive chemoradiation either:
 - in combination with platinum- and fluoropyrimidine-based chemotherapy, or
 - as a single agent after one or more prior lines of systemic therapy for patients with tumors of squamous cell histology that express PD-L1 (CPS ≥ 10) as determined by an FDA-approved test.

Cervical Cancer

- for the treatment of patients with recurrent or metastatic cervical cancer with disease progression on or after chemotherapy whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 1] as determined by an FDA-approved test.¹

Hepatocellular Carcinoma (HCC)

- for the treatment of patients with HCC who have been previously treated with sorafenib.¹

Merkel Cell Carcinoma (MCC)

- for the treatment of adult and pediatric patients with recurrent locally advanced or metastatic Merkel cell carcinoma.¹

Renal Cell Carcinoma (RCC)

- in combination with axitinib, for the first-line treatment of adult patients with advanced RCC.
- in combination with lenvatinib, for the first-line treatment of adult patients with advanced RCC.

Endometrial Carcinoma

- in combination with lenvatinib, for the treatment of patients with advanced endometrial carcinoma that is not MSI-H or dMMR, who have disease progression following prior systemic therapy in any setting and are not candidates for curative surgery or radiation.

Tumor Mutational Burden-High (TMB-H) Cancer

- for the treatment of adult and pediatric patients with unresectable or metastatic tumor mutational burden-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] solid tumors, as determined by an FDA-approved test, that have progressed following prior treatment and who have no satisfactory alternative treatment options.¹
- Limitations of Use: The safety and effectiveness of KEYTRUDA® in pediatric patients with TMB-H central nervous system cancers have not been established.

Cutaneous Squamous Cell Carcinoma (cSCC)

- for the treatment of patients with recurrent or metastatic cSCC or locally advanced cSCC that is not curable by surgery or radiation.

Triple-Negative Breast Cancer (TNBC)

Tumor Mutational Burden (continued)

- for the treatment of patients with high-risk early-stage TNBC in combination with chemotherapy as neoadjuvant treatment, and then continued as a single agent as adjuvant treatment after surgery.
- in combination with chemotherapy, for the treatment of patients with locally recurrent unresectable or metastatic TNBC whose tumors express PD-L1 [Combined Positive Score (CPS) ≥ 10] as determined by an FDA approved test.

Adult Indications: Additional Dosing Regimen of 400 mg Every 6 Weeks

- for use at an additional recommended dosage of 400 mg every 6 weeks for all approved adult indications.²

¹This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

²This indication is approved under accelerated approval based on pharmacokinetic data, the relationship of exposure to efficacy, and the relationship of exposure to safety. Continued approval for this dosing may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125514s117,s118lbl.pdf

Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2021-10-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

Tumor Mutational Burden

☐ pembrolizumab

Cancer type: Chondrosarcoma, Osteosarcoma Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2022]

☐ pembrolizumab

Cancer type: Breast Cancer Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IV; Invasive, Unresectable, Metastatic, Progression (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 8.2021]

☐ pembrolizumab

Cancer type: Cervical Cancer Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Metastatic, Progression (Second-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2021]

☐ pembrolizumab

Cancer type: Esophageal Cancer,
Gastroesophageal Junction Adenocarcinoma Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Squamous Cell; Unresectable, Locally Advanced, Recurrent, Metastatic (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 4.2021]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Gastric Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Locally Advanced, Recurrent, Metastatic (Second-line therapy, Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Gastric Cancer [Version 4.2021]

○ pembrolizumab

Cancer type: Head and Neck Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Nasopharyngeal; Recurrent, Unresectable, Metastatic (Subsequent therapy); Useful in certain circumstances
- Salivary Gland Neoplasm; Recurrent, Unresectable, Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 3.2021]

○ pembrolizumab

Cancer type: Extrahepatic Cholangiocarcinoma,
Gallbladder Carcinoma, Intrahepatic
Cholangiocarcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Hepatobiliary Cancers [Version 5.2021]

○ pembrolizumab

Cancer type: Ovarian Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Epithelial, Less Common Ovarian Cancers, Fallopian Tube, Primary Peritoneal; Recurrent (Recurrence therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Ovarian Cancer [Version 3.2021]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Castration-Resistant Prostate Cancer **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma; Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 1.2022]

○ pembrolizumab

Cancer type: Testicular Cancer

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Germ Cell Tumor; Metastatic (Third-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Testicular Cancer [Version 2.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Follicular Carcinoma, Thyroid Gland Hurthle Cell Carcinoma, Thyroid Gland Papillary Carcinoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Locally Recurrent, Advanced, Metastatic (Line of therapy not specified); Consider

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Medullary Carcinoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Recurrent, Persistent, Local, Distant Metastases, Regional (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2021]

○ pembrolizumab

Cancer type: Thyroid Gland Anaplastic Carcinoma **Variant class:** Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Stage IVC; Metastatic (Second-line therapy); Useful in certain circumstances, Consider

Reference: NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2021]

Tumor Mutational Burden (continued)

○ pembrolizumab

Cancer type: Endometrial Carcinoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Second-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 4.2021]

○ pembrolizumab

Cancer type: Uterine Sarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Unresectable, Metastatic, Progression (Second-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 4.2021]

○ pembrolizumab

Cancer type: Ewing Sarcoma

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Metastatic (Line of therapy not specified); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2022]

○ pembrolizumab

Cancer type: Castration-Resistant Prostate Cancer Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- Adenocarcinoma; Visceral Metastases, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Prostate Cancer [Version 1.2022]

Clinical Trials in Taiwan region:

Clinical Trials Summary

Tumor Mutational Burden

| NCT ID | Title | Phase |
|-------------|---|-------|
| NCT04589845 | Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial | II |
| NCT02628067 | A Clinical Trial of Pembrolizumab (MK-3475) Evaluating Predictive Biomarkers in Subjects With Advanced Solid Tumors (KEYNOTE 158). | II |
| NCT03516981 | A Phase II Precision Oncology Study of Biomarker-Directed, Pembrolizumab-(MK-3475, SCH 900475) Based Combination Therapy for Advanced Non-Small Cell Lung Cancer (KEYNOTE-495; KeyImPaCT) | II |

NF1 p.(R816*) c.2446C>T

| NCT ID | Title | Phase |
|-------------|--|-------|
| NCT04418661 | A Phase 1/2, Open-label, Multicenter, Dose Escalation and Dose Expansion Study of SAR442720 in Combination With Pembrolizumab in Patients With Advanced Malignancies | I/II |

TP53 p.(R273H) c.818G>A

| NCT ID | Title | Phase |
|-------------|--|-------|
| NCT04634877 | A Phase III, Randomized, Double-Blind Study of Pembrolizumab Versus Placebo in Combination With Adjuvant Chemotherapy With or Without Radiotherapy for the Treatment of Newly Diagnosed High-Risk Endometrial Cancer After Surgery With Curative Intent (KEYNOTE-B21/ ENGOT-en11 / GOG-3053) | III |

Alerts Informed By Public Data Sources

Current NCCN Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

NCCN information is current as of 2021-10-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

Tumor Mutational Burden

pembrolizumab

Cancer type: Giant Cell Tumor of Soft Tissue

Variant class: Tumor Mutational Burden

NCCN Recommendation category: 2A

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "NCCN does not recommend this systemic treatment for GCTB since it is not technically a malignant tumor."

Reference: NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2022]

Current ESMO Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

ESMO information is current as of 2021-10-01. For the most up-to-date information, search www.esmo.org.

NF1 p.(R816*) c.2446C>T

imatinib

Cancer type: Gastrointestinal Stromal Tumor

Variant class: NF1 mutation

Other criteria: SDH underexpression

ESMO Level of Evidence/Grade of Recommendation: IV / D

Summary:

ESMO™ Clinical Practice Guidelines include the following supporting statement:

- "Adjuvant treatment should be avoided in NF1-related and SDH expression negative GISTs [IV, D]."

Reference: ESMO Clinical Practice Guidelines - ESMO-EUROCAN-Gastrointestinal Stromal Tumours [Annals of Oncology (2021), doi: <https://doi.org/10.1016/j.annonc.2021.09.005>.]

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Li. Mechanisms and functions of DNA mismatch repair. *Cell Res.* 2008 Jan;18(1):85-98. PMID: 18157157
3. Zhao et al. Mismatch Repair Deficiency/Microsatellite Instability-High as a Predictor for anti-PD-1/PD-L1 Immunotherapy Efficacy. *J Hematol Oncol.* 12(1),54. PMID: 31151482
4. Berends et al. MLH1 and MSH2 protein expression as a pre-screening marker in hereditary and non-hereditary endometrial hyperplasia and cancer. *Int. J. Cancer.* 2001 May 1;92(3):398-403. PMID: 11291077
5. Lynch et al. Review of the Lynch syndrome: history, molecular genetics, screening, differential diagnosis, and medicolegal ramifications. *Clin. Genet.* 2009 Jul;76(1):1-18. PMID: 19659756
6. Baudrin et al. Molecular and Computational Methods for the Detection of Microsatellite Instability in Cancer. *Front Oncol.* 2018 Dec 12;8:621. doi: 10.3389/fonc.2018.00621. eCollection 2018. PMID: 30631754
7. Saeed et al. Microsatellites in Pursuit of Microbial Genome Evolution. *Front Microbiol.* 2016 Jan 5;6:1462. doi: 10.3389/fmicb.2015.01462. eCollection 2015. PMID: 26779133
8. Nojadeh et al. Microsatellite instability in colorectal cancer. *EXCLI J.* 2018;17:159-168. PMID: 29743854
9. Imai et al. Carcinogenesis and microsatellite instability: the interrelationship between genetics and epigenetics. *Carcinogenesis.* 2008 Apr;29(4):673-80. PMID: 17942460
10. Latham et al. Microsatellite Instability Is Associated With the Presence of Lynch Syndrome Pan-Cancer. *J. Clin. Oncol.* 2019 Feb 1;37(4):286-295. PMID: 30376427
11. NCCN-Genetic/Familial High-Risk Assessment:Breast, Ovarian, and Pancreatic. NCCN-Genetic/Familial High-Risk Assessment:Breast, Ovarian, and Pancreatic
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
13. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
14. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125514s117,s118lbl.pdf
15. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125554s097bledt.pdf
16. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125377s121lbl.pdf
17. Philpott et al. The NF1 somatic mutational landscape in sporadic human cancers. 2017 Jun 21;11(1):13. doi: 10.1186/s40246-017-0109-3. PMID: 28637487
18. Scheffzek et al. The Ras-RasGAP complex: structural basis for GTPase activation and its loss in oncogenic Ras mutants. *Science.* 1997 Jul 18;277(5324):333-8. PMID: 9219684
19. Fioretos et al. Isochromosome 17q in blast crisis of chronic myeloid leukemia and in other hematologic malignancies is the result of clustered breakpoints in 17p11 and is not associated with coding TP53 mutations. *Blood.* 1999 Jul 1;94(1):225-32. PMID: 10381517
20. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2021]
21. Brems et al. Mechanisms in the pathogenesis of malignant tumours in neurofibromatosis type 1. *Lancet Oncol.* 2009 May;10(5):508-15. PMID: 19410195
22. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
23. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2021]
24. Pan et al. The TET2 interactors and their links to hematological malignancies. *IUBMB Life.* 2015 Jun;67(6):438-45. PMID: 26099018
25. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature.* 2010 Dec 9;468(7325):839-43. PMID: 21057493
26. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. *Leukemia.* 2014 Mar;28(3):485-96. PMID: 24220273
27. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. *Exp. Mol. Med.* 2017 Apr 28;49(4):e323. PMID: 28450733
28. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood.* 2009 Oct 8;114(15):3285-91. PMID: 19666869
29. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2021]

References (continued)

30. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood*. 2014 Apr 3;123(14):2220-8. PMID: 24478400
31. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell*. 2014 Mar 17;25(3):304-17. PMID: 24651012
32. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol*. 2010 Jan;2(1):a001008. PMID: 20182602
33. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med*. 2017 Apr 3;7(4). PMID: 28270529
34. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature*. 2012 Sep 27;489(7417):519-25. PMID: 22960745
35. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
36. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet*. 2016 Jun;48(6):607-16. PMID: 27158780
37. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature*. 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
38. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat*. 2002 Jun;19(6):607-14. PMID: 12007217
39. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer*. 2011 Apr;2(4):466-74. PMID: 21779514
40. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene*. 2007 Apr 2;26(15):2157-65. PMID: 17401424
41. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat*. 2014 Jun;35(6):766-78. PMID: 24729566
42. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
43. <https://ir.aprea.com//news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>
44. <http://vp280.alertir.com/en/pressreleases/karolinska-development%27s-portfolio-company-aprea-therapeutics-receives-fda-breakthrough-therapy-designation-1769167>
45. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol*. 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
46. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci*. 2017 Nov;74(22):4171-4187. PMID: 28643165
47. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2021]
48. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 1.2022]
49. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 5.2021]
50. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med*. 2020 Aug 3. PMID: 32747829
51. Rosset et al. TSC1 and TSC2 gene mutations and their implications for treatment in Tuberous Sclerosis Complex: a review. *Genet Mol Biol*. 2017 Jan-Mar;40(1):69-79. PMID: 28222202
52. Henske et al. Tuberous sclerosis complex. *Nat Rev Dis Primers*. 2016 May 26;2:16035. PMID: 27226234
53. Santiago et al. Identification of regions critical for the integrity of the TSC1-TSC2-TBC1D7 complex. *PLoS ONE*. 2014;9(4):e93940. PMID: 24714658
54. Yuan et al. Novel technologies and emerging biomarkers for personalized cancer immunotherapy. 4:3. PMID: 26788324
55. Temko et al. Somatic POLE exonuclease domain mutations are early events in sporadic endometrial and colorectal carcinogenesis, determining driver mutational landscape, clonal neoantigen burden and immune response. *J. Pathol*. 2018 Jul;245(3):283-296. PMID: 29604063
56. Mehnert et al. Immune activation and response to pembrolizumab in POLE-mutant endometrial cancer. 126(6):2334-40. PMID: 27159395

References (continued)

57. Bouffet et al. Immune Checkpoint Inhibition for Hypermutant Glioblastoma Multiforme Resulting From Germline Biallelic Mismatch Repair Deficiency. *J. Clin. Oncol.* 2016 Jul 1;34(19):2206-11. PMID: 27001570
58. Humphris et al. Hypermutation In Pancreatic Cancer. *Gastroenterology.* 2017 Jan;152(1):68-74.e2. PMID: 27856273
59. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
60. Snyder et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N. Engl. J. Med.* 2014 Dec 4;371(23):2189-2199. PMID: 25409260
61. Alexandrov et al. Signatures of mutational processes in human cancer. *Nature.* 2013 Aug 22;500(7463):415-21. PMID: 23945592
62. Rizvi et al. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science.* 2015 Apr 3;348(6230):124-8. PMID: 25765070
63. Van et al. Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science.* 2015 Oct 9;350(6257):207-211. PMID: 26359337
64. Chalmers et al. Analysis of 100,000 human cancer genomes reveals the landscape of tumor mutational burden. *9:34.* PMID: 28420421
65. Govindan et al. Genomic landscape of non-small cell lung cancer in smokers and never-smokers. *Cell.* 2012 Sep 14;150(6):1121-34. PMID:22980976
66. Endris et al. Measurement of tumor mutational burden (TMB) in routine molecular diagnostics: in silico and real-life analysis of three larger gene panels. *Int. J. Cancer.* 2019 May 1;144(9):2303-2312. PMID: 30446996
67. Meléndez et al. Methods of measurement for tumor mutational burden in tumor tissue. *Transl Lung Cancer Res.* 2018 Dec;7(6):661-667. PMID: 30505710
68. Hellmann et al. Nivolumab plus Ipilimumab in Lung Cancer with a High Tumor Mutational Burden. *N. Engl. J. Med.* 2018 May 31;378(22):2093-2104. PMID: 29658845
69. Ready et al. First-Line Nivolumab Plus Ipilimumab in Advanced Non-Small-Cell Lung Cancer (CheckMate 568): Outcomes by Programmed Death Ligand 1 and Tumor Mutational Burden as Biomarkers. *J. Clin. Oncol.* 2019 Apr 20;37(12):992-1000. PMID: 30785829
70. Alborelli et al. Tumor mutational burden assessed by targeted NGS predicts clinical benefit from immune checkpoint inhibitors in non-small cell lung cancer. *J. Pathol.* 2020 Jan;250(1):19-29. PMID: 31471895
71. Heeke et al. In-house Implementation of Tumor Mutational Burden Testing to Predict Durable Clinical Benefit in Non-small Cell Lung Cancer and Melanoma Patients. *Cancers (Basel).* 2019 Aug 29;11(9). PMID: 31470674
72. F. et al. CheckMate 817: First-Line Nivolumab + Ipilimumab in Patients with ECOG PS 2 and Other Special Populations with Advanced NSCLC. Volume 14, Issue 10, Supplement, Pages S214–S215. Abstract OA04.02
73. Zhu et al. Association Between Tumor Mutation Burden (TMB) and Outcomes of Cancer Patients Treated With PD-1/PD-L1 Inhibitions: A Meta-Analysis. *Front Oncol.* 9:1161, 04 Nov 2019. PMID: 31258479
74. Castellanos et al. Tumor mutational burden (TMB) and PD-L1 expression as predictors of response to immunotherapy (IO) in NSCLC. *Journal of Clinical Oncology* 37, no. 15_suppl (May 20, 2019) 2630-2630. Abstract 2630
75. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 6.2021]
76. NCCN Guidelines® - NCCN-Bone Cancer [Version 1.2022]
77. NCCN Guidelines® - NCCN-Head and Neck Cancers [Version 3.2021]
78. NCCN Guidelines® - NCCN-Testicular Cancer [Version 2.2021]
79. NCCN Guidelines® - NCCN-Cervical Cancer [Version 1.2021]
80. NCCN Guidelines® - NCCN-Uterine Neoplasms [Version 4.2021]
81. NCCN Guidelines® - NCCN-Thyroid Carcinoma [Version 2.2021]
82. NCCN Guidelines® - NCCN-Esophageal and Esophagogastric Junction Cancers [Version 4.2021]
83. NCCN Guidelines® - NCCN-Ovarian Cancer [Version 3.2021]
84. NCCN Guidelines® - NCCN-Breast Cancer [Version 8.2021]
85. <https://www.focr.org/tmb>
86. <http://www.iqnpa.org/category/tmb>
87. Stenzinger et al. Tumor mutational burden standardization initiatives: Recommendations for consistent tumor mutational burden assessment in clinical samples to guide immunotherapy treatment decisions. *Genes Chromosomes Cancer.* 2019 Aug;58(8):578-588. PMID: 30664300

References (continued)

88. Merino et al. Establishing guidelines to harmonize tumor mutational burden (TMB): in silico assessment of variation in TMB quantification across diagnostic platforms: phase I of the Friends of Cancer Research TMB Harmonization Project. *J Immunother Cancer*. 2020 Mar;8(1). PMID: 32217756