



Sample Information

Patient Name: 張瑞勳
Gender: Male
ID No.: K120885036
History No.: 17821544
Age: 64

Ordering Doctor: DOC1878G 沈佳儀
Ordering REQ.: 0BJGZSN
Signing in Date: 2021/07/28

Path No.: S110-99166
MP No.: F21062
Assay: Oncomine Focus Assay
Sample Type: FFPE
Block No.: S110-18979A+B
Percentage of tumor cells: 30%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	4
Clinical Trials Summary	5
Alert Details	5

Report Highlights
1 Relevant Biomarkers
1 Therapies Available
6 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	EGFR exon 20 insertion	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	EGFR exon 20 insertion epidermal growth factor receptor Allele Frequency: 17.53%	amivantamab ¹	None	6

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

 Alerts informed by public data sources:  Contraindicated,  Resistance

EGFR exon 20 insertion  **gefitinib**²
 afatinib, dacomitinib, erlotinib, gefitinib

Public data sources included in alerts: FDA¹, NCCN, EMA², ESMO

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
EGFR	p.(A767_V769dup)	c.2308_2309insCCA GCGTGG	COSM12376	chr7:55248998	17.53%	NM_005228.5	nonframeshift Insertion	1945
MTOR	p.(A1792=)	c.5376G>A	.	chr1:11190823	48.35%	NM_004958.4	synonymous	2000

Biomarker Descriptions

EGFR (epidermal growth factor receptor)

Background: The EGFR gene encodes the epidermal growth factor receptor (EGFR) tyrosine kinase, a member of the ERBB/human epidermal growth factor receptor (HER) family. In addition to EGFR/ERBB1/HER1, other members of the ERBB/HER family include ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4¹. EGFR ligand induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways. Activation of these pathways promote cell proliferation, differentiation, and survival^{2,3}.

Alterations and prevalence: Recurrent somatic mutations in the tyrosine kinase domain (TKD) of EGFR are observed in approximately 10-20% of lung adenocarcinoma, and at higher frequencies in never-smoker, female, and Asian populations^{4,5,6,7}. The most common mutations occur near the ATP-binding pocket of the TKD and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21⁸. These mutations constitutively activate EGFR resulting in downstream signaling, and represent 80% of the EGFR mutations observed in lung cancer. A second group of less prevalent activating mutations include E709K, G719X, S768I, L861Q, and short in-frame insertion mutations in exon 20^{9,10,11,12}. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations¹³. In contrast, a different set of recurrent activating EGFR mutations in the extracellular domain include R108K, A289V and G598V and are primarily observed in glioblastoma^{8,14}. Amplification of EGFR is observed in several cancer types including 30% of glioblastoma, 12% of esophageal cancer, 10% of head and neck cancer, 5% of bladder cancer, and 5% of lung squamous cell carcinoma^{5,6,7,14,15}. Deletion of exons 2-7, encoding the extracellular domain of EGFR (EGFRvIII), results in overexpression of a ligand-independent constitutively active protein and is observed in approximately 30% of glioblastoma^{16,17,18}.

Potential relevance: Approved first-generation EGFR tyrosine kinase inhibitors (TKIs) include erlotinib¹⁹ (2004) and gefitinib²⁰ (2015), which block the activation of downstream signaling by reversible interaction with the ATP-binding site. Although initially approved for advanced lung cancer, the discovery that drug sensitivity was associated with exon 19 and exon 21 activating mutations allowed first-generation TKIs to become subsequently approved for front-line therapy in lung cancer tumors containing exon 19 or exon 21 activating mutations. Second-generation TKIs afatinib²¹ (2013) and dacomitinib²² (2018) bind EGFR and other ERBB/HER gene family members irreversibly and were subsequently approved. First- and second-generation TKIs afatinib, dacomitinib, erlotinib, and gefitinib are recommended for the treatment NSCLC harboring EGFR exon 19 insertions, exon 19 deletions, point mutations L861Q, L858R, S768I, and codon 719 mutations, whereas most EGFR exon 20 insertions, except p.A763_Y764insFQEA, confer resistance to the same

Biomarker Descriptions (continued)

therapies^{23,24,25,26}. In lung cancer containing EGFR exon 19 or 21 activating mutations, treatment with TKIs is eventually associated with the emergence of drug resistance²⁷. The primary resistance mutation that emerges following treatment with first-generation TKI is T790M, accounting for 50-60% of resistant cases⁸. Third generation TKIs were developed to maintain sensitivity in the presence of T790M. Osimertinib²⁸ (2015) is an irreversible inhibitor indicated for metastatic EGFR T790M positive lung cancer and for the first-line treatment of metastatic NSCLC containing EGFR exon 19 deletions or exon 21 L858R mutations. Like first-generation TKIs, treatment with osimertinib is associated with acquired resistance. In this case, resistance is associated with the C797S mutation, and occurs in 22-44% of cases²⁷. The T790M and C797S mutations may be each selected following sequential treatment with a first-generation TKI followed by a third-generation TKI or vice versa²⁹. T790M and C797S can occur in either cis or trans allelic orientation²⁹. If C797S is observed following progression after treatment with a third-generation TKI in the first-line setting, sensitivity may be retained to first-generation TKIs²⁹. If C797S co-occurs in trans with T790M following sequential treatment with first- and third-generation TKIs, patients may exhibit sensitivity to combination first- and third-generation TKIs, but resistance to third-generation TKIs alone^{29,30}. However, C797S occurring in cis conformation with T790M, confers resistance to first- and third-generation TKIs²⁹. Fourth-generation TKIs are in development to overcome acquired C797S and T790M resistance mutations after osimertinib treatment. EGFR targeting antibodies including cetuximab (2004), panitumumab (2006), and necitumumab (2016) are under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations. The bispecific antibody, JNJ-61186372³¹, targeting EGFR and MET, and the TKI mobocertinib³², each received a breakthrough designation from the FDA (2020) for NSCLC tumors harboring EGFR exon 20 insertion mutations. The OncoPrex immunogene therapy CNVN-202³³ in combination with osimertinib received a fast track designation from the FDA (2020) for NSCLC tumors harboring EGFR mutations that progressed on osimertinib alone. BDTX-189³⁴ was granted a fast track designation (2020) for the treatment of solid tumors harboring an EGFR exon 20 insertion mutation.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

EGFR exon 20 insertion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
amivantamab	●	×	×	×	● (I)
durvalumab, chemotherapy	×	×	×	×	● (III)
mobocertinib	×	×	×	×	● (III)
DZD-9008	×	×	×	×	● (I/II)
TPC-064	×	×	×	×	● (I/II)
lazertinib, amivantamab	×	×	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type ☐ In other cancer type ☐ In this cancer type and other cancer types

FDA information is current as of 2021-06-09. For the most up-to-date information, search www.fda.gov.

EGFR exon 20 insertion

☒ amivantamab

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-05-21

Variant class: EGFR exon 20 insertion

Indications and usage:

RYBREVENT™ is a bispecific EGF receptor-directed and MET receptor directed antibody indicated for the treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 20 insertion mutations, as detected by an FDA-approved test, whose disease has progressed on or after platinum-based chemotherapy.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/761210s000lbl.pdf

Clinical Trials in Taiwan region:

Clinical Trials Summary

EGFR exon 20 insertion

NCT ID	Title	Phase
NCT04129502	A Randomized Phase III Multicenter Open-label Study to Compare the Efficacy of TAK-788 as First-line Treatment Versus Platinum-Based Chemotherapy in Patients With Non-Small Cell Lung Cancer With EGFR Exon 20 Insertion Mutations	III
NCT03974022	A Phase I/II, Open-Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumor Efficacy of DZD9008 in Patients With Advanced Non-Small Cell Lung Cancer (NSCLC) With EGFR or HER2 Mutation	I/II
NCT04036682	A Phase I/IIa, Open-Label, Multi-Center Trial To Assess Safety, Tolerability, Pharmacokinetics, Pharmacodynamics, And Efficacy Of CLN-081 In Patients With Non-Small Cell Lung Cancer Harboring EGFR Exon 20 Insertion Mutations	I/II
NCT02609776	A Phase I, First-in-Human, Open-Label, Dose Escalation Study of JNJ-61186372, a Human Bispecific EGFR and cMet Antibody, in Subjects With Advanced Non-Small Cell Lung Cancer.	I
NCT04077463	An Open-label Phase I/Ib Study to Evaluate the Safety and Pharmacokinetics of JNJ-73841937 (Lazertinib), a Third Generation EGFR-TKI, as Monotherapy or in Combinations With JNJ-61186372, a Human Bispecific EGFR and cMet Antibody in Participants With Advanced Non-Small Cell Lung Cancer	I
NCT03800134	A Phase III, Double-blind, Placebo-controlled, Multi-center International Study of Neoadjuvant/Adjuvant Durvalumab for the Treatment of Patients With Resectable Stages II and III Non-small Cell Lung Cancer (AEGEAN)	III

Alerts Informed By Public Data Sources

Current FDA Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

FDA information is current as of 2021-06-09. For the most up-to-date information, search www.fda.gov.

EGFR exon 20 insertion

mobocertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 20 insertion

Supporting Statement:

The FDA has granted Breakthrough Therapy Designation to the small molecule inhibitor, mobocertinib, for EGFR exon 20 insertion mutant non-small cell lung cancer (NSCLC) that has progressed on or after platinum-based chemotherapy.

Reference:

<https://www.takeda.com/newsroom/newsreleases/2020/takeda-announces-u.s.-fda-breakthrough-therapy-designation-for-mobocertinib-tak-788-for-the-treatment-of-nsclc-patients-with-egfr-exon-20-insertion-mutations/>

EGFR exon 20 insertion (continued)

BDTX-189

Cancer type: Solid Tumor

Variant class: EGFR exon 20 insertion

Supporting Statement:

The FDA has granted Fast Track Designation to BDTX-189 for solid tumors harboring a HER2 mutation or an EGFR or HER2 exon 20 insertion after progression on prior therapy.

Reference:

<https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>

osimertinib + quaratusugene ozeplasmid

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation


Supporting Statement:

The FDA has granted Fast Track Designation to the immunogene therapy, quaratusugene ozeplasmid, in combination with EGFR inhibitor osimertinib for the treatment of non-small cell lung cancer (NSCLC) with EGFR mutations that progressed after treatment with osimertinib alone.

Reference:


<https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>

Current NCCN Information

 Contraindicated

 Not recommended

 Resistance

 Breakthrough

 Fast Track

NCCN information is current as of 2021-06-01. For the most up-to-date information, search www.nccn.org. For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

EGFR exon 20 insertion

atezolizumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

nivolumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

EGFR exon 20 insertion (continued)

pembrolizumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

afatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 20 insertion

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with EGFR exon 20 insertion mutations are usually resistant to erlotinib, gefitinib, afatinib, or dacomitinib, although there are rare exceptions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

dacomitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 20 insertion

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with EGFR exon 20 insertion mutations are usually resistant to erlotinib, gefitinib, afatinib, or dacomitinib, although there are rare exceptions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 20 insertion

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with EGFR exon 20 insertion mutations are usually resistant to erlotinib, gefitinib, afatinib, or dacomitinib, although there are rare exceptions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR exon 20 insertion

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with EGFR exon 20 insertion mutations are usually resistant to erlotinib, gefitinib, afatinib, or dacomitinib, although there are rare exceptions."

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]

Current EMA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

EMA information is current as of 2021-06-09. For the most up-to-date information, search www.ema.europa.eu/ema.

EGFR exon 20 insertion

gefitinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-03-05

Variant class: EGFR exon 20 insertion

Reference:

https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information_en.pdf

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985 Sep 6;229(4717):974-6. PMID: 2992089
2. Zhixiang. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
3. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
4. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
5. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
9. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
10. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
11. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
12. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
13. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
14. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
15. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
16. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469
17. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene.* 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
18. Gan et al. The EGFRvIII variant in glioblastoma multiforme. *J Clin Neurosci.* 2009 Jun;16(6):748-54. PMID: 19324552
19. https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf
20. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/206995s004lbl.pdf
21. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/201292s015lbl.pdf
22. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/211288s003lbl.pdf
23. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 4.2021]
24. Naidoo et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer.* 2015 Sep 15;121(18):3212-3220. PMID: 26096453
25. Vyse et al. Targeting EGFR exon 20 insertion mutations in non-small cell lung cancer. *Signal Transduct Target Ther.* 2019;4:5. PMID: 30854234
26. Yi et al. A comparison of epidermal growth factor receptor mutation testing methods in different tissue types in non-small cell lung cancer. *Int J Mol Med.* 2014 Aug;34(2):464-74. PMID: 24891042
27. Madic et al. EGFR C797S, EGFR T790M and EGFR sensitizing mutations in non-small cell lung cancer revealed by six-color crystal digital PCR. *Oncotarget.* 2018 Dec 21;9(100):37393-37406. PMID: 30647840
28. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208065s021lbl.pdf
29. Niederst et al. The Allelic Context of the C797S Mutation Acquired upon Treatment with Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin. Cancer Res.* 2015 Sep 1;21(17):3924-33. PMID: 25964297

References (continued)

30. Wang et al. Lung Adenocarcinoma Harboring EGFR T790M and In Trans C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol*. 2017 Nov;12(11):1723-1727. PMID: 28662863
31. <https://www.jnj.com/janssen-announces-u-s-fda-breakthrough-therapy-designation-granted-for-jnj-6372-for-the-treatment-of-non-small-cell-lung-cancer>
32. <https://www.takeda.com/newsroom/newsreleases/2020/takeda-announces-u.s.-fda-breakthrough-therapy-designation-for-mobocertinib-tak-788-for-the-treatment-of-nscl-patients-with-egfr-exon-20-insertion-mutations/>
33. <https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>
34. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>