

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Head and neck adenocarcinoma	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Head and Neck
	NAME Chen, Wei-Cheng		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S110-38254B (PF22053)
	DATE OF BIRTH 02 January 1966		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Male		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 29 October 2021
	MEDICAL RECORD # 36421600		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 18 April 2022

Biomarker Findings

Tumor Mutational Burden - 18 Muts/Mb

Microsatellite status - MS-Stable

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

PIK3CA E726K

APC S2223*

CCND1 amplification

CDK6 amplification - equivocal[†]

MTAP loss exons 6-8

CDKN2A/B CDKN2B loss, CDKN2A loss

EPHB4 amplification - equivocal[†]

FGF19 amplification

FGF3 amplification

FGF4 amplification

NOTCH1 P1443fs*36

TP53 R273H

[†] See About the Test in appendix for details.

Report Highlights

- Targeted therapies with potential clinical benefit **approved in this patient's tumor type**: Dostarlimab (p. 12), Pembrolizumab (p. 12)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 17)

BIOMARKER FINDINGS

Tumor Mutational Burden - 18 Muts/Mb

10 Trials see p. 17

Microsatellite status - MS-Stable

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

Dostarlimab

Pembrolizumab

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Atezolizumab

Avelumab

Cemiplimab

Durvalumab

Nivolumab

Nivolumab +
Ipilimumab

No therapies or clinical trials. see Biomarker Findings section

GENOMIC FINDINGS	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
PIK3CA - E726K	none	Everolimus
10 Trials see p. 24		Temsirolimus
APC - S2223*	none	none
3 Trials see p. 19		
CCND1 - amplification	none	none
5 Trials see p. 20		
CDK6 - amplification - equivocal	none	none
9 Trials see p. 21		
MTAP - loss exons 6-8	none	none
1 Trial see p. 23		

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CDKN2A/B - CDKN2B loss, CDKN2A loss	p. 7	FGF4 - amplification	p. 9
EPHB4 - amplification - equivocal	p. 8	NOTCH1 - P1443fs*36	p. 10
FGF19 - amplification	p. 8	TP53 - R273H	p. 11
FGF3 - amplification	p. 9		

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

ORDERED TEST # ORD-1346479-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

18 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1 therapies¹⁻⁴, and combination nivolumab and ipilimumab⁵⁻¹⁰. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors^{1-4,11}. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment for patients with 9 types of advanced tumors¹. Analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with chemotherapy¹² or those with lower TMB treated

with PD-1 or PD-L1-targeting agents². However, the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors found significant improvement in ORR for patients with TMB ≥ 10 Muts/Mb (based on this assay or others) compared to those with TMB < 10 Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials^{4,11}. Together, these studies suggest that patients with TMB ≥ 10 Muts/Mb may derive clinical benefit from PD-1 or PD-L1 inhibitors. In the first-line setting for patients with advanced nasopharyngeal carcinoma (NPC), the Phase 3 JUPITER-02 study of the anti-PD-1 antibody toripalimab in combination with chemotherapy reported significantly improved median PFS relative to chemotherapy plus placebo (11.7 vs. 8.0 months, HR=0.52); median PFS improved regardless of PD-L1 status¹³. For patients with previously-treated NPC, the Phase 2 POLARIS-02 study of single-agent toripalimab reported a 20% (39/190) ORR, a 12.8-month median duration of response, a 1.9-month median PFS, and a 17.4-month median OS; activity and efficacy was not associated with either PD-L1 status or tumor mutational burden¹⁴.

FREQUENCY & PROGNOSIS

Head and neck carcinoma, including squamous cell carcinoma and adenocarcinoma, harbors a

median TMB of 3.8 mutations per megabase (mut/Mb), with a higher median TMB of 5 mut/Mb in head and neck squamous cell carcinoma (HNSCC) specifically, and 5.8% of cases [10.1% of HNSCC cases specifically] have high TMB (> 20 mut/Mb)¹⁵. Published data investigating the prognostic implications of TMB in nasopharyngeal and sinonasal carcinoma are limited (PubMed, Oct 2021).

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma¹⁶⁻¹⁷ and cigarette smoke in lung cancer¹⁸⁻¹⁹, treatment with temozolomide-based chemotherapy in glioma²⁰⁻²¹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes²²⁻²⁶, and microsatellite instability (MSI)^{22,25-26}. This sample harbors a TMB level that may be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors in multiple solid tumor types^{2-4,11}.

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors²⁷⁻²⁹, including approved therapies nivolumab and pembrolizumab³⁰. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)³¹.

FREQUENCY & PROGNOSIS

MSI-high (MSI-H) has been observed at high frequency in endometrial cancers (14-33%)³²⁻³⁹, colorectal cancers (CRCs; 10-15%)^{25,29,40-42}, and gastric cancers (12-35%)⁴³⁻⁴⁶ and at lower frequencies in many other tumor types, including esophageal⁴⁷, small bowel⁴⁸⁻⁵², hepatobiliary⁵³⁻⁵⁹, prostate⁶⁰⁻⁶², and urinary tract carcinomas⁶³⁻⁶⁵. In one study, MSI-H status was associated with a positive prognostic effect in patients with gastric cancer treated with surgery alone and a negative predictive effect in patients treated with chemotherapy⁶⁶. Data regarding the role of MSI-H on prognosis and survival in endometrial cancer are conflicting^{32,35-36,38,67-69}. However, studies specifically analyzing early stage endometrial cancer have reported a correlation between MSI-H and decreased survival^{34,37,39,68}, thereby suggesting that MSI-H predicts for poor prognosis in this subset of endometrial tumors.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor⁴². Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2^{42,70-71}. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers^{41,72-73}. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{41-42,71,73}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS
GENE
PIK3CA
ALTERATION

E726K

TRANSCRIPT ID

NM_006218

CODING SEQUENCE EFFECT

2176G>A

VARIANT ALLELE FREQUENCY (% VAF)

18.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Clinical and preclinical data in various tumor types indicate that PIK3CA activating alterations may predict sensitivity to therapies targeting PI3K⁷⁴⁻⁸¹, AKT⁸²⁻⁸³, or mTOR⁸⁴⁻⁹¹. The Phase 2 NCI-MATCH study of copanlisib for patients with refractory solid tumors harboring PIK3CA mutations with or without PTEN loss met its primary endpoint with an ORR of 16% (4/25 PRs); responses (PR or SD >6 months) were seen in patients with ameloblastoma, liposarcoma, and carcinomas of the endometrium, ovary, esophagus, lung, and prostate⁸¹. However, the Phase 2 study of copanlisib for patients with endometrial

carcinoma harboring PIK3CA hotspot mutations failed to report any objective responses (n=11)⁸⁰. Two other studies of copanlisib for patients with genomically unselected tumors reported 1 CR and 2 PRs (1 unconfirmed) among 16 total patients with PIK3CA-mutated solid tumors with or without PTEN alterations⁷⁸⁻⁷⁹. In the Phase 2 MATCH trial for patients with PIK3CA-mutated solid tumors, 28% (18/65) of patients experienced PFS lasting at least 6 months after treatment with taselisib; however, no ORs were observed in this study⁹². A separate Phase 1b study of taselisib in combination with the CDK4/6 inhibitor palbociclib for patients with PIK3CA-mutated solid tumors reported an ORR of 0% (n=12) and a DCR of 17% (2/12)⁹³. In a Phase 1 trial of the dual PI3K/mTOR kinase inhibitor apitolisib, 79% (11/14) of patients with PIK3CA-mutated advanced solid tumors experienced disease control (3 PRs, 8 SDs)⁹⁴. The PI3K inhibitor alpelisib is approved as a single agent for the treatment of patients with PIK3CA-related overgrowth spectrum (PROS)⁹⁵, but has shown limited activity as monotherapy for PIK3CA-mutated solid tumors with a Phase 1a study reporting an ORR of 6.0% (8/134) and a DCR of 58% (78/134)⁹⁶.

FREQUENCY & PROGNOSIS

Two studies have reported PIK3CA mutation in

4.9-9.6% of nasopharyngeal carcinomas (NPC)⁹⁷⁻⁹⁸, whereas several studies have reported that PIK3CA amplification and overexpression occur frequently in NPC, but that PIK3CA mutations are infrequent⁹⁹⁻¹⁰⁴. PIK3CA amplification and expression have been reported in 20% and 44% of NPCs, respectively¹⁰⁵. PIK3CA mutations were detected in 7.1% of head and neck carcinomas (COSMIC, Jun 2021)¹⁰⁶. In one study, PI3K mutations were prevalent in advanced Stage 4 HNSCC tumors and associated with tumor progression¹⁰⁷. PIK3CA amplification has been found to be associated with poor prognosis in HNSCC¹⁰⁸⁻¹¹⁰. PIK3CA amplification has been associated with more aggressive disease in NPC¹⁰¹, although another study found that PIK3CA mutation did not affect disease-specific survival⁹⁷.

FINDING SUMMARY

PIK3CA encodes p110- α , which is the catalytic subunit of phosphatidylinositol 3-kinase (PI3K). The PI3K pathway is involved in cell signaling that regulates a number of critical cellular functions, including cell growth, proliferation, differentiation, motility, and survival¹¹¹⁻¹¹². PIK3CA alterations that have been characterized as activating, such as observed here, are predicted to be oncogenic^{107,113-133}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS

GENE

APC

ALTERATION

S2223*

TRANSCRIPT ID

NM_000038

CODING SEQUENCE EFFECT

6668C>A

VARIANT ALLELE FREQUENCY (% VAF)

24.7%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved drugs targeting APC inactivation in cancer. Loss of APC function leads to accumulation of beta-catenin and upregulation of WNT pathway transcription programs¹³⁴, and potential therapeutic approaches to target this pathway include CBP/beta-catenin antagonists, which interfere with the ability of beta-catenin to interact with transcriptional co-activator CBP¹³⁵⁻¹³⁶. In a Phase 1 trial of the CBP/beta-

catenin antagonist E7386, 1 patient with APC-mutated small bowel adenocarcinoma achieved a PR with tumor shrinkage of -69% and response duration of 165 days¹³⁷; preclinical data support sensitivity of APC-deficient gastric or colorectal cancer models to E7386¹³⁸⁻¹³⁹.

FREQUENCY & PROGNOSIS

APC mutations have been reported in 8.2% (8/97) of sinonasal and nasal cavity tumors analyzed in COSMIC (Jan 2022)¹⁰⁶. Small studies of sinonasal adenocarcinomas in the literature have reported APC mutations in 0-30% of samples¹⁴⁰⁻¹⁴¹. Hypermethylation of the APC promoter, likely to result in downregulation of Apc expression, has been frequently reported various types of head and neck carcinomas, including laryngeal papilloma, nasopharyngeal carcinoma, squamous cell carcinoma (HNSCC), and mixed exocrine-neuroendocrine carcinoma of the nasal cavity¹⁴²⁻¹⁴⁵. An analysis of a mixed exocrine-neuroendocrine carcinoma of the nasal cavity found reduced Apc protein expression in the neuroendocrine component only, correlating with APC promoter methylation and copy number

loss¹⁴⁵. Studies on the prognostic effect of APC alterations in sinonasal and nasal cavity tumors are limited (PubMed, Mar 2021). Solid tumors with WNT/beta-catenin pathway alterations, as seen here, were observed to have significantly less T-cell inflammation in one study¹⁴⁶.

FINDING SUMMARY

APC (adenomatous polyposis coli) encodes a tumor suppressor with critical roles in regulating cell division and adhesion. APC interacts with beta-catenin and controls signaling in the WNT pathway, which regulates embryonic development and cell differentiation¹⁴⁷. Alterations such as seen here may disrupt APC function or expression¹⁴⁸⁻¹⁵².

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in APC are found in more than 90% of patients with familial adenomatous polyposis (FAP)¹⁵³⁻¹⁵⁵. The prevalence for FAP in the general population is estimated to be 1:8,300 from birth¹⁵⁶, and in the appropriate clinical context germline testing of APC is recommended.

GENE

CCND1

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Amplification or overexpression of CCND1 may predict sensitivity to CDK4/6 inhibitors, such as abemaciclib, palbociclib, and ribociclib¹⁵⁷⁻¹⁶², although as monotherapy these agents have shown limited activity in tumor types other than breast cancer^{161,163}. In refractory advanced solid tumors with CCND1 (n=39) or CCND3 (n=1) amplification and retinoblastoma protein expression, palbociclib resulted in SD for 39% (14/36) of patients and a median PFS of 1.8 months in the NCI-MATCH trial¹⁶⁴; 4 patients (13%, 4/36 overall) with squamous cell carcinomas (lung,

esophageal, or laryngeal) or adenoid cystic carcinoma experienced prolonged SD in this study¹⁶⁴. Among 9 patients with CCND1-amplified advanced solid tumors, 1 patient with bladder cancer responded to ribociclib in a Phase 2 trial¹⁶⁵. Two of 3 patients with advanced solid tumors and concurrent alterations in CCND1/3 and NOTCH1, as seen here, achieved prolonged SD on palbociclib in the exploratory analysis of NCI-MATCH¹⁶⁴.

— Potential Resistance —

CCND1 amplification may predict worse outcomes on immune checkpoint inhibitors (anti-PD-1/PD-L1/CTLA-4) in solid tumors on the basis of 2 meta-analyses¹⁶⁶⁻¹⁶⁷; in these studies, CCND1 amplification was associated with significantly decreased response rate¹⁶⁷ and OS (HR=1.6-2.0)¹⁶⁶⁻¹⁶⁷ across various tumor types and significantly shorter OS specifically in urothelial carcinoma (HR=2.2-3.6), melanoma (HR=1.6-2.5), and solid tumors harboring elevated TMB (HR=2.8)¹⁶⁶⁻¹⁶⁷.

FREQUENCY & PROGNOSIS

Amplification of CCND1 has been observed in 28-45% of head and neck squamous cell carcinoma (HNSCC) cases, including 63% of pharyngeal carcinomas, 37% of laryngeal, and 25% of oral carcinomas¹⁶⁸⁻¹⁷⁰. A study reported cyclin D1 expression in 66% of HNSCC samples¹⁷¹. CCND1 amplification or overexpression has been associated with advanced disease, lymph node metastasis, and decreased overall and disease-free survival in patients with HNSCC¹⁷⁰⁻¹⁷⁴. CCND1 or CCND3 protein expression may also associate with cisplatin resistance in HNSCC¹⁷⁵.

FINDING SUMMARY

CCND1 encodes cyclin D1, a binding partner of the kinases CDK4 and CDK6, that regulates RB activity and cell cycle progression. Amplification of CCND1 has been positively correlated with cyclin D1 overexpression¹⁷⁶ and may lead to excessive proliferation¹⁷⁷⁻¹⁷⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS

GENE

CDK6

ALTERATION

amplification - equivocal

palbociclib, and ribociclib^{158-159,161,179}. Clinical benefit has been reported for patients with CDK6-amplified or mutated solid tumors in response to treatment with ribociclib^{165,180}.

FREQUENCY & PROGNOSIS

No alterations in CDK6 were reported in 56 samples analyzed in the Singapore Nasopharynx TCGA dataset⁹⁹. Published data investigating the prognostic implications of CDK6 alterations in head and neck carcinomas are limited (PubMed, Jan 2021).

FINDING SUMMARY

CDK6 encodes cyclin-dependent kinase 6, which regulates the cell cycle, differentiation, senescence, and apoptosis¹⁸¹⁻¹⁸³. CDK6 and its functional homolog CDK4 are activated by D-type cyclins and promote cell cycle progression by inactivating the tumor suppressor Rb¹⁸⁴⁻¹⁸⁵. Amplification of the chromosomal region that includes CDK6 has been reported in multiple cancer types, and has been associated with overexpression of CDK6 protein¹⁸⁶⁻¹⁸⁷.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Tumors with CDK6 activation may be sensitive to CDK4/6 inhibitors, such as abemaciclib,

GENE

MTAP

ALTERATION

loss exons 6-8

sensitivity to inhibitors of purine synthesis or purine analogs, especially upon addition of exogenous MTA, which is converted to adenine in normal cells, thereby providing competition to purine poisons lacking in MTAP-deficient cells¹⁹⁶⁻²⁰⁶. A Phase 2 study of L-alanosine, an inhibitor of adenine synthesis, as a monotherapy for 65 patients with MTAP-deficient cancers reported no responses and stable disease in 23.6% (13/55) of patients²⁰⁷.

FREQUENCY & PROGNOSIS

MTAP loss/homozygous deletion as well as loss of expression has been reported in a wide variety of solid tumors and hematologic cancers²⁰⁸⁻²⁰⁹; such events have been correlated with poor prognosis in a variety of cancer types, including hepatocellular carcinoma²¹⁰, gastrointestinal stromal tumors²¹¹, mantle cell lymphoma (MCL)²¹², melanoma²¹³⁻²¹⁴, gastric cancer²¹⁵, myxofibrosarcoma²¹⁶, nasopharyngeal carcinoma²¹⁷, ovarian carcinoma²⁰⁸ and non-small cell lung cancer²¹⁸. MTAP loss was not prognostic in pediatric B-cell acute lymphocytic leukemia²¹⁹ or in astrocytoma²²⁰. However, MTAP has also

been reported to be overexpressed in colorectal cancer (CRC) samples²²¹, and MTAP retention is thought to be important for prostate cancer growth due to continuous supply of SAM²²². Germline SNPs in MTAP have been correlated with the development of cutaneous melanoma²²³⁻²²⁴, esophageal cancer²²⁵⁻²²⁶, osteosarcoma²²⁷, and CRC²²⁸.

FINDING SUMMARY

MTAP encodes S-methyl-5'-thioadenosine (MTA) phosphorylase, a tumor suppressor involved in polyamine metabolism and methionine synthesis, although its enzymatic function is dispensable for its tumor suppressor activity²²⁹⁻²³⁰. Decreased expression of MTAP leads to MTA accumulation within tumor cells and their microenvironment^{210,231-232}, thereby reducing intracellular arginine methylation^{188,190,233} and altering cell signaling^{232,234}. MTAP is located at 9p21, adjacent to CDKN2A and CDKN2B, with which it is frequently co-deleted in various cancers. Other alterations in MTAP are rare and have not been extensively characterized.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical and limited clinical evidence indicate that MTAP inactivation produces specific metabolic vulnerabilities. MTAP inactivation may confer sensitivity to MAT2A inhibitors¹⁸⁸. A Phase 1 trial of MAT2A inhibitor AG-270 reported 1 PR and 2 SDs lasting longer than 6 months for patients with advanced solid tumors displaying MTAP loss¹⁸⁹. Although preclinical data have suggested that MTAP loss sensitizes cells to PRMT5 inhibition^{188,190-191}, MTAP loss may not be a biomarker of response to previously developed small-molecule SAM-uncompetitive PRMT5 inhibitors¹⁹²; dual PRMT1 and PRMT5 inhibition may be more effective¹⁹³⁻¹⁹⁵. In preclinical cancer models, MTAP inactivation showed increased

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS

GENE

CDKN2A/B

ALTERATION

CDKN2B loss, CDKN2A loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical data suggest that tumors with loss of p16INK4a function may be sensitive to CDK4/6 inhibitors, such as abemaciclib, ribociclib, and palbociclib²³⁵⁻²³⁸. Although case studies have reported that patients with breast cancer or uterine leiomyosarcoma harboring CDKN2A loss responded to palbociclib treatment²³⁹⁻²⁴⁰, multiple other clinical studies have shown no significant correlation between p16INK4a loss or inactivation and therapeutic benefit of these agents^{159-160,165,180,241-243}; it is not known whether CDK4/6 inhibitors would be beneficial in this case. Although preclinical studies have suggested that loss of p14ARF function may be associated with reduced sensitivity to MDM2 inhibitors²⁴⁴⁻²⁴⁵, the clinical relevance of p14ARF as a predictive biomarker is not clear. There are no drugs that directly target the mutation or loss of CDKN2B in cancer. Because the p15INK4b protein encoded by CDKN2B is known to inhibit CDK4, tumors with CDKN2B mutation or loss may predict sensitivity to CDK4/6 inhibitors, such as ribociclib, abemaciclib, and palbociclib^{158-159,165,242,246-247}.

FREQUENCY & PROGNOSIS

Mutation in CDKN2A or CDKN2B has been detected in 1.1% (1/90) and 0% (0/21) sinonasal carcinomas analyzed in COSMIC, respectively (Jan 2022)¹⁰⁶. In a genomic analysis of 56 nasopharyngeal carcinoma cases, mutation of CDKN2B was detected in 2% (1/56) of samples, while CDKN2A mutation was absent^{99,248-249}. In another study of nasopharyngeal carcinoma, p16INK4a homozygous deletion was observed in 35% (7/20) of cases²⁵⁰. CDKN2A and CDKN2B were identified as susceptibility loci in nasopharyngeal carcinoma²⁵¹. Inactivation of CDKN2A and CDKN2B by gene methylation was reported to be common in undifferentiated nasopharyngeal carcinoma, with CDKN2A methylation occurring in 23-42% of cases and CDKN2B methylation occurring in 20-50% of cases²⁵²⁻²⁵³. CDKN2A is also often inactivated in head and neck squamous cell carcinoma (HNSCC) through deletion, promoter methylation, or mutation²⁵⁴⁻²⁵⁶. In the context of nasopharyngeal carcinoma, loss of p16INK4a correlated with worse overall survival²⁵⁷ and local recurrence²⁵⁸.

FINDING SUMMARY

CDKN2A encodes two different, unrelated tumor suppressor proteins, p16INK4a and p14ARF, whereas CDKN2B encodes the tumor suppressor p15INK4b²⁵⁹⁻²⁶⁰. Both p15INK4b and p16INK4a bind to and inhibit CDK4 and CDK6, thereby maintaining the growth-suppressive activity of the Rb tumor suppressor; loss or inactivation of either p15INK4b or p16INK4a contributes to

dysregulation of the CDK4/6-cyclin-Rb pathway and loss of cell cycle control²⁶¹⁻²⁶². The tumor suppressive functions of p14ARF involve stabilization and activation of p53, via a mechanism of MDM2 inhibition²⁶³⁻²⁶⁴. One or more alterations observed here are predicted to result in p16INK4a loss of function²⁶⁵⁻²⁸⁶. One or more alterations seen here are predicted to result in p14ARF loss of function^{269,286-289}. CDKN2B alterations such as seen here are predicted to inactivate p15INK4b²⁹⁰.

POTENTIAL GERMLINE IMPLICATIONS

Germline CDKN2A mutation is associated with melanoma-pancreatic cancer syndrome, a condition marked by increased risk of developing malignant melanoma and/or pancreatic cancer²⁹¹. Mutation carriers within families may develop either or both types of cancer, and melanoma cases may be referred to as familial or hereditary melanoma²⁹²⁻²⁹³. CDKN2A is the most implicated gene in familial melanoma, with germline mutations present in 16% to 20% of familial melanoma cases²⁹⁴⁻²⁹⁶. CDKN2A alteration has also been implicated in familial melanoma-astrocytoma syndrome, an extremely rare tumor association characterized by dual predisposition to melanoma and nervous system tumors²⁹⁷⁻²⁹⁹. In the appropriate clinical context, germline testing of CDKN2A is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS

GENE

EPHB4

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies available to target EPHB4 alterations in cancer. sEPHB4 is a soluble monomeric extracellular domain of EPHB4 that functions as an antagonist of EphrinB2-EPHB4 interaction³⁰⁰, and fusion of sEPHB4 with human serum albumin (HSA) increases its stability³⁰¹. Recombinant sEPHB4-HSA is under investigation in clinical trials. Preclinical studies have demonstrated that sEPHB4-HSA inhibits cell proliferation and xenograft tumor growth, including for cells expressing cancer-associated EPHB4 mutants or overexpressing wild-type EPHB4^{300,302-306}. In addition, small-molecule

inhibitors targeting multiple tyrosine kinases including EPHB4, such as JI-101 and XL647, have been under preclinical and clinical investigation³⁰⁷⁻³⁰⁹.

FREQUENCY & PROGNOSIS

Increased EPHB4 mRNA and/or protein expression has been reported in a variety of cancer types, including head and neck squamous cell carcinoma (HNSCC)³¹⁰⁻³¹³, gastric and esophageal³¹⁴⁻³¹⁸, colorectal carcinoma (CRC)³¹⁹⁻³²⁵, breast³²⁶⁻³³⁰, ovarian³³⁰⁻³³², endometrial³³³⁻³³⁵, thyroid³³⁶⁻³³⁸, lung³³⁹⁻³⁴⁰, glioma³⁴¹⁻³⁴², and other solid tumors^{302,343-350}. In several of these studies, increased EPHB4 expression has been associated with clinicopathologic features, including disease stage^{302,310,326,331-332,340,343,345}, histological grade^{316,326,333,342}, and hormone receptor status^{329,334}. High EPHB4 expression has been associated with inferior survival in multivariate analyses for patients with CRC treated with bevacizumab [hazard ratio (HR) = 5.95]³²⁴, HNSCC (HR = 2.95)³¹³, epithelial ovarian cancer (HR =

4.53)³³⁰, or glioma (HR = 3.21)³⁴².

FINDING SUMMARY

EPHB4 encodes a member of the EPH family of receptor tyrosine kinases³⁵¹. Ephrin signaling has been implicated in multiple processes, including cell adhesion, cytoskeletal organization, and cell migration³⁵², and signaling between EPHB4 and its ligand EphrinB2 is particularly important for angiogenesis³⁵³⁻³⁵⁴. EPHB receptors, including EPHB4, have been shown to undergo dysregulation (amplification, mutation, under- or overexpression) in a number of different cancer types³⁵⁵. EPHB4 amplification has been reported in several solid tumor types^{310-311,316,356-357} and was associated with advanced disease stage in head and neck squamous cell carcinoma (HNSCC)³¹⁰. Activating missense mutations in or near the tyrosine kinase domain, including G723S, A742V, and P881S, have also been identified in lung cancer³⁰⁶.

GENE

FGF19

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

A Phase 1 study of the FGFR4 inhibitor fsgatinib (BLU-554) for patients with advanced hepatocellular carcinoma (HCC) reported a 17% ORR (11/66, 1 CR, ongoing for >1.5 years) and 3.3-month PFS for FGF19 IHC-positive patients; patients with negative or unknown FGF19 IHC scores experienced poorer outcomes (0% ORR, 2.3-month PFS)³⁵⁸. A Phase 1/2 study evaluating another FGFR4 inhibitor, FGF401, demonstrated an ORR of 7.5% (4/53) and SD rate of 53% (28/53) for patients with HCC³⁵⁹. A Phase 1 study of the FGFR4 inhibitor H3B-6527 reported a 17% ORR (OS of 10.3 months, 46% clinical benefit rate)

among patients with HCC; enrollment of patients with intrahepatic cholangiocarcinoma (ICC) was suspended due to efficacy³⁶⁰. A retrospective analysis reported that 50% (2/4) of patients with HCC harboring FGF19 amplification experienced a CR to sorafenib³⁶¹, though another retrospective study found patients with higher pretreatment serum levels of FGF19 experienced reduced benefit from sorafenib compared with those with lower serum FGF19 (PFS of 86 vs. 139 days, OS of 353 vs. 494 days); no difference was observed for lenvatinib³⁶². A patient with head and neck squamous cell carcinoma (HNSCC) with 11q13 (FGF3, FGF4, FGF19) and 12p13 (FGF6 and FGF23) amplification experienced a CR lasting 9 months from a pan-FGFR inhibitor³⁶³.

FREQUENCY & PROGNOSIS

For patients with solid tumors, FGF19 amplification has been reported most frequently in breast cancer (17%), head and neck cancer (12%), lung squamous cell carcinoma (SCC; 12%), and urothelial carcinoma cancer (11%)³⁶⁴⁻³⁶⁶. FGF19

mutations are rare in solid tumors³⁶⁴. FGF19 expression or amplification has been associated with poor prognosis in hepatocellular carcinoma (HCC)³⁶⁷⁻³⁶⁸, and in prostate cancer following radical prostatectomy³⁶⁹. Studies suggest FGF19 expression may also be a poor prognostic indicator in head and neck squamous cell carcinoma (HNSCC)³⁷⁰ and lung SCC³⁷¹.

FINDING SUMMARY

FGF19 encodes fibroblast growth factor 19, an FGFR4 ligand involved with bile acid synthesis and hepatocyte proliferation in the liver³⁷²⁻³⁷³. FGF19 lies in a region of chromosome 11q13 that also contains FGF3, FGF4, and CCND1; this region is frequently amplified in a diverse range of malignancies³⁷⁴. Correlation between FGF19 amplification and protein expression has been reported in hepatocellular carcinoma (HCC)³⁷⁵, lung squamous cell carcinoma^{371,376}, and head and neck squamous cell carcinoma (HNSCC)³⁷⁰, but was not observed in other cancers^{362,377}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS
GENE
FGF3
ALTERATION
 amplification

trials in a number of different cancers. Limited data suggest that pan-FGFR inhibitors show activity in FGF amplified cancers; following treatment with a selective pan-FGFR inhibitor, a patient with head and neck squamous cell carcinoma (HNSCC) and amplification of 11q13 (FGF3, FGF4, FGF19) and 12p13 (FGF6 and FGF23) experienced a radiologic CR³⁷⁸.

cycle progression. This chromosomal region is frequently amplified in a diverse range of malignancies¹⁷⁷.

FINDING SUMMARY

FGF3 encodes fibroblast growth factor 3, a growth factor that plays a central role in development of the inner ear. Germline mutations in FGF3 give rise to an autosomal recessive syndrome characterized by microdontia, deafness, and complete lack of inner ear structures³⁷⁹.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies that directly address genomic alterations in FGF3. Inhibitors of FGF receptors, however, are undergoing clinical

FREQUENCY & PROGNOSIS

FGF3 lies in a region of chromosome 11q13 that also contains FGF19, FGF4, and CCND1, the latter gene encoding cyclin D1, a key regulator of cell

GENE
FGF4
ALTERATION
 amplification

inhibitor, a patient with head and neck squamous cell carcinoma (HNSCC) and amplification of 11q13 (FGF3, FGF4, FGF19) and 12p13 (FGF6 and FGF23) experienced a radiologic CR³⁷⁸.

(HCC; 5%), however FGF4 amplification is rare in hematopoietic and lymphoid malignancies, reported in less than 1% of samples analyzed (cBioPortal, Jan 2022)²⁴⁸⁻²⁴⁹.

FREQUENCY & PROGNOSIS

FGF4 lies in a region of chromosome 11q13 that also contains FGF19, FGF3, and CCND1, the latter gene encoding cyclin D1, a key regulator of cell cycle progression. This chromosomal region is frequently amplified in a diverse range of malignancies¹⁷⁷ including esophageal carcinoma (35%), head and neck squamous cell carcinoma (HNSCC; 24%), breast invasive carcinoma (14%), lung squamous cell carcinoma (13%), cholangiocarcinoma (11%), bladder urothelial carcinoma (10%), stomach adenocarcinoma (7%), skin melanoma (5%), and hepatocellular carcinoma

FINDING SUMMARY

FGF4 encodes fibroblast growth factor 4, which plays a central role in development of the teeth³⁸² and acts synergistically with other FGFs and SHH (sonic hedgehog) to regulate limb outgrowth in vertebrate development³⁸³. FGF4 lies in a region of chromosome 11q13 that also contains FGF19, FGF3, and CCND1, the latter gene encoding cyclin D1, a key regulator of cell cycle progression. Amplification of FGF4, along with that of FGF3, FGF19, and CCND1, has been reported in a variety of cancers^{177,380,384-387} and may confer sensitivity to the multi-kinase inhibitor sorafenib³⁸⁰.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

FGF4 amplification and overexpression was associated with cell sensitivity to the multikinase inhibitor sorafenib in preclinical studies³⁸⁰⁻³⁸¹ and amplification of FGF4/FGF3 in HCC significantly correlated with patient response to sorafenib ($p=0.006$)³⁸⁰. Limited data suggest that pan-FGFR inhibitors show activity in FGF amplified cancers; following treatment with a selective pan-FGFR

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS
GENE

NOTCH1

ALTERATION

P1443fs*36

TRANSCRIPT ID

NM_017617

CODING SEQUENCE EFFECT

4319_4320insT

VARIANT ALLELE FREQUENCY (% VAF)

56.6%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

NOTCH1 inhibitors and gamma-secretase inhibitors (GSIs) may be potential therapeutic approaches in the case of NOTCH1 activating mutations³⁸⁸⁻³⁹⁶. In a Phase 2 study, the GSI AL101 (BMS-906024) elicited PRs in 15% (6/39) and SDs in 54% (21/39) of patients with metastatic adenoid cystic carcinoma (ACC) harboring NOTCH activating alterations³⁹⁷. Additional responses to AL101 have been reported in a patient with gastroesophageal junction adenocarcinoma harboring multiple NOTCH1 mutations, a patient with T-cell acute lymphoblastic leukemia (T-ALL) harboring a NOTCH1 HD domain mutation, and a patient with ACC harboring a single NOTCH1 mutation³⁹⁸. A Phase 1 study of the pan-NOTCH inhibitor CB-103 for patients with advanced or

recurrent solid tumors reported a preliminary mPFS of 21.7 weeks for patients with ACC, with 2 patients harboring NOTCH1-mutated ACC demonstrating SD > 6 months as best response³⁹⁹. On the basis of clinical data in non-Hodgkin lymphoma, NOTCH1 activating alterations may be associated with sensitivity to the approved PI3K inhibitor copanlisib⁴⁰⁰; this is further supported by limited preclinical data that suggest that NOTCH1 may be a negative regulator of PTEN⁴⁰¹⁻⁴⁰². Two of 3 patients with advanced solid tumors and concurrent alterations in CCND1/3 and NOTCH1, as seen here, achieved prolonged SD on palbociclib in the exploratory analysis of NCI-MATCH¹⁶⁴. While activating mutations may be targeted via gamma-secretase inhibitors or PI3K inhibitors, there are no therapies available to address NOTCH1 inactivation, as seen here.

FREQUENCY & PROGNOSIS

NOTCH1 mutation has been reported in 13% (4/30) of upper aerodigestive tract adenocarcinoma cases (COSMIC, Mar 2021)¹⁰⁶. Studies have reported NOTCH1 mutations in 15-26% of head and neck squamous cell carcinomas (HNSCCs) analyzed, suggesting that NOTCH1 acts as a tumor suppressor in HNSCC⁴⁰³⁻⁴⁰⁴. NOTCH1 mutations have been reported in 3% of nasopharyngeal carcinoma and 22% (2/9) of sinonasal adenocarcinomas analyzed in COSMIC (Aug 2021)¹⁰⁶. Published data investigating the prognostic implications of NOTCH1 alterations in

head and neck adenocarcinoma are limited (PubMed, Mar 2021). Published data investigating the prognostic implications of NOTCH1 alteration in head and neck carcinomas outside of the context of HNSCC are limited (PubMed, Aug 2021).

FINDING SUMMARY

NOTCH1 encodes a member of the NOTCH family of receptors, which are involved in cell fate determination and various developmental processes. Depending on cellular context, NOTCH1 can act as either a tumor suppressor or an oncogene⁴⁰⁵⁻⁴⁰⁶. Upon binding of membrane-bound ligands, the NOTCH1 intracellular domain (NICD) is cleaved and forms part of a transcription factor complex that regulates downstream target genes involved in cell fate determination, proliferation, and apoptosis⁴⁰⁷⁻⁴⁰⁸. NOTCH1 alterations that disrupt ligand binding⁴⁰⁹⁻⁴¹¹ or remove the transmembrane domain (amino acids 1736-1756), RAM domain (amino acids 1757-1926), ankyrin repeats (amino acids 1927-2122) and/or transactivation domain (amino acids 2123-2374) that are necessary for NOTCH1 function, such as observed here, are predicted to be inactivating^{408,412-414}. Several point mutations, including D469G, A465T, C478F, R1594Q, and P1770S, have also been reported to inactivate NOTCH1^{405,415-416}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

R273H

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

818G>A

VARIANT ALLELE FREQUENCY (% VAF)

47.2%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib⁴¹⁷⁻⁴²⁰, or p53 gene therapy and immunotherapeutics such as SGT-53⁴²¹⁻⁴²⁵ and ALT-801⁴²⁶. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype⁴²⁷. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer⁴²⁸. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer⁴²⁹. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone⁴³⁰. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel⁴³¹. A Phase 1

trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations⁴³². The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring⁴³³. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage⁴²⁵. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutated p53 such as APR-246⁴³⁴⁻⁴³⁶. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR⁴³⁷. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies⁴³⁸⁻⁴³⁹; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies⁴⁴⁰⁻⁴⁴¹. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

Mutations of TP53 have been reported in 36-51% of head and neck carcinomas, across numerous subtypes including in 40% of adenocarcinomas⁴⁴²⁻⁴⁴⁵. In one study of 78 patients with undifferentiated nasopharyngeal carcinoma, p53 protein expression was associated with worse disease-free survival⁴⁴⁶.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers⁴⁴⁷. Alterations such

as seen here may disrupt TP53 function or expression⁴⁴⁸⁻⁴⁵².

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Sep 2021)⁴⁵³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers⁴⁵⁴⁻⁴⁵⁶, including sarcomas⁴⁵⁷⁻⁴⁵⁸. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000⁴⁵⁹ to 1:20,000⁴⁵⁸. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30⁴⁶⁰. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁴⁶¹⁻⁴⁶⁶. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy⁴⁶¹⁻⁴⁶². Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease⁴⁶⁷. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{465,468-469}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

ORDERED TEST # ORD-1346479-01

THERAPIES WITH CLINICAL BENEFIT
IN PATIENT'S TUMOR TYPE

Dostarlimab

Assay findings association
Tumor Mutational Burden
 18 Muts/Mb

AREAS OF THERAPEUTIC USE

Dostarlimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor response. It is FDA approved to treat patients with mismatch repair deficient recurrent or advanced endometrial cancer or solid tumors. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher

TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Clinical data on the efficacy of dostarlimab for the treatment of head and neck cancer are limited (PubMed, Mar 2022). Dostarlimab has been studied primarily in recurrent and advanced mismatch repair-deficient (dMMR) endometrial and non-endometrial cancers⁴⁷¹⁻⁴⁷³. In the Phase 1 GARNET trial, single-agent dostarlimab elicited an ORR of 39% (41/106) and an immune-related ORR of 46% (50/110) for patients with non-endometrial dMMR solid tumors^{471,474}.

Pembrolizumab

Assay findings association
Tumor Mutational Burden
 18 Muts/Mb

AREAS OF THERAPEUTIC USE

Pembrolizumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved for patients with tumor mutational burden (TMB)-high (≥ 10 Muts/Mb), microsatellite instability-high (MSI-H), or mismatch repair-deficient (dMMR) solid tumors; as monotherapy for PD-L1-positive non-small cell lung cancer (NSCLC), head and neck squamous cell cancer (HNSCC), cervical cancer, or esophageal cancer; and in combination with chemotherapy for PD-L1-positive triple-negative breast cancer (TNBC) or cervical cancer. It is also approved in various treatment settings as monotherapy for patients with melanoma, HNSCC, urothelial carcinoma, hepatocellular carcinoma, Merkel cell carcinoma, cutaneous squamous cell carcinoma, endometrial carcinoma that is MSI-H or dMMR, classical Hodgkin lymphoma, or primary mediastinal large B-cell lymphoma; and in combination with chemotherapy or targeted therapy for NSCLC, HNSCC, esophageal or gastroesophageal junction cancer, renal cell carcinoma, TNBC, or endometrial carcinoma that is not MSI-H or dMMR. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

A Phase 1b trial for 27 patients with previously treated PD-L1-positive advanced nasopharyngeal carcinoma achieved an ORR of 25.9% (7/27), a median PFS of 6.5 months, and a median OS of 16.5 months on pembrolizumab⁴⁷⁵. In the Phase 2 KEYNOTE 158 multi-solid tumor trial, treatment with the PD-1 inhibitor pembrolizumab led to improved ORR for patients with TMB of 10 Muts/Mb or higher compared those with TMB < 10 Muts/Mb (28.3% [34/120] vs. 6.5% [41/635])¹¹. In the KEYNOTE 028/012 pan-solid tumor trials, a similar improvement in ORR was reported for patients with > 103 non-synonymous mutations/exome (~equivalency > 8 Muts/Mb as measured by this assay) compared to those with < 103 non-synonymous mutations/exome (30.6% [11/36] vs. 6.5% [5/77])⁴.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Atezolizumab

Assay findings association

Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Atezolizumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC) and urothelial carcinoma, depending on treatment setting. Atezolizumab is also approved in combination with other therapies to treat patients with non-squamous NSCLC lacking EGFR or ALK alterations, small cell lung cancer, hepatocellular carcinoma, and BRAF V600-positive melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

A Phase 1 trial of atezolizumab for patients with previously treated advanced head and neck cancer reported an ORR of 22% (7/32), a median duration of response of 7.4 months, median PFS of 2.6 months, and median OS of 6.0 months. Responses were achieved for

patients with primary tumors of the oral cavity (1/7), oropharynx (3/18), larynx (2/2), and nasopharynx (1/4), and did not correlate with PD-L1 expression or HPV status⁴⁷⁶. A 16% (3/19) ORR was reported for a dose-expansion cohort in a Phase 1b trial of atezolizumab combined with selicrelumab to treat head and neck cancer⁴⁷⁷. A Phase 1 study of atezolizumab combined with the IDO1 inhibitor navoximod reported 1 PR (in head and neck squamous cell carcinoma) and 1 prolonged SD outcome among 6 patients with advanced head and neck cancer⁴⁷⁸. A case study reported a patient with both metastatic lung adenocarcinoma and locally advanced epipharyngeal carcinoma who experienced a PR to atezolizumab in combination with several chemotherapy agents⁴⁷⁹. In the prospective Phase 2a MyPathway basket study evaluating atezolizumab for patients with TMB-High solid tumors, patients with TMB ≥ 16 Muts/Mb achieved improved ORR (38% [16/42] vs. 2.1% [1/48]), DCR (62% [26/42] vs. 23% [11/48]), mPFS (5.7 vs. 1.8 months, HR 0.34), and mOS (19.8 vs. 11.4, HR 0.53) as compared to those with TMB ≥ 10 and < 16 Muts/Mb⁴⁸⁰. In a retrospective analysis of patients with 17 solid tumor types (comprised of 47% NSCLC, 40% urothelial carcinoma, and 13% encompassing 15 other solid tumors), TMB of 16 Muts/Mb or greater was reported to be associated with an improved ORR to atezolizumab compared to chemotherapy (30% vs. 14%)¹².

Avelumab

Assay findings association

Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Avelumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 in order to enhance antitumor immune responses. It is FDA approved to treat patients 12 years and older with Merkel cell carcinoma, or for urothelial carcinoma in various treatment settings. The combination of avelumab and axitinib is FDA approved for patients with renal cell carcinoma (RCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Clinical data on the efficacy of avelumab for the treatment

of head and neck adenocarcinoma are limited (PubMed, Apr 2022). The JAVELIN Phase 1b study has demonstrated clinical benefit from single-agent avelumab in a variety of solid tumor types, including non-small cell lung carcinoma (NSCLC)⁴⁸¹, gastric carcinoma and gastroesophageal junction (GEJ) adenocarcinoma⁴⁸², urothelial carcinoma⁴⁸³, mesothelioma⁴⁸⁴, ovarian carcinoma⁴⁸⁵, and breast cancer⁴⁸⁶, and from avelumab combined with axitinib in renal cell carcinoma⁴⁸⁷. Emerging clinical data show a positive trend toward the association of tumor cell PD-L1 expression and improved objective response rate, progression-free survival, or overall survival in NSCLC in the first-line setting and in ovarian and breast cancer^{481,485-486}. Limited clinical data indicate activity of avelumab in adrenocortical carcinoma, metastatic castration-resistant prostate cancer, and thymic cancer⁴⁸⁸⁻⁴⁹⁰. Phase 3 studies are evaluating avelumab with chemoradiotherapy alone (NCT02952586) or in combination with cetuximab (NCT02999087) in patients with locally advanced head and neck squamous cell carcinoma (Jun 2021).

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Cemiplimab

Assay findings association
Tumor Mutational Burden
18 Muts/Mb

AREAS OF THERAPEUTIC USE

Cemiplimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved to treat patients with NSCLC with high PD-L1 expression (TPS \geq 50%), cutaneous squamous cell carcinoma (CSCC), or basal cell carcinoma (BCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of \geq 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Cemiplimab has been studied primarily in advanced cutaneous squamous cell carcinoma (CSCC), where it elicited a combined ORR of 48% (41/85) in Phase 1 and 2 studies⁴⁹¹. A Phase 2 trial of cemiplimab in patients with basal cell carcinoma (BCC) reported ORRs of 31% (5 CRs and 21 PRs) in patients with locally advanced BCC and 21% (6 PRs) in patients with metastatic BCC⁴⁹²⁻⁴⁹³. The Phase 3 EMPOWER-Lung 1 trial for advanced non-small cell lung cancer (NSCLC) with PD-L1 expression \geq 50% reported that cemiplimab is associated with improved PFS (8.2 vs. 5.7 months), OS (not reached vs. 14.2 months), and ORR (37% vs. 21%) compared with chemotherapy⁴⁹⁴. Clinical data on the efficacy of cemiplimab for the treatment of head and neck adenocarcinoma are limited (PubMed, Apr 2022).

Durvalumab

Assay findings association
Tumor Mutational Burden
18 Muts/Mb

AREAS OF THERAPEUTIC USE

Durvalumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of \geq 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Clinical data on the efficacy of durvalumab for the treatment of head and neck adenocarcinoma are limited (PubMed, Apr 2022). Single-agent durvalumab has demonstrated efficacy in non-small cell lung cancer⁴⁹⁵⁻⁴⁹⁶, and head and neck squamous cell carcinoma⁴⁹⁷⁻⁴⁹⁸. In patients with advanced solid tumors, durvalumab monotherapy has elicited disease control rates (DCRs) of

36.8–46.2% (7/19 to 12/26) in Phase 1/2 studies⁴⁹⁹⁻⁵⁰⁰. Durvalumab is also under investigation in combination with other agents in Phase 1/2 trials. In advanced melanoma, durvalumab in combination with trametinib and dabrafenib elicited ORRs and DCRs of 76.2% (16/21) and 100% (21/21) in patients with BRAF-mutant tumors, and durvalumab with trametinib elicited ORRs and DCRs of 21.4% (3/14) and 64.3% (9/14) in patients whose tumors were BRAF wild-type⁵⁰¹. Durvalumab in combination with the PARP inhibitor olaparib has shown activity in patients with metastatic castration-resistant prostate cancer and progression on enzalutamide and/or abiraterone⁵⁰² and in patients with BRCA-wild-type breast or gynecological cancer⁵⁰³. Durvalumab in combination with the anti-CTLA4 antibody tremelimumab, but not durvalumab as a single-agent, has shown activity in patients with previously treated advanced germ cell tumors⁵⁰⁴. Responses have also been reported for patients with solid tumors treated with durvalumab in combination with the anti-PD-1 antibody MEDI680⁵⁰⁵, the CXCR2 antagonist AZD5069⁵⁰⁶, or the ATR inhibitor AZD6738⁵⁰⁷. In patients with treatment-refractory solid tumors, concurrent durvalumab and radiotherapy achieved an ORR of 60% (6/10) for in-field evaluable lesions, including 2 CRs and 4 PRs⁵⁰⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Everolimus

Assay findings association
PIK3CA
E726K

AREAS OF THERAPEUTIC USE

Everolimus is an orally available mTOR inhibitor that is FDA approved to treat renal cell carcinoma (RCC) following antiangiogenic therapy; pancreatic neuroendocrine tumors; and well-differentiated non-functional neuroendocrine tumors of the lung or gastrointestinal tract. Everolimus is also approved to treat either renal angiomyolipoma or subependymal giant cell astrocytoma in association with tuberous sclerosis complex (TSC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence⁸⁴⁻⁹¹, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{88-91,509-513}.

SUPPORTING DATA

Clinical data on the efficacy of everolimus for the treatment of non-squamous head and neck carcinoma are

limited (PubMed, Feb 2022). A patient with lacrimal gland ductal adenocarcinoma achieved an 8 month PR after treatment with everolimus⁸⁵. A Phase 2 study of everolimus therapy has reported no objective responses in any of nine patients with refractory head and neck squamous cell carcinoma (HNSCC)⁵¹⁴. A Phase 1 trial in patients with advanced solid tumors reported that everolimus in combination with low dose weekly cisplatin showed activity in several tumor types, with three partial responses and prolonged stable disease observed in five patients out of 28 evaluable patients; one patient with oropharyngeal squamous cell carcinoma obtained stable disease after more than 6 cycles of treatments⁵¹⁵. Another Phase 1 trial of everolimus in combination with docetaxel and cisplatin reported progression-free survival rate of 87.5% at one year and 76.6% at two years in patients with advanced HNSCC⁵¹⁶. Whereas frequent adverse events precluded a recommended Phase 2 dose and schedule for the combination of trametinib and everolimus in a Phase 1b trial for solid tumors⁵¹⁷, a retrospective study for heavily pretreated patients with solid tumors reported tolerable regimens of the combination for 23/31 patients, with 16 patients treated >3 months and evaluable patients achieving a median PFS of 6.5 months⁵¹⁸.

Nivolumab

Assay findings association
Tumor Mutational Burden
18 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response. It is FDA approved as a monotherapy in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), urothelial carcinoma, colorectal cancer (CRC), classical Hodgkin lymphoma (cHL), gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma or squamous cell carcinoma (ESCC). It is also approved in combination with cabozantinib to treat RCC, and in combination with relatlimab to treat melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,470}, TMB of ≥10 Muts/Mb (based on this assay or others) may

predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

A Phase 2 study of single-agent nivolumab treatment in recurrent and metastatic nasopharyngeal carcinoma showed that patients had an ORR of 20.5% and a DCR of 54.5%, with 1 CR, 8 PRs, and 15 SD out of 44 patients; the median PFS was 2.8 months and the median OS was 17.1 months, with similar PFS and OS for patients with PD-L1-negative and PD-L1-positive tumors⁵¹⁹. A Phase 1/2 study of nivolumab for the treatment of patients with virus-associated tumors reported an ORR of 21% (5/24) and median PFS of 2.4 months in patients with recurrent/metastatic nasopharyngeal carcinoma; 88% (21/24) of cases were EBV-positive⁵²⁰.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Nivolumab + Ipilimumab

Assay findings association
Tumor Mutational Burden
18 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response, and ipilimumab is a cytotoxic T-lymphocyte antigen 4 (CTLA-4)-blocking antibody. The combination is FDA approved in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), hepatocellular carcinoma (HCC), and pleural mesothelioma. Furthermore, nivolumab is approved in combination with ipilimumab to treat patients with mismatch repair-deficient (dMMR) or microsatellite instability-high (MSI-H) colorectal cancer (CRC). Please see the drug label for full prescribing

information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{5-6,521}, a TMB score of ≥ 10 Muts/Mb (as measured by this assay) may predict sensitivity to combination nivolumab and ipilimumab treatment.

SUPPORTING DATA

A Phase 2 trial of nivolumab plus ipilimumab for patients with EBV-associated advanced nasopharyngeal carcinoma in Asia reported an ORR of 30% (12/40 PRs), median PFS of 5.3 months, and median OS of 17.6 months⁵²².

Temsirolimus

Assay findings association
PIK3CA
E726K

AREAS OF THERAPEUTIC USE

Temsirolimus is an intravenous mTOR inhibitor that is FDA approved for the treatment of advanced renal cell carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence⁸⁴⁻⁹¹, PIK3CA activating mutations may predict sensitivity to mTOR inhibitors such as everolimus and temsirolimus. Studies have reported modest activity of these therapies as single agents (ORRs of 0-4%), but improved activity has been observed when they are combined with other agents such as bevacizumab and doxorubicin (ORRs of 25-44%), for patients with PIK3CA-mutated solid tumors^{88-91,509-513}.

SUPPORTING DATA

Clinical data on the efficacy of temsirolimus for the treatment of non-squamous head and neck carcinoma are limited (PubMed, Nov 2021). A Phase 2 study evaluated temsirolimus for patients with recurrent or metastatic

head and neck squamous cell carcinoma (HNSCC) after failure of platinum and cetuximab and reported a median PFS of 1.9 months and overall survival of 5.1 months⁵²³. Temsirolimus has been tested preclinically and in clinical trials for HNSCC, in combination with the VEGF antibody bevacizumab, and has shown significant efficacy⁵²⁴. A study assessing temsirolimus in combination with metformin in patients with advanced solid tumors reported a partial response for one patient with HNSCC, despite disease progression after treatment with docetaxel and cisplatin and subsequent treatment with zalutumumab⁵²⁵. A Phase 1 study of temsirolimus in combination with carboplatin and paclitaxel in 18 patients with HNSCC reported a partial response rate of 22% and recommended Phase 2 testing⁵²⁶. However, a Phase 2 study of temsirolimus and erlotinib in patients with recurrent and/or metastatic, platinum-refractory HNSCC has reported that this combination therapy was poorly tolerated, with the trial ending early after 50% (6/12) of patients withdrew from the study⁵²⁷.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.clinicaltrials.gov). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

BIOMARKER

Tumor Mutational Burden

RESULT

18 Muts/Mb

RATIONALE

Increased tumor mutational burden may predict response to anti-PD-1 (alone or in combination

with anti-CTLA-4) or anti-PD-L1 immune checkpoint inhibitors.

NCT04237649
PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS

ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Sunto Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China), Beijing (China), Chengdu City (China), Changchun (China)

NCT04521621
PHASE 1/2

A Study of V937 in Combination With Pembrolizumab (MK-3475) in Participants With Advanced/Metastatic Solid Tumors (V937-013)

TARGETS

PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Tainan (Taiwan), Kaohsiung (Taiwan), Kashiwa (Japan), Afula (Israel), Jerusalem (Israel), Tel Aviv (Israel), Warszawa (Poland), Oslo (Norway)

NCT03674567
PHASE 1/2

Dose Escalation and Expansion Study of FLX475 Monotherapy and in Combination With Pembrolizumab

TARGETS

PD-1, CCR4

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Busan (Korea, Republic of), Shatin (Hong Kong), High West (Hong Kong), Ulsan (Korea, Republic of), Chungbuk (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Bangkok (Thailand)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
NCT04261439
PHASE 1

A Phase I/Ib Study of NIZ985 Alone and in Combination With Spartalizumab

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Chuo ku (Japan), Essen (Germany), Napoli (Italy), Leuven (Belgium), Barcelona (Spain), California, Texas

NCT03861793
PHASE 1/2

A Dose Escalation and Cohort Expansion Study of Subcutaneously-Administered Cytokine (ALKS 4230) as a Single Agent and in Combination With Anti-PD-1 Antibody (Pembrolizumab) in Subjects With Select Advanced or Metastatic Solid Tumors (ARTISTRY-2)

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Suwon (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Edmonton (Canada), Badalona (Spain), Rotterdam (Netherlands), Valencia (Spain), Madrid (Spain)

NCT04047862
PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS
 PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Guangdong (China), Changsha (China), Wuhan (China), Jinju-si (Korea, Republic of)

NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
 PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Amsterdam (Netherlands), Ravenna (Italy), Meldola (Italy)

NCT03891953
PHASE 1

Study of Safety and Efficacy of DKY709 Alone or in Combination With PDR001 in Patients With Advanced Solid Tumors.

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Chuo ku (Japan), Dresden (Germany), Essen (Germany), Barcelona (Spain), Massachusetts, Tennessee

NCT03396445
PHASE 1

Safety and Pharmacokinetics Study of MK-5890 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Adults With Advanced Solid Tumors (MK-5890-001)

TARGETS
 PD-1, CD27

LOCATIONS: Taipei (Taiwan), Seoul (Korea, Republic of), Jerusalem (Israel), Ramat Gan (Israel), Be'er Sheva (Israel), Amsterdam (Netherlands), Rotterdam (Netherlands), Barcelona (Spain), Madrid (Spain), Pozuelo de Alarcon (Spain)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

GENE
APC
ALTERATION

S2223*

RATIONALE

Based on preclinical and limited clinical data, APC inactivation may be associated with sensitivity to CBP/beta-catenin interaction inhibitors.

NCT03833700
PHASE 1

A Study of E7386 in Participants With Advanced Solid Tumor Including Colorectal Cancer (CRC)

TARGETS
 CBP, Beta-catenin

LOCATIONS: Fukuoka (Japan), Nagaizumi-cho (Japan), Chuo Ku (Japan), Kashiwa (Japan)

NCT04008797
PHASE 1

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

TARGETS
 CBP, Beta-catenin, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Osakasayama (Japan), Chuo-Ku (Japan), Kashiwa (Japan)

NCT03264664
PHASE 1

Study of E7386 in Participants With Selected Advanced Neoplasms

TARGETS
 CBP, Beta-catenin

LOCATIONS: Glasgow (United Kingdom), Manchester (United Kingdom), Sutton (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
GENE
CCND1
RATIONALE

CCND1 amplification or overexpression may activate CDK4/6 and may predict sensitivity to

single-agent CDK4/6 inhibitors.

ALTERATION

amplification

NCT04282031
PHASE 1/2

A Study of BPI-1178 in Patients With Advanced Solid Tumor and HR+/HER2- Breast Cancer

TARGETS

CDK6, CDK4, ER, Aromatase

LOCATIONS: Shanghai (China)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT05252416
PHASE 1/2

(VELA) Study of BLU-222 in Advanced Solid Tumors

TARGETS

ER, CDK4, CDK6, CDK2

LOCATIONS: Massachusetts

NCT02896335
PHASE 2

Palbociclib In Progressive Brain Metastases

TARGETS

CDK4, CDK6

LOCATIONS: Massachusetts

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
GENE
CDK6
RATIONALE

Tumors with CDK6 amplification may be sensitive to CDK4/6 inhibitors.

ALTERATION

amplification - equivocal

NCT04282031
PHASE 1/2

A Study of BPI-1178 in Patients With Advanced Solid Tumor and HR+/HER2- Breast Cancer

TARGETS

CDK6, CDK4, ER, Aromatase

LOCATIONS: Shanghai (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04594005
PHASE 1/2

CDK4/6 Tumor, Abemaciclib, Paclitaxel

TARGETS

CDK4, CDK6

LOCATIONS: Seoul (Korea, Republic of)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT02693535
PHASE 2

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, CDK4, CDK6, FLT3, CSF1R, KIT, RET, mTOR, EGFR, ERBB2, MEK, BRAF, SMO, DDR2, PARP, PD-1, CTLA-4, ERBB4

LOCATIONS: Hawaii, Washington, Oregon, California

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
NCT03994796
PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, CDK4, CDK6, PI3K, mTOR

LOCATIONS: Washington, Oregon, Idaho, Montana

NCT05252416
PHASE 1/2

(VELA) Study of BLU-222 in Advanced Solid Tumors

TARGETS

ER, CDK4, CDK6, CDK2

LOCATIONS: Massachusetts

NCT02896335
PHASE 2

Palbociclib In Progressive Brain Metastases

TARGETS

CDK4, CDK6

LOCATIONS: Massachusetts

NCT03310879
PHASE 2

Study of the CDK4/6 Inhibitor Abemaciclib in Solid Tumors Harboring Genetic Alterations in Genes Encoding D-type Cyclins or Amplification of CDK4 or CDK6

TARGETS

CDK4, CDK6

LOCATIONS: Massachusetts

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
GENE
MTAP
RATIONALE

MTAP loss may predict sensitivity to MAT2A inhibitors.

ALTERATION

loss exons 6-8

NCT03435250
PHASE 1

Study of AG-270 in Participants With Advanced Solid Tumors or Lymphoma With MTAP Loss

TARGETS
MAT2A
LOCATIONS: Villejuif Cedex (France), Barcelona (Spain), Massachusetts, Connecticut, New York, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
GENE
PIK3CA
ALTERATION
E726K
RATIONALE
 PIK3CA activating mutations may lead to activation of the PI3K-AKT-mTOR pathway and may therefore indicate sensitivity to inhibitors of

this pathway. Strong clinical data support sensitivity of PIK3CA-mutated solid tumors to the PI3K-alpha inhibitor alpelisib.

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS
 TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China), Beijing (China), Chengdu City (China), Changchun (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
 EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS
 mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
 mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS
 mTOR

LOCATIONS: Tianjin (China)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

CLINICAL TRIALS
NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT04632992
PHASE 2

A Study Evaluating Targeted Therapies in Participants Who Have Advanced Solid Tumors With Genomic Alterations or Protein Expression Patterns Predictive of Response

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, PD-L1, ERBB2, PI3K-alpha, RET, AKTs

LOCATIONS: Alaska, Washington, Oregon, California, Idaho

NCT04770246
PHASE 2

TAS-117 in Patients With Advanced Solid Tumors Harboring Germline PTEN Mutations

TARGETS

AKT2, AKT1, AKT3

LOCATIONS: Vienna (Austria), London (United Kingdom), Villejuif (France), California, Ohio, Pennsylvania, Texas

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

AKT1

R15Q

ATM

E2347Q

ATR

S1616L

ATRX

L1517F

BRCA1

E75K

DAXX

R91Q

INPP4B

S219R

IRS2

W1220*

KDM5A

G284D

KMT2A (MLL)

E1992Q

MED12

S1123C

MLL2

S504C

MSH6

K1358fs*2

NOTCH1

R2549C

P2RY8

E323G

PRKCI

I156M

SGK1

E46A

TEK

E434K

TYRO3

S648F

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKKN1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV1
ETV4	ETVS	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT
KMT2A (MLL)	MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA
RAF1	RARA	RET	ROS1	RSPO2	SDC4	SLC34A2	TERC*	TERT**
TPRSS2								

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.



ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 - Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 291656669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 6.1.0

The median exon coverage for this sample is 721x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX
References

1. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
2. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
3. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
4. Cristescu R, et al. Science (2018) PMID: 30309915
5. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
6. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
7. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
8. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
9. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
10. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
11. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
12. Legrand et al., 2018; ASCO Abstract 12000
13. Mai HQ, et al. Nat Med (2021) PMID: 34341578
14. Wang FH, et al. J Clin Oncol (2021) PMID: 33492986
15. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
16. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
17. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
18. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
19. Rizvi NA, et al. Science (2015) PMID: 25765070
20. Johnson BE, et al. Science (2014) PMID: 24336570
21. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
22. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
23. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
24. Heitzner E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
25. Nature (2012) PMID: 22810696
26. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
27. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
28. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
29. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
30. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
31. Ayers et al., 2016; ASCO-SITC Abstract P60
32. Zigelboim I, et al. J. Clin. Oncol. (2007) PMID: 17513808
33. Hampel H, et al. Cancer Res. (2006) PMID: 16885385
34. Stelloo E, et al. Clin. Cancer Res. (2016) PMID: 27006490
35. Kanopienė D, et al. Medicina (Kaunas) (2014) PMID: 25458958
36. Black D, et al. J. Clin. Oncol. (2006) PMID: 16549821
37. Nout RA, et al. Gynecol. Oncol. (2012) PMID: 22609107
38. Steinbakk A, et al. Cell Oncol (Dordr) (2011) PMID: 21547578
39. Bilbao C, et al. Int. J. Radiat. Oncol. Biol. Phys. (2010) PMID: 20005452
40. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
41. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
42. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
43. Hiyama T, et al. J. Gastroenterol. Hepatol. (2004) PMID: 15209621
44. Wu MS, et al. Cancer Res. (1998) PMID: 9537253
45. dos Santos NR, et al. Gastroenterology (1996) PMID: 8536886
46. Fang WL, et al. Biomed Res Int (2013) PMID: 23555086
47. Farris AB, et al. Am. J. Surg. Pathol. (2011) PMID: 21422910
48. Agaram NP, et al. Am. J. Clin. Pathol. (2010) PMID: 20395525
49. Ruemmele P, et al. Am. J. Surg. Pathol. (2009) PMID: 19252434
50. Planck M, et al. Cancer (2003) PMID: 12627520
51. Hibi K, et al. Jpn. J. Cancer Res. (1995) PMID: 7775257
52. Muneyuki T, et al. Dig. Dis. Sci. (2000) PMID: 11117578
53. Zhang SH, et al. World J. Gastroenterol. (2005) PMID: 15918185
54. Chiappini F, et al. Carcinogenesis (2004) PMID: 14656944
55. Suto T, et al. J Surg Oncol (2001) PMID: 11223838
56. Momoi H, et al. J. Hepatol. (2001) PMID: 11580146
57. Liengswangwong U, et al. Int. J. Cancer (2003) PMID: 14506736
58. Moy AP, et al. Virchows Arch. (2015) PMID: 25680569
59. Yoshida T, et al. J. Gastroenterol. (2000) PMID: 11063221
60. Pritchard CC, et al. Nat Commun (2014) PMID: 25255306
61. Azzouzi AR, et al. BJU Int. (2007) PMID: 17233803
62. Burger M, et al. J. Mol. Med. (2006) PMID: 16924473
63. Bai S, et al. Am. J. Clin. Pathol. (2013) PMID: 23690119
64. Giedl J, et al. Am. J. Clin. Pathol. (2014) PMID: 25319978
65. Yamamoto Y, et al. Clin. Cancer Res. (2006) PMID: 16675567
66. Smyth et al., 2015; ASCO Gastrointestinal Cancers Symposium Abstract 62
67. Bilbao-Sieyro C, et al. Oncotarget (2014) PMID: 25026289
68. Mackay HJ, et al. Eur. J. Cancer (2010) PMID: 20304627
69. Arabi H, et al. Gynecol. Oncol. (2009) PMID: 19275958
70. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
71. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
72. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
73. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
74. Fritsch C, et al. Mol. Cancer Ther. (2014) PMID: 24608574
75. Juric D, et al. J. Clin. Oncol. (2018) PMID: 29401002
76. Gallant JN, et al. NPJ Precis Oncol (2019) PMID: 30793038
77. Delestre F, et al. Sci Transl Med (2021) PMID: 34613809
78. Morschhauser F, et al. Mol Cancer Ther (2020) PMID: 31619463
79. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
80. Santin AD, et al. Gynecol Oncol Rep (2020) PMID: 31934607
81. Damodaran S, et al. J Clin Oncol (2022) PMID: 35133871
82. André F, et al. N. Engl. J. Med. (2019) PMID: 31091374
83. Smyth LM, et al. NPJ Breast Cancer (2021) PMID: 33863913
84. Park HS, et al. PLoS ONE (2016) PMID: 27105424
85. Lim SM, et al. Oncotarget (2016) PMID: 26859683
86. Hou MM, et al. Oncotarget (2014) PMID: 25426553
87. Varnier R, et al. Eur J Cancer (2019) PMID: 31351267
88. Janku F, et al. Cell Rep (2014) PMID: 24440717
89. Moroney J, et al. Clin. Cancer Res. (2012) PMID: 22927482
90. Basho RK, et al. JAMA Oncol (2017) PMID: 27893038
91. Moroney JW, et al. Clin. Cancer Res. (2011) PMID: 21890452
92. Krop et al., 2018; ASCO Abstract 101
93. Pascual J, et al. Cancer Discov (2021) PMID: 32958578
94. Dolly SO, et al. Clin. Cancer Res. (2016) PMID: 26787751
95. Canaud et al., 2021; ESMO Abstract LBA23
96. Aust Fam Physician (1986) PMID: 2941002
97. Chou CC, et al. Med. Oncol. (2009) PMID: 19012001
98. Zhang ZC, et al. Onco Targets Ther (2014) PMID: 24672248
99. Lin DC, et al. Nat. Genet. (2014) PMID: 24952746
100. Or YY, et al. Int. J. Cancer (2006) PMID: 16114017
101. Fendri A, et al. Cancer Sci. (2009) PMID: 19735264
102. Jiang N, et al. Oncol. Rep. (2014) PMID: 25109408
103. Zhang JW, et al. Chin J Cancer (2015) PMID: 25963410
104. Hui AB, et al. Int. J. Oncol. (2002) PMID: 11836556
105. Yip WK, et al. Pathol. Oncol. Res. (2016) PMID: 26581613
106. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
107. Lui VW, et al. Cancer Discov (2013) PMID: 23619167
108. Suda T, et al. BMC Cancer (2012) PMID: 22994622
109. Fenic I, et al. Oncol. Rep. (2007) PMID: 17549376
110. Sticht C, et al. Br. J. Cancer (2005) PMID: 15700036
111. Samuels Y, et al. Cancer Cell (2005) PMID: 15950905
112. Nat. Rev. Cancer (2009) PMID: 19629070
113. Kang S, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15647370
114. Ikenoue T, et al. Cancer Res. (2005) PMID: 15930273
115. Gymnopoulos M, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17376864
116. Horn S, et al. Oncogene (2008) PMID: 18317450
117. Rudd ML, et al. Clin. Cancer Res. (2011) PMID: 21266528
118. Hon WC, et al. Oncogene (2012) PMID: 22120714
119. Burke JE, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22949682
120. Wu H, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19915146
121. Laurenti R, et al. Rev Saude Publica (1990) PMID: 2103068
122. Dan S, et al. Cancer Res. (2010) PMID: 20530683
123. Oda K, et al. Cancer Res. (2008) PMID: 18829572
124. Zhao L, et al. Oncogene (2008) PMID: 18794883
125. Ross RL, et al. Oncogene (2013) PMID: 22430209
126. Rivière JB, et al. Nat. Genet. (2012) PMID: 22729224
127. Shibata T, et al. Cancer Lett. (2009) PMID: 19394761
128. Dogruluk T, et al. Cancer Res. (2015) PMID: 26627007
129. Croessmann S, et al. Clin. Cancer Res. (2018) PMID: 29284706
130. Ng PK, et al. Cancer Cell (2018) PMID: 29533785
131. Spangle JM, et al. (2020) PMID: 32929011
132. Chen L, et al. Nat Commun (2018) PMID: 29636477
133. Jin N, et al. J Clin Invest (2021) PMID: 34779417
134. Zhan T, et al. Oncogene (2017) PMID: 27617575
135. Jung YS, et al. Exp Mol Med (2020) PMID: 32037398
136. Krishnamurthy N, et al. Cancer Treat Rev (2018) PMID: 29169144
137. Kawazoe et al., 2021; ESMO Abstract 473P
138. Yamada K, et al. Cancer Res (2021) PMID: 33408116
139. Kanda Y, et al. Biochem Biophys Res Commun (2022) PMID: 34837838
140. Yom SS, et al. Mod. Pathol. (2005) PMID: 15492756
141. Frattini M, et al. Head Neck (2006) PMID: 16906516
142. Stephen JK, et al. Int J Head Neck Surg (2010) PMID: 21603083
143. Loyo M, et al. Int. J. Cancer (2011) PMID: 20473931
144. Chen K, et al. Arch. Otolaryngol. Head Neck Surg. (2007) PMID: 18025318
145. La Rosa S, et al. Head Neck Pathol (2013) PMID: 22740238
146. Luke JJ, et al. Clin Cancer Res (2019) PMID: 30635339
147. Logan CY, et al. Annu. Rev. Cell Dev. Biol. (2004) PMID: 15647370

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX **References**

- 15473860
148. Eklof Spink K, et al. EMBO J. (2001) PMID: 11707392
149. Liu J, et al. J. Mol. Biol. (2006) PMID: 16753179
150. Dikovskaya D, et al. J. Cell. Sci. (2010) PMID: 20144988
151. Murphy SJ, et al. Dig. Dis. Sci. (2007) PMID: 17410430
152. Aretz S, et al. Hum. Mutat. (2004) PMID: 15459959
153. Kerr SE, et al. J Mol Diagn (2013) PMID: 23159591
154. Annu Rev Pathol (2011) PMID: 21090969
155. Kastritis E, et al. Int. J. Cancer (2009) PMID: 18844223
156. Half E, et al. Orphanet J Rare Dis (2009) PMID: 19822006
157. Morschhauser F, et al. Haematologica (2020) PMID: 32381571
158. Flaherty KT, et al. Clin. Cancer Res. (2012) PMID: 22090362
159. Finn RS, et al. Lancet Oncol. (2015) PMID: 25524798
160. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
161. Patnaik A, et al. Cancer Discov (2016) PMID: 27217383
162. Leonard JP, et al. Blood (2012) PMID: 22383795
163. Dickler MN, et al. Clin. Cancer Res. (2017) PMID: 28533223
164. Clark et al., 2019; AACR Abstract LB-010/2
165. Peguero et al., 2016; ASCO Abstract 2528
166. Chen Y, et al. Front Immunol (2020) PMID: 32903763
167. Litchfield K, et al. Cell (2021) PMID: 33508232
168. Nature (2015) PMID: 25631445
169. Hanken H, et al. Clin Oral Investig (2014) PMID: 23494454
170. Rodrigo JP, et al. J. Pathol. (2009) PMID: 18991334
171. Sakashita T, et al. Acta Otolaryngol. (2013) PMID: 23692432
172. Maruyama S, et al. Cancer Genet. Cytogenet. (2010) PMID: 19963131
173. Rasamny JJ, et al. Otolaryngol Head Neck Surg (2012) PMID: 22323434
174. Sugahara K, et al. Int. J. Oncol. (2011) PMID: 21701773
175. Zhang P, et al. BMC Cancer (2006) PMID: 16978399
176. Elsheikh S, et al. Breast Cancer Res. Treat. (2008) PMID: 17653856
177. Fu M, et al. Endocrinology (2004) PMID: 15331580
178. Takahashi-Yanaga F, et al. Cell. Signal. (2008) PMID: 18023328
179. Turner NC, et al. N. Engl. J. Med. (2015) PMID: 26030518
180. Konecny et al., 2016; ASCO Abstract 5557
181. Meyerson M, et al. Mol. Cell. Biol. (1994) PMID: 8114739
182. Gossel MJ, et al. J. Cell. Biochem. (2006) PMID: 16294322
183. Choi YJ, et al. Oncogene (2014) PMID: 23644662
184. Cell (1995) PMID: 7736585
185. Musgrave EA, et al. Nat. Rev. Cancer (2011) PMID: 21734724
186. Ismail A, et al. Clin. Cancer Res. (2011) PMID: 21593195
187. van Dekken H, et al. Cancer Genet. Cytogenet. (2009) PMID: 19167610
188. Marjon K, et al. Cell Rep (2016) PMID: 27068473
189. Heist et al., 2019; AACR-NCI-EORTC Abstract B116
190. Mavrakis KJ, et al. Science (2016) PMID: 26912361
191. Endoscopy (1989) PMID: 2691236
192. Guccione E, et al. Nat. Rev. Mol. Cell Biol. (2019) PMID: 31350521
193. Fedoriv A, et al. Cancer Cell (2019) PMID: 31257072
194. Srour N, et al. Cancer Cell (2019) PMID: 31287990
195. Gao G, et al. Nucleic Acids Res. (2019) PMID: 30916320
196. Hansen LJ, et al. Cancer Res. (2019) PMID: 31040154
197. Tang B, et al. Cancer Res. (2018) PMID: 29844120
198. Munshi PN, et al. Oncologist (2014) PMID: 24928612
199. de Oliveira SF, et al. PLoS ONE (2016) PMID: 26751376
200. Lubin M, et al. PLoS ONE (2009) PMID: 19478948
201. Tang B, et al. Cancer Biol. Ther. (2012) PMID: 22825330
202. Collins CC, et al. Mol. Cancer Ther. (2012) PMID: 22252602
203. Bertino JR, et al. Cancer Biol. Ther. (2011) PMID: 21301207
204. Coulthard SA, et al. Mol. Cancer Ther. (2011) PMID: 21282358
205. Miyazaki S, et al. Int. J. Oncol. (2007) PMID: 17912432
206. Efferth T, et al. Blood Cells Mol. Dis. () PMID: 11987241
207. Kindler HL, et al. Invest New Drugs (2009) PMID: 18618081
208. Wei R, et al. Sci Rep (2016) PMID: 27929028
209. Zhao M, et al. BMC Genomics (2016) PMID: 27556634
210. Kirovski G, et al. Am. J. Pathol. (2011) PMID: 21356366
211. Huang HY, et al. Clin. Cancer Res. (2009) PMID: 19887491
212. Marcé S, et al. Clin. Cancer Res. (2006) PMID: 16778103
213. Meyer S, et al. Exp. Dermatol. (2010) PMID: 20500769
214. Wild PJ, et al. Arch Dermatol (2006) PMID: 16618867
215. Kim J, et al. Genes Chromosomes Cancer (2011) PMID: 21412930
216. Li CF, et al. Oncotarget (2014) PMID: 25426549
217. He HL, et al. Medicine (Baltimore) (2015) PMID: 26656376
218. Su CY, et al. Eur J Surg Oncol (2014) PMID: 24969958
219. Mirebeau D, et al. Haematologica (2006) PMID: 16818274
220. Becker AP, et al. Pathobiology (2015) PMID: 26088413
221. Snezhkina AV, et al. Oxid Med Cell Longev (2016) PMID: 27433286
222. Bistulfi G, et al. Oncotarget (2016) PMID: 26910893
223. Antonopoulou K, et al. J. Invest. Dermatol. (2015) PMID: 25407435
224. Maccioni L, et al. BMC Cancer (2013) PMID: 23816148
225. Hyland PL, et al. Int J Epidemiol (2016) PMID: 26635288
226. Lin X, et al. Cancer Sci. (2017) PMID: 27960044
227. Zhi L, et al. J Cancer (2016) PMID: 27994653
228. Gu F, et al. Br. J. Cancer (2013) PMID: 23361049
229. Limm K, et al. PLoS ONE (2016) PMID: 27479139
230. Tang B, et al. G3 (Bethesda) (2014) PMID: 25387827
231. Limm K, et al. Eur. J. Cancer (2013) PMID: 23265702
232. Stevens AP, et al. J. Cell. Biochem. (2009) PMID: 19097084
233. Kryukov GV, et al. Science (2016) PMID: 26912360
234. Limm K, et al. Eur. J. Cancer (2014) PMID: 25087184
235. Konecny GE, et al. Clin. Cancer Res. (2011) PMID: 21278246
236. Katsumi Y, et al. Biochem. Biophys. Res. Commun. (2011) PMID: 21871868
237. Cen L, et al. Neuro-oncology (2012) PMID: 22711607
238. Logan JE, et al. Anticancer Res. (2013) PMID: 23898052
239. Elvin JA, et al. Oncologist (2017) PMID: 28283584
240. Gao J, et al. Curr Oncol (2015) PMID: 26715889
241. Gopalan et al., 2014; ASCO Abstract 8077
242. DeMichele A, et al. Clin. Cancer Res. (2015) PMID: 25501126
243. Johnson DB, et al. Oncologist (2014) PMID: 24797823
244. Van Maerken T, et al. Mol. Cancer Ther. (2011) PMID: 21460101
245. Gamble LD, et al. Oncogene (2012) PMID: 21725357
246. Shapiro et al., 2013; ASCO Abstract 2500
247. Dickson MA, et al. J. Clin. Oncol. (2013) PMID: 23569312
248. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
249. Gao J, et al. Sci Signal (2013) PMID: 23550210
250. Lo KW, et al. Cancer Res. (1995) PMID: 7743498
251. Bei JX, et al. Nat. Genet. (2010) PMID: 20512145
252. Wong TS, et al. Int. J. Oncol. (2003) PMID: 12632081
253. Wong TS, et al. Clin. Cancer Res. (2004) PMID: 15073117
254. Pierini S, et al. Head Neck (2014) PMID: 23804521
255. Lechner M, et al. Genome Med (2013) PMID: 23718828
256. Reed AL, et al. Cancer Res. (1996) PMID: 8705996
257. Mäkitie AA, et al. Clin. Cancer Res. (2003) PMID: 12796384
258. Hwang CF, et al. Ann. Oncol. (2002) PMID: 12181248
259. Quelle DE, et al. Cell (1995) PMID: 8521522
260. Mutat. Res. (2005) PMID: 15878778
261. Gazzeri S, et al. Oncogene (1998) PMID: 9484839
262. Oncogene (1999) PMID: 10498883
263. Sherr CJ, et al. Cold Spring Harb. Symp. Quant. Biol. (2005) PMID: 16869746
264. Ozenne P, et al. Int. J. Cancer (2010) PMID: 20549699
265. Ruas M, et al. Oncogene (1999) PMID: 10498896
266. Jones R, et al. Cancer Res. (2007) PMID: 17909018
267. Haferkamp S, et al. Aging Cell (2008) PMID: 18843795
268. Huot TJ, et al. Mol. Cell. Biol. (2002) PMID: 12417717
269. Rizo H, et al. J. Biol. Chem. (2001) PMID: 11518711
270. Gombart AF, et al. Leukemia (1997) PMID: 9324288
271. Yang R, et al. Cancer Res. (1995) PMID: 7780957
272. Parry D, et al. Mol. Cell. Biol. (1996) PMID: 8668202
273. Greenblatt MS, et al. Oncogene (2003) PMID: 12606942
274. Yarbrough WG, et al. J. Natl. Cancer Inst. (1999) PMID: 10491434
275. Poi MJ, et al. Mol. Carcinog. (2001) PMID: 11255261
276. Byeon IJ, et al. Mol. Cell (1998) PMID: 9660926
277. Kannengiesser C, et al. Hum. Mutat. (2009) PMID: 19260062
278. Lal G, et al. Genes Chromosomes Cancer (2000) PMID: 10719365
279. Koh J, et al. Nature (1995) PMID: 7777061
280. McKenzie HA, et al. Hum. Mutat. (2010) PMID: 20340136
281. Miller PJ, et al. Hum. Mutat. (2011) PMID: 21462282
282. Kutscher CL, et al. Physiol. Behav. (1977) PMID: 905385
283. Scaini MC, et al. Hum. Mutat. (2014) PMID: 24659262
284. Jenkins NC, et al. J. Invest. Dermatol. (2013) PMID: 23190892
285. Walker GJ, et al. Int. J. Cancer (1999) PMID: 10389768
286. Rutter JL, et al. Oncogene (2003) PMID: 12853981
287. Itahana K, et al. Cancer Cell (2008) PMID: 18538737
288. Zhang Y, et al. Mol. Cell (1999) PMID: 10360174
289. Zhang Y, et al. Cell (1998) PMID: 9529249
290. Jaffr M, et al. Cancer Discov (2015) PMID: 25873077
291. Whelan AJ, et al. N Engl J Med (1995) PMID: 7666917
292. Adv Exp Med Biol (2010) PMID: 20687502
293. Hogg D, et al. J Cutan Med Surg (1998) PMID: 9479083
294. De Unamuno B, et al. Melanoma Res (2018) PMID: 29543703
295. Soura E, et al. J Am Acad Dermatol (2016) PMID: 26892650
296. Huerta C, et al. Acta Derm Venereol (2018) PMID: 29405243
297. Kaufman DK, et al. Neurology (1993) PMID: 8414022
298. Bahuau M, et al. Cancer Res (1998) PMID: 9622062
299. Chan AK, et al. Clin Neuropathol () PMID: 28699883
300. Kertesz N, et al. Blood (2006) PMID: 16322467
301. Shi S, et al. J Pharm Sci (2012) PMID: 22411527
302. Li X, et al. PLoS ONE (2014) PMID: 25148033
303. Liu R, et al. BMC Cancer (2013) PMID: 23721559

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX **References**

304. Bhatia S, et al. Sci Rep (2016) PMID: 27941840
305. Sehn JS, et al. Blood (2009) PMID: 18836096
306. Ferguson BD, et al. Sci Rep (2015) PMID: 26073592
307. Werner TL, et al. Invest New Drugs (2015) PMID: 26365907
308. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22011666
309. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22722787
310. Sinha UK, et al. Arch. Otolaryngol. Head Neck Surg. (2006) PMID: 17043250
311. Masood R, et al. Int. J. Cancer (2006) PMID: 16615113
312. Ferguson BD, et al. Growth Factors (2014) PMID: 25391996
313. Yavrouian EJ, et al. Arch. Otolaryngol. Head Neck Surg. (2008) PMID: 18794445
314. Liersch-Löhn B, et al. Int. J. Cancer (2016) PMID: 26414866
315. Hu F, et al. Tumour Biol. (2014) PMID: 24771266
316. Hasina R, et al. Cancer Res. (2013) PMID: 23100466
317. Li M, et al. Dig. Dis. Sci. (2011) PMID: 20686847
318. Yin J, et al. Anticancer Res. (2017) PMID: 28739744
319. Stephenson SA, et al. BMC Mol. Biol. (2001) PMID: 11801186
320. Liu W, et al. Cancer (2002) PMID: 11920461
321. McCall JL, et al. Mol. Cell. Biol. (2016) PMID: 27273865
322. Stremtitz S, et al. Mol. Cancer Ther. (2016) PMID: 27535973
323. Lv J, et al. Exp. Mol. Pathol. (2016) PMID: 27072105
324. Guíjarro-Muñoz I, et al. Med. Oncol. (2013) PMID: 23579861
325. Kumar SR, et al. Cancer Res. (2009) PMID: 19366806
326. Wu Q, et al. Pathol. Oncol. Res. (2004) PMID: 15029258
327. Berclaz G, et al. Oncol. Rep. () PMID: 12168060
328. Brantley-Sieders DM, et al. PLoS ONE (2011) PMID: 21935409
329. Huang G, et al. Int J Clin Exp Pathol (2015) PMID: 26191333
330. Pradeep S, et al. Cancer Cell (2015) PMID: 26481148
331. Alam SM, et al. Br. J. Cancer (2008) PMID: 18231102
332. Kumar SR, et al. Br. J. Cancer (2007) PMID: 17353927
333. Takai N, et al. Oncol. Rep. () PMID: 11295082
334. Dong LD, et al. Oncol Lett (2017) PMID: 28454369
335. Berclaz G, et al. Ann. Oncol. (2003) PMID: 12562648
336. Sharma GK, et al. Head Neck (2015) PMID: 24634162
337. Giaginis C, et al. Pathol. Oncol. Res. (2016) PMID: 26220827
338. Xuqing W, et al. Tumour Biol. (2012) PMID: 22528941
339. Ferguson BD, et al. PLoS ONE (2013) PMID: 23844053
340. Zheng MF, et al. Mol Med Rep (2012) PMID: 22684742
341. Chen T, et al. Tumour Biol. (2013) PMID: 23138393
342. Tu Y, et al. Clin Transl Oncol (2012) PMID: 22374425
343. Li M, et al. Mol. Biol. Rep. (2013) PMID: 23079712
344. Xia G, et al. Cancer Res. (2005) PMID: 15930280
345. Alam SM, et al. Gynecol. Oncol. (2009) PMID: 19356789
346. Özgür E, et al. Urol. Oncol. () PMID: 19272799
347. Pierscianek D, et al. Neuropathology (2017) PMID: 27388534
348. Pierscianek D, et al. Brain Tumor Pathol (2016) PMID: 26951238
349. Becerikli M, et al. Int. J. Cancer (2015) PMID: 25274141
350. Xia G, et al. Clin. Cancer Res. (2005) PMID: 15958611
351. Noren NK, et al. Cancer Res. (2007) PMID: 17483308
352. Cell (2008) PMID: 18394988
353. Salvucci O, et al. Adv. Cancer Res. (2012) PMID: 22588055
354. Pitulescu ME, et al. Genes Dev. (2010) PMID: 21078817
355. Nat. Rev. Cancer (2010) PMID: 20179713
356. Boberg DR, et al. Chem. Biol. Interact. (2013) PMID: 23063927
357. Cromer A, et al. Oncogene (2004) PMID: 14676830
358. Kim RD, et al. Cancer Discov (2019) PMID: 31575541
359. Chan et al., 2017; AACR Abstract CT106
360. Macarulla et al., 2021; ASCO Abstract 4090
361. Kaibori M, et al. Oncotarget (2016) PMID: 27384874
362. Kanzaki H, et al. Sci Rep (2021) PMID: 33674622
363. Dumbrava EI, et al. JCO Precis Oncol (2018) PMID: 31123723
364. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
365. Nature (2012) PMID: 22960745
366. Robertson AG, et al. Cell (2017) PMID: 28988769
367. Miura S, et al. BMC Cancer (2012) PMID: 22309595
368. Kang HJ, et al. Liver Cancer (2019) PMID: 30815392
369. Nagamatsu H, et al. Prostate (2015) PMID: 25854696
370. Gao L, et al. Oncogene (2019) PMID: 30518874
371. Li F, et al. Oncogene (2020) PMID: 32111983
372. Xie MH, et al. Cytokine (1999) PMID: 10525310
373. Hagel M, et al. Cancer Discov (2015) PMID: 25776529
374. Int. J. Oncol. (2002) PMID: 12429977
375. Kan Z, et al. Genome Res. (2013) PMID: 23788652
376. Caruso S, et al. Gastroenterology (2019) PMID: 31063779
377. Sawey ET, et al. Cancer Cell (2011) PMID: 21397858
378. Dumbrava et al., 2018; doi/full/10.1200/PO.18.00100
379. Tekin M, et al. Am. J. Hum. Genet. (2007) PMID: 17236138
380. Arai T, et al. Hepatology (2013) PMID: 22890726
381. Yamada T, et al. BMC Cancer (2015) PMID: 25885470
382. Kratochwil K, et al. Genes Dev. (2002) PMID: 12502739
383. Scherz PJ, et al. Science (2004) PMID: 15256670
384. Zaharieva BM, et al. J. Pathol. (2003) PMID: 14648664
385. Arai H, et al. Cancer Genet. Cytogenet. (2003) PMID: 14499691
386. Ribeiro IP, et al. Tumour Biol. (2014) PMID: 24477574
387. Schulze K, et al. Nat. Genet. (2015) PMID: 25822088
388. Debeb BG, et al. Breast Cancer Res. Treat. (2012) PMID: 22547109
389. Fouladi M, et al. J. Clin. Oncol. (2011) PMID: 21825264
390. Groth C, et al. Semin. Cell Dev. Biol. (2012) PMID: 22309842
391. Kamstrup MR, et al. Blood (2010) PMID: 20538790
392. Kridel R, et al. Blood (2012) PMID: 22210878
393. Krop I, et al. J. Clin. Oncol. (2012) PMID: 22547604
394. Rosati E, et al. Int. J. Cancer (2013) PMID: 23001755
395. Samon JB, et al. Mol. Cancer Ther. (2012) PMID: 22504949
396. Lehal R, et al. Proc Natl Acad Sci U S A (2020) PMID: 32601208
397. Ferrarotto et al., 2020; ESMO Abstract 919MO
398. Knoechel B, et al. Cold Spring Harb Mol Case Stud (2015) PMID: 27148573
399. Lopez-Miranda et al., 2021; ASCO Abstract 3020
400. Dreyling M, et al. Ann. Oncol. (2017) PMID: 28633365
401. Palomero T, et al. Nat. Med. (2007) PMID: 17873882
402. Liu S, et al. Urol. Oncol. (2013) PMID: 21993533
403. Agrawal N, et al. Science (2011) PMID: 21798897
404. Stransky N, et al. Science (2011) PMID: 21798893
405. Wang NJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2011) PMID: 22006338
406. Klinakis A, et al. Nature (2011) PMID: 21562564
407. Penton AL, et al. Semin. Cell Dev. Biol. (2012) PMID: 22306179
408. Kopan R, et al. Cell (2009) PMID: 19379690
409. Andrawes MB, et al. J. Biol. Chem. (2013) PMID: 23839946
410. Rebay I, et al. Cell (1991) PMID: 1657403
411. Ge C, et al. BMC Dev. Biol. (2008) PMID: 18445292
412. Aster JC, et al. Mol. Cell. Biol. (2000) PMID: 11003647
413. Weng AP, et al. Science (2004) PMID: 15472075
414. Deregowski V, et al. J. Bone Miner. Res. (2006) PMID: 16869730
415. Uchibori M, et al. Oncol. Rep. (2017) PMID: 28791383
416. Liu J, et al. Proc Natl Acad Sci U S A (2013) PMID: 24277854
417. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
418. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
419. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
420. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
421. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
422. Xu L, et al. Mol. Med. (2001) PMID: 11713371
423. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
424. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
425. Pirolo KF, et al. Mol. Ther. (2016) PMID: 27357628
426. Hajdenberg et al., 2012; ASCO Abstract e15010
427. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
428. Moore et al., 2019; ASCO Abstract 5513
429. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
430. Oza et al., 2015; ASCO Abstract 5506
431. Lee J, et al. Cancer Discov (2019) PMID: 31315834
432. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
433. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
434. Lehmann S, et al. J. Clin. Oncol. (2012) PMID: 22965953
435. Mohell N, et al. Cell Death Dis (2015) PMID: 26086967
436. Fransson Å, et al. J Ovarian Res (2016) PMID: 27179933
437. Gourley et al., 2016; ASCO Abstract 5571
438. Kwok M, et al. Blood (2016) PMID: 26563132
439. Boudny M, et al. Haematologica (2019) PMID: 30975914
440. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
441. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
442. Bolt J, et al. Oral Oncol. (2005) PMID: 16139561
443. Yamazaki Y, et al. Oral Oncol. (2003) PMID: 12509970
444. McBride SM, et al. Head Neck (2014) PMID: 23852799
445. Pickering CR, et al. Cancer Discov (2013) PMID: 23619168
446. Ma BB, et al. Head Neck (2003) PMID: 12966511
447. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
448. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
449. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
450. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
451. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
452. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
453. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
454. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
455. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
456. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # ORD-1346479-01

APPENDIX
References

457. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
458. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
459. Lalloo F, et al. Lancet (2003) PMID: 12672316
460. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713
461. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
462. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
463. Xie M, et al. Nat. Med. (2014) PMID: 25326804
464. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
465. Severson EA, et al. Blood (2018) PMID: 29678827
466. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
467. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
468. Chabon JJ, et al. Nature (2020) PMID: 32269342
469. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
470. Marabelle et al., 2019; ESMO Abstract 11920
471. Andre et al., 2021; ASCO GI Abstract 9
472. Oaknin A, et al. JAMA Oncol (2020) PMID: 33001143
473. Berton et al., 2021; ASCO Abstract 2564
474. Andre et al., 2021; ESMO GI Abstract SO-9
475. Hsu C, et al. J. Clin. Oncol. (2017) PMID: 28837405
476. Colevas AD, et al. Ann. Oncol. (2018) PMID: 30219915
477. Barlesi et al., 2020; SITC Abstract 291
478. Jung KH, et al. Clin. Cancer Res. (2019) PMID: 30770348
479. Okauchi S, et al. In Vivo () PMID: 31882503
480. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
481. Verschraegen et al., 2016; ASCO Abstract 9036
482. Chung et al., 2016; ASCO Abstract 4009
483. Patel et al., 2016; ESMO Abstract 777PD
484. Hassan et al., 2016; ASCO Abstract 8503
485. Disis et al., 2016; ASCO Abstract 5533
486. Dirix et al., 2016; SABCS Abstract S1-04
487. Larkin et al., 2016; ESMO Abstract 775PD
488. Le Tourneau et al., 2016; ASCO Abstract 4516
489. Fakhrejahani et al., 2017; ASCO GU Abstract 159
490. Rajan et al., 2016; ASCO Abstract e20106
491. Migden MR, et al. N. Engl. J. Med. (2018) PMID: 29863979
492. Stratigos et al., 2020; EMSO Abstract LBA47
493. Lewis et al. 2020; doi: 10.1136/jitc-2020-SITC2020.0428
494. Sezer et al., 2020; ESMO Abstract LBA52
495. Bais et al., 2017; AACR Abstract 3720/5
496. Garassino et al., 2016; IASLC Abstract PLO4a.03
497. Segal et al., 2016; ESMO Abstract 9490
498. Segal et al., 2015; ASCO Abstract 3011
499. Lutzky et al., 2014; ASCO Abstract 3001
500. Iguchi et al., 2015; ASCO Abstract 3039
501. Ribas et al., 2015; ASCO Abstract 3003
502. Karzai et al., 2017; ASCO Genitourinary Abstract 162
503. Lee et al., 2016; ASCO Abstract 3015
504. Necchi A, et al. Eur. Urol. (2019) PMID: 30243800
505. Hamid et al., 2016; ESMO Abstract 1050PD
506. Hong et al., 2016; ESMO 2016 Abstract 1049PD
507. Yap et al., 2016; EORTC-NCI-AACR Abstract 1LBA
508. Levy A, et al. Eur. J. Cancer (2016) PMID: 27764686
509. Janku F, et al. Cancer Res. (2013) PMID: 23066039
510. Janku F, et al. J. Clin. Oncol. (2012) PMID: 22271473
511. Janku F, et al. Mol. Cancer Ther. (2011) PMID: 21216929
512. Moulder S, et al. Ann. Oncol. (2015) PMID: 25878190
513. Byeon et al., 2020; doi: 10.21037/tcr.2020.04.07
514. Varadarajan et al., 2012; ASCO Abstract 5541
515. Fury MG, et al. Cancer Chemother. Pharmacol. (2012) PMID: 21913034
516. Fury MG, et al. Cancer (2013) PMID: 23408298
517. Tolcher AW, et al. Ann. Oncol. (2015) PMID: 25344362
518. Patterson et al., 2018; AACR Abstract 3891
519. Ma BBY, et al. J. Clin. Oncol. (2018) PMID: 29584545
520. Delord et al., 2017; ASCO Abstract 6025
521. Hodi et al., 2019; AACR abstract CT037
522. Kao et al., 2020; ESMO Abstract 2660
523. Grünwald V, et al. Ann. Oncol. (2015) PMID: 25527417
524. Trafalis DT, et al. Anticancer Drugs (2012) PMID: 22510794
525. MacKenzie MJ, et al. Invest New Drugs (2012) PMID: 20978924
526. Fury MG, et al. Cancer Chemother. Pharmacol. (2012) PMID: 22644799
527. Bauman JE, et al. Oral Oncol. (2013) PMID: 23384718

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 25 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531