



## Sample Information

**Patient Name:** 朱森堯  
**Gender:** Male  
**ID No.:** A124992382  
**History No.:** 40887318  
**Age:** 42

**Ordering Doctor:** DOC8716K 李怡緻  
**Ordering REQ.:** OBNMFRH  
**Signing in Date:** 2021/11/18

**Path No.:** S110-89554  
**MP No.:** F21096  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S110-04165C  
**Percentage of tumor cells:** 40%

**Reporting Doctor:** DOC5466K 葉奕成 (Phone: 8#5466)

**Note:**

## Sample Cancer Type: Gastric Cancer

Table of Contents	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Alert Details	3

**Report Highlights**  
1 Relevant Biomarkers  
0 Therapies Available  
0 Clinical Trials

## Relevant Gastric Cancer Variants

Gene	Finding
ERBB2	None detected
NTRK1	None detected
NTRK2	None detected
NTRK3	None detected

## Relevant Biomarkers

No relevant biomarkers found in this sample.

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### Copy Number Variations

Gene	Locus	Copy Number
FGFR2	chr10:123247505	78.93

## Biomarker Descriptions

### FGFR2 (fibroblast growth factor receptor 2)

**Background:** The FGFR2 gene encodes fibroblast growth receptor 2, a member of the fibroblast growth- factor receptor (FGFR) family that also includes FGFR1, 3, and 4. These proteins are single-transmembrane receptors composed of three extracellular immunoglobulin (Ig)-type domains and an intracellular kinase domain. Upon FGF-mediated stimulation, FGFRs activate several oncogenic signaling pathways, including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, PLC/PKC, and JAK/STAT pathways influencing cell proliferation, migration, and survival<sup>1,2,3</sup>.

**Alterations and prevalence:** Aberrations most common to the FGFR family are amplifications, followed by mutations and fusions. The majority of these aberrations result in gain of function<sup>4</sup>. Missense mutations are the most prevalent alterations in FGFR2 and are observed in up to 15% of uterine carcinomas<sup>5,6,7</sup>. These mutations are predominantly activating, most often involve substitutions at S252 and P253, and confer sensitivity to pan-FGFR2 inhibitors<sup>5,8</sup>. FGFR2 amplification occurs in up to 4% of gastric carcinoma, and is associated with poor prognosis as well as tumor invasion and metastasis<sup>6,9,10,11</sup>. FGFR2 fusions have also been reported in up to 14% of cholangiocarcinoma and confer sensitivity to select FGFR inhibitors<sup>6,12,13</sup>.

**Potential relevance:** The pan-FGFR inhibitor, infigratinib, has been granted accelerated approval (2021) for previously treated, unresectable locally advanced or metastatic cholangiocarcinoma positive for FGFR2 fusion or other rearrangement<sup>14</sup>. The pan-FGFR inhibitor, erdafitinib<sup>15</sup>, received FDA approval (2019) for the treatment of locally advanced or metastatic urothelial cancer that is positive for FGFR2 fusions including, FGFR2-BICC1 and FGFR2-CASP7, FGFR3 fusions, or FGFR3 mutation. Additionally, the Pan-FGFR inhibitor, futibatinib<sup>16</sup>, has been granted Breakthrough Therapy Designation (2021) for FGFR2 rearrangement or fusion-positive locally advanced or metastatic cholangiocarcinoma. The FGFR kinase inhibitor, pemigatinib<sup>17</sup>, received FDA approval (2020), for previously treated, advanced or unresectable cholangiocarcinoma harboring FGFR2 fusions or other FGFR2 rearrangements. The FDA also granted fast-track designation (2018) to Debio 1347<sup>18</sup> for solid tumors harboring FGFR1, FGFR2, or FGFR3 aberrations. Additional FGFR inhibitors are under clinical evaluation for FGFR2 aberrations. In a phase II study of patients with FGFR2 fusion-positive intrahepatic cholangiocarcinoma, the pan-kinase inhibitor derazantinib, demonstrated an overall response rate (ORR) of 20.7% with progression-free survival (PFS) of 5.7 months<sup>19</sup>. Likewise, results of a phase II trial testing the pan-FGFR inhibitor, infigratinib (BGJ398) demonstrated an ORR of 14.8% (18.8% FGFR2 fusions only), disease control rate (DCR) of 75.4% (83.3% FGFR2 fusions only), and a median PFS of 5.8 months<sup>20</sup>.

## Clinical Trials in Taiwan region:

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2021-08-18. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### FGFR2 amplification

#### Debio 1347

**Cancer type:** Solid Tumor

**Variant class:** FGFR2 aberration

**Supporting Statement:**

The FDA has granted Fast Track Designation to the FGFR 1-3 inhibitor, debio 1347, for FGFR1/2/3 alterations in unresectable or metastatic solid tumors.

**Reference:**

<https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. Babina et al. Advances and challenges in targeting FGFR signalling in cancer. *Nat. Rev. Cancer*. 2017 May;17(5):318-332. PMID: 28303906
2. Ahmad et al. Mechanisms of FGFR-mediated carcinogenesis. *Biochim. Biophys. Acta*. 2012 Apr;1823(4):850-60. PMID: 22273505
3. Sarabipour et al. Mechanism of FGF receptor dimerization and activation. *Nat Commun*. 2016 Jan 4;7:10262. doi: 10.1038/ncomms10262. PMID: 26725515
4. Helsten et al. The FGFR Landscape in Cancer: Analysis of 4,853 Tumors by Next-Generation Sequencing. *Clin. Cancer Res*. 2016 Jan 1;22(1):259-67. PMID: 26373574
5. Touat et al. Targeting FGFR Signaling in Cancer. *Clin. Cancer Res*. 2015 Jun 15;21(12):2684-94. PMID: 26078430
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
8. Byron et al. The N550K/H mutations in FGFR2 confer differential resistance to PD173074, dovitininib, and ponatinib ATP-competitive inhibitors . *Neoplasia*. 2013 Aug;15(8):975-88. PMID: 23908597
9. Chae et al. Inhibition of the fibroblast growth factor receptor (FGFR) pathway: the current landscape and barriers to clinical application. *Oncotarget*. 2017 Feb 28;8(9):16052-16074. PMID: 28030802
10. Ahn et al. FGFR2 in gastric cancer: protein overexpression predicts gene amplification and high H-index predicts poor survival. *Mod. Pathol*. 2016 Sep;29(9):1095-103. PMID: 27230412
11. Jung et al. Fibroblast growth factor receptor 2 gene amplification status and its clinicopathologic significance in gastric carcinoma. *Hum. Pathol*. 2012 Oct;43(10):1559-66. PMID: 22440694
12. Borad et al. Fibroblast growth factor receptor 2 fusions as a target for treating cholangiocarcinoma. *Curr. Opin. Gastroenterol*. 2015 May;31(3):264-8. PMID: 25763789
13. Ghedini et al. Future applications of FGF/FGFR inhibitors in cancer. *Expert Rev Anticancer Ther*. 2018 Sep;18(9):861-872. PMID: 29936878
14. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/214622s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214622s000lbl.pdf)
15. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/212018s001lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/212018s001lbl.pdf)
16. [https://www.taihooncology.com/us/news/2021-04-01\\_toi\\_tpc\\_futibatinib\\_btd/](https://www.taihooncology.com/us/news/2021-04-01_toi_tpc_futibatinib_btd/)
17. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/213736s001lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/213736s001lbl.pdf)
18. <https://www.debiopharm.com/drug-development/press-releases/fda-grants-fast-track-designation-to-debiopharm-internationals-debio-1347-for-the-treatment-of-patients-with-unresectable-or-metastatic-tumors-with-a-specific-fgfr-gene-alteration/>
19. Mazzaferro et al. Derazantinib (ARQ 087) in advanced or inoperable FGFR2 gene fusion-positive intrahepatic cholangiocarcinoma. *Br. J. Cancer*. 2019 Jan;120(2):165-171. PMID: 30420614
20. Javle et al. Phase II Study of BGJ398 in Patients With FGFR-Altered Advanced Cholangiocarcinoma. *J. Clin. Oncol*. 2018 Jan 20;36(3):276-282. PMID: 29182496