

# ACT Onco<sup>®</sup> + Report

PATIENT		
Name: 楊麗珠		Patient ID: 10739727
Date of Birth: Jun 29, 1955		Gender: Female
Diagnosis: Metastatic lung adenocarcinoma		
ORDERING PHYSICIAN		
Name: 周德盈醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11170966A		Type: FFPE tissue
Collection site: Liver		
Date received: Aug 12, 2022	Lab ID: AA-22-04727	D/ID: NA

## ABOUT ACT Onco<sup>®</sup>+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) ( $\leq 15$  nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

## SUMMARY FOR ACTIONABLE VARIANTS

### VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
EGFR L858R	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	-	-

### VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
CCND1 Amplification	Abemaciclib, Palbociclib, Ribociclib	-
EGFR Amplification	Afatinib, Cetuximab, Erlotinib, Gefitinib, Necitumumab, Osimertinib, Panitumumab	-
IL6 Amplification	-	Erlotinib, Gefitinib, Trastuzumab
LYN Amplification	Dasatinib	-

#### Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

# ACTOnco® + Report

## TESTING RESULTS

### VARIANT(S) WITH CLINICAL RELEVANCE

#### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
EGFR	L858R	30.0%
TP53	Q104*	67.4%

#### - Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr13	BRCA2	Heterozygous deletion	1
Chr17	FLCN	Heterozygous deletion	1
Chr3	MLH1	Heterozygous deletion	1
Chr9	TSC1	Heterozygous deletion	1
Chr11	CCND1, VEGFB	Amplification	6
Chr7	CARD11, EGFR, IL6	Amplification	6
Chr8	LYN	Amplification	6

#### - Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

#### - Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	2.6 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

#### Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 61% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at  $\geq 7.5$  mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is  $< 30\%$ .

# ACT Onco<sup>®</sup> + Report

## THERAPEUTIC IMPLICATIONS

### TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
<b>Level 1</b>		
<b>EGFR</b> L858R	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	<b>sensitive</b>
<b>Level 3B</b>		
<b>CCND1</b> Amplification	Abemaciclib, Palbociclib, Ribociclib	<b>sensitive</b>
<b>EGFR</b> Amplification	Afatinib, Cetuximab, Erlotinib, Gefitinib, Necitumumab, Osimertinib, Panitumumab	<b>sensitive</b>
<b>Level 4</b>		
<b>LYN</b> Amplification	Dasatinib	<b>sensitive</b>
<b>IL6</b> Amplification	Erlotinib, Gefitinib, Trastuzumab	<b>resistant</b>

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
<b>1</b>	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
<b>2</b>	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
<b>3A</b>	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
<b>3B</b>	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
<b>4</b>	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

# ACT Onco<sup>®</sup> + Report

## IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

### - Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
EGFR aberration	Likely associated with WORSE response to ICIs

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

## CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

## HORMONAL THERAPIES

Genomic Alterations	Therapies	Effect	Level of Evidence	Cancer Type
CCND1 Amplification	Anastrozole	Less sensitive	Clinical	Breast cancer
	Tamoxifen	Less sensitive	Clinical	Breast cancer

## OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

### Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

# ACT Onco<sup>®</sup> + Report

## VARIANT INTERPRETATION

### EGFR L858R, Amplification

#### Biological Impact

The EGFR gene encodes for the Epidermal Growth Factor Receptor, a receptor tyrosine kinase which binds to its ligands, including Epidermal Growth Factor (EGF) and Transforming Growth Factor- $\alpha$  (TGF- $\alpha$ ), activates downstream signaling pathways, including the canonical oncogenic MAPK and PI3K/AKT/mTOR signaling cascades<sup>[1]</sup>. Increased EGFR activity by mutations and/or amplification of the EGFR gene has been described in a wide range of cancers, such as lung, brain, colorectal and head and neck cancer<sup>[2]</sup>. Mutations in the kinase domain of EGFR are commonly observed in non-small cell lung cancer (NSCLC), resulting in a constitutively activated form of the receptor<sup>[3]</sup>. On the other hand, in the brain and colorectal cancers, the most prevalent EGFR alteration is copy number amplification that results in receptor overexpression<sup>[4]</sup>.

EGFR L858R is a missense mutation at position 858, located in exon 21, which encodes part of the kinase domain, from a leucine to an arginine residue<sup>[5]</sup>. The two most common EGFR alterations, L858R mutation and exon 19 deletions can result in constitutive activation of signal transduction pathways, leading to cell proliferation or anti-apoptosis without ligand binding<sup>[6]</sup>.

#### Therapeutic and prognostic relevance

There is accumulated clinical evidence suggested that patients with MDM2/MDM4 amplification or EGFR aberrations exhibited poor clinical outcome and demonstrated a significantly increased rate of tumor growth (hyper-progression) after receiving immune checkpoint (PD-1/PD-L1) inhibitors therapies<sup>[7]</sup> (Annals of Oncology (2017) 28 (suppl\_5): v403-v427. 10.1093/annonc/mdx376).

The first- and second-generation EGFR tyrosine kinase inhibitors (EGFR-TKIs), dacomitinib, erlotinib, gefitinib and afatinib have been approved by the U.S. Food and Drug Administration (FDA) as the first-line treatment in non-small cell lung cancer (NSCLC) patients whose tumor carries EGFR exon 19 deletion or L858R mutation<sup>[8][9][10]</sup>, as detected by a U.S. FDA-approved test. A Phase III clinical trial (NCT01774721) show that dacomitinib significantly improved progression-free survival over gefitinib in first-line treatment of patients with EGFR-mutation-positive NSCLC<sup>[8]</sup>. Another Phase III clinical trial (NCT00949650) demonstrated that median progression-free survival (PFS) among lung cancer patients with exon 19 deletion or L858R EGFR mutation (n=308) was 13.6 months for afatinib and 6.9 months for chemotherapy. The EGFR T790M mutation has been demonstrated to confer resistance to TKIs (dacomitinib, gefitinib, erlotinib, and afatinib) in preclinical and clinical studies<sup>[11][12][13][14]</sup>.

Osimertinib, a third-generation irreversible EGFR-TKI that selectively inhibits both EGFR-TKI-sensitizing and EGFR T790M resistance mutations, has been approved by the U.S. FDA for NSCLC patient harboring T790M mutation-positive tumor<sup>[15][16][17]</sup>. Results from a double-blind, Phase 3 trial further showed that osimertinib significantly demonstrated longer PFS than standard EGFR-TKIs (18.9 months vs. 10.2 months) in previously untreated EGFR mutation-positive (exon 19 deletion or L858R) advanced NSCLC<sup>[18]</sup>.

Increased EGFR copy number is associated with tumor response to panitumumab, an EGFR-targeted antibody, in colorectal cancer patients, based on data from a phase III study<sup>[19]</sup>. A recent Phase II trial of cetuximab (another approved anti-EGFR antibody) oxaliplatin/leucovorin/5-fluorouracil therapy in first-line setting also demonstrated an association between higher EGFR copy number and better overall survival in gastric cancer patients<sup>[20]</sup>. The addition of cetuximab to chemotherapy reduced the risk of death by 44% for advanced squamous non-small cell lung cancer (NSCLC) patients with EGFR-amplified tumor, according to clinical trial findings presented at the 2015 World Conference on Lung Cancer. Preclinical data of gastric cancer (GC)-derived xenograft also showed that EGFR amplification or overexpression is associated with response to cetuximab<sup>[21]</sup>. Besides, a phase III study of necitumumab showed squamous cell lung cancer patients with EGFR amplification had improved overall survival (14.8 versus 7.6 months, p = 0.033) (NCT00981058)<sup>[22]</sup>.

# ACT Onco<sup>®</sup> + Report

Increased EGFR copy number has been shown to be associated with better response and survival in gefitinib or erlotinib treatment for NSCLC<sup>[23][24][25][26][27][28]</sup>, esophageal cancer<sup>[29]</sup>, and mucinous urethral adenocarcinoma<sup>[30]</sup>. Concurrent amplification of EGFR and ERBB2 is associated with response to afatinib in patients with trastuzumab-refractory esophagogastric cancer<sup>[31]</sup>. However, dacomitinib has been reported with a limited single-agent activity in recurrent glioblastoma with EGFR amplification in a phase II trial<sup>[32]</sup>. EGFR amplification has been determined as an inclusion criterion for the trials evaluating erlotinib, afatinib, and osimertinib efficacy in PDAC with co-expressing EGFR and c-Met (NCT03213626), glioblastoma (NCT03732352), urothelial tract carcinoma (NCT02780687), and brain cancer (NCT02423525).

## TP53 Q104\*

### Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis<sup>[33]</sup>. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation<sup>[34]</sup>.

Q104\* mutation results in a premature truncation of the p53 protein at amino acid 104 (UniProtKB). This mutation is predicted to lead to a loss of p53 function, despite not having characterized in the literature.

### Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)<sup>[35]</sup>.

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib<sup>[36]</sup>. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat<sup>[37]</sup>.

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53<sup>[38][39][40]</sup>. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)<sup>[41]</sup>. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy<sup>[42][43]</sup>. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53<sup>[44]</sup>.

## BRCA2 Heterozygous deletion

### Biological Impact

The BRCA2 gene encodes a tumor suppressor involved in the homologous recombination pathway for double-strand DNA repair<sup>[45]</sup>. BRCA2 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions<sup>[46]</sup>. BRCA2 germline mutations confer an increased lifetime risk of developing breast, ovarian, prostate and pancreatic cancer, limited reports of related gastric cancer, and Fanconi anemia subtype D1-associated risk of brain cancer, medulloblastoma, pharyngeal cancer, chronic lymphocytic leukemia and acute myeloid leukemia<sup>[47]</sup>. Somatic mutations in BRCA2 are highest in colorectal, non-small cell lung cancer (NSCLC), and ovarian cancers<sup>[48]</sup>.



# ACT Onco<sup>®</sup> + Report

## Therapeutic and prognostic relevance

The U.S. FDA has approved olaparib in advanced ovarian cancer under several settings including (1) first-line maintenance treatment for patients with deleterious or suspected deleterious germline or somatic BRCA mutation who are in complete or partial response to first-line platinum-based chemotherapy<sup>[49]</sup>; (2) in combination with bevacizumab as first-line maintenance treatment for patients with homologous recombination deficiency (HRD)-positive status<sup>[50]</sup>; (3) maintenance treatment for patients with germline BRCA-mutated recurrent ovarian cancer who are in complete or partial response to platinum-based chemotherapy<sup>[51][52]</sup>; (4) treatment for patients with germline BRCA-mutated advanced ovarian cancer who have been treated with three or more prior lines of chemotherapy<sup>[53]</sup>. In addition, olaparib has also been approved in patients with deleterious or suspected deleterious germline BRCA-mutated, HER2-negative metastatic breast cancer who have been treated with chemotherapy in either neoadjuvant, adjuvant, or metastatic setting<sup>[54]</sup> and germline BRCA-mutated metastatic pancreatic cancer<sup>[55]</sup>. Of note, in May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate<sup>[56]</sup>.

Rucaparib has been approved for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy and patients with BRCA-mutated epithelial ovarian, fallopian tube, or primary peritoneal cancer, who have been treated with two or more chemotherapies<sup>[57][58]</sup>. In May 2020, the U.S. FDA also approved rucaparib to treat adult patients with a deleterious BRCA mutation-associated metastatic castration-resistant prostate cancer (mCRPC) who have been treated with androgen receptor-directed therapy and a taxane-based chemotherapy (TRITON2, NCT02952534).

The U.S. FDA also approved niraparib for the maintenance treatment of patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in response to platinum-based chemotherapy and patients who have been treated with three or more prior lines of chemotherapy and associated with HRD positive status<sup>[59][60][61]</sup>. In addition, talazoparib for patients with deleterious or suspected deleterious germline BRCA-mutated, HER2 negative locally advanced or metastatic breast cancer<sup>[62]</sup>.

## CARD11 Amplification

### Biological Impact

CARD11 (caspase recruitment domain 11) gene encodes a cytoplasmic scaffold protein of the CARD11/BCL10/MALT1 (CBM) complex which plays essential roles in regulating apoptosis and NF-κB activation in response to upstream stimuli<sup>[63][64]</sup>. CARD11 gain-of-function mutations are frequently detected in human diffuse large B-cell lymphoma (DLBCL)<sup>[65]</sup> and cutaneous squamous cell carcinoma<sup>[66]</sup>. Moreover, CARD11 gene amplification has been observed in a significant proportion of DLBCL<sup>[67]</sup>. Biochemical assays revealed that enforced expression of CARD11/BCL10/MALT1 is essential for transformation of B-cell and survival of DLBCL cell<sup>[68]</sup>.

## Therapeutic and prognostic relevance

Retrospective studies have shown that high CARD11 expression or CARD11 gene amplification was associated with poor survival in diffuse large B cell lymphoma (DLBCL)<sup>[69][70]</sup>.

## CCND1 Amplification

### Biological Impact

The cyclin D1 (CCND1) gene encodes a protein involved in the control of cell growth, proliferation, transcription, and DNA repair<sup>[70]</sup>. CCND1 forms a complex with CDK4 and CDK6, leading to G1-S cell-cycle transition by inhibiting the retinoblastoma (RB) protein<sup>[70]</sup>. Amplification or overexpression of CCND1 could be oncogenic and is associated with carcinogenesis of various cancer types<sup>[71]</sup>.

# ACT Onco<sup>®</sup> + Report

## Therapeutic and prognostic relevance

Several CDK4 inhibitors, including palbociclib (PD0332991), LEE011, and LY2835219 have entered clinical trials for tumors with CCND1 amplification<sup>[72][73]</sup>. In the Phase II study of palbociclib and letrozole in patients with ER-positive HER2-negative metastatic breast cancer, patient selection based on CCND1 amplification or p16 loss did not further improve patient outcome<sup>[74]</sup>. Preclinical studies also demonstrated conflicting results regarding the correlation between high-level CCND1 and palbociclib sensitivity<sup>[75][76][77]</sup>.

CCND1 amplification has been implicated in predicting poor clinical outcomes in postmenopausal breast cancer patients treated with either anastrozole or tamoxifen<sup>[78]</sup>. In lung cancer patients, the increased CCND1 copy number is associated with poorer overall survival<sup>[79]</sup>. A retrospective study showed that melanoma patients whose tumor harboring CCND1, cRAF or KRAS gene copy number gain had better treatment response with CPS (carboplatin, paclitaxel, and sorafenib)<sup>[80]</sup>. Of note, 3 of 4 patients treated with ribociclib for the longest duration had CCND1 amplification in a phase I trial<sup>[81]</sup>.

Amplification of CCND1 are frequent and contributes to dedifferentiation and cellular proliferative activity of intrahepatic cholangiocarcinoma (ICC), and also indicates a poor prognosis for ICC patients<sup>[82]</sup>. Of note, CCND1 amplification has been selected as an inclusion criterion for the trial examining CDK4/6 inhibitors in different types of malignant solid tumors (NCT02187783, NCT02896335, NCT03526250, NCT02693535, NCT01037790, NCT03454919, NCT03310879, and NCT03356223).

## FLCN Heterozygous deletion

### Biological Impact

The FLCN gene encodes the tumor suppressor, Folliculin, a GTPase activating protein (GAP) for RagC/D GTPase proteins involved in amino acid sensing and signaling to mTORC1<sup>[83]</sup>. FLCN has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions<sup>[84][85]</sup>. Inactivation of the FLCN gene by mutation or deletion results in the activation of the mTOR pathway and AKT signaling<sup>[86][87]</sup>. Germline mutation of the FLCN gene causes the Birt-Hogg-Dubé syndrome, a rare disorder that is characterized by benign hamartomatous skin lesions and an increased risk of pneumothorax and renal tumors<sup>[88]</sup>.

## Therapeutic and prognostic relevance

In a prospective Phase 2 study, four anaplastic thyroid cancer (ATC)/ poorly differentiated thyroid cancer (PDTC) patients who had PI3K/mTOR/AKT alterations, including TSC2, FLCN or NF1, showed impressive progression-free survival (PFS) of 15.2 months after receiving everolimus<sup>[89]</sup>. mTOR inhibition via rapamycin also demonstrated potential in inhibiting the growth of renal cells deficient in FLCN in the preclinical setting<sup>[90]</sup>.

## IL6 Amplification

### Biological Impact

The Interleukin 6 (IL6) gene encodes a key cytokine produced by various cell types including monocytes, macrophages, fibroblasts, keratinocytes, endothelial cells, B cells, T cells, and also several tumor cells during infection and inflammation. It is a pleiotropic cytokine that regulates the immune response, tissue regeneration, and promoting tumor growth and survival<sup>[91][92][93][94]</sup>. In cancers, IL6 is implicated in inflammation to malignant transformation by activating the NF-κB pathway, and the induction of an epithelial-mesenchymal transition (EMT)<sup>[95]</sup>.

## Therapeutic and prognostic relevance

Preclinical studies showed that IL6 is sufficient to modify the sensitivity of EGFR mutant non-small cell lung cancer (NSCLC) cell line to erlotinib treatment<sup>[96]</sup>. Inhibition of IL6 signaling by metformin can potentiate gefitinib-induced antitumor activity<sup>[97]</sup>.



# ACT Onco<sup>®</sup> + Report

A preclinical study showed that increased expression of the cytokines IL6 was associated with trastuzumab resistance in HER2+ breast cancer<sup>[98]</sup>.

Amplification of IL6 was associated with shortened survival in glioblastoma patients<sup>[99]</sup>.

## LYN Amplification

### Biological Impact

The LYN Proto-Oncogene, Src Family Tyrosine Kinase (LYN) gene encodes a non-receptor tyrosine protein kinase of the Src family (SFK)<sup>[100]</sup>. LYN plays an important role in the regulation of immune responses, hematopoiesis, signal transduction of growth factors and cytokines and is activated in the cellular response to DNA damage and genotoxic agents<sup>[101][102][103][104]</sup>. LYN has been described to promote tumor growth, invasion, epithelial to mesenchymal transition (EMT) and ERK signaling in different cancer types<sup>[105][106][107]</sup>. Amplification of LYN has been identified in prostate cancer, breast cancer and ovarian cancer (cBioPortal).

### Therapeutic and prognostic relevance

LYN could be pharmacologically targeted with Src-kinase inhibitors. Results of preclinical studies showed that dasatinib, a dual-specificity tyrosine kinase inhibitor of ABL and the Src-family tyrosine kinases, exerted antitumor activity in LYN-expressing breast cancer cells<sup>[105]</sup>.

## MLH1 Heterozygous deletion

### Biological Impact

The MutL protein homolog 1 (MLH1) gene encodes a tumor suppressor that dimerizes with PMS2 protein to form a component of the DNA mismatch repair (MMR) system<sup>[108]</sup>. Deletion of one copy of the MLH1 gene resulted in haploinsufficiency in the correction of small insertions/deletions (indels), and could be a driving force in pancreatic and renal carcinogenesis<sup>[109]</sup>. Genetic alterations such as mutation, loss of heterozygosity or epigenetic silencing could lead to inactivation of MLH1 and are associated with a broad spectrum of cancers, including a subset of sporadic colon, gastric and endometrial cancers, as well as the hereditary non-polyposis colon cancer (HNPCC, also known as Lynch syndrome)<sup>[110][111][112]</sup>.

### Therapeutic and prognostic relevance

Currently, there are no FDA-approved medications specifically targeting MLH1. A screening test for microsatellite instability (MSI) is commonly used to identify an MMR-deficient tumor in the clinic<sup>[113][114]</sup>. Pembrolizumab (KEYTRUDA), an inhibitor targeting programmed cell death 1 (PD-1), has been approved by the U.S. FDA for adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI-H) or mismatch repair-deficient cancer. Down-regulation of genes involved in the MMR pathway such as MLH1, MSH2 and, MSH6 in high-grade serous epithelial ovarian cancer cell lines rendered cells sensitive to PARP inhibitors<sup>[115]</sup>.

## TSC1 Heterozygous deletion

### Biological Impact

The tuberous sclerosis complex 1 (TSC1) gene encodes a tumor suppressor, hamartin, a key negative regulator of the mammalian target of rapamycin (mTOR) pathway<sup>[116][117]</sup>. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis<sup>[118][119][120]</sup>, while LOH in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)<sup>[121]</sup> and endometrial cancer<sup>[122]</sup>. Loss of single TSC1 allele (haploinsufficiency) may provide a growth advantage to bladder epithelial cells, contributing to bladder cancer development<sup>[123]</sup>. Both TSC1 and TSC2 mutations cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC), in which individuals develop a variety of benign but often progressive neoplasms<sup>[124]</sup>.

# ACT Onco<sup>®</sup> + Report

## Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple neoplasms, such as bladder tumors<sup>[125]</sup>, gastric, sarcoma, thyroid cancer, and HNSCC<sup>[126]</sup>. There were case reports demonstrated the efficacy of sirolimus in malignant uterine perivascular epithelioid cell tumors (PEComa) patients harboring mutations/deletions in TSC1 and TSC2 genes, and temsirolimus in PEComa patients with hyperactivated mTOR pathway. Genomic profiling analysis of GOG248, a Phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer showed that mutations in AKT1, TSC1 and TSC2 may predict clinical benefit from temsirolimus<sup>[127]</sup>. Recent studies indicate that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets<sup>[128]</sup>.

Everolimus has been approved by the U.S. FDA for Tuberous Sclerosis Complex (TSC)-associated renal angiomyolipoma and Tuberous Sclerosis Complex (TSC)-associated subependymal giant cell astrocytoma (SEGA). This approval is based on the results from EXIST-1, EXIST-2, and Study 2485 trials (NCT00789828, NCT00790400, and NCT00411619).

## VEGFB Amplification

### Biological Impact

The VEGF gene family is constituted of VEGF-A, VEGF-B, VEGF-C, VEGF-D and placenta growth factor (PlGF)<sup>[129]</sup>. Bindings of VEGFs with their receptors, such as Flt-1 (VEGFR-1), Flk-1/KDR (VEGFR-2) and VEGFR-3, lead to endothelial cell proliferation and new blood vessel growth and thus plays a key role in angiogenesis<sup>[130][131]</sup>. VEGFs and their receptors are expressed in the vast majority of human solid tumors, including those of the lung<sup>[132]</sup>, breast<sup>[133]</sup>, gastrointestinal tract<sup>[134]</sup>, kidney<sup>[135]</sup>, bladder<sup>[135]</sup>, ovary<sup>[136]</sup>, endometrium<sup>[136]</sup>, brain<sup>[137]</sup> and endocrine<sup>[138]</sup> system.

In contrast to VEGFA, VEGF-B plays a relatively less pronounced role in the vascular system.

## Therapeutic and prognostic relevance

Aflibercept (VEGF trap), a soluble decoy receptor that binds and inactivates members of the VEGF family, including VEGF-A, VEGF-B, and PlGF, has been shown to inhibit ascites, and causes dramatic vascular remodeling in an ovarian cancer model<sup>[139]</sup>. Moreover, Aflibercept plus paclitaxel significantly exerted antitumor activity and inhibited ascites while prolonging survival in human ovarian cancer model<sup>[140]</sup> and demonstrated promising antitumor activity in combination with docetaxel in patients with recurrent gynecologic malignancies, including ovarian cancer<sup>[141]</sup>.

High VEGFB expression is associated with significantly shortened survival in lung squamous cell carcinoma (LSCC) and melanoma<sup>[142]</sup>. In ovarian cancer, overexpression of intratumoral VEGF, found to correlate with poorer prognosis<sup>[143][144][145]</sup>.

# ACT Onco<sup>®</sup> + Report

## US FDA-APPROVED DRUG(S)

### Abemaciclib (VERZENIO)

Abemaciclib is a cyclin-dependent kinase 4/6 (CDK4/6) inhibitor. Abemaciclib is developed and marketed by Eli Lilly under the trade name VERZENIO.

#### - FDA Approval Summary of Abemaciclib (VERZENIO)

<b>monarchE</b> NCT03155997	<b>Breast cancer</b> (Approved on 2021/10/12)
	<b>HR-positive, HER2-negative</b> Abemaciclib + tamoxifen/aromatase inhibitor vs. Tamoxifen/aromatase inhibitor [IDFS at 36 months(%): 86.1 vs. 79.0]
<b>MONARCH 3</b> <sup>[146]</sup> NCT02246621	<b>Breast cancer</b> (Approved on 2018/02/26)
	<b>HR-positive, HER2-negative</b> Abemaciclib + anastrozole/letrozole vs. Placebo + anastrozole/letrozole [PFS(M): 28.2 vs. 14.8]
<b>MONARCH 2</b> <sup>[147]</sup> NCT02107703	<b>Breast cancer</b> (Approved on 2017/09/28)
	<b>HR-positive, HER2-negative</b> Abemaciclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 16.4 vs. 9.3]
<b>MONARCH 1</b> <sup>[148]</sup> NCT02102490	<b>Breast cancer</b> (Approved on 2017/09/28)
	<b>HR-positive, HER2-negative</b> Abemaciclib [ORR(%): 19.7 vs. 17.4]

### Afatinib (GILOTRIF)

Afatinib acts as an irreversible covalent inhibitor of the ErbB family of receptor tyrosine kinases, including epidermal growth factor receptor (EGFR) and erbB-2 (HER2). Afatinib is developed and marketed by Boehringer Ingelheim under the trade name GILOTRIF (United States) and GIOTRIF (Europe).

#### - FDA Approval Summary of Afatinib (GILOTRIF)

<b>LUX-Lung 8</b> <sup>[149]</sup> NCT01523587	<b>Non-small cell lung carcinoma</b> (Approved on 2016/04/15)
	<b>EGFR Del19/L858R</b> Afatinib vs. Erlotinib [PFS(M): 2.4 vs. 1.9]
<b>LUX-Lung 3</b> <sup>[150]</sup> NCT00949650	<b>Non-small cell lung carcinoma</b> (Approved on 2013/07/13)
	<b>EGFR Del19/L858R</b> Afatinib vs. Pemetrexed + cisplatin [PFS(M): 11.1 vs. 6.9]

### Cetuximab (ERBITUX)

Cetuximab is a recombinant, chimeric (human/mouse) monoclonal antibody that binds to the extracellular domain and inhibits epidermal growth factor receptor (EGFR). Cetuximab is developed by ImClone and marketed by Eli Lilly under the trade name ERBITUX.

#### - FDA Approval Summary of Cetuximab (ERBITUX)

<b>CRYSTAL</b> <sup>[151]</sup> NCT00154102	<b>Colorectal cancer</b> (Approved on 2012/07/06)
	<b>EGFR-expressing, K-Ras Wild-type</b> Cetuximab + 5-fluorouracil + folinic acid + irinotecan vs. 5-fluorouracil + folinic acid + irinotecan [PFS(M): 8.9 vs. 8.1]

# ACT Onco<sup>®</sup> + Report

EXTREME <sup>[152]</sup> NCT00122460	Head and neck cancer (Approved on 2011/11/07)
	-
	Cetuximab + cisplatin/carboplatin + 5-FU vs. Cisplatin/carboplatin + 5-FU [OS(M): 10.1 vs. 7.4]
<sup>[153]</sup> NCT00004227	Head and neck cancer (Approved on 2006/03/01)
	-
	Cetuximab + radiation vs. Radiation [OS(M): 49 vs. 29.3]
<sup>[154]</sup> NCT00063141	Colorectal cancer (Approved on 2004/02/12)
	EGFR-expressing
	Cetuximab + irinotecan vs. Irinotecan [DOR(M): 5.7 vs. 4.2]

## Dacomitinib (VIZIMPRO)

Dacomitinib is an oral kinase inhibitor that targets EGFR. Dacomitinib is developed and marketed by Pfizer under the trade name VIZIMPRO.

### - FDA Approval Summary of Dacomitinib (VIZIMPRO)

ARCHER 1050 <sup>[8]</sup> NCT01774721	Non-small cell lung carcinoma (Approved on 2018/09/27)
	EGFR Del 19/ L858R
	Dacomitinib vs. Gefitinib [PFS(M): 14.7 vs. 9.2]

## Dasatinib (SPRYCEL)

Dasatinib is an oral Bcr-Abl tyrosine kinase inhibitor (inhibits the "Philadelphia chromosome") and Src family tyrosine kinase inhibitor. Dasatinib is produced by Bristol-Myers Squibb and sold under the trade name SPRYCEL.

### - FDA Approval Summary of Dasatinib (SPRYCEL)

DASISION <sup>[155]</sup> NCT00481247	Chronic myeloid leukemia (Approved on 2010/10/28)
	-
	Dasatinib vs. Imatinib [ORR(%): 76.8 vs. 66.2]
<sup>[156]</sup> NCT00123474	Chronic myeloid leukemia (Approved on 2007/11/08)
	-
	Dasatinib [ORR(%): 63.0]
<sup>[157]</sup> NCT00123487	Acute lymphocytic leukemia (Approved on 2006/06/28)
	-
	Dasatinib [ORR(%): 38.0]

# ACT Onco<sup>®</sup> + Report

## Erlotinib (TARCEVA)

Erlotinib is a small molecule, reversible inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Erlotinib is developed by OSI Pharmaceuticals, Genentech and Roche, and marketed by Astellas Pharm Global Development under the trade name TARCEVA.

### - FDA Approval Summary of Erlotinib (TARCEVA)

<b>RELAY</b> NCT02411448	<b>Non-small cell lung carcinoma</b> (Approved on 2020/05/29)
	<b>EGFR exon 19 deletion or exon 21 (L858R)</b> Erlotinib + ramucirumab vs. Erlotinib + placebo [PFS(M): 19.4 vs. 12.4]
<b>EURTAC<sup>[158]</sup></b> NCT00446225	<b>Non-small cell lung carcinoma</b> (Approved on 2013/05/14)
	<b>Exon 19 Del/Exon 21 substitution (L858R)</b> Erlotinib vs. Cisplatin + gemcitabine or cisplatin + docetaxel or carboplatin + gemcitabine or carboplatin + docetaxel [PFS(M): 10.4 vs. 5.2]
<b>PA.3<sup>[159]</sup></b> NCT00026338	<b>Pancreatic cancer</b> (Approved on 2005/11/02)
	- Gemcitabine vs. Placebo [OS(M): 6.4 vs. 6]

## Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

### - FDA Approval Summary of Everolimus (AFINITOR)

<b>RADIANT-4<sup>[160]</sup></b> NCT01524783	<b>Lung or gastrointestinal neuroendocrine tumor</b> (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
<b>BOLERO-2<sup>[161]</sup></b> NCT00863655	<b>Breast cancer</b> (Approved on 2012/07/20)
	<b>ER+/HER2-</b> Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
<b>EXIST-2</b> NCT00790400	<b>Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma</b> (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
<b>RADIANT-3<sup>[162]</sup></b> NCT00510068	<b>Pancreatic neuroendocrine tumor</b> (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
<b>EXIST-1<sup>[163]</sup></b> NCT00789828	<b>Subependymal giant cell astrocytoma</b> (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
<b>RECORD-1<sup>[164]</sup></b> NCT00410124	<b>Renal cell carcinoma</b> (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]



# ACT Onco® + Report

## Gefitinib (IRESSA)

Gefitinib is a small molecule inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Gefitinib is developed and marketed by AstraZeneca under the trade name IRESSA.

### - FDA Approval Summary of Gefitinib (IRESSA)

<b>IFUM</b> <sup>[165]</sup> NCT01203917	<b>Non-small cell lung carcinoma</b> (Approved on 2015/07/13)
	<b>Exon 19 Del/Exon 21 substitution (L858R)</b>
	Gefitinib [ORR(%): 50.0]

## Necitumumab (PORTRAZZA)

Necitumumab is a recombinant human IgG1 monoclonal antibody against the human epidermal growth factor receptor (EGFR) and blocks the binding of EGFR to its ligands. Necitumumab is developed and marketed by Eli Lilly under the trade name PORTRAZZA.

### - FDA Approval Summary of Necitumumab (PORTRAZZA)

<b>SQUIRE</b> <sup>[166]</sup> NCT00981058	<b>Lung squamous cell carcinoma</b> (Approved on 2015/11/14)
	-
	Gemcitabine + cisplatin vs. Placebo [OS(M): 11.5 vs. 9.9]

## Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

### - FDA Approval Summary of Niraparib (ZEJULA)

<b>PRIMA</b> NCT02655016	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2020/04/29)
	-
	Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
<b>QUADRA</b> <sup>[61]</sup> NCT02354586	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2019/10/23)
	<b>HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)</b>
	Niraparib [ORR(%): 24.0, DOR(M): 8.3]
<b>NOVA</b> <sup>[60]</sup> NCT01847274	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2017/03/27)
	-
	Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

# ACT Onco<sup>®</sup> + Report

## Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

### - FDA Approval Summary of Olaparib (LYNPARZA)

<b>OlympiA</b> NCT02032823	<b>Her2-negative high-risk early breast cancer</b> (Approved on 2022/03/11)
	<b>gBRCA</b> Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M): ]
<b>PROfound</b> <sup>[56]</sup> NCT02987543	<b>Prostate cancer</b> (Approved on 2020/05/19)
	<b>ATMm, BRCA1m, BRCA2m, BARD1m, BRIP1m, CDK12m, CHEK1m, CHEK2m, FANCLm, PALB2m, RAD51Bm, RAD51Cm, RAD51Dm, RAD54Lm</b> Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
<b>PAOLA-1</b> <sup>[50]</sup> NCT02477644	<b>Ovarian cancer</b> (Approved on 2020/05/08)
	<b>HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)</b> Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
<b>POLO</b> <sup>[55]</sup> NCT02184195	<b>Pancreatic adenocarcinoma</b> (Approved on 2019/12/27)
	<b>Germline BRCA mutation (deleterious/suspected deleterious)</b> Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
<b>SOLO-1</b> <sup>[49]</sup> NCT01844986	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2018/12/19)
	<b>Germline or somatic BRCA-mutated (gBRCAm or sBRCAm)</b> Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
<b>OlympiAD</b> <sup>[54]</sup> NCT02000622	<b>Breast cancer</b> (Approved on 2018/02/06)
	<b>Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative</b> Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
<b>SOLO-2/ENGOT-Ov21</b> <sup>[167]</sup> NCT01874353	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2017/08/17)
	<b>gBRCA+</b> Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
<b>Study19</b> <sup>[168]</sup> NCT00753545	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2017/08/17)
	- Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]
<b>Study 42</b> <sup>[169]</sup> NCT01078662	<b>Ovarian cancer</b> (Approved on 2014/12/19)
	<b>Germline BRCA mutation (deleterious/suspected deleterious)</b> Olaparib [ORR(%): 34.0, DOR(M): 7.9]

## Osimertinib (TAGRISSO)

Osimertinib is a third-generation tyrosine kinase inhibitor (TKI) for patients with tumors harboring EGFR T790M mutation. Osimertinib is developed and marketed by AstraZeneca under the trade name TAGRISSO.

### - FDA Approval Summary of Osimertinib (TAGRISSO)

<b>ADAURA</b> NCT02511106	<b>Non-small cell lung carcinoma</b> (Approved on 2020/12/18)
	<b>EGFR exon 19 deletions or exon 21 L858R mutations</b> Osimertinib vs. Placebo + adjuvant chemotherapy [DFS(M): NR vs. 19.6]

# ACT Onco<sup>®</sup> + Report

<b>FLAURA<sup>[18]</sup></b> NCT02296125	<b>Non-small cell lung carcinoma</b> (Approved on 2018/04/18)
	<b>EGFR Del19/L858R</b>
	Osimertinib vs. Gefitinib or erlotinib [PFS(M): 18.9 vs. 10.2]
<b>AURA3<sup>[170]</sup></b> NCT02151981	<b>Non-small cell lung carcinoma</b> (Approved on 2017/03/30)
	<b>EGFR T790M+</b>
	Osimertinib vs. Chemotherapy [PFS(M): 10.1 vs. 4.4]
<b>AURA<sup>[17]</sup></b> NCT01802632	<b>Non-small cell lung carcinoma</b> (Approved on 2015/11/13)
	<b>EGFR T790M+</b>
	Osimertinib [ORR(%): 59.0]

## Palbociclib (IBRANCE)

Palbociclib is an oral, cyclin-dependent kinase (CDK) inhibitor specifically targeting CDK4 and CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Palbociclib is developed and marketed by Pfizer under the trade name IBRANCE.

### - FDA Approval Summary of Palbociclib (IBRANCE)

<b>PALOMA-2<sup>[171]</sup></b> NCT01740427	<b>Breast cancer</b> (Approved on 2017/03/31)
	<b>ER+, HER2-</b>
	Palbociclib + letrozole vs. Placebo + letrozole [PFS(M): 24.8 vs. 14.5]
<b>PALOMA-3<sup>[172]</sup></b> NCT01942135	<b>Breast cancer</b> (Approved on 2016/02/19)
	<b>ER+, HER2-</b>
	Palbociclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 9.5 vs. 4.6]

## Panitumumab (VECTIBIX)

Panitumumab is a fully human monoclonal antibody against the human epidermal growth factor receptor (EGFR) and binds to the extracellular domain to prevent its activation. Panitumumab is developed by Abgenix and Amgen, and marketed by the latter under the trade name VECTIBIX.

### - FDA Approval Summary of Panitumumab (VECTIBIX)

<b>Study 20050203<sup>[173]</sup></b> NCT01412957	<b>Colorectal cancer</b> (Approved on 2017/06/29)
	-
	Panitumumab + BSC vs. BSC [OS(M): 10 vs. 6.9]
<b>PRIME<sup>[174]</sup></b> NCT00364013	<b>Colorectal cancer</b> (Approved on 2014/05/23)
	-
	Panitumumab + FOLFOX vs. FOLFOX [PFS(M): 9.6 vs. 8]
<b>ASPECCT<sup>[175]</sup></b> NCT01001377	<b>Colorectal cancer</b> (Approved on 2014/05/23)
	-
	Panitumumab vs. Cetuximab [OS(M): 10.4 vs. 10]
<b>Study 20080763<sup>[176]</sup></b> NCT00113763	<b>Colorectal cancer</b> (Approved on 2006/09/27)
	-
	Panitumumab + BSC vs. BSC [PFS(M): 3.2 vs. 2]

# ACT Onco<sup>®</sup> + Report

## Ribociclib (KISQALI)

Ribociclib is a cyclin-dependent kinase (CDK) inhibitor specifically targeting cyclin D1/CDK4 and cyclin D3/CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Ribociclib is developed by Novartis and Astex Pharmaceuticals and marketed by Novartis under the trade name KISQALI.

### - FDA Approval Summary of Ribociclib (KISQALI)

<b>MONALEESA-2</b> <sup>[177]</sup> NCT01958021	<b>Breast cancer</b> (Approved on 2017/03/13)
	<b>HR+, HER2-</b>
	Ribociclib vs. Letrozole [PFS(M): NR vs. 14.7]

## Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

### - FDA Approval Summary of Rucaparib (RUBRACA)

<b>TRITON2</b> NCT02952534	<b>Prostate cancer</b> (Approved on 2020/05/15)
	<b>gBRCA+, sBRCA</b>
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
<b>ARIEL3</b> <sup>[57]</sup> NCT01968213	<b>Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma</b> (Approved on 2018/04/06)
	<b>All HRD tBRCA</b>
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]
<b>ARIEL2</b> <sup>[178]</sup> NCT01482715, NCT01891344	<b>Ovarian cancer</b> (Approved on 2016/12/19)
	<b>Germline and/or somatic BRCA mutation</b>
	Rucaparib [ORR(%): 54.0]

## Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

### - FDA Approval Summary of Talazoparib (TALZENNA)

<b>EMBRACA</b> <sup>[62]</sup> NCT01945775	<b>Breast cancer</b> (Approved on 2018/10/16)
	<b>Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative</b>
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

# ACT Onco<sup>®</sup> + Report

## Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

## - FDA Approval Summary of Temsirolimus (TORISEL)

[179] NCT00065468	<b>Renal cell carcinoma</b> (Approved on 2007/05/30) - Temsirolimus vs. IFN- $\alpha$ [OS(M): 10.9 vs. 7.3]
----------------------	---

D=day; W=week; M=month



# ACT Onco<sup>®</sup> + Report

## ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

# ACT Onco<sup>®</sup> + Report


## SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

### - Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
EGFR	L858R	21	c.2573T>G	NM_005228	COSM6224	30.0%	3971
TP53	Q104*	4	c.310C>T	NM_000546	COSM10886	67.4%	727

### - Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



# ACT Onco<sup>®</sup> + Report

## OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ABL2	P1044S	12	c.3130C>T	NM_007314	-	17.7%	1395
ADAMTS9	A117T	2	c.349G>A	NM_182920	-	21.2%	1025
CARD11	A1033V	23	c.3098C>T	NM_032415	COSM1088959	7.2%	1746
CARD11	A687V	16	c.2060C>T	NM_032415	-	61.0%	751
CDK12	V605A	2	c.1814T>C	NM_016507	COSM9248152	20.8%	2566
CTNNA1	V702A	15	c.2105T>C	NM_001903	-	6.9%	3293
EPHA5	V537A	7	c.1610T>C	NM_001281765	COSM6929960	37.4%	1093
ERCC5	A185V	6	c.554C>T	NM_000123	-	72.1%	305
FAT1	V2654I	10	c.7960G>A	NM_005245	COSM260847	34.9%	1551
FGFR4	Splice region	8	c.1056A>T	NM_213647	-	44.7%	445
FLT1	A1319E	30	c.3956C>A	NM_002019	-	61.5%	728
HIST1H1C	A24V	1	c.71C>T	NM_005319	-	46.2%	1185
MSH2	I169V	3	c.505A>G	NM_000251	COSM1684714	82.5%	766
NFE2L2	R449H	5	c.1346G>A	NM_006164	COSM1009927	58.2%	622
NFKB1	R534H	15	c.1601G>A	NM_003998	-	37.0%	3419
NPM1	E245Q	9	c.733G>C	NM_002520	-	49.7%	963
PIK3C2B	A1441V	30	c.4322C>T	NM_002646	-	29.3%	1254
RUNX1T1	Splice region	10	c.1281C>T	NM_175634	COSM2790695	29.6%	2056
USH2A	T3635N	55	c.10904C>A	NM_206933	-	81.1%	848


### Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

# ACT Onco<sup>®</sup> + Report

## TEST DETAILS

### SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Jul 2022
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11170966A
- Collection site: Liver
- Examined by: Dr. Chien-Ta Chiang
  1. The percentage of viable tumor cells in total cells in the whole slide (%): 10%
  2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 60%
  3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 5%
  4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
  5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

## RUN QC

- Panel: ACTOnco<sup>®</sup>+

### DNA test

- Mean Depth: 1033x
- Target Base Coverage at 100x: 95%

### RNA test

- Average unique RNA Start Sites per control GSP2: 124

## LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.

# ACT Onco<sup>®</sup> + Report

2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

## NEXT-GENERATION SEQUENCING (NGS) METHODS

### DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage  $\geq 20$ , allele frequency  $\geq 5\%$  and actionable variants with allele frequency  $\geq 2\%$  were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at  $100\times \geq 85\%$  with a mean coverage  $\geq 500\times$ .

Variants reported in Genome Aggregation database with  $> 1\%$  minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation  $\geq 0.3$  were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco<sup>®</sup> to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to  $\geq 7.5$  mutations per megabase (Muts/Mb); TMB-Low corresponds to  $< 7.5$  Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is  $< 30\%$ .

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in  $> 400$  genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

### RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be  $\geq 10$ .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2  $\geq 3$ ; (2) Number of supporting



# ACT Onco<sup>®</sup> + Report

reads spanning the fusion junction  $\geq 5$ ; (3) Percentage of supporting reads spanning the fusion junction  $\geq 10\%$ ; (4) Fusions annotated in Quiver Gene Fusion Database.

## DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

### Variant Analysis:

醫檢師黃靖婷 博士  
Ching-Ting Huang Ph.D.  
檢字第 016511 號

CT Huang

### Sign Off

解剖病理專科醫師王業翰  
Yeh-Han Wang M.D.  
病解字第 000545 號

Yeh

# ACT Onco<sup>®</sup> + Report

## GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLCO1B1*
SLCO1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

\*Analysis of copy number alterations NOT available.

## FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


# ACT Onco<sup>®</sup> + Report

## APPENDIX

### POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Gene	Therapies	Possible effect
<i>FLCN</i>	Everolimus, Temsirolimus	<b>sensitive</b>
<i>TSC1</i>	Everolimus, Temsirolimus	<b>sensitive</b>
<i>BRCA2</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	<b>sensitive</b>
<i>MLH1</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	<b>sensitive</b>

### SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Olaparib, Niraparib, Rucaparib, Talazoparib



# ACT Onco<sup>®</sup> + Report

## Receptor Tyrosine Kinase/Growth Factor Signalling



1: Gefitinib, Afatinib, Erlotinib, Osimertinib, Dacomitinib; 2: Cetuximab, Panitumumab, Necitumumab; 3: Dasatinib; 4: Dasatinib; 5: Afatinib



# ACT Onco<sup>®</sup> + Report

## DISCLAIMER

### 法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

### 醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

### 基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

### 證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

### 責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

# ACT Onco<sup>®</sup> + Report

## REFERENCE

1. PMID: 18045542; 2007, Cell;131(5):1018  
SnapShot: EGFR signaling pathway.
2. PMID: 10880430; 2000, EMBO J;19(13):3159-67  
The ErbB signaling network: receptor heterodimerization in development and cancer.
3. PMID: 15329413; 2004, Proc Natl Acad Sci U S A;101(36):13306-11  
EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib.
4. PMID: 11426640; 2000, Oncogene;19(56):6550-65  
The EGF receptor family as targets for cancer therapy.
5. PMID: 17318210; 2007, Nat Rev Cancer;7(3):169-81  
Epidermal growth factor receptor mutations in lung cancer.
6. PMID: 22263017; 2010, J Thorac Dis;2(1):48-51  
Epidermal growth factor receptor (EGFR) in lung cancer: an overview and update.
7. PMID: 28351930; 2017, Clin Cancer Res;23(15):4242-4250  
Hyperprogressors after Immunotherapy: Analysis of Genomic Alterations Associated with Accelerated Growth Rate.
8. PMID: 28958502; 2017, Lancet Oncol;18(11):1454-1466  
Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial.
9. PMID: 24868098; 2014, Oncologist;19(7):774-9  
U.S. Food and Drug Administration approval summary: Erlotinib for the first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor exon 19 deletions or exon 21 (L858R) substitution mutations.
10. PMID: 23982599; 2013, Drugs;73(13):1503-15  
Afatinib: first global approval.
11. PMID: 27912760; 2016, J Biomed Sci;23(1):86  
Update on recent preclinical and clinical studies of T790M mutant-specific irreversible epidermal growth factor receptor tyrosine kinase inhibitors.
12. PMID: 15728811; 2005, N Engl J Med;352(8):786-92  
EGFR mutation and resistance of non-small-cell lung cancer to gefitinib.
13. PMID: 26862733; 2016, Oncotarget;7(11):12404-13  
The mechanism of acquired resistance to irreversible EGFR tyrosine kinase inhibitor-afatinib in lung adenocarcinoma patients.
14. PMID: 29410323; 2018, J Thorac Oncol;13(5):727-731  
EGFR T790M and C797S Mutations as Mechanisms of Acquired Resistance to Dacomitinib.
15. PMID: 24893891; 2014, Cancer Discov;4(9):1046-61  
AZD9291, an irreversible EGFR TKI, overcomes T790M-mediated resistance to EGFR inhibitors in lung cancer.
16. PMID: 27071706; 2016, J Hematol Oncol;9():34  
Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer.
17. PMID: 25923549; 2015, N Engl J Med;372(18):1689-99  
AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer.
18. PMID: 29151359; 2018, N Engl J Med;378(2):113-125  
Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer.

# ACT Onco<sup>®</sup> + Report

19. PMID: 17664472; 2007, J Clin Oncol;25(22):3238-45  
Epidermal growth factor receptor gene copy number and clinical outcome of metastatic colorectal cancer treated with panitumumab.
20. PMID: 22152101; 2011, BMC Cancer;11():509  
Biomarker analysis of cetuximab plus oxaliplatin/leucovorin/5-fluorouracil in first-line metastatic gastric and oesophago-gastric junction cancer: results from a phase II trial of the Arbeitsgemeinschaft Internistische Onkologie (AIO).
21. PMID: 24141978; 2013, Sci Rep;3():2992  
A subset of gastric cancers with EGFR amplification and overexpression respond to cetuximab therapy.
22. PMID: 29158193; 2018, J Thorac Oncol;13(2):228-236  
EGFR Gene Copy Number by FISH May Predict Outcome of Necitumumab in Squamous Lung Carcinomas: Analysis from the SQUIRE Study.
23. PMID: 19102716; 2009, Pharmacogenomics;10(1):59-68  
EGFR-targeted therapies in lung cancer: predictors of response and toxicity.
24. PMID: 15870435; 2005, J Natl Cancer Inst;97(9):643-55  
Epidermal growth factor receptor gene and protein and gefitinib sensitivity in non-small-cell lung cancer.
25. PMID: 15998906; 2005, J Clin Oncol;23(28):6838-45  
Increased epidermal growth factor receptor gene copy number detected by fluorescence in situ hybridization associates with increased sensitivity to gefitinib in patients with bronchioloalveolar carcinoma subtypes: a Southwest Oncology Group Study.
26. PMID: 16204011; 2005, J Clin Oncol;23(31):8081-92  
Epidermal growth factor receptor mutations and gene amplification in non-small-cell lung cancer: molecular analysis of the IDEAL/INTACT gefitinib trials.
27. PMID: 23557218; 2013, J Transl Med;11():90  
High EGFR copy number predicts benefits from tyrosine kinase inhibitor treatment for non-small cell lung cancer patients with wild-type EGFR.
28. PMID: 26141217; 2015, Lung Cancer;89(3):337-42  
Concurrence of EGFR amplification and sensitizing mutations indicate a better survival benefit from EGFR-TKI therapy in lung adenocarcinoma patients.
29. PMID: 28537764; 2017, J Clin Oncol;35(20):2279-2287  
Gefitinib and EGFR Gene Copy Number Aberrations in Esophageal Cancer.
30. PMID: 28057415; 2017, Clin Genitourin Cancer;15(4):e727-e734  
Comprehensive Genomic Analysis of Metastatic Mucinous Urethral Adenocarcinoma Guides Precision Oncology Treatment: Targetable EGFR Amplification Leading to Successful Treatment With Erlotinib.
31. PMID: 30463996; 2019, Cancer Discov;9(2):199-209  
EGFR and MET Amplifications Determine Response to HER2 Inhibition in ERBB2-Amplified Esophagogastric Cancer.
32. PMID: 28575464; 2017, Neuro Oncol;19(11):1522-1531  
Phase II trial of dacomitinib, a pan-human EGFR tyrosine kinase inhibitor, in recurrent glioblastoma patients with EGFR amplification.
33. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70  
Unravelling mechanisms of p53-mediated tumour suppression.
34. PMID: 21125671; 2011, J Pathol;223(2):137-46  
Haplo-insufficiency: a driving force in cancer.
35. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361  
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months.
36. PMID: 26646755; 2016, Ann Oncol;27(3):539-43  
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.

# ACT Onco<sup>®</sup> + Report

37. PMID: 25669829; 2015, Ann Oncol;26(5):1012-8  
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
38. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485  
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
39. PMID: 23670029; 2013, Oncotarget;4(5):705-14  
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
40. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14  
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
41. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52  
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
42. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21  
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
43. PMID: 10786679; 2000, Cancer Res;60(8):2155-62  
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
44. PMID: 25672981; 2015, Cancer Res;75(7):1187-90  
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
45. PMID: 11239455; 2001, Mol Cell;7(2):263-72  
BRCA2 is required for homology-directed repair of chromosomal breaks.
46. PMID: 17597348; 2007, Ann Surg Oncol;14(9):2510-8  
Heterogenic loss of the wild-type BRCA allele in human breast tumorigenesis.
47. PMID: 22193408; 2011, Nat Rev Cancer;12(1):68-78  
BRCA1 and BRCA2: different roles in a common pathway of genome protection.
48. PMID: 27283171; 2016, J Natl Compr Canc Netw;14(6):795-806  
The Relevance of Hereditary Cancer Risks to Precision Oncology: What Should Providers Consider When Conducting Tumor Genomic Profiling?
49. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505  
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
50. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428  
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
51. PMID: 28884698; 2017, Lancet Oncol;18(9):e510  
Correction to Lancet Oncol 2017; 18: 1274-84.
52. PMID: 22452356; 2012, N Engl J Med;366(15):1382-92  
Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer.
53. PMID: 26187614; 2015, Clin Cancer Res;21(19):4257-61  
FDA Approval Summary: Olaparib Monotherapy in Patients with Deleterious Germline BRCA-Mutated Advanced Ovarian Cancer Treated with Three or More Lines of Chemotherapy.
54. PMID: 28578601; 2017, N Engl J Med;377(6):523-533  
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
55. PMID: 31157963; 2019, N Engl J Med;381(4):317-327

# ACT Onco<sup>®</sup> + Report

Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.

56. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102  
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
57. PMID: 28916367; 2017, Lancet;390(10106):1949-1961  
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
58. PMID: 28882436; 2017, Gynecol Oncol;147(2):267-275  
Antitumor activity and safety of the PARP inhibitor rucaparib in patients with high-grade ovarian carcinoma and a germline or somatic BRCA1 or BRCA2 mutation: Integrated analysis of data from Study 10 and ARIEL2.
59. PMID: 31562799; 2019, N Engl J Med;381(25):2391-2402  
Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
60. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164  
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
61. PMID: 30948273; 2019, Lancet Oncol;20(5):636-648  
Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial.
62. PMID: 30110579; 2018, N Engl J Med;379(8):753-763  
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
63. PMID: 11278692; 2001, J Biol Chem;276(15):11877-82  
CARD11 and CARD14 are novel caspase recruitment domain (CARD)/membrane-associated guanylate kinase (MAGUK) family members that interact with BCL10 and activate NF-kappa B.
64. PMID: 26260210; 2015, Mol Immunol;68(2 Pt C):546-57  
TCR signaling to NF-kB and mTORC1: Expanding roles of the CARMA1 complex.
65. PMID: 18323416; 2008, Science;319(5870):1676-9  
Oncogenic CARD11 mutations in human diffuse large B cell lymphoma.
66. PMID: 26212909; 2015, Am J Pathol;185(9):2354-63  
Novel CARD11 Mutations in Human Cutaneous Squamous Cell Carcinoma Lead to Aberrant NF-kB Regulation.
67. PMID: 22397314; 2012, Leuk Lymphoma;53(10):1971-7  
Role of nuclear factor-kB regulators TNFAIP3 and CARD11 in Middle Eastern diffuse large B-cell lymphoma.
68. PMID: 26668357; 2015, Proc Natl Acad Sci U S A;112(52):E7230-8  
Lymphomagenic CARD11/BCL10/MALT1 signaling drives malignant B-cell proliferation via cooperative NF-kB and JNK activation.
69. PMID: 26876250; 2016, Zhonghua Xue Ye Xue Za Zhi;37(1):30-4  
[Expression and prognostic value of CARD11 in diffuse large B cell lymphoma].
70. PMID: 8114739; 1994, Mol Cell Biol;14(3):2077-86  
Identification of G1 kinase activity for cdk6, a novel cyclin D partner.
71. PMID: 12432268; 2002, Cancer Biol Ther;1(3):226-31  
Cycling to cancer with cyclin D1.
72. PMID: 24795392; 2014, Clin Cancer Res;20(13):3379-83  
Molecular pathways: CDK4 inhibitors for cancer therapy.
73. PMID: 27030077; 2016, Nat Rev Clin Oncol;13(7):417-30  
Treating cancer with selective CDK4/6 inhibitors.
74. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35

# ACT Onco<sup>®</sup> + Report

The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.

75. PMID: 27334220; 2016, Ann Surg Oncol;23(9):2745-52  
Targeting Cyclin-Dependent Kinases in Synovial Sarcoma: Palbociclib as a Potential Treatment for Synovial Sarcoma Patients.
76. PMID: 19874578; 2009, Breast Cancer Res;11(5):R77  
PD 0332991, a selective cyclin D kinase 4/6 inhibitor, preferentially inhibits proliferation of luminal estrogen receptor-positive human breast cancer cell lines in vitro.
77. PMID: 23898052; 2013, Anticancer Res;33(8):2997-3004  
PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity.
78. PMID: 22475046; 2012, Breast Cancer Res;14(2):R57  
Effects of cyclin D1 gene amplification and protein expression on time to recurrence in postmenopausal breast cancer patients treated with anastrozole or tamoxifen: a TransATAC study.
79. PMID: 17070615; 2007, Lung Cancer;55(1):1-14  
Cyclin D1 in non-small cell lung cancer: a key driver of malignant transformation.
80. PMID: 26307133; 2016, Clin Cancer Res;22(2):374-82  
Copy Number Changes Are Associated with Response to Treatment with Carboplatin, Paclitaxel, and Sorafenib in Melanoma.
81. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705  
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
82. PMID: 11495045; 2001, J Hepatol;35(1):74-9  
The role of overexpression and gene amplification of cyclin D1 in intrahepatic cholangiocarcinoma.
83. PMID: 24095279; 2013, Mol Cell;52(4):495-505  
The folliculin tumor suppressor is a GAP for the RagC/D GTPases that signal amino acid levels to mTORC1.
84. PMID: 26342594; 2016, Fam Cancer;15(1):127-32  
Birt-Hogg-Dubé syndrome: novel FLCN frameshift deletion in daughter and father with renal cell carcinomas.
85. PMID: 23223565; 2013, J Clin Pathol;66(3):178-86  
Birt-Hogg-Dubé syndrome: clinicopathological features of the lung.
86. PMID: 19850877; 2009, Proc Natl Acad Sci U S A;106(44):18722-7  
Homozygous loss of BHD causes early embryonic lethality and kidney tumor development with activation of mTORC1 and mTORC2.
87. PMID: 24908670; 2014, Hum Mol Genet;23(21):5706-19  
Folliculin (Flcn) inactivation leads to murine cardiac hypertrophy through mTORC1 deregulation.
88. PMID: 15956655; 2005, J Natl Cancer Inst;97(12):931-5  
High frequency of somatic frameshift BHD gene mutations in Birt-Hogg-Dubé-associated renal tumors.
89. PMID: 29301825; 2018, Clin Cancer Res;24(7):1546-1553  
Genomic Correlates of Response to Everolimus in Aggressive Radioiodine-refractory Thyroid Cancer: A Phase II Study.
90. PMID: 26418749; 2015, Oncotarget;6(32):32761-73  
Flcn-deficient renal cells are tumorigenic and sensitive to mTOR suppression.
91. PMID: 9237710; 1997, J Infect Dis;176(2):439-44  
Interleukin-6 gene-deficient mice show impaired defense against pneumococcal pneumonia.
92. PMID: 14581334; 2003, Clin Cancer Res;9(13):4653-65  
Targeted anti-interleukin-6 monoclonal antibody therapy for cancer: a review of the rationale and clinical evidence.



# ACT Onco<sup>®</sup> + Report

93. PMID: 16199153; 2005, Eur J Cancer;41(16):2502-12  
The role of IL-6 and STAT3 in inflammation and cancer.
94. PMID: 28778705; 2017, Autoimmun Rev;16(10):1079-1089  
Targeting interleukin-6 in autoimmune uveitis.
95. PMID: 21220315; 2011, Proc Natl Acad Sci U S A;108(4):1397-402  
Inducible formation of breast cancer stem cells and their dynamic equilibrium with non-stem cancer cells via IL6 secretion.
96. PMID: 20713723; 2010, Proc Natl Acad Sci U S A;107(35):15535-40  
TGF-beta IL-6 axis mediates selective and adaptive mechanisms of resistance to molecular targeted therapy in lung cancer.
97. PMID: 24644001; 2014, Clin Cancer Res;20(10):2714-26  
Metformin sensitizes EGFR-TKI-resistant human lung cancer cells in vitro and in vivo through inhibition of IL-6 signaling and EMT reversal.
98. PMID: 22819326; 2012, Mol Cell;47(4):570-84  
Activation of an IL6 inflammatory loop mediates trastuzumab resistance in HER2+ breast cancer by expanding the cancer stem cell population.
99. PMID: 17224923; 2007, Br J Cancer;96(3):474-6  
Interleukin-6 gene amplification and shortened survival in glioblastoma patients.
100. PMID: 9202419; 1997, Int J Biochem Cell Biol;29(3):397-400  
Lyn, a src-like tyrosine kinase.
101. PMID: 19290919; 2009, Immunol Rev;228(1):23-40  
Multiple roles of Lyn kinase in myeloid cell signaling and function.
102. PMID: 11825908; 2002, J Biol Chem;277(16):13732-8  
Phosphatidylinositol 3-kinase and Src family kinases are required for phosphorylation and membrane recruitment of Dok-1 in c-Kit signaling.
103. PMID: 11435302; 2001, Blood;98(2):343-50  
Lyn is required for normal stem cell factor-induced proliferation and chemotaxis of primary hematopoietic cells.
104. PMID: 10574731; 1999, J Exp Biol;202 Pt 24():3541-53  
Redox control in development and evolution: evidence from colonial hydroids
105. PMID: 20215510; 2010, Cancer Res;70(6):2296-306  
LYN is a mediator of epithelial-mesenchymal transition and a target of dasatinib in breast cancer.
106. PMID: 22490227; 2012, Clin Cancer Res;18(10):2850-60  
Lyn kinase mediates cell motility and tumor growth in EGFRvIII-expressing head and neck cancer.
107. PMID: 22731636; 2012, Mol Cancer;11():43  
Lyn is involved in CD24-induced ERK1/2 activation in colorectal cancer.
108. PMID: 16873062; 2006, Cell;126(2):297-308  
Endonucleolytic function of MutLalpha in human mismatch repair.
109. PMID: 22156295; 2012, Genome Res;22(2):208-19  
Whole-exome sequencing of human pancreatic cancers and characterization of genomic instability caused by MLH1 haploinsufficiency and complete deficiency.
110. PMID: 8484122; 1993, Science;260(5109):816-9  
Microsatellite instability in cancer of the proximal colon.
111. PMID: 8040889; 1994, J Natl Cancer Inst;86(16):1216-21  
Microsatellite instability in sporadic endometrial carcinoma.
112. PMID: 8261393; 1993, Cancer Res;53(24):5853-5  
Microsatellite instability is associated with tumors that characterize the hereditary non-polyposis colorectal carcinoma syndrome.

# ACT Onco<sup>®</sup> + Report

113. PMID: 9823339; 1998, Cancer Res;58(22):5248-57  
A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer.
114. PMID: 12454837; 2002, Gastroenterology;123(6):1804-11  
Evaluation of tumor microsatellite instability using five quasimonomorphic mononucleotide repeats and pentaplex PCR.
115. PMID: 27374179; 2017, Oncotarget;8(25):40152-40168  
Cumulative defects in DNA repair pathways drive the PARP inhibitor response in high-grade serous epithelial ovarian cancer cell lines.
116. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35  
mTOR: from growth signal integration to cancer, diabetes and ageing.
117. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6  
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
118. PMID: 9242607; 1997, Science;277(5327):805-8  
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
119. PMID: 8269512; 1993, Cell;75(7):1305-15  
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
120. PMID: 1303246; 1992, Nat Genet;2(1):37-41  
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
121. PMID: 18538015; 2008, BMC Cancer;8():163  
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
122. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784  
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
123. PMID: 20610279; 2010, Urol Oncol;28(4):409-28  
Bladder cancer or bladder cancers? Genetically distinct malignant conditions of the urothelium.
124. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56  
The tuberous sclerosis complex.
125. PMID: 22923433; 2012, Science;338(6104):221  
Genome sequencing identifies a basis for everolimus sensitivity.
126. PMID: 26859683; 2016, Oncotarget;7(9):10547-56  
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
127. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8  
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
128. PMID: 26412398; 2015, Sci Rep;5():14534  
PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.
129. PMID: 21756109; 2011, Annu Rev Cell Dev Biol;27():563-84  
Developmental and pathological angiogenesis.
130. PMID: 15585754; 2005, J Clin Oncol;23(5):1011-27  
Role of the vascular endothelial growth factor pathway in tumor growth and angiogenesis.
131. PMID: 10815805; 2000, Oncogene;19(17):2138-46  
Roles of two VEGF receptors, Flt-1 and KDR, in the signal transduction of VEGF effects in human vascular endothelial cells.

# ACT Onco<sup>®</sup> + Report

132. PMID: 9066636; 1997, Anticancer Res;17(1A):99-103  
Angiogenic growth factors and their receptors in non-small cell lung carcinomas and their relationships to drug response in vitro.
133. PMID: 7821921; 1995, Hum Pathol;26(1):86-91  
Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in breast cancer.
134. PMID: 8402650; 1993, Cancer Res;53(19):4727-35  
Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract.
135. PMID: 8238242; 1993, Am J Pathol;143(5):1255-62  
Increased expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in kidney and bladder carcinomas.
136. PMID: 8697391; 1996, Cancer;78(3):454-60  
Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in endometrial carcinoma.
137. PMID: 15949234; 2005, Neurol Res;27(4):371-7  
Expression of VEGF and its receptors in different brain tumors.
138. PMID: 14570746; 2003, Endocr Rev;24(5):600-32  
Angiogenesis in endocrine tumors.
139. PMID: 14654557; 2003, Clin Cancer Res;9(15):5721-8  
Vascular endothelial growth factor-trap decreases tumor burden, inhibits ascites, and causes dramatic vascular remodeling in an ovarian cancer model.
140. PMID: 16203789; 2005, Clin Cancer Res;11(19 Pt 1):6966-71  
Vascular endothelial growth factor trap combined with paclitaxel strikingly inhibits tumor and ascites, prolonging survival in a human ovarian cancer model.
141. PMID: 21992853; 2011, Lancet Oncol;12(12):1109-17  
Phase 1-2 study of docetaxel plus aflibercept in patients with recurrent ovarian, primary peritoneal, or fallopian tube cancer.
142. PMID: 25991856; 2015, Proc Natl Acad Sci U S A;112(22):E2900-9  
VEGF-B promotes cancer metastasis through a VEGF-A-independent mechanism and serves as a marker of poor prognosis for cancer patients.
143. PMID: 9570355; 1997, Cancer Lett;121(2):169-75  
Vascular endothelial growth factor (VEGF) expression and survival in human epithelial ovarian carcinomas.
144. PMID: 9210714; 1997, Cancer;80(1):98-106  
Vascular endothelial growth factor expression in early stage ovarian carcinoma.
145. PMID: 15087251; 2004, Clin Biochem;37(5):363-9  
Vascular endothelial growth factor and interleukin-8 are associated with poor prognosis in epithelial ovarian cancer patients.
146. PMID: 28968163; 2017, J Clin Oncol;35(32):3638-3646  
MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer.
147. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884  
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
148. PMID: 28533223; 2017, Clin Cancer Res;23(17):5218-5224  
MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR+/HER2- Metastatic Breast Cancer.
149. PMID: 26156651; 2015, Lancet Oncol;16(8):897-907  
Afatinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label

# ACT Onco<sup>®</sup> + Report

randomised controlled phase 3 trial.

150. PMID: 23816960; 2013, J Clin Oncol;31(27):3327-34  
Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations.
151. PMID: 19339720; 2009, N Engl J Med;360(14):1408-17  
Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer.
152. PMID: 18784101; 2008, N Engl J Med;359(11):1116-27  
Platinum-based chemotherapy plus cetuximab in head and neck cancer.
153. PMID: 16467544; 2006, N Engl J Med;354(6):567-78  
Radiotherapy plus cetuximab for squamous-cell carcinoma of the head and neck.
154. PMID: 15269313; 2004, N Engl J Med;351(4):337-45  
Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer.
155. PMID: 20525995; 2010, N Engl J Med;362(24):2260-70  
Dasatinib versus imatinib in newly diagnosed chronic-phase chronic myeloid leukemia.
156. PMID: 18541900; 2008, J Clin Oncol;26(19):3204-12  
Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia.
157. PMID: 17496201; 2007, Blood;110(7):2309-15  
Dasatinib induces rapid hematologic and cytogenetic responses in adult patients with Philadelphia chromosome positive acute lymphoblastic leukemia with resistance or intolerance to imatinib: interim results of a phase 2 study.
158. PMID: 22285168; 2012, Lancet Oncol;13(3):239-46  
Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTAC): a multicentre, open-label, randomised phase 3 trial.
159. PMID: 17452677; 2007, J Clin Oncol;25(15):1960-6  
Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group.
160. PMID: 26703889; 2016, Lancet;387(10022):968-977  
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
161. PMID: 22149876; 2012, N Engl J Med;366(6):520-9  
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
162. PMID: 21306238; 2011, N Engl J Med;364(6):514-23  
Everolimus for advanced pancreatic neuroendocrine tumors.
163. PMID: 23158522; 2013, Lancet;381(9861):125-32  
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
164. PMID: 18653228; 2008, Lancet;372(9637):449-56  
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
165. PMID: 24263064; 2014, Br J Cancer;110(1):55-62  
First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study.
166. PMID: 26045340; 2015, Lancet Oncol;16(7):763-74  
Necitumumab plus gemcitabine and cisplatin versus gemcitabine and cisplatin alone as first-line therapy in patients with stage IV squamous non-small-cell lung cancer (SQUIRE): an open-label, randomised, controlled phase 3 trial.

# ACT Onco<sup>®</sup> + Report

167. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284  
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
168. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589  
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
169. PMID: 25366685; 2015, J Clin Oncol;33(3):244-50  
Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation.
170. PMID: 27959700; 2017, N Engl J Med;376(7):629-640  
Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer.
171. PMID: 27959613; 2016, N Engl J Med;375(20):1925-1936  
Palbociclib and Letrozole in Advanced Breast Cancer.
172. PMID: 26030518; 2015, N Engl J Med;373(3):209-19  
Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer.
173. PMID: 27736842; 2016, Br J Cancer;115(10):1206-1214  
A phase 3 trial evaluating panitumumab plus best supportive care vs best supportive care in chemorefractory wild-type KRAS or RAS metastatic colorectal cancer.
174. PMID: 24024839; 2013, N Engl J Med;369(11):1023-34  
Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer.
175. PMID: 24739896; 2014, Lancet Oncol;15(6):569-79  
Panitumumab versus cetuximab in patients with chemotherapy-refractory wild-type KRAS exon 2 metastatic colorectal cancer (ASPECCT): a randomised, multicentre, open-label, non-inferiority phase 3 study.
176. PMID: 17470858; 2007, J Clin Oncol;25(13):1658-64  
Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer.
177. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748  
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
178. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87  
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
179. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81  
Temsilimus, interferon alfa, or both for advanced renal-cell carcinoma.