

Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C. Tel: 02-2875-7449

Date: 09 Apr 2020 1 of 6

Indicated Contraindicated

Sample Information

Patient Name: 藍嘉齊

Gender: Male **ID No.:** G121456429 **History No.:** 43719411

Age: 38

Ordering Doctor: DOC8340G 劉品均

Ordering REQ.: 0AQWDXP Signing in Date: 2020/04/09

Path No.: \$109-99340 **MP No.:** F2009

Assay: Oncomine Focus Assay

Sample Type: FFPE Block No.: S109-75857A Percentage of tumor cells: 90%

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	1 Clinically Significant Biomarkers
Biomarker Descriptions	3	0 Therapies Available
Relevant Therapy Summary	3	6 Clinical Trials

Relevant Non-Small Cell Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	Not detected	NTRK1	Not detected
BRAF	Not detected	NTRK2	Not detected
EGFR	Not detected	NTRK3	Not detected
ERBB2	Not detected	RET	Not detected
KRAS	Not detected	ROS1	Not detected
MET	Not detected		

Clinically Significant Biomarkers

Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
CDK4 amplification cyclin dependent kinase 4	None	None	6
Tier: IIC			

Sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO



Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

Date: 09 Apr 2020 2 of 6

Prevalent cancer biomarkers without clinical significance based on included data sources CTNNB1 p.(D32H) c.94G>C

Tier Criteria Met

Genomic Alteration	Tier Classification for Non-Small Cell Lung Cancer		
CDK4 amplification Tier: IIC	IIC: Biomarker is an inclusion criteria for clinical trials		

Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
CTNNB1	p.(D32H)	c.94G>C	COSM5668	chr3:41266097	15.71%	NM_001904.3	missense	1999
JAK1	p.(=)	c.2199A>G		chr1:65310489	99.09%	NM_002227.3	synonymous	1983
ALK	p.(D1529E)	c.4587C>G		chr2:29416366	100.00%	NM_004304.4	missense	1998
ALK	p.(I1461V)	c.4381A>G		chr2:29416572	99.95%	NM_004304.4	missense	1998
ALK	p.(=)	c.3375C>A		chr2:29445458	100.00%	NM_004304.4	synonymous	1998
FGFR3	p.(=)	c.1953G>A		chr4:1807894	99.90%	NM_000142.4	synonymous	1999
PDGFRA	p.(=)	c.1701A>G		chr4:55141055	99.95%	NM_006206.5	synonymous	1998
PDGFRA	p.(=)	c.2472C>T		chr4:55152040	38.20%	NM_006206.5	synonymous	2000
FGFR4	p.(P136L)	c.407C>T		chr5:176517797	99.45%	NM_213647.2	missense	1999
EGFR	p.(H773fs)	c.2317_2318insC		chr7:55249014	5.44%	NM_005228.4	frameshift Insertion	1968
FGFR1	p.(=)	c.2178T>G		chr8:38271771	8.57%	NM_001174067.1	synonymous	1353
MYC	p.(G123R)	c.367G>A		chr8:128750830	4.95%	NM_002467.4	missense	2000
RET	p.(=)	c.2307G>T		chr10:43613843	99.75%	NM_020975.4	synonymous	1997
MAP2K1	p.(Y134F)	c.401A>T		chr15:66729193	5.71%	NM_002755.3	missense	1998

Copy Number Variations Gene Locus Copy Number CDK4 chr12:58142052 5.62



Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

(II)

×

Date: 09 Apr 2020 3 of 6

Biomarker Descriptions

CDK4 (cyclin dependent kinase 4)

Background: The CDK4 gene encodes the cyclin-dependent kinase-4 protein, a homologue of CDK6. Both proteins are serine/threonine protein kinases that are involved in the regulation of the G1/S phase transition of the mitotic cell cycle^{1,2}. CDK4 kinase is activated by complex formation with D-type cyclins (e.g., CCND1, CCND2, or CCND3), which leads to the phosphorylation of retinoblastoma protein (RB), followed by E2F activation, DNA replication, and cell-cycle progression³. Germline mutations in CDK4 are associated with familial melanoma^{4,5,6}.

Alterations and prevalence: Recurrent somatic mutations of CDK4 codon K22 and R24 are observed in melanoma (1-2%) and lung cancer (approximately 0.1%). Codons K22 and R24 are necessary for binding and inhibition by p16/CDKN2A^{7,8,9}. CDK4 is recurrently amplified in several cancer types, most notably in sarcomas (15-20%), glioma (10-15%), adrenocortical carcinoma (5%), lung adenocarcinoma (5%), and melanoma (3%)^{10,11,12,13}.

Potential clinical relevance: Currently, no therapies are approved for CDK4 aberrations. Small molecule inhibitors targeting CDK4/6-including palbociclib (2015), abemaciclib (2017), and ribociclib (2017)— are FDA approved in combination with an aromatase inhibitor or fulvestrant for the treatment of hormone receptor-positive, HER2-negative advanced or metastatic breast cancer.

CTNNB1 (catenin beta 1)

Background: The CTNNB1 gene encodes catenin beta-1 (β -catenin), an integral component of cadherin-based adherens junctions involved in maintaining adhesion and regulating the growth of epithelial cell layers¹⁴. CTNNB1 binds to the APC protein in the cytoplasm and also interacts with TCF and LEF transcription factors in the nucleus to regulate WNT signaling¹⁵. Steady state levels of CTNNB1 are regulated by ubiquitin-dependent proteolysis^{16,17,18}.

Alterations and prevalence: Recurrent somatic mutations leading to CTNNB1 activation are common in cancer. The most prevalent alterations include missense mutations in exon 3 at codons S33, S37, T41, and S45 that block phosphorylation by GSK- β and inhibit CTNNB1 degradation^{19,20,21,22}. These activating mutations are observed in diverse solid tumors and have a prevalence of 20-30% in hepatocellular carcinoma, 20% of uterine carcinoma, and 15% of adrenocortical carcinoma^{10,11,23,24,25,26,27}.

Potential clinical relevance: Currently, no therapies have been approved for CTNNB1 alterations. CTNNB1 alterations in EGFR positive lung cancer have been proposed to promote cancer progression and limit the response to EGFR tyrosine kinase inhibitors²⁸.

Relevant Therapy Summary

In this cancer type	In other cancer (type	In this cancer type and other cancer types	⊘ Contraindicated	Both for use and contraindicated		No evidence
CDK4 amplificati	ion					
Relevant Therapy		FDA	NCCN	EMA	ESMO	Clinical Trials*
abemaciclib		×	×	×	×	(II)

×

×

Signatures

palbociclib

Testing Personnel:

^{*} Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Taipei Veterans General Hospital



Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C. Tel: 02-2875-7449

Date: 09 Apr 2020 4 of 6

Laboratory Supervisor:

Pathologist:

Taipei Veterans General Hospital



Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

Date: 09 Apr 2020 5 of 6

References

- 1. Malumbres et al. Cell cycle, CDKs and cancer: a changing paradigm. Nat. Rev. Cancer. 2009 Mar;9(3):153-66. PMID: 19238148
- 2. Sherr et al. Targeting CDK4 and CDK6: From Discovery to Therapy. Cancer Discov. 2016 Apr;6(4):353-67. PMID: 26658964
- 3. Weinberg. The retinoblastoma protein and cell cycle control. Cell. 1995 May 5;81(3):323-30. PMID: 7736585
- 4. Rane et al. Germ line transmission of the Cdk4(R24C) mutation facilitates tumorigenesis and escape from cellular senescence. Mol. Cell. Biol. 2002 Jan;22(2):644-56. PMID: 11756559
- 5. Zuo et al. Germline mutations in the p16INK4a binding domain of CDK4 in familial melanoma. Nat. Genet. 1996 Jan;12(1):97-9. PMID: 8528263
- 6. Molven et al. A large Norwegian family with inherited malignant melanoma, multiple atypical nevi, and CDK4 mutation. Genes Chromosomes Cancer. 2005 Sep;44(1):10-8. PMID: 15880589
- 7. Ceha et al. Several noncontiguous domains of CDK4 are involved in binding to the P16 tumor suppressor protein. Biochem. Biophys. Res. Commun. 1998 Aug 19;249(2):550-5. PMID: 9712735
- 8. Tsao et al. Novel mutations in the p16/CDKN2A binding region of the cyclin-dependent kinase-4 gene. Cancer Res. 1998 Jan 1;58(1):109-13. PMID: 9426066
- 9. Sotillo et al. Invasive melanoma in Cdk4-targeted mice. Proc. Natl. Acad. Sci. U.S.A. 2001 Nov 6;98(23):13312-7. PMID: 11606789
- 10. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4. PMID: 22588877
- 11. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
- 12. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. Nature. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
- 13. Brennan et al. The somatic genomic landscape of glioblastoma. Cell. 2013 Oct 10;155(2):462-77. PMID: 24120142
- 14. Valenta et al. The many faces and functions of β-catenin. EMBO J. 2012 Jun 13;31(12):2714-36. PMID: 22617422
- 15. Korinek et al. Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma. Science. 1997 Mar 21;275(5307):1784-7. PMID: 9065401
- 16. Aberle et al. beta-catenin is a target for the ubiquitin-proteasome pathway. EMBO J. 1997 Jul 1;16(13):3797-804. PMID: 9233789
- 17. Winston et al. The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha ubiquitination in vitro. Genes Dev. 1999 Feb 1;13(3):270-83. PMID: 9990852
- 18. Kitagawa et al. An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of beta-catenin. EMBO J. 1999 May 4;18(9):2401-10. PMID: 10228155
- 19. Liu et al. Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. Cell. 2002 Mar 22;108(6):837-47. PMID: 11955436
- 20. Miyoshi et al. Activation of the beta-catenin gene in primary hepatocellular carcinomas by somatic alterations involving exon 3. Cancer Res. 1998 Jun 15;58(12):2524-7. PMID: 9635572
- 21. Gao et al. Exon 3 mutations of CTNNB1 drive tumorigenesis: a review. Oncotarget. 2018 Jan 12;9(4):5492-5508. PMID: 29435196
- 22. Morin et al. Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC. Science. 1997 Mar 21;275(5307):1787-90. PMID: 9065402
- 23. Schulze et al. Exome sequencing of hepatocellular carcinomas identifies new mutational signatures and potential therapeutic targets. Nat. Genet. 2015 May;47(5):505-511. PMID: 25822088
- 24. Ahn et al. Genomic portrait of resectable hepatocellular carcinomas: implications of RB1 and FGF19 aberrations for patient stratification. Hepatology. 2014 Dec;60(6):1972-82. PMID: 24798001
- 25. Harding et al. Prospective Genotyping of Hepatocellular Carcinoma: Clinical Implications of Next-Generation Sequencing for Matching Patients to Targeted and Immune Therapies. Clin. Cancer Res. 2018 Oct 29. PMID: 30373752
- 26. Cancer et al. Integrated genomic characterization of endometrial carcinoma. Nature. 2013 May 2;497(7447):67-73. PMID: 23636398

Taipei Veterans General Hospital



Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

Date: 09 Apr 2020 6 of 6

References (continued)

- 27. Soumerai et al. Clinical Utility of Prospective Molecular Characterization in Advanced Endometrial Cancer. Clin. Cancer Res. 2018 Dec 1;24(23):5939-5947. PMID: 30068706
- 28. Blakely et al. Evolution and clinical impact of co-occurring genetic alterations in advanced-stage EGFR-mutant lung cancers. Nat. Genet. 2017 Dec;49(12):1693-1704. PMID: 29106415