

ACT Onco[®] + Report

PATIENT		
Name: 史玉琴		Patient ID: 48538267
Date of Birth: Mar 24, 1968		Gender: Female
Diagnosis: Thymic carcinoma		
ORDERING PHYSICIAN		
Name: 顏厥全醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11120097A	Collection site: Soft tissue	Type: FFPE tissue
Date received: Jul 22, 2022	Lab ID: AA-22-04243	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
Not detected			

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
BRAF D594G	Binimetinib, Encorafenib, Sorafenib, Vemurafenib	-
CAPZA2(1)-MET(2) fusion	-	Cabozantinib, Crizotinib

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACT Onco[®] + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
<i>BRAF</i>	D594G	32.5%
<i>MUC4</i>	P4858fs	48.1%
<i>TP53</i>	R248Q	51.8%

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr10	<i>PTEN</i>	Heterozygous deletion	1
Chr11	<i>CHEK1</i>	Heterozygous deletion	1
Chr16	<i>PALB2, TSC2</i>	Heterozygous deletion	1
Chr17	<i>RAD51C</i>	Heterozygous deletion	1
Chr18	<i>SMAD4</i>	Heterozygous deletion	1
Chr22	<i>CHEK2, NF2</i>	Heterozygous deletion	1
Chr4	<i>FBXW7</i>	Heterozygous deletion	1
Chr9	<i>CDKN2A</i>	Heterozygous deletion	1
Chr17	<i>ERBB2</i>	Amplification	6*

* Increased gene copy number was observed.

- Fusions

Fusion Gene & Exon	Transcript ID
<i>CAPZA2(1)-MET(2) fusion</i>	CAPZA2(NM_006136.2), MET(NM_000245.3)

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	6.3 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 61% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- The fusion gene reported above is confirmed to be in-frame and includes the kinase/functional domain. Such alteration may indicate potential benefits from kinase inhibitors. However, for a novel fusion, its functional significance and response to kinase inhibitors are undetermined.
- TMB was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 3B		
BRAF D594G	Binimetinib, Encorafenib	sensitive
Level 4		
BRAF D594G	Sorafenib, Vemurafenib	sensitive
CAPZA2(1)-MET(2) fusion	Cabozantinib, Crizotinib	resistant

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
Not detected	

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

Genomic Alterations	Therapies	Effect	Level of Evidence	Cancer Type
TP53 R248Q	Platinum- and taxane-based regimens	Less sensitive	Clinical	Ovarian cancer

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

CAPZA2(1)-MET(2) fusion

Biological Impact

The Mesenchymal-Epithelial Transition (MET) is an oncogene that encodes the MET receptor tyrosine kinase (c-MET, also called HGFR, hepatocyte growth factor receptor). Binding of HGF leads to autophosphorylation and activation of MET and downstream effectors through the PI3K/AKT and RAS/RAF/MEK pathways, which regulates cell growth, proliferation, migration, and angiogenesis^{[1][2]}. Gene amplification or overexpression of the MET occur in a wide range of cancers, including breast cancer^[3], non-small cell lung cancer (NSCLC)^[4], prostate cancer^[5], renal papillary carcinoma^{[6][7]}, glioblastoma^[8], hepatocellular carcinoma^[9], and gastric cancer^[10].

CAPZA2-MET fusion has been identified in glioblastoma^[11] (DOI: 10.1200/JCO.2017.35.15_suppl.2019), non-small cell lung cancer (<https://doi.org/10.1016/j.jtho.2018.08.451>) and endometrial cancer^[12].

MET gene rearrangement, which resulting in constitutively active MET fusion kinase, has been identified in various tumor types (DOI: 10.1200/JCO.2019.37.15_suppl.3078)^{[13][14]}.

Therapeutic and prognostic relevance

Preclinical data showed that 5-8F cell line which harbors CAPZA2-MET fusion gene is sensitive to tivantinib but resistance to crizotinib and cabozantinib (Int J Clin Exp Pathol 2016;9(8):8308-8317). Results from a MET-amplified glioblastoma patient-derived xenograft (PDX) model study revealed that capmatinib, a selective c-MET inhibitor, can improve overall survival of PDX models carrying MET amplification and in-frame fusion of exon 1 of CAPZA2 to exon 2 of MET, which resulted in full-length MET coding region, with altered 5' cis-regulatory sequences. However, no improvement of survival was observed in PDXs harboring MET amplification and in-frame fusion of exon 1 of CAPZA2 to exon 6 of MET (Abstract 2519, 2018 AACR).

BRAF D594G

Biological Impact

BRAF is a serine/threonine kinase that belongs to the RAF family. The protein plays an essential role in the regulation of mitogen-activated protein kinase (MAPK) cascade, which affects a range of cellular response including cell division, differentiation, and secretion^{[15][16]}. Mutations in the BRAF gene, most commonly the V600 residue, are the most frequently identified oncogenic mutations in melanomas, and have been identified in several types of cancers including non-Hodgkin lymphoma, thyroid cancers, non-small cell lung carcinoma, hairy cell leukemia, glioma, gastrointestinal stromal tumor, and colorectal cancers (CRCs)^{[17][18]}. Of note, in the vast majority of cases, BRAF mutations are non-overlapping with other oncogenic mutations (e.g., NRAS mutations, KIT mutations, etc.) found in melanoma. V600E has been determined to be an activating mutation, which results in enhanced BRAF kinase activity and constitutive activation of downstream MEK/ERK signaling cascade^{[19][20]}.

BRAF D594G mutation is located within the protein kinase domain of the BRAF protein, resulting in impaired BRAF kinase activity. Meanwhile, expression of this mutant in melanoma cells models induces constitutive levels of pERK through CRAF^{[21][22]}.

Therapeutic and prognostic relevance

In the phase II NCI-MATCH trial (NCT02465060), trametinib treatment resulted in stable disease in two patients with prostate cancer and lung adenocarcinoma respectively, both harboring BRAF D594G. However, four gastrointestinal system cancer patients harboring BRAF D594G resulted in progression disease under trametinib treatment^[23]. In the phase II BEAVER trial (NCT03839342), an ampullary cancer patient harboring BRAF D594G achieved a confirmed partial response to combination treatment with binimetinib and encorafenib, and continued to be on treatment for 5.4 months (Annals of Oncology 32 (2021): S596).

ACT Onco[®] + Report

Data from cell-based and xenograft studies demonstrated that the CRAF inhibitor, sorafenib, induced apoptosis and reduces tumor growth in melanoma cell lines harboring BRAF D594G mutation^[21]. Besides, a preclinical study revealed that cells harboring BRAF D594N/G/A mutant are sensitive to trametinib but are insensitive to vemurafenib^[22]. However, there is a case report showed that responses to vemurafenib were observed in a patient with advanced melanoma harboring D594G^[24].

The NCCN guidelines for central nervous system cancers recommended selumetinib for pilocytic astrocytoma patients with BRAF fusion or BRAF V600E activating mutation. BRAF activating mutations have been determined as an inclusion criterion for the trials evaluating selumetinib efficacies in cancers (NCT01089101, NCT00888134, NCT00866177, and NCT00936221).

MUC4 P4858fs

Biological Impact

Mucin 4 (MUC4) gene encodes a membrane protein of the mucin family which is expressed in normal stomach, ovary, salivary gland, colon, and lung^[25]. Because of its large size and extended conformation, MUC4 can protect the epithelium from invasion by microbes^[26]. Similarly, MUC4 also can protect tumor cells from killing mechanisms by immune cells and anti-HER2 therapy^{[27][28]}. In cancer, MUC4 has been proposed to promote cell differentiation and proliferation via interaction with ERBB2 and activating downstream signaling pathway^[29]. Of note, it is able to disrupt tumor cell-cell interaction and promote invasion and metastasis^[30]. Overexpression of MUC4 has been reported in many cancers such as pancreatic cancer^{[31][32]}, ovarian cancer^[33], breast cancer^[34], and colorectal cancer^[35].

P4858fs mutation results in a change in the amino acid sequence beginning at 4858, likely to cause premature truncation of the functional MUC4 protein (UniProtKB). This mutation is predicted to lead to a loss of MUC4 protein function, despite not being characterized in the literature.

Therapeutic and prognostic relevance

Relatively low expression of MUC4 was associated with better survival in patients with resectable pancreatic cancer receiving adjuvant gemcitabine^[36]. However, low MUC4 expression was correlated with shorter overall survival in clear-cell renal cell carcinoma^[25].

TP53 R248Q

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[37]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[38].

R248Q is a missense mutation located in the DNA-binding domain (DBD) of the p53 protein (UniProtKB). This mutation causes oncogenic gain-of-function phenotypes in a breast cancer cell line^[39] and has been reported to target the proteasome activator REG gamma to promote endometrial cancer progression^[40].

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[41].

ACT Onco[®] + Report

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[42]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[43].

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[44][45][46]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[47]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[48][49]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[50].

TP53 oncomorphic mutations, including P151S, Y163C, R175H, L194R, Y220C, R248Q, R248W, R273C, R273H, R273L and R282W have been shown to predict resistance to platinum- and taxane-based chemotherapy in advanced serous ovarian carcinoma patients^[51].

CDKN2A Heterozygous deletion

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^{[52][53][54]}. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[55]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[56][57]}.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[58][59]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[60][61][62]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[63][64][65]}. However, CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[59][66][67]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[61]. Administration of combined palbociclib and MEK inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[68].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[69].

ACT Onco[®] + Report

CHEK1 Heterozygous deletion

Biological Impact

The checkpoint kinase 1 (CHEK1 or CHK1) gene encodes a protein kinase involved in transducing DNA damage signals and is required for both the intra-S phase and G2/M checkpoints^[70]. CHEK1 heterozygosity has been shown to cause haploinsufficient phenotypes that can contribute to tumorigenesis through inappropriate S phase entry, accumulation of DNA damage during replication, and failure to restrain mitotic entry^{[71][72]}. Despite acting as a tumor suppressor, homozygous loss-of-function mutations in CHEK1 have not been identified in tumors^[73], and CHEK1 mutations are extremely rare^[70]. Overexpression of CHEK1 has been observed in a variety of tumors, including liver cancer^[74], breast cancer^[75], colorectal cancer^[76], non-small cell lung (NSCLC) cancer^[77], and nasopharyngeal cancer^[78].

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[79].

In addition, CHEK1 has been determined as an inclusion criterion for the trials evaluating olaparib efficacy in advanced solid tumors (NCT03297606; CAPTUR trial), rucaparib efficacy in ovarian cancer^[80], prostate cancer (NCT03533946), niraparib efficacy in pancreatic cancer (NCT03553004), and any malignancy, except prostate (NCT03207347), and talazoparib efficacy in lung cancer (NCT03377556), respectively.

Selective inhibitors for CHEK1 and CHEK2 alone or in combination with other agents are currently being investigated in clinical trials^[81].

CHEK2 Heterozygous deletion

Biological Impact

The checkpoint kinase 2 (CHEK2 or CHK2) gene encodes a serine/threonine protein kinase involved in transducing DNA damage signals that are required for both the intra-S phase and G2/M checkpoints^[82]. CHEK2 heterozygosity has been shown to cause haploinsufficient phenotypes that can contribute to tumorigenesis through inappropriate S phase entry, accumulation of DNA damage during replication, and failure to restrain mitotic entry^{[71][72]}. CHEK2 aberrations are associated with glioblastoma, breast, ovarian, prostate, colorectal, gastric, thyroid, and lung cancers^{[83][84][85][86][87]}.

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[79].

In addition, CHEK2 has been determined as an inclusion criterion for the trials evaluating rucaparib efficacy in ovarian cancer or prostate cancer (NCT03533946)^{[80][88]}, niraparib efficacy in melanoma (NCT03925350), pancreatic cancer (NCT03553004), prostate cancer (NCT02854436), and any malignancy, except prostate (NCT03207347), and talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795), and lung cancer (NCT03377556), respectively.

In a phase 2 trial, two prostate cancer patients harboring CHEK2 homozygous deletion was enrolled. One of the two patients had a response to olaparib^[89].

ACT Onco[®] + Report

ERBB2 Amplification

Biological Impact

The epidermal growth factor receptor 2 (HER2, or ERBB2) gene encodes a transmembrane receptor tyrosine kinase that belongs to the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases^[90]. Amplification or activating mutations of ERBB2 can lead to aberrant activation of downstream pathways, such as phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase (MAPK) cascades, which are involved in cell survival and proliferation, respectively^{[90][91]}. ERBB2 mutations are mostly observed in HER2-negative (non-overexpressed or non-amplified) samples^[92].

Therapeutic and prognostic relevance

Early Phase clinical trials and case reports have reported clinical benefits to afatinib and dacomitinib treatment in non-small cell lung cancer (NSCLC) patients harboring ERBB2 exon 20 in-frame insertion mutations^{[93][94][95][96][97][98]}. A preclinical study in lung cancer cell lines demonstrated that ERBB2 exon 20 in-frame insertion mutations resulted in constitutive phosphorylation and activation of ERBB2 and resistance to EGFR tyrosine kinase inhibitors, erlotinib and gefitinib^{[99][96]}. ERBB2 exon 20 in-frame insertion mutations in NSCLC cells were shown to be sensitive to neratinib and poziotinib^{[100][101]}.

There were case reports showing NSCLC patients with ERBB2 exon 20 in-frame insertion mutations derived benefit from trastuzumab-based therapies^{[94][102][103][104]}, or trastuzumab emtansine^{[104][105]}.

In NCCN guidelines for NSCLC, ERBB2 (HER2) mutations have been suggested as an emerging biomarker for ado-trastuzumab emtansine and fam-trastuzumab deruxtecan-nxki (T-DXd; DS-8201) in patients with NSCLC (DOI: 10.1200/JCO.2020.38.15_suppl.9504)^[106]. ERBB2 mutations have been determined as an inclusion criterion for the trials evaluating neratinib, lapatinib, afatinib, dacomitinib, and trastuzumab efficacies in multiple types of solid tumors (NCT01670877, NCT01953926, NCT01306045, NCT02780687, NCT00818441, and NCT02693535).

FDA has approved several HER2 targeting drugs including trastuzumab (NCT00004067, NCT00005970)^{[107][108]}, pertuzumab (NCT01358877)^[109], ado-trastuzumab emtansine (NCT01772472)^[110], fam-trastuzumab deruxtecan-nxki (DS-8201) (NCT03248492)^[111], lapatinib (NCT00078572)^[112], neratinib (NCT01808573, NCT00878709) (DOI 10.1200/JCO.2019.37.15_suppl.1002)^{[113][114][115]}, and tucatinib (NCT02614794)^[116] for HER2-positive breast cancer under different clinical settings.

Besides, fam-trastuzumab deruxtecan-nxki also has been approval by the US FDA for adult patients with locally advanced or metastatic HER2-positive gastric or gastroesophageal (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen.

Afatinib also demonstrates remarkable anti-tumor activity against HER2-amplified uterine serous endometrial cancer in vitro and in vivo^[117].

Aberrant activation of ERBB2 by gene amplification or activating mutations may confer increased sensitivity to several ERBB-directed agents, such as trastuzumab^[118], ado-trastuzumab emtansine (T-DM1)^{[119][120]}, lapatinib^{[121][122]}, and neratinib^{[123][124][125]} in breast cancer and ovarian cancer^[126] (DOI 10.3816/COC.2008.n.005).

Of note, combined neoadjuvant therapy consisting of pertuzumab and trastuzumab achieved a significantly higher response rate than with trastuzumab alone^{[120][127][128]}. FDA has approved trastuzumab for metastatic gastric or gastroesophageal junction adenocarcinoma (NCT01041404)^{[107][129]}.

In NCCN guidelines for colon cancer, HER2 amplification has been suggested as an emerging biomarker for fam-trastuzumab deruxtecan-nxki or trastuzumab plus either pertuzumab or lapatinib in mCRC patients^{[130][131]}. For salivary

ACT Onco[®] + Report

gland tumors, the NCCN guidelines for head and neck cancer also suggested that HER2 amplification is an emerging biomarker for trastuzumab, T-DM1, pertuzumab, and fam-trastuzumab deruxtecan-nxki^{[132][126][133][134]}(DOI: 10.1200/JCO.2021.39.15_suppl.6079). For esophageal cancer, the NCCN guidelines suggested that HER2 amplification is an emerging biomarker for fam-trastuzumab deruxtecan-nxki, trastuzumab and FDA-approved biosimilar of trastuzumab^{[129][111]}. For patients with stage III/IV or recurrent HER2-positive uterine serous carcinoma, the NCCN guidelines suggested that HER2 amplification is an emerging biomarker for carboplatin/paclitaxel plus trastuzumab^[135].

Clinical studies have shown that neither trastuzumab nor lapatinib exhibited significant activity in HER2-overexpressing advanced salivary gland tumors^{[136][137]}. However, although no response of lapatinib was observed, it was well tolerated, with prolonged tumor stabilization of more than 6 months in 36% (95% CI, 21% to 54%) of patients^[137]. In addition, trastuzumab in combination with paclitaxel and carboplatin showed a durable clinical response in a patient with HER2-positive metastatic submandibular salivary ductal carcinoma^[138].

In ovarian cancers, clinical evaluation of trastuzumab and pertuzumab in Phase 2 studies showed clinical benefit of in 7.3% and 14.5%, respectively in unselected patients^{[139][140]}.

Beyond EGFR T790M, several mechanisms of acquired resistance to first- and second-generation EGFR-TKIs have been identified in lung cancer patients, including rare EGFR resistance-associated point mutations and HER2 amplification^[141]. The NCCN guidelines for NSCLC suggests that acquired resistance can also be mediated by the acquisition of ERBB2 amplification^[142].

As reviewed in Nahta et al. and Rexer^{[143][144]}, accumulated evidence have suggested several intrinsic or acquired mechanisms that could contribute to trastuzumab and lapatinib resistance, such as alternations of the drug-binding domain of the HER2 gene that abrogate trastuzumab binding, overexpression of mucins that block the binding of trastuzumab, loss-of-function mutations of the phosphatase and tensin homolog (PTEN) gene or activating mutations of PIK3CA gene, and upregulation of IGF-1 or MET^{[145][146][147][148][149]}. However, there is conflicting data suggested that gain-of-function PIK3CA mutations may not be associated with anti-HER2 resistance^[150]. Several clinical trials of inhibitors targeting PI3K/AKT pathway combined with HER2-targeted treatments are underway^{[151][152]}.

ERBB2 gene amplification is associated with shorter overall survival and time to disease recurrence in breast cancers^{[153][154][155]}. ERBB2 amplification has been implicated in colorectal cancer patients who exhibited either de novo or acquired resistance to cetuximab-based therapy^[156]. Studies also suggested that overexpression of the ERBB2 gene is associated with resistance to tamoxifen therapy in patients with advanced breast cancer^{[157][158][159]}. HER2 positivity was correlated with a worse survival outcome in endometrial cancer^{[160][161][162]}.

FBXW7 Heterozygous deletion

Biological Impact

The F-box/WD repeat-containing protein 7 (FBXW7) gene encodes a protein that belongs to the SCF (SKP1-CUL1-F-box protein) E3 ligase complex. FBXW7 is recognized as a tumor suppressor which is involved in the negative regulation of oncogenes such as c-Myc^{[163][164]}, c-Jun^[165], cyclin E^[166], Notch family members^{[167][168]}, Aurora-A^[169], mTOR^[170], KLF5^[171], and MCL-1^[172]. Inactivating FBXW7 mutation or copy number loss may result in the accumulation of oncoproteins and therefore lead to malignant transformation^[173]. FBXW7 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[171][172][174]}.

Therapeutic and prognostic relevance

Clinical efficacy of mTOR inhibitors was seen in patients harboring aberrations in the FBXW7 gene (one patient with refractory fibrolamellar hepatocellular carcinoma, and one patient with lung adenocarcinoma)^{[175][176]}. Moreover, in vitro

ACT Onco[®] + Report

assay also suggested that loss or inactivation of FBXW7 may confer sensitivity to mTOR inhibitor^[170].

Preclinical studies suggested that mutations or loss of FBXW7 were associated with regorafenib and oxaliplatin resistance in CRC cell lines and gefitinib resistance in lung cancer cells^{[177][178][179][180]}.

Retrospective studies have indicated that a relatively low expression level of FBXW7 is an independent prognostic marker of poor survival for patients with hepatocellular carcinoma, lung adenocarcinoma and squamous cell carcinoma^{[181][179]}.

NF2 Heterozygous deletion

Biological Impact

The neurofibromin (NF2) gene encodes the protein Merlin, a tumor suppressor that functions as a negative regulator of the PI3K/AKT/mTOR pathway^{[182][183][184]}. NF2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[185]. Inactivation germline mutations in the NF2 are associated with the hereditary neurofibromatosis type 2, a disorder characterized by the growth of noncancerous tumors in the nervous system^{[182][186]}. Somatic mutations or deletion of NF2 are frequently observed in human cancers, including 20-50% of pleural mesotheliomas^[187], 6% papillary renal cell carcinoma, 5% pancreas cancer, and 4% melanoma (cbioPortal; June 2015), and less frequently in other cancers^[188].

Therapeutic and prognostic relevance

Genomic alterations with activating effects on the mTOR signaling pathway have been identified to confer sensitivity to everolimus across multiple cancer types^{[189][190][191][192]}. There are at least two case studies indicating the clinical efficacy of everolimus in bladder cancer and urothelial carcinoma^{[193][194]}, both harboring NF2 truncating mutations. Preclinical evidence has shown the efficacy of MEK1/2 inhibitor selumetinib in KRAS-mutant thyroid cancer model with NF2 loss^[195].

Analysis of afatinib-plus-cetuximab-resistant biopsy specimens revealed a loss-of-function alteration in genes that modulate mTOR signaling pathway, including NF2 and TSC1^[196].

PALB2 Heterozygous deletion

Biological Impact

The partner and localizer of BRCA2 (PALB2) gene encodes a protein that plays a critical role in homologous recombination repair (HRR) through its ability to interact with BRCA2 in nuclear foci, promoting its localization and stability in key nuclear structures^[197]. The Fanconi anemia complementation group (FANC) which includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM and FANCN (also called PALB2) are involved in the repair of DNA double-strand breaks (DSBs) by homologous recombination (HR)^{[198][199][200]}. PALB2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological function^[201]. Biallelic germline loss-of-function mutations in PALB2 cause Fanconi anemia, whereas monoallelic loss-of-function mutations are associated with an increased risk of breast cancer and pancreatic cancer^[202]. Fanconi Anemia is an autosomal recessive disease characterized by hematological abnormalities, bone marrow failure, limb deformities, skin hyperpigmentation, and susceptibility to hematologic and solid malignancies, such as acute myeloid leukemia and head and neck carcinoma^{[203][204]}.

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and

ACT Onco[®] + Report

progressed following prior treatment with enzalutamide or abiraterone acetate^[79].

PALB2 loss of function mutation has been determined as an inclusion criterion for the trial evaluating rucaparib efficacy in ovarian cancer^[80] or prostate cancer^[88]; talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795) or lung cancer (NCT03377556); niraparib efficacy in pancreatic cancer (NCT03553004), prostate cancer (NCT02854436), or any malignancy (except prostate) cancer (NCT03207347).

A case report demonstrated an exceptional response to mitomycin C and cisplatin treatment in a gemcitabine-resistant pancreatic cancer patient with biallelic inactivation of PALB2^[205].

PTEN Heterozygous deletion

Biological Impact

The phosphatase and tensin homolog deleted on chromosome ten (PTEN) gene encodes a lipid/protein phosphatase that is important for the regulation of cell proliferation, survival, homologous recombination and maintenance of genomic integrity^{[206][207]}. PTEN acts as an essential tumor suppressor by antagonizing the PI3K/AKT/mTOR signaling pathway^[208]. PTEN is a haploinsufficient tumor suppressor gene, in which having only one copy of the wild-type allele does not produce enough protein product to execute wild-type functions^{[38][209][210]}. Germline loss-of-function PTEN mutations are found in approximately 80% of patients with Cowden syndrome, a disorder that is associated with high-penetrance breast and thyroid cancer^{[211][212][213]}. Somatic mutations or monoallelic loss of PTEN is regularly observed in a significant fraction of human cancers, including sporadic breast cancer, colon cancer, endometrial cancer, prostate cancer, and glioblastoma^{[214][215][216][217][218]}.

Therapeutic and prognostic relevance

Somatic loss of PTEN results in aberrant activation of PI3K/AKT/mTOR signaling pathway and provides a mechanistic rationale for PI3K pathway inhibitors treatment^{[219][220]}. Preclinical studies demonstrated that PTEN deficiency was associated with increased sensitivity to PI3K pathway inhibitors in selected cancer subtypes^{[221][222][223][224][225][226]}. Moreover, early clinical data also indicated that PTEN loss was associated with improved response and longer PFS in patients with advanced breast cancer^[189], advanced pancreatic neuroendocrine tumors^[227], and metastatic castration-resistant prostate cancer treated with mTORC1 inhibitor, everolimus^[228].

Several groups found that PTEN loss was generally associated with poor response to trastuzumab therapy, whether this agent was administered in the neoadjuvant, adjuvant, or metastatic settings^{[145][147][229][148][149]}.

Loss of PTEN expression in advanced colorectal cancer (CRC) has been linked with resistance to anti-EGFR mAbs like cetuximab and panitumumab^{[230][231][232][233][234][235]}. However, encouraging anti-tumor activity of the combination of an EGFR antibody and a mTORC1 inhibitor (everolimus or temsirolimus) have been reported in early-phase clinical studies (J Clin Oncol. 2011;29 (suppl): abstr 3587; J Clin Oncol. 2013;31 (suppl): abstr 608). Ongoing phase I/II studies testing combinations of EGFR antibodies and PI3K/AKT/mTOR pathway inhibitors (e.g., NCT01816984, NCT01252628, NCT01719380) will provide larger numbers of patients to assess the role of PTEN status in therapeutic response.

Preclinical studies showed that loss of PTEN expression in EGFR mutant cells was associated with decreased sensitivity to EGFR TKIs, erlotinib and gefitinib^{[236][237]}. Inhibition of the PI3K/AKT/mTOR signal pathway has been shown to be an effective strategy to radiosensitize NSCLC cells harboring the EGFR activating mutation that acquires resistance to both TKIs due to PTEN loss or inactivation mutations^[238].

Loss or biallelic inactivation of PTEN is associated with resistance to anti-PD-1 checkpoint blockade therapies, including pembrolizumab and nivolumab in melanoma and leiomyosarcoma patients^{[239][240][241]}.

PTEN loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy

ACT Onco[®] + Report

in metastatic biliary tract cancer (NCT04042831); talazoparib efficacy in HER2-negative breast cancer (NCT02401347), and niraparib efficacy in breast cancer (NCT04508803) or any malignancy (except prostate) cancer (NCT03207347). Clinical data also suggested that PTEN deficient cancers may be sensitive to olaparib^[242].

RAD51C Heterozygous deletion

Biological Impact

The RAD51C (RAD51 paralog C) encodes a member of the RAD51 protein family involved in the late phase of homologous recombination DNA repair. Germline mutations in RAD51C have been shown to confer increased susceptibility to ovarian cancer and head and neck squamous cell carcinoma (HNSCC)^{[243][244][245][246][247]}. Amplification of RAD51C has been implicated in tumor progression^{[248][249]}. RAD51C is a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological function^[250].

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[79].

A preclinical study using gastric cancer xenograft model showed that RAD51C deficiency caused sensitivity to PARP inhibitor olaparib^[251].

RAD51C loss of function mutation has been determined as an inclusion criterion for the trials evaluating rucaparib efficacy in ovarian cancer or prostate cancer^{[80][88]}; talazoparib efficacy in HER2-negative breast cancer (NCT02401347) or prostate cancer (NCT03148795), and niraparib efficacies in pancreatic cancer (NCT03553004).

SMAD4 Heterozygous deletion

Biological Impact

The SMAD family member 4 (SMAD4) gene encodes a transcription factor that acts as a downstream effector in the TGF- β signaling pathway. Upon phosphorylated and activated by serine-threonine receptor kinase, Smad4 is the Co-Smad which recruits other activated R-Smad proteins to the Smad transcriptional complex and regulate TGF- β -targeted genes^[252]. Smad4 has been identified as a haploinsufficient gene with one copy loss may lead to a weak protein expression and is insufficient to execute its original physiological function^[253]. SMAD4 germline mutations are associated with juvenile polyposis syndrome (JPS)^{[254][255][256][257]}. Somatic mutations of SMAD4 are commonly observed in pancreatic cancer^[258], colorectal cancer (CRC)^{[256][259][260]}, and less frequently seen in other cancers such as lung adenocarcinoma^[261], head and neck cancer^{[262][263]}, and cutaneous squamous cell carcinoma^[264].

Therapeutic and prognostic relevance

In Chinese patients with metastatic colorectal cancer, SMAD4 or NF1 mutations are suggested as a potential biomarker for poor prognosis to cetuximab-based therapy^[265]. Preclinical data demonstrated that depletion of SMAD4 by shRNA knockdown increased clonogenic survival and cetuximab resistance in HPV-negative head and neck squamous cell carcinoma cells^[266].

SMAD4 is also suggested as a predictive marker for 5-fluorouracil-based chemotherapy in colorectal cancer (CRC)^{[267][268]}. CRC patients with normal SMAD4 diploidy exhibited three-fold higher benefit of 5-FU/mitomycin-based adjuvant therapy when compared with those with SMAD4 deletion^[269].

ACT Onco[®] + Report

Results from clinical and meta-analyses showed that loss of SMAD4 in CRC, pancreatic cancer was correlated with poor prognosis^{[270][271][272][273][274][275][276][277]}. In cervical cancer patients, weak cytoplasmic SMAD4 expression and absent nuclear SMAD4 expression were shown to be significantly associated with poor disease-free and overall 5-year survival^[278].

TSC2 Heterozygous deletion

Biological Impact

The tuberous sclerosis complex 2 (TSC2) gene encodes a protein called tuberin, which interact with a protein called hamartin (encoded by the TSC1 gene). This hamartin-tuberin tumor suppressor complex plays a critical role in growth control as a negative regulator of the mammalian target of rapamycin (mTOR) pathway^{[279][280]}. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis complex^{[281][282][283]}, while the loss of heterozygosity (LOH) in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)^[284] and endometrial cancer^[285]. TSC2 deletion, splicing-mutant, and inactivating mutations such as A1141T, G305V, S1514X, and R1032X, has been identified in TSC2-null hepatocellular carcinoma (HCC) cell lines, patient-derived xenograft, and primary tumors. Mutations in the TSC1 and TSC2 genes cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC)^[286].

Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple cancer types, such as bladder cancer, gastric cancer, sarcoma, thyroid cancer, hepatocellular carcinoma (HCC) as well as head and neck squamous cell carcinoma (HNSCC)^{[193][192][287]}. Results from one Phase II study of advanced endometrial cancer showed that mutations in AKT1, TSC1, and TSC2 might predict sensitivity to temsirolimus^[288]. Recent studies indicated that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets^[289].

Everolimus has been approved by the U.S. FDA for Tuberous Sclerosis Complex (TSC)-associated renal angiomyolipoma and Tuberous Sclerosis Complex (TSC)-associated subependymal giant cell astrocytoma (SEGA). This approval is based on the results from EXIST-1, EXIST-2, and Study 2485 trials (NCT00789828, NCT00790400, and NCT00411619).

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Abemaciclib (VERZENIO)

Abemaciclib is a cyclin-dependent kinase 4/6 (CDK4/6) inhibitor. Abemaciclib is developed and marketed by Eli Lilly under the trade name VERZENIO.

- FDA Approval Summary of Abemaciclib (VERZENIO)

monarchE NCT03155997	Breast cancer (Approved on 2021/10/12)
	HR-positive, HER2-negative Abemaciclib + tamoxifen/aromatase inhibitor vs. Tamoxifen/aromatase inhibitor [IDFS at 36 months(%): 86.1 vs. 79.0]
MONARCH 3 ^[290] NCT02246621	Breast cancer (Approved on 2018/02/26)
	HR-positive, HER2-negative Abemaciclib + anastrozole/letrozole vs. Placebo + anastrozole/letrozole [PFS(M): 28.2 vs. 14.8]
MONARCH 2 ^[67] NCT02107703	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative Abemaciclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 16.4 vs. 9.3]
MONARCH 1 ^[291] NCT02102490	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative Abemaciclib [ORR(%): 19.7 vs. 17.4]

Binimetinib (MEKTOVI)

Binimetinib is an oral kinase inhibitor that targets MEK. Binimetinib is developed and marketed by Array BioPharma under the trade name MEKTOVI.

- FDA Approval Summary of Binimetinib (MEKTOVI)

MEKTOVI ^[292] NCT01909453	Melanoma (Approved on 2018/06/27)
	BRAF V600E/K Encorafenib + binimetinib vs. Vemurafenib [PFS(M): 14.9 vs. 7.3]

Encorafenib (BRAFTOVI)

Encorafenib is an oral kinase inhibitor that targets BRAF. Encorafenib is developed and marketed by Array BioPharma under the trade name BRAFTOVI.

- FDA Approval Summary of Encorafenib (BRAFTOVI)

BEACON CRC ^[293] NCT02928224	Colorectal cancer (Approved on 2020/04/08)
	BRAF V600E Encorafenib in combination with cetuximab vs. Irinotecan or folfiri with cetuximab [OS(M): 8.4 vs. 5.4]
COLUMBUS ^[292] NCT01909453	Melanoma (Approved on 2018/06/27)
	BRAF V600E/K Encorafenib + binimetinib vs. Vemurafenib [PFS(M): 14.9 vs. 7.3]

ACT Onco[®] + Report

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4 ^[294] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
BOLERO-2 ^[295] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
EXIST-2 NCT00790400	Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
RADIANT-3 ^[227] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1 ^[296] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
RECORD-1 ^[297] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (ZEJULA)

PRIMA NCT02655016	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29)
	- Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
QUADRA ^[298] NCT02354586	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2019/10/23)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability) Niraparib [ORR(%): 24.0, DOR(M): 8.3]
NOVA ^[299] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	- Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

ACT Onco[®] + Report

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (LYNPARZA)

OlympiA NCT02032823	Her2-negative high-risk early breast cancer (Approved on 2022/03/11)
	gBRCA Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):]
PROfound ^[79] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	ATMm, BRCA1m, BRCA2m, BARD1m, BRIP1m, CDK12m, CHEK1m, CHEK2m, FANCLm, PALB2m, RAD51Bm, RAD51Cm, RAD51Dm, RAD54Lm Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1 ^[300] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability) Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
POLO ^[301] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	Germline BRCA mutation (deleterious/suspected deleterious) Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1 ^[302] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	Germline or somatic BRCA-mutated (gBRCAm or sBRCAm) Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
OlympiAD ^[303] NCT02000622	Breast cancer (Approved on 2018/02/06)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21 ^[304] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA+ Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
Study19 ^[305] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	- Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]
Study 42 ^[306] NCT01078662	Ovarian cancer (Approved on 2014/12/19)
	Germline BRCA mutation (deleterious/suspected deleterious) Olaparib [ORR(%): 34.0, DOR(M): 7.9]

Palbociclib (IBRANCE)

Palbociclib is an oral, cyclin-dependent kinase (CDK) inhibitor specifically targeting CDK4 and CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Palbociclib is developed and marketed by Pfizer under the trade name IBRANCE.

- FDA Approval Summary of Palbociclib (IBRANCE)

PALOMA-2 ^[307] NCT01740427	Breast cancer (Approved on 2017/03/31)
	ER+, HER2- Palbociclib + letrozole vs. Placebo + letrozole [PFS(M): 24.8 vs. 14.5]

ACT Onco[®] + Report

PALOMA-3 ^[308] NCT01942135	Breast cancer (Approved on 2016/02/19)
	ER+, HER2-
	Palbociclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 9.5 vs. 4.6]

Ribociclib (KISQALI)

Ribociclib is a cyclin-dependent kinase (CDK) inhibitor specifically targeting cyclin D1/CDK4 and cyclin D3/CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Ribociclib is developed by Novartis and Astex Pharmaceuticals and marketed by Novartis under the trade name KISQALI.

- FDA Approval Summary of Ribociclib (KISQALI)

MONALEESA-2 ^[66] NCT01958021	Breast cancer (Approved on 2017/03/13)
	HR+, HER2-
	Ribociclib vs. Letrozole [PFS(M): NR vs. 14.7]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA+, sBRCA
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
ARIEL3 ^[80] NCT01968213	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	All HRD tBRCA
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]
ARIEL2 ^[309] NCT01482715, NCT01891344	Ovarian cancer (Approved on 2016/12/19)
	Germline and/or somatic BRCA mutation
	Rucaparib [ORR(%): 54.0]

Sorafenib (NEXAVAR)

Sorafenib is a small molecule multi-kinase inhibitor that targets multiple kinase families including VEGFR, PDGFRB, and the RAF family kinases. Sorafenib is co-developed and co-marketed by Bayer HealthCare Pharmaceuticals and Onyx Pharmaceuticals under the trade name NEXAVAR.

- FDA Approval Summary of Sorafenib (NEXAVAR)

DECISION ^[310] NCT00984282	Differentiated thyroid carcinoma (Approved on 2013/11/22)
	-
	Sorafenib vs. Placebo [PFS(M): 10.8 vs. 5.8]
SHARP ^[311] NCT00105443	Hepatocellular carcinoma (Approved on 2007/11/16)
	-
	Sorafenib vs. Placebo [OS(M): 10.7 vs. 7.9]

ACT Onco[®] + Report

TARGET ^[312] NCT00073307	Renal cell carcinoma (Approved on 2005/12/20)
	-
	Sorafenib vs. Placebo [PFS(D): 167 vs. 84]

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

EMBRACA ^[313] NCT01945775	Breast cancer (Approved on 2018/10/16)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

^[314] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	-
	Temsirolimus vs. Ifn- α [OS(M): 10.9 vs. 7.3]

Vemurafenib (ZELBORAF)

Vemurafenib is an anti-cancer inhibitor which targets B-Raf. Vemurafenib is developed and marketed by Genentech under the trade name ZELBORAF.

- FDA Approval Summary of Vemurafenib (ZELBORAF)

VE-BASKET ^[315] NCT01524978	Erdheim-cheester disease (Approved on 2017/11/06)
	BRAF V600
	Vemurafenib [ORR(%): 54.5]
BRIM 3 ^[316] NCT01006980	Melanoma (Approved on 2011/08/17)
	BRAF V600E
	Vemurafenib vs. Dacarbazine [PFS(M): 5.3 vs. 1.6, OS(M): 13.6 vs. 10.3]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
BRAF	D594G	15	c.1781A>G	NM_004333	COSM467	32.5%	1478
MUC4	P485fs	15	c.14573del	NM_018406	-	48.1%	954
TP53	R248Q	7	c.743G>A	NM_000546	COSM10662	51.8%	2120

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
AMER1	R641*	2	c.1921C>T	NM_152424	COSM174910	37.7%	1150
AR	A597T	3	c.1789G>A	NM_000044	COSM1468978	28.3%	1697
ARID2	Q1462*	15	c.4384C>T	NM_152641	COSM2067865	31.8%	2444
ARID2	Q995*	15	c.2983C>T	NM_152641	COSM7318561	30.7%	2119
ATM	R2443Q	50	c.7328G>A	NM_000051	COSM20404	29.6%	778
BCOR	D1133Y	7	c.3397G>T	NM_001123385	-	37.8%	757
BRIP1	N196S	6	c.587A>G	NM_032043	COSM249456	71.6%	578
CREBBP	G447E	6	c.1340G>A	NM_004380	-	51.6%	1555
ERBB2	R143Q	3	c.428G>A	NM_004448	COSM1382867	20.4%	3164
FAT1	T1679I	10	c.5036C>T	NM_005245	-	72.1%	1095
FLT4	V528M	12	c.1582G>A	NM_182925	-	52.7%	431
INSR	K294R	3	c.881A>G	NM_000208	COSM2728577	47.6%	1146
LRP1B	E1327Q	25	c.3979G>C	NM_018557	-	35.0%	1303
NTRK1	R748W	17	c.2242C>T	NM_002529	-	54.5%	861
RAD51D	I311N	10	c.932T>A	NM_002878	-	73.4%	2154
RARA	P440R	9	c.1319C>G	NM_000964	-	21.6%	222
SMAD4	V128_Y131del	3	c.382_393del	NM_005359	-	46.8%	808



Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco® + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: May 2022
- Facility retrieved: 臺北榮總桃園分院
- H&E-stained section No.: S11120097A
- Collection site: Soft tissue
- Examined by: Dr. Chien-Ta Chiang
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 25%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 30%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco®+

DNA test

- Mean Depth: 1298x
- Target Base Coverage at 100x: 95%

RNA test

- Average unique RNA Start Sites per control GSP2: 49

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

ACT Onco[®] + Report

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師張筑芃 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



Sign Off

醫檢師張筑芃 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTX	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC1B1*
SLC1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION


Gene	Therapies	Possible effect
<i>CDKN2A</i>	Abemaciclib, Palbociclib, Ribociclib	sensitive
<i>PTEN</i>	Everolimus, Temsirolimus, Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>FBXW7</i>	Everolimus, Temsirolimus	sensitive
<i>NF2</i>	Everolimus, Temsirolimus	sensitive
<i>TSC2</i>	Everolimus, Temsirolimus	sensitive
<i>CHEK1</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>CHEK2</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>PALB2</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>RAD51C</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>SMAD4</i>	Cetuximab	resistant
<i>PTEN</i>	Erlotinib, Gefitinib, Panitumumab, Cetuximab, Trastuzumab	resistant
<i>FBXW7</i>	Gefitinib, Regorafenib	resistant

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Everolimus, Temsirolimus; 2: Sorafenib, Encorafenib, Vemurafenib; 3: Binimetinib

ACT Onco[®] + Report



1: Abemaciclib, Palbociclib, Ribociclib



1: Olaparib, Niraparib, Rucaparib, Talazoparib

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 25770121; 2015, J Biochem;157(5):271-84
Hepatocyte growth factor and Met in drug discovery.
2. PMID: 23867513; 2013, Cancer J;19(4):316-23
Targeting the hepatocyte growth factor/c-Met signaling pathway in renal cell carcinoma.
3. PMID: 15455388; 2005, Int J Cancer;113(4):678-82
C-Met overexpression in node-positive breast cancer identifies patients with poor clinical outcome independent of Her2/neu.
4. PMID: 9699182; 1998, Lung Cancer;20(1):1-16
Differential expression of Met/hepatocyte growth factor receptor in subtypes of non-small cell lung cancers.
5. PMID: 10454259; 1999, Cancer Lett;141(1-2):173-8
Progression-linked overexpression of c-Met in prostatic intraepithelial neoplasia and latent as well as clinical prostate cancers.
6. PMID: 24812413; 2014, Clin Cancer Res;20(13):3361-3
MET as a target in papillary renal cell carcinoma.
7. PMID: 24658158; 2014, Clin Cancer Res;20(13):3411-21
MET is a potential target across all papillary renal cell carcinomas: result from a large molecular study of pRCC with CGH array and matching gene expression array.
8. PMID: 18772890; 2008, Nature;455(7216):1061-8
Comprehensive genomic characterization defines human glioblastoma genes and core pathways.
9. PMID: 24222167; 2013, Anticancer Res;33(11):5179-86
A survey of c-MET expression and amplification in 287 patients with hepatocellular carcinoma.
10. PMID: 9759658; 1998, Lab Invest;78(9):1143-53
Amplification of c-myc, K-sam, and c-met in gastric cancers: detection by fluorescence in situ hybridization.
11. PMID: 29718398; 2018, J Neuropathol Exp Neurol;77(6):437-442
Targetable Gene Fusions Associate With the IDH Wild-Type Astrocytic Lineage in Adult Gliomas.
12. PMID: 26689674; 2015, Sci Rep;5():18657
Novel kinase fusion transcripts found in endometrial cancer.
13. PMID: 25135958; 2014, Genome Res;24(11):1765-73
RNA-seq of 272 gliomas revealed a novel, recurrent PTPRZ1-MET fusion transcript in secondary glioblastomas.
14. PMID: 24240688; 2014, Oncogene;33(47):5434-41
Novel fusion transcripts in human gastric cancer revealed by transcriptome analysis.
15. PMID: 15520807; 2004, Nat Rev Mol Cell Biol;5(11):875-85
The RAF proteins take centre stage.
16. PMID: 24737949; 2014, J Carcinog;13():1
BRAF and beyond: Tailoring strategies for the individual melanoma patient.
17. PMID: 12068308; 2002, Nature;417(6892):949-54
Mutations of the BRAF gene in human cancer.
18. PMID: 24071849; 2013, Nat Genet;45(10):1113-20
The Cancer Genome Atlas Pan-Cancer analysis project.
19. PMID: 20179705; 2010, Nature;464(7287):427-30
RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF.

ACT Onco[®] + Report

20. PMID: 21388974; 2011, Mol Cancer Ther;10(3):385-94
BRAFV600E: implications for carcinogenesis and molecular therapy.
21. PMID: 18794803; 2009, Oncogene;28(1):85-94
CRF inhibition induces apoptosis in melanoma cells with non-V600E BRAF mutations.
22. PMID: 28783719; 2017, Nature;548(7666):234-238
Tumours with class 3 BRAF mutants are sensitive to the inhibition of activated RAS.
23. PMID: 31924734; 2020, Clin Cancer Res;26(8):1812-1819
Trametinib Activity in Patients with Solid Tumors and Lymphomas Harboring BRAF Non-V600 Mutations or Fusions: Results from NCI-MATCH (EAY131).
24. PMID: 29076950; 2017, Melanoma Res;27(6):585-590
Vemurafenib treatment for patients with locally advanced, unresectable stage IIIC or metastatic melanoma and activating exon 15 BRAF mutations other than V600E.
25. PMID: 27124015; 2016, Medicine (Baltimore);95(17):e3225
Low Expression of Mucin-4 Predicts Poor Prognosis in Patients With Clear-Cell Renal Cell Carcinoma.
26. PMID: 20001800; 2009, Future Oncol;5(10):1631-40
Muc4/MUC4 functions and regulation in cancer.
27. PMID: 12115478; 2002, Int J Cancer;99(6):783-91
Rat Muc4 (sialomucin complex) reduces binding of anti-ErbB2 antibodies to tumor cell surfaces, a potential mechanism for herceptin resistance.
28. PMID: 15695389; 2005, Cancer Res;65(2):473-82
Decreased accessibility and lack of activation of ErbB2 in JIMT-1, a herceptin-resistant, MUC4-expressing breast cancer cell line.
29. PMID: 12102554; 2002, Prog Nucleic Acid Res Mol Biol;71():149-85
Muc4/sialomucin complex, the intramembrane ErbB2 ligand, in cancer and epithelia: to protect and to survive.
30. PMID: 16891313; 2006, J Biol Chem;281(39):29411-20
Membrane mucin Muc4 induces density-dependent changes in ERK activation in mammary epithelial and tumor cells: role in reversal of contact inhibition.
31. PMID: 16049287; 2005, J Clin Pathol;58(8):845-52
MUC4 expression is a novel prognostic factor in patients with invasive ductal carcinoma of the pancreas.
32. PMID: 28460571; 2017, Expert Opin Ther Targets;21(7):657-669
MUC4 mucin- a therapeutic target for pancreatic ductal adenocarcinoma.
33. PMID: 18665193; 2008, Br J Cancer;99(3):520-6
MUC4 activates HER2 signalling and enhances the motility of human ovarian cancer cells.
34. PMID: 23408941; 2013, PLoS One;8(2):e54455
MUC4 overexpression augments cell migration and metastasis through EGFR family proteins in triple negative breast cancer cells.
35. PMID: 20564074; 2010, Cancer;116(15):3577-86
Prognostic value of mucin 4 expression in colorectal adenocarcinomas.
36. PMID: 28270046; 2017, Scand J Gastroenterol;52(5):595-600
Low MUC4 expression is associated with survival benefit in patients with resectable pancreatic cancer receiving adjuvant gemcitabine.
37. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
38. PMID: 21125671; 2011, J Pathol;223(2):137-46

ACT Onco[®] + Report

Haplo-insufficiency: a driving force in cancer.

39. PMID: 26703669; 2015, Int J Environ Res Public Health;13(1):ijerph13010022
Hot Spot Mutation in TP53 (R248Q) Causes Oncogenic Gain-of-Function Phenotypes in a Breast Cancer Cell Line Derived from an African American patient.
40. PMID: 25697482; 2015, Cancer Lett;360(2):269-79
Mutant p53 (p53-R248Q) functions as an oncogene in promoting endometrial cancer by up-regulating REGy.
41. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months.
42. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
43. PMID: 25669829; 2015, Ann Oncol;26(5):1012-8
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
44. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
45. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
46. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
47. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
48. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
49. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
50. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
51. PMID: 25385265; 2015, Int J Oncol;46(2):607-18
TP53 oncomorphic mutations predict resistance to platinum and taxane based standard chemotherapy in patients diagnosed with advanced serous ovarian carcinoma.
52. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
53. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
54. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
55. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.
56. PMID: 7550353; 1995, Nat Genet;11(2):210-2

ACT Onco[®] + Report

Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.

57. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
58. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
59. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
60. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
61. PMID: 27217383; 2016, Cancer Discov;6(7):740-53
Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.
62. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
63. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
64. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
65. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.
66. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
67. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
68. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
69. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
70. PMID: 12781359; 2003, Cancer Cell;3(5):421-9
Chk1 and Chk2 kinases in checkpoint control and cancer.
71. PMID: 15261141; 2004, Cancer Cell;6(1):45-59
Chk1 is haploinsufficient for multiple functions critical to tumor suppression.
72. PMID: 15539958; 2005, Cell Cycle;4(1):131-9
Chk1 is essential for tumor cell viability following activation of the replication checkpoint.
73. PMID: 15459660; 2004, Nat Rev Mol Cell Biol;5(10):792-804
Checking on DNA damage in S phase.
74. PMID: 22585575; 2012, J Clin Invest;122(6):2165-75
CHK1 targets spleen tyrosine kinase (L) for proteolysis in hepatocellular carcinoma.

ACT Onco[®] + Report

75. PMID: 17638866; 2007, Cancer Res;67(14):6574-81
The E2F-regulated gene Chk1 is highly expressed in triple-negative estrogen receptor /progesterone receptor /HER-2 breast carcinomas.
76. PMID: 17848589; 2007, Mol Cell Proteomics;6(12):2150-64
A proteomics analysis of cell signaling alterations in colorectal cancer.
77. PMID: 24418519; 2014, J Surg Res;187(1):6-13
Checkpoint kinase 1 protein expression indicates sensitization to therapy by checkpoint kinase 1 inhibition in non-small cell lung cancer.
78. PMID: 15297395; 2004, Clin Cancer Res;10(15):4944-58
Global gene expression profile of nasopharyngeal carcinoma by laser capture microdissection and complementary DNA microarrays.
79. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
80. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
81. PMID: 21458083; 2011, Trends Pharmacol Sci;32(5):308-16
Anticancer therapy with checkpoint inhibitors: what, where and when?
82. PMID: 21088254; 2011, Clin Cancer Res;17(3):401-5
Tumor suppressor CHK2: regulator of DNA damage response and mediator of chromosomal stability.
83. PMID: 23296741; 2013, Fam Cancer;12(3):473-8
The risk of gastric cancer in carriers of CHEK2 mutations.
84. PMID: 24713400; 2014, Hered Cancer Clin Pract;12(1):10
A risk of breast cancer in women - carriers of constitutional CHEK2 gene mutations, originating from the North - Central Poland.
85. PMID: 25583358; 2015, Int J Cancer;137(3):548-52
CHEK2 mutations and the risk of papillary thyroid cancer.
86. PMID: 12052256; 2002, Breast Cancer Res;4(3):R4
Mutation analysis of the CHK2 gene in breast carcinoma and other cancers.
87. PMID: 15125777; 2004, Mol Cancer;3():14
CHK2 kinase expression is down-regulated due to promoter methylation in non-small cell lung cancer.
88. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
89. PMID: 26510020; 2015, N Engl J Med;373(18):1697-708
DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer.
90. PMID: 11252954; 2001, Nat Rev Mol Cell Biol;2(2):127-37
Untangling the ErbB signalling network.
91. PMID: 21204711; 2011, Arch Pathol Lab Med;135(1):55-62
HER2: biology, detection, and clinical implications.
92. PMID: 25480824; 2015, Oncologist;20(1):7-12
Oncogenic alterations in ERBB2/HER2 represent potential therapeutic targets across tumors from diverse anatomic sites of origin.
93. PMID: 22325357; 2012, Lung Cancer;76(1):123-7
Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu.

ACT Onco[®] + Report

94. PMID: 23610105; 2013, J Clin Oncol;31(16):1997-2003
Lung cancer that harbors an HER2 mutation: epidemiologic characteristics and therapeutic perspectives.
95. PMID: 26964772; 2016, J Thorac Oncol;11(6):918-23
Pulse Afatinib for ERBB2 Exon 20 Insertion-Mutated Lung Adenocarcinomas.
96. PMID: 26545934; 2016, Cancer Sci;107(1):45-52
Antitumor effect of afatinib, as a human epidermal growth factor receptor 2-targeted therapy, in lung cancers harboring HER2 oncogene alterations.
97. PMID: 32162827; 2020, Oncologist;25(3):e545-e554
Mutation Variants and Co-Mutations as Genomic Modifiers of Response to Afatinib in HER2-Mutant Lung Adenocarcinoma.
98. PMID: 25899785; 2015, Ann Oncol;26(7):1421-7
Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors.
99. PMID: 16843263; 2006, Cancer Cell;10(1):25-38
HER2 kinase domain mutation results in constitutive phosphorylation and activation of HER2 and EGFR and resistance to EGFR tyrosine kinase inhibitors.
100. PMID: 16818618; 2006, Cancer Res;66(13):6487-91
Non-small-cell lung cancer and Ba/F3 transformed cells harboring the ERBB2 G776insV_G/C mutation are sensitive to the dual-specific epidermal growth factor receptor and ERBB2 inhibitor HKI-272.
101. PMID: 31588020; 2019, Cancer Cell;36(4):444-457.e7
Pan-Cancer Landscape and Analysis of ERBB2 Mutations Identifies Pozotinib as a Clinically Active Inhibitor and Enhancer of T-DM1 Activity.
102. PMID: 16775247; 2006, N Engl J Med;354(24):2619-21
HER2 mutation and response to trastuzumab therapy in non-small-cell lung cancer.
103. PMID: 23328556; 2013, J Thorac Oncol;8(2):e19-20
Non-small-cell lung cancer with HER2 exon 20 mutation: regression with dual HER2 inhibition and anti-VEGF combination treatment.
104. PMID: 26598547; 2016, Ann Oncol;27(2):281-6
Lung cancer patients with HER2 mutations treated with chemotherapy and HER2-targeted drugs: results from the European EUHER2 cohort.
105. PMID: 25789838; 2015, J Thorac Oncol;10(4):e16-7
Rapid response to trastuzumab emtansine in a patient with HER2-driven lung cancer.
106. PMID: 29989854; 2018, J Clin Oncol;36(24):2532-2537
Ado-Trastuzumab Emtansine for Patients With HER2-Mutant Lung Cancers: Results From a Phase II Basket Trial.
107. PMID: 11248153; 2001, N Engl J Med;344(11):783-92
Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2.
108. PMID: 25332249; 2014, J Clin Oncol;32(33):3744-52
Trastuzumab plus adjuvant chemotherapy for human epidermal growth factor receptor 2-positive breast cancer: planned joint analysis of overall survival from NSABP B-31 and NCCTG N9831.
109. PMID: 28581356; 2017, N Engl J Med;377(2):122-131
Adjuvant Pertuzumab and Trastuzumab in Early HER2-Positive Breast Cancer.
110. PMID: 30516102; 2019, N Engl J Med;380(7):617-628
Trastuzumab Emtansine for Residual Invasive HER2-Positive Breast Cancer.
111. PMID: 31825192; 2020, N Engl J Med;382(7):610-621
Trastuzumab Deruxtecan in Previously Treated HER2-Positive Breast Cancer.
112. PMID: 18188694; 2008, Breast Cancer Res Treat;112(3):533-43

ACT Onco[®] + Report

A phase III randomized comparison of lapatinib plus capecitabine versus capecitabine alone in women with advanced breast cancer that has progressed on trastuzumab: updated efficacy and biomarker analyses.

113. PMID: 26874901; 2016, Lancet Oncol;17(3):367-77
Neratinib after trastuzumab-based adjuvant therapy in patients with HER2-positive breast cancer (ExteNET): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial.
114. PMID: 31140297; 2019, Future Oncol;15(21):2489-2501
Neratinib after trastuzumab-based adjuvant therapy in patients from Asia with early stage HER2-positive breast cancer.
115. PMID: 29146401; 2017, Lancet Oncol;18(12):1688-1700
Neratinib after trastuzumab-based adjuvant therapy in HER2-positive breast cancer (ExteNET): 5-year analysis of a randomised, double-blind, placebo-controlled, phase 3 trial.
116. PMID: 31825569; 2020, N Engl J Med;382(7):597-609
Tucatinib, Trastuzumab, and Capecitabine for HER2-Positive Metastatic Breast Cancer.
117. PMID: 25268372; 2014, Br J Cancer;111(9):1750-6
Afatinib demonstrates remarkable activity against HER2-amplified uterine serous endometrial cancer in vitro and in vivo.
118. PMID: 28520362; 2012, ;():
Trastuzumab (Herceptin) Therapy and ERBB2 (HER2) Genotype
119. PMID: 23020162; 2012, N Engl J Med;367(19):1783-91
Trastuzumab emtansine for HER2-positive advanced breast cancer.
120. PMID: 24793816; 2014, Lancet Oncol;15(7):689-99
Trastuzumab emtansine versus treatment of physician's choice for pretreated HER2-positive advanced breast cancer (TH3RESA): a randomised, open-label, phase 3 trial.
121. PMID: 20124187; 2010, J Clin Oncol;28(7):1124-30
Randomized study of Lapatinib alone or in combination with trastuzumab in women with ErbB2-positive, trastuzumab-refractory metastatic breast cancer.
122. PMID: 22689807; 2012, J Clin Oncol;30(21):2585-92
Overall survival benefit with lapatinib in combination with trastuzumab for patients with human epidermal growth factor receptor 2-positive metastatic breast cancer: final results from the EGF104900 Study.
123. PMID: 20142587; 2010, J Clin Oncol;28(8):1301-7
Neratinib, an irreversible ErbB receptor tyrosine kinase inhibitor, in patients with advanced ErbB2-positive breast cancer.
124. PMID: 24323026; 2014, J Clin Oncol;32(2):68-75
Phase I study of neratinib in combination with temsirolimus in patients with human epidermal growth factor receptor 2-dependent and other solid tumors.
125. PMID: 26358790; 2015, J Natl Compr Canc Netw;13(9):1061-4
HER2-Mutated Breast Cancer Responds to Treatment With Single-Agent Neratinib, a Second-Generation HER2/EGFR Tyrosine Kinase Inhibitor.
126. PMID: 31504139; 2019, Ann Oncol;30(11):1821-1830
Ado-trastuzumab emtansine (T-DM1) in patients with HER2-amplified tumors excluding breast and gastric/gastroesophageal junction (GEJ) adenocarcinomas: results from the NCI-MATCH trial (EAY131) subprotocol Q.
127. PMID: 22149875; 2012, N Engl J Med;366(2):109-19
Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer.
128. PMID: 23602601; 2013, Lancet Oncol;14(6):461-71
Pertuzumab, trastuzumab, and docetaxel for HER2-positive metastatic breast cancer (CLEOPATRA study): overall survival results from a randomised, double-blind, placebo-controlled, phase 3 study.

ACT Onco[®] + Report

129. PMID: 20728210; 2010, Lancet;376(9742):687-97
Trastuzumab in combination with chemotherapy versus chemotherapy alone for treatment of HER2-positive advanced gastric or gastro-oesophageal junction cancer (ToGA): a phase 3, open-label, randomised controlled trial.
130. PMID: 27108243; 2016, Lancet Oncol;17(6):738-746
Dual-targeted therapy with trastuzumab and lapatinib in treatment-refractory, KRAS codon 12/13 wild-type, HER2-positive metastatic colorectal cancer (HERACLES): a proof-of-concept, multicentre, open-label, phase 2 trial.
131. PMID: 30857956; 2019, Lancet Oncol;20(4):518-530
Pertuzumab plus trastuzumab for HER2-amplified metastatic colorectal cancer (MyPathway): an updated report from a multicentre, open-label, phase 2a, multiple basket study.
132. PMID: 28006087; 2017, Head Neck;39(3):E40-E44
Significant and durable clinical benefit from trastuzumab in 2 patients with HER2-amplified salivary gland cancer and a review of the literature.
133. PMID: 32067683; 2020, Ann Oncol;31(3):412-421
Targeted therapy for advanced salivary gland carcinoma based on molecular profiling: results from MyPathway, a phase IIa multiple basket study.
134. PMID: 30452336; 2019, J Clin Oncol;37(2):125-134
Phase II Trial of Trastuzumab and Docetaxel in Patients With Human Epidermal Growth Factor Receptor 2-Positive Salivary Duct Carcinoma.
135. PMID: 29584549; 2018, J Clin Oncol;36(20):2044-2051
Randomized Phase II Trial of Carboplatin-Paclitaxel Versus Carboplatin-Paclitaxel-Trastuzumab in Uterine Serous Carcinomas That Overexpress Human Epidermal Growth Factor Receptor 2/neu.
136. PMID: 12907212; 2003, Oral Oncol;39(7):724-7
Herceptin in patients with advanced or metastatic salivary gland carcinomas. A phase II study.
137. PMID: 17761983; 2007, J Clin Oncol;25(25):3978-84
Phase II study of lapatinib in recurrent or metastatic epidermal growth factor receptor and/or erbB2 expressing adenoid cystic carcinoma and non adenoid cystic carcinoma malignant tumors of the salivary glands.
138. PMID: 17972317; 2008, Head Neck;30(5):680-3
Successful treatment of pulmonary metastatic salivary ductal carcinoma with trastuzumab-based therapy.
139. PMID: 12525520; 2003, J Clin Oncol;21(2):283-90
Evaluation of monoclonal humanized anti-HER2 antibody, trastuzumab, in patients with recurrent or refractory ovarian or primary peritoneal carcinoma with overexpression of HER2: a phase II trial of the Gynecologic Oncology Group.
140. PMID: 16896006; 2006, J Clin Oncol;24(26):4324-32
Clinical activity of pertuzumab (rhuMab 2C4), a HER dimerization inhibitor, in advanced ovarian cancer: potential predictive relationship with tumor HER2 activation status.
141. PMID: 22956644; 2012, Cancer Discov;2(10):922-33
HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation.
142. PMID: 29530932; 2018, Clin Cancer Res;24(13):3108-3118
Concurrent Alterations in EGFR-Mutant Lung Cancers Associated with Resistance to EGFR Kinase Inhibitors and Characterization of MTOR as a Mediator of Resistance.
143. PMID: 16683005; 2006, Nat Clin Pract Oncol;3(5):269-80
Mechanisms of disease: understanding resistance to HER2-targeted therapy in human breast cancer.
144. PMID: 22471661; 2012, Crit Rev Oncog;17(1):1-16
Intrinsic and acquired resistance to HER2-targeted therapies in HER2 gene-amplified breast cancer: mechanisms and clinical implications.
145. PMID: 15324695; 2004, Cancer Cell;6(2):117-27
PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients.

ACT Onco[®] + Report

146. PMID: 25559818; 2015, J Clin Oncol;33(12):1334-9
PIK3CA mutations are associated with decreased benefit to neoadjuvant human epidermal growth factor receptor 2-targeted therapies in breast cancer.
147. PMID: 20813970; 2010, Am J Pathol;177(4):1647-56
PTEN, PIK3CA, p-AKT, and p-p70S6K status: association with trastuzumab response and survival in patients with HER2-positive metastatic breast cancer.
148. PMID: 21594665; 2011, Breast Cancer Res Treat;128(2):447-56
Evaluation of the association of PIK3CA mutations and PTEN loss with efficacy of trastuzumab therapy in metastatic breast cancer.
149. PMID: 17936563; 2007, Cancer Cell;12(4):395-402
A functional genetic approach identifies the PI3K pathway as a major determinant of trastuzumab resistance in breast cancer.
150. PMID: 23739063; 2013, J Natl Cancer Inst;105(13):960-7
Somatic mutation profiling and associations with prognosis and trastuzumab benefit in early breast cancer.
151. PMID: 23918797; 2013, Cancer Res;73(19):6013-23
Combination of antibody that inhibits ligand-independent HER3 dimerization and a p110 α inhibitor potently blocks PI3K signaling and growth of HER2+ breast cancers.
152. PMID: 20007781; 2009, Proc Natl Acad Sci U S A;106(52):22299-304
Specific apoptosis induction by the dual PI3K/mTor inhibitor NVP-BEZ235 in HER2 amplified and PIK3CA mutant breast cancer cells.
153. PMID: 3798106; 1987, Science;235(4785):177-82
Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene.
154. PMID: 9469329; 1998, J Clin Oncol;16(2):462-9
Prognostic and predictive value of c-erbB-2 overexpression in primary breast cancer, alone and in combination with other prognostic markers.
155. PMID: 9552035; 1998, J Clin Oncol;16(4):1340-9
neu/erbB-2 amplification identifies a poor-prognosis group of women with node-negative breast cancer. Toronto Breast Cancer Study Group.
156. PMID: 21900593; 2011, Sci Transl Med;3(99):99ra86
Activation of ERBB2 signaling causes resistance to the EGFR-directed therapeutic antibody cetuximab.
157. PMID: 24841429; 2014, Oncol Rep;32(1):3-15
Mechanisms associated with resistance to tamoxifen in estrogen receptor-positive breast cancer (review).
158. PMID: 10098763; 1999, Br J Cancer;79(7-8):1220-6
Overexpression of c-erbB2 is an independent marker of resistance to endocrine therapy in advanced breast cancer.
159. PMID: 12618500; 2003, J Natl Cancer Inst;95(5):353-61
Role of the estrogen receptor coactivator AIB1 (SRC-3) and HER-2/neu in tamoxifen resistance in breast cancer.
160. PMID: 16710036; 2006, J Clin Oncol;24(15):2376-85
HER-2 is an independent prognostic factor in endometrial cancer: association with outcome in a large cohort of surgically staged patients.
161. PMID: 19088718; 2009, Br J Cancer;100(1):89-95
HER2 gene amplification and EGFR expression in a large cohort of surgically staged patients with nonendometrioid (type II) endometrial cancer.
162. PMID: 15284264; 2004, J Clin Oncol;22(15):3126-32
Her-2/neu overexpression and amplification in uterine papillary serous carcinoma.
163. PMID: 15498494; 2004, Curr Biol;14(20):1852-7
A nucleolar isoform of the Fbw7 ubiquitin ligase regulates c-Myc and cell size.
164. PMID: 15103331; 2004, EMBO J;23(10):2116-25

ACT Onco[®] + Report

Phosphorylation-dependent degradation of c-Myc is mediated by the F-box protein Fbw7.

165. PMID: 16023596; 2005, Cancer Cell;8(1):25-33
The v-Jun point mutation allows c-Jun to escape GSK3-dependent recognition and destruction by the Fbw7 ubiquitin ligase.
166. PMID: 11533444; 2001, Science;294(5540):173-7
Phosphorylation-dependent ubiquitination of cyclin E by the SCFFbw7 ubiquitin ligase.
167. PMID: 11461910; 2001, J Biol Chem;276(38):35847-53
The Notch intracellular domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog.
168. PMID: 11425854; 2001, J Biol Chem;276(37):34371-8
Functional interaction between SEL-10, an F-box protein, and the nuclear form of activated Notch1 receptor.
169. PMID: 16863506; 2006, Cancer Sci;97(8):729-36
Fbxw7 contributes to tumor suppression by targeting multiple proteins for ubiquitin-dependent degradation.
170. PMID: 18787170; 2008, Science;321(5895):1499-502
FBXW7 targets mTOR for degradation and cooperates with PTEN in tumor suppression.
171. PMID: 20484041; 2010, Cancer Res;70(11):4728-38
The Fbw7 tumor suppressor targets KLF5 for ubiquitin-mediated degradation and suppresses breast cell proliferation.
172. PMID: 21368833; 2011, Nature;471(7336):104-9
SCF(FBW7) regulates cellular apoptosis by targeting MCL1 for ubiquitylation and destruction.
173. PMID: 18094723; 2008, Nat Rev Cancer;8(2):83-93
FBW7 ubiquitin ligase: a tumour suppressor at the crossroads of cell division, growth and differentiation.
174. PMID: 23032637; 2012, Cancer Inform;11():157-71
Haploinsufficiency of Tumor Suppressor Genes is Driven by the Cumulative Effect of microRNAs, microRNA Binding Site Polymorphisms and microRNA Polymorphisms: An In silico Approach.
175. PMID: 24586741; 2014, PLoS One;9(2):e89388
FBXW7 mutations in patients with advanced cancers: clinical and molecular characteristics and outcomes with mTOR inhibitors.
176. PMID: 24360397; 2014, Lung Cancer;83(2):300-1
Temozolomide therapy in a patient with lung adenocarcinoma harboring an FBXW7 mutation.
177. PMID: 27399335; 2017, Oncogene;36(6):787-796
FBW7 mutations mediate resistance of colorectal cancer to targeted therapies by blocking Mcl-1 degradation.
178. PMID: 25860929; 2015, Oncotarget;6(11):9240-56
FBXW7-mutated colorectal cancer cells exhibit aberrant expression of phosphorylated-p53 at Serine-15.
179. PMID: 29633504; 2018, Mol Oncol;12(6):883-895
FBXW7 deletion contributes to lung tumor development and confers resistance to gefitinib therapy.
180. PMID: 28522751; 2017, Cancer Res;77(13):3527-3539
Targeting FBW7 as a Strategy to Overcome Resistance to Targeted Therapy in Non-Small Cell Lung Cancer.
181. PMID: 24884509; 2014, Mol Cancer;13():110
Fbxw7 is an independent prognostic marker and induces apoptosis and growth arrest by regulating YAP abundance in hepatocellular carcinoma.
182. PMID: 25893302; 2016, Oncogene;35(5):537-48
Role of Merlin/NF2 inactivation in tumor biology.
183. PMID: 19451229; 2009, Mol Cell Biol;29(15):4235-49
Loss of the tumor suppressor gene NF2, encoding merlin, constitutively activates integrin-dependent mTORC1 signaling.

ACT Onco[®] + Report

184. PMID: 19451225; 2009, Mol Cell Biol;29(15):4250-61
NF2/merlin is a novel negative regulator of mTOR complex 1, and activation of mTORC1 is associated with meningioma and schwannoma growth.
185. PMID: 17655741; 2007, Brain Pathol;17(4):371-6
Role of NF2 haploinsufficiency in NF2-associated polyneuropathy.
186. PMID: 19545378; 2009, Orphanet J Rare Dis;4():16
Neurofibromatosis type 2 (NF2): a clinical and molecular review.
187. PMID: 21642991; 2011, Nat Genet;43(7):668-72
The nuclear deubiquitinase BAP1 is commonly inactivated by somatic mutations and 3p21.1 losses in malignant pleural mesothelioma.
188. PMID: 24393766; 2014, Oncotarget;5(1):67-77
NF2/merlin in hereditary neurofibromatosis 2 versus cancer: biologic mechanisms and clinical associations.
189. PMID: 27091708; 2016, J Clin Oncol;34(18):2115-24
Molecular Alterations and Everolimus Efficacy in Human Epidermal Growth Factor Receptor 2-Overexpressing Metastatic Breast Cancers: Combined Exploratory Biomarker Analysis From BOLERO-1 and BOLERO-3.
190. PMID: 26503204; 2016, J Clin Oncol;34(5):419-26
Correlative Analysis of Genetic Alterations and Everolimus Benefit in Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: Results From BOLERO-2.
191. PMID: 24833916; 2014, Breast Cancer (Dove Med Press);6():43-57
Use of mTOR inhibitors in the treatment of breast cancer: an evaluation of factors that influence patient outcomes.
192. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
193. PMID: 22923433; 2012, Science;338(6104):221
Genome sequencing identifies a basis for everolimus sensitivity.
194. PMID: 25630452; 2015, Eur Urol;67(6):1195-1196
Exceptional Response on Addition of Everolimus to Taxane in Urothelial Carcinoma Bearing an NF2 Mutation.
195. PMID: 26359368; 2015, Cancer Discov;5(11):1178-93
NF2 Loss Promotes Oncogenic RAS-Induced Thyroid Cancers via YAP-Dependent Transactivation of RAS Proteins and Sensitizes Them to MEK Inhibition.
196. PMID: 24813888; 2014, Cell Rep;7(4):999-1008
Acquired resistance of EGFR-mutant lung adenocarcinomas to afatinib plus cetuximab is associated with activation of mTORC1.
197. PMID: 16793542; 2006, Mol Cell;22(6):719-29
Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2.
198. PMID: 23325218; 2013, Nature;493(7432):356-63
Fanconi anaemia and the repair of Watson and Crick DNA crosslinks.
199. PMID: 15905196; 2005, Carcinogenesis;26(10):1731-40
The Fanconi anemia group A protein modulates homologous repair of DNA double-strand breaks in mammalian cells.
200. PMID: 27037238; 2016, EMBO J;35(9):909-23
Interplay between Fanconi anemia and homologous recombination pathways in genome integrity.
201. PMID: 24153426; 2013, Nat Commun;4():2578
Heterozygous mutations in PALB2 cause DNA replication and damage response defects.
202. PMID: 20858716; 2010, Cancer Res;70(19):7353-9

ACT Onco[®] + Report

PALB2/FANCN: recombining cancer and Fanconi anemia.

203. PMID: 25754594; 2015, Hum Mutat;36(5):562-8
Loss-of-Function FANCL Mutations Associate with Severe Fanconi Anemia Overlapping the VACTERL Association.
204. PMID: 28678401; 2017, Cancer;123(20):3943-3954
Assessing the spectrum of germline variation in Fanconi anemia genes among patients with head and neck carcinoma before age 50.
205. PMID: 21135251; 2011, Mol Cancer Ther;10(1):3-8
Personalizing cancer treatment in the age of global genomic analyses: PALB2 gene mutations and the response to DNA damaging agents in pancreatic cancer.
206. PMID: 17218262; 2007, Cell;128(1):157-70
Essential role for nuclear PTEN in maintaining chromosomal integrity.
207. PMID: 18794879; 2008, Oncogene;27(41):5443-53
PTEN: a new guardian of the genome.
208. PMID: 18767981; 2009, Annu Rev Pathol;4():127-50
PTEN and the PI3-kinase pathway in cancer.
209. PMID: 11553783; 2001, Proc Natl Acad Sci U S A;98(20):11563-8
Haploinsufficiency of the Pten tumor suppressor gene promotes prostate cancer progression.
210. PMID: 20400965; 2010, Nat Genet;42(5):454-8
Subtle variations in Pten dose determine cancer susceptibility.
211. PMID: 9467011; 1998, Hum Mol Genet;7(3):507-15
Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation.
212. PMID: 24136893; 2013, J Natl Cancer Inst;105(21):1607-16
Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria.
213. PMID: 21430697; 2011, Nat Rev Cancer;11(4):289-301
PTEN loss in the continuum of common cancers, rare syndromes and mouse models.
214. PMID: 18455982; 2008, Cell;133(3):403-14
Tenets of PTEN tumor suppression.
215. PMID: 9393738; 1997, Cancer Res;57(23):5221-5
MMAC1/PTEN mutations in primary tumor specimens and tumor cell lines.
216. PMID: 9829719; 1998, Clin Cancer Res;4(11):2577-83
Loss of heterozygosity and mutational analysis of the PTEN/MMAC1 gene in synchronous endometrial and ovarian carcinomas.
217. PMID: 9582022; 1998, Oncogene;16(13):1743-8
Analysis of PTEN and the 10q23 region in primary prostate carcinomas.
218. PMID: 9671321; 1998, Oncogene;17(1):123-7
Allelic loss of chromosome 10q23 is associated with tumor progression in breast carcinomas.
219. PMID: 11504908; 2001, Proc Natl Acad Sci U S A;98(18):10314-9
Enhanced sensitivity of PTEN-deficient tumors to inhibition of FRAP/mTOR.
220. PMID: 23714559; 2013, Am Soc Clin Oncol Educ Book;():
Targeting the PI3K/AKT/mTOR pathway: biomarkers of success and tribulation.
221. PMID: 20231295; 2010, J Biol Chem;285(20):14980-9
Phosphoinositide 3-kinase pathway activation in phosphate and tensin homolog (PTEN)-deficient prostate cancer cells is independent of

ACT Onco[®] + Report

receptor tyrosine kinases and mediated by the p110beta and p110delta catalytic subunits.

222. PMID: 23287563; 2013, Clin Cancer Res;19(7):1760-72
Targeting activated Akt with GDC-0068, a novel selective Akt inhibitor that is efficacious in multiple tumor models.
223. PMID: 17047067; 2006, Cancer Res;66(20):10040-7
Inhibition of mammalian target of rapamycin or apoptotic pathway induces autophagy and radiosensitizes PTEN null prostate cancer cells.
224. PMID: 22422409; 2012, Clin Cancer Res;18(6):1777-89
PIK3CA/PTEN mutations and Akt activation as markers of sensitivity to allosteric mTOR inhibitors.
225. PMID: 22662154; 2012, PLoS One;7(5):e37431
Genotype-dependent efficacy of a dual PI3K/mTOR inhibitor, NVP-BEZ235, and an mTOR inhibitor, RAD001, in endometrial carcinomas.
226. PMID: 23136191; 2012, Clin Cancer Res;18(24):6771-83
Phosphoinositide 3-kinase (PI3K) pathway alterations are associated with histologic subtypes and are predictive of sensitivity to PI3K inhibitors in lung cancer preclinical models.
227. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
228. PMID: 23582881; 2013, Eur Urol;64(1):150-8
Phase 2 trial of single-agent everolimus in chemotherapy-naïve patients with castration-resistant prostate cancer (SAKK 08/08).
229. PMID: 21135276; 2011, J Clin Oncol;29(2):166-73
Loss of phosphatase and tensin homolog or phosphoinositide-3 kinase activation and response to trastuzumab or lapatinib in human epidermal growth factor receptor 2-overexpressing locally advanced breast cancers.
230. PMID: 18700047; 2008, BMC Cancer;8():234
Potential value of PTEN in predicting cetuximab response in colorectal cancer: an exploratory study.
231. PMID: 17940504; 2007, Br J Cancer;97(8):1139-45
PTEN loss of expression predicts cetuximab efficacy in metastatic colorectal cancer patients.
232. PMID: 19398573; 2009, J Clin Oncol;27(16):2622-9
PTEN expression and KRAS mutations on primary tumors and metastases in the prediction of benefit from cetuximab plus irinotecan for patients with metastatic colorectal cancer.
233. PMID: 19953097; 2010, Br J Cancer;102(1):162-4
PTEN status in advanced colorectal cancer treated with cetuximab.
234. PMID: 27605871; 2016, World J Gastroenterol;22(28):6345-61
Mechanisms of resistance to anti-epidermal growth factor receptor inhibitors in metastatic colorectal cancer.
235. PMID: 24666267; 2014, Acta Oncol;53(7):852-64
The predictive value of KRAS, NRAS, BRAF, PIK3CA and PTEN for anti-EGFR treatment in metastatic colorectal cancer: A systematic review and meta-analysis.
236. PMID: 19351834; 2009, Cancer Res;69(8):3256-61
PTEN loss contributes to erlotinib resistance in EGFR-mutant lung cancer by activation of Akt and EGFR.
237. PMID: 23133538; 2012, PLoS One;7(10):e48004
Modeling of tumor progression in NSCLC and intrinsic resistance to TKI in loss of PTEN expression.
238. PMID: 23592446; 2013, J Cell Biochem;114(6):1248-56
mTOR inhibitors radiosensitize PTEN-deficient non-small-cell lung cancer cells harboring an EGFR activating mutation by inducing autophagy.
239. PMID: 26645196; 2016, Cancer Discov;6(2):202-16
Loss of PTEN Promotes Resistance to T Cell-Mediated Immunotherapy.

ACT Onco[®] + Report

240. PMID: 28228279; 2017, Immunity;46(2):197-204
Loss of PTEN Is Associated with Resistance to Anti-PD-1 Checkpoint Blockade Therapy in Metastatic Uterine Leiomyosarcoma.
241. PMID: 30150660; 2018, Nat Genet;50(9):1271-1281
Genomic correlates of response to immune checkpoint blockade in microsatellite-stable solid tumors.
242. PMID: 21468130; 2011, Nat Rev Clin Oncol;8(5):302-6
Treatment with olaparib in a patient with PTEN-deficient endometrioid endometrial cancer.
243. PMID: 20400964; 2010, Nat Genet;42(5):410-4
Germline mutations in breast and ovarian cancer pedigrees establish RAD51C as a human cancer susceptibility gene.
244. PMID: 21990120; 2012, Hum Mutat;33(1):95-9
Analysis of RAD51C germline mutations in high-risk breast and ovarian cancer families and ovarian cancer patients.
245. PMID: 21616938; 2011, Hum Mol Genet;20(16):3278-88
RAD51C is a susceptibility gene for ovarian cancer.
246. PMID: 22538716; 2012, Nat Genet;44(5):475-6; author reply 476
Germline RAD51C mutations confer susceptibility to ovarian cancer.
247. PMID: 24315737; 2014, Oral Oncol;50(3):196-9
RAD51C--a new human cancer susceptibility gene for sporadic squamous cell carcinoma of the head and neck (HNSCC).
248. PMID: 11034073; 2000, Cancer Res;60(19):5371-5
17q23 amplifications in breast cancer involve the PAT1, RAD51C, PS6K, and SIGMa1B genes.
249. PMID: 11034067; 2000, Cancer Res;60(19):5340-4
Multiple genes at 17q23 undergo amplification and overexpression in breast cancer.
250. PMID: 20471405; 2010, Mutat Res;689(1-2):50-8
Rad51C is essential for embryonic development and haploinsufficiency causes increased DNA damage sensitivity and genomic instability.
251. PMID: 23512992; 2013, Mol Cancer Ther;12(6):865-77
RAD51C-deficient cancer cells are highly sensitive to the PARP inhibitor olaparib.
252. PMID: 25935112; 2015, Trends Biochem Sci;40(6):296-308
Structural determinants of Smad function in TGF- β signaling.
253. PMID: 19014666; 2008, Pathogenetics;1(1):2
Smad4 haploinsufficiency: a matter of dosage.
254. PMID: 9545410; 1998, Am J Hum Genet;62(5):1129-36
A gene for familial juvenile polyposis maps to chromosome 18q21.1.
255. PMID: 8553070; 1996, Science;271(5247):350-3
DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1.
256. PMID: 8673134; 1996, Nat Genet;13(3):343-6
Evaluation of candidate tumour suppressor genes on chromosome 18 in colorectal cancers.
257. PMID: 18662538; 2008, Cell;134(2):215-30
TGFbeta in Cancer.
258. PMID: 9135016; 1997, Cancer Res;57(9):1731-4
Tumor-suppressive pathways in pancreatic carcinoma.
259. PMID: 23139211; 2013, Cancer Res;73(2):725-35
SMAD2, SMAD3 and SMAD4 mutations in colorectal cancer.

ACT Onco[®] + Report

260. PMID: 22810696; 2012, Nature;487(7407):330-7
Comprehensive molecular characterization of human colon and rectal cancer.
261. PMID: 25890228; 2015, World J Surg Oncol;13():128
Clinical outcome and expression of mutant P53, P16, and Smad4 in lung adenocarcinoma: a prospective study.
262. PMID: 19841540; 2009, J Clin Invest;119(11):3208-11
Smad4: gatekeeper gene in head and neck squamous cell carcinoma.
263. PMID: 15867212; 2005, Clin Cancer Res;11(9):3191-7
Differences in Smad4 expression in human papillomavirus type 16-positive and human papillomavirus type 16-negative head and neck squamous cell carcinoma.
264. PMID: 25589618; 2015, Clin Cancer Res;21(6):1447-56
Genomic analysis of metastatic cutaneous squamous cell carcinoma.
265. PMID: 29703253; 2018, BMC Cancer;18(1):479
SMAD4 and NF1 mutations as potential biomarkers for poor prognosis to cetuximab-based therapy in Chinese metastatic colorectal cancer patients.
266. PMID: 28522603; 2017, Clin Cancer Res;23(17):5162-5175
SMAD4 Loss Is Associated with Cetuximab Resistance and Induction of MAPK/JNK Activation in Head and Neck Cancer Cells.
267. PMID: 16144935; 2005, Clin Cancer Res;11(17):6311-6
SMAD4 levels and response to 5-fluorouracil in colorectal cancer.
268. PMID: 24384683; 2014, Br J Cancer;110(4):946-57
Loss of Smad4 in colorectal cancer induces resistance to 5-fluorouracil through activating Akt pathway.
269. PMID: 12237773; 2002, Br J Cancer;87(6):630-4
SMAD4 is a predictive marker for 5-fluorouracil-based chemotherapy in patients with colorectal cancer.
270. PMID: 25749173; 2015, Transl Oncol;8(1):18-24
A Meta-Analysis of SMAD4 Immunohistochemistry as a Prognostic Marker in Colorectal Cancer.
271. PMID: 19478385; 2009, Cell Oncol;31(3):169-78
Presence of a high amount of stroma and downregulation of SMAD4 predict for worse survival for stage I-II colon cancer patients.
272. PMID: 25681512; 2015, J Clin Pathol;68(5):341-5
Smad4 inactivation predicts for worse prognosis and response to fluorouracil-based treatment in colorectal cancer.
273. PMID: 26861460; 2016, Clin Cancer Res;22(12):3037-47
Reduced Expression of SMAD4 Is Associated with Poor Survival in Colon Cancer.
274. PMID: 26947875; 2016, Transl Oncol;9(1):1-7
Prognostic Value of SMAD4 in Pancreatic Cancer: A Meta-Analysis.
275. PMID: 25760429; 2015, Pancreas;44(4):660-4
SMAD4 expression predicts local spread and treatment failure in resected pancreatic cancer.
276. PMID: 22504380; 2012, Pancreas;41(4):541-6
SMAD4 genetic alterations predict a worse prognosis in patients with pancreatic ductal adenocarcinoma.
277. PMID: 19584151; 2009, Clin Cancer Res;15(14):4674-9
SMAD4 gene mutations are associated with poor prognosis in pancreatic cancer.
278. PMID: 18425078; 2008, Mod Pathol;21(7):866-75
Expression of Smad2 and Smad4 in cervical cancer: absent nuclear Smad4 expression correlates with poor survival.
279. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35

ACT Onco[®] + Report

mTOR: from growth signal integration to cancer, diabetes and ageing.

280. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
281. PMID: 9242607; 1997, Science;277(5327):805-8
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
282. PMID: 8269512; 1993, Cell;75(7):1305-15
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
283. PMID: 1303246; 1992, Nat Genet;2(1):37-41
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
284. PMID: 18538015; 2008, BMC Cancer;8():163
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
285. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
286. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56
The tuberous sclerosis complex.
287. PMID: 25724664; 2015, Mol Cancer Ther;14(5):1224-35
Loss of Tuberous Sclerosis Complex 2 (TSC2) Is Frequent in Hepatocellular Carcinoma and Predicts Response to mTORC1 Inhibitor Everolimus.
288. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
289. PMID: 26412398; 2015, Sci Rep;5():14534
PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.
290. PMID: 28968163; 2017, J Clin Oncol;35(32):3638-3646
MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer.
291. PMID: 28533223; 2017, Clin Cancer Res;23(17):5218-5224
MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR+/HER2-Metastatic Breast Cancer.
292. PMID: 29573941; 2018, Lancet Oncol;19(5):603-615
Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial.
293. PMID: 33503393; 2021, J Clin Oncol;39(4):273-284
Encorafenib Plus Cetuximab as a New Standard of Care for Previously Treated BRAF V600E-Mutant Metastatic Colorectal Cancer: Updated Survival Results and Subgroup Analyses from the BEACON Study.
294. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
295. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
296. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.

ACT Onco[®] + Report

297. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
298. PMID: 30948273; 2019, Lancet Oncol;20(5):636-648
Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial.
299. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
300. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
301. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
302. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
303. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
304. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
305. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
306. PMID: 25366685; 2015, J Clin Oncol;33(3):244-50
Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation.
307. PMID: 27959613; 2016, N Engl J Med;375(20):1925-1936
Palbociclib and Letrozole in Advanced Breast Cancer.
308. PMID: 26030518; 2015, N Engl J Med;373(3):209-19
Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer.
309. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
310. PMID: 24768112; 2014, Lancet;384(9940):319-28
Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial.
311. PMID: 18650514; 2008, N Engl J Med;359(4):378-90
Sorafenib in advanced hepatocellular carcinoma.
312. PMID: 17189398; 2006, Clin Cancer Res;12(24):7271-8
Sorafenib for the treatment of advanced renal cell carcinoma.
313. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
314. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsilolimus, interferon alfa, or both for advanced renal-cell carcinoma.
315. PMID: 29188284; 2018, JAMA Oncol;4(3):384-388

ACT Onco[®] + Report

Vemurafenib for BRAF V600-Mutant Erdheim-Chester Disease and Langerhans Cell Histiocytosis: Analysis of Data From the Histology-Independent, Phase 2, Open-label VE-BASKET Study.

316. PMID: 21639808; 2011, N Engl J Med;364(26):2507-16
Improved survival with vemurafenib in melanoma with BRAF V600E mutation.