

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE	Brain glioblastoma (GBM)	PHYSICIAN	ORDERING PHYSICIAN	Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE	Brain
	NAME	Hua, Shu-Chen		MEDICAL FACILITY	Taipei Veterans General Hospital		SPECIMEN ID	S110-38114B (PF21072)
	DATE OF BIRTH	12 February 1970		ADDITIONAL RECIPIENT	None		SPECIMEN TYPE	Slide Deck
	SEX	Female		MEDICAL FACILITY ID	205872		DATE OF COLLECTION	04 December 2021
	MEDICAL RECORD #	47897723		PATHOLOGIST	Not Provided		SPECIMEN RECEIVED	18 December 2021

Biomarker Findings

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

FGFR3 FGFR3-TACC3 fusion

CDK4 amplification

MDM2 amplification

QKI Y204*

TERT promoter -124C>T

3 Disease relevant genes with no reportable alterations: **EGFR**, **IDH1**, **PDGFRA**

† See About the Test in appendix for details.

Report Highlights

- Variants with **diagnostic implications** that may indicate a specific cancer type: **TERT** promoter -124C>T (p. 6)
- Targeted therapies with potential clinical benefit **approved in another tumor type**: **Erdafeitinib** (p. 7)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 8)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **TERT** promoter -124C>T (p. 6)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

GENOMIC FINDINGS

FGFR3 - FGFR3-TACC3 fusion

10 Trials see p. 10

CDK4 - amplification

10 Trials see p. 8

MDM2 - amplification

6 Trials see p. 12

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
none	Erdafeitinib
none	none
none	none

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

QKI - Y204* p. 5 **TERT** - promoter -124C>T p. 6

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

Low-level MSI has been reported in 5-9% of glioblastoma (GBM) samples⁶⁻⁸. A large-scale study did not find high-level microsatellite instability (MSI-H) in any of 129 GBM samples⁶, although a small-scale study reported MSI-H in 4 of 15 pediatric GBMs and 1 of 12 adult GBMs⁹. The frequency of MSI has been reported to be increased in relapsed compared to primary GBM⁶, in GBMs with a previous lower grade astrocytoma⁷, and in giant cell GBM compared to classic GBM⁸.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁰⁻¹². This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹³⁻¹⁵. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{10,12,14-15}.

BIOMARKER

Tumor Mutational Burden

RESULT

1 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁶⁻¹⁸, anti-PD-1 therapies¹⁶⁻¹⁹, and combination nivolumab and ipilimumab²⁰⁻²⁵. In glioma, a lack of association between TMB and clinical benefit from immune checkpoint inhibitors has been reported^{16,26-27}. However, multiple case studies have reported that patients with ultramutated gliomas driven by POLE

mutations have benefited from treatment with anti-PD-1²⁸⁻²⁹ or anti-PD-L1³⁰ therapies. Therefore, although increased TMB alone may not be a strong biomarker for PD-1 or PD-L1 inhibitors in this cancer type, these agents may have efficacy for patients with glioma harboring both high TMB and POLE mutation.

FREQUENCY & PROGNOSIS

Glioblastoma (GBM) harbors a median TMB of 2.7 mutations per megabase (mut/Mb), and 4.2% of cases have high TMB (>20 muts/Mb)³¹. For pediatric patients, high TMB has been reported in a subset of high-grade gliomas, frequently in association with mutations in mismatch repair or proofreading genes and in TP53, whereas BRAF alterations or other oncogene fusions were observed more frequently in brain tumors harboring low TMB³²⁻³³. Increased TMB has been reported to correlate with higher tumor grade in glioma³⁴ and glioblastoma (GBM) tissue samples with biallelic mismatch repair deficiency

(bMMRD)²⁸, as well as with shorter OS of patients with diffuse glioma³⁵.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁶⁻³⁷ and cigarette smoke in lung cancer³⁸⁻³⁹, treatment with temozolomide-based chemotherapy in glioma⁴⁰⁻⁴¹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁴²⁻⁴⁶, and microsatellite instability (MSI)^{42,45-46}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{16,26-30}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

GENOMIC FINDINGS

GENE

FGFR3

ALTERATION

FGFR3-TACC3 fusion

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Alterations that activate FGFR3 may predict sensitivity to selective FGFR kinase inhibitors, including erdafitinib⁴⁷⁻⁴⁹, pemigatinib⁵⁰, infigratinib⁵¹⁻⁵², rogaratinib⁵³, Debio 1347⁵⁴⁻⁵⁵, and derazantinib⁵⁶; multikinase inhibitors such as pazopanib⁵⁷⁻⁵⁸ and ponatinib⁵⁹⁻⁶⁰; and vofatamab, an antibody targeting FGFR3⁶¹⁻⁶³. In the context of FGFR3 alterations, FGFR inhibitors, such as erdafitinib⁴⁷, pemigatinib⁵⁰, infigratinib⁵¹, rogaratinib⁵³, and Debio 1347⁶⁴, have predominantly been studied in the context of urothelial carcinoma, resulting in ORRs of 25-40% and DCRs of 64-80%. Clinical benefit has been reported in patients with gliomas harboring FGFR3 fusions treated in a Phase 1 trial of erdafitinib^{48,65}, and a prolonged SD has been observed in a case study treated with Debio 1347⁶⁶. For infigratinib, activity against non-urothelial tumors harboring FGFR3 alterations are limited⁶⁷, with responses reported in individuals with an FGFR3-amplified and -rearranged glioma or FGFR3-mutated head and neck squamous cell carcinoma with co-occurring FGF

amplifications⁶⁸.

FREQUENCY & PROGNOSIS

FGFR3-TACC3 fusions have been reported in 1-8% of glioblastoma (GBM) cases^{65,69-75} and 0-1% of lower-grade glioma cases^{65,69,74}. In both GBM and lower-grade glioma, FGFR3-TACC3 fusions have been observed only in IDH1/2 wildtype cases (3.6% [21/588] of GBM cases in one study; 3/85 [3.5%] and 3/80 [3.8%] lower-grade glioma cases in 2 studies) but not in any samples harboring IDH1 or IDH2 mutations (0/126 lower-grade glioma cases in 1 study; 0/187 combined GBM and lower-grade glioma cases in another study)^{65,74}. A case report described a patient with polymorphous low-grade neuroepithelial tumor harboring an FGFR3-TACC3 fusion⁷⁶. FGFR3 has been reported to be expressed in GBM cell lines⁷⁷. FGFR3 fusion was reported to be an independent predictor of longer OS in patients with IDH-wildtype glioma (stage II-IV combined cohort) or glioblastoma (GBM) specifically; however, OS of patients with IDH-wildtype, FGFR3 fusion-positive GBM was found to be significantly shorter than that of patients with IDH-mutated GBM⁷⁴. Additionally, IDH-wildtype gliomas with FGFR3 fusions were reported to have a significantly lower rate of MGMT methylation than IDH-wildtype cases lacking such fusions⁷⁴. In other studies, FGFR3 fusions were reported to be associated with recurrent histologic features in both low-grade and high-grade glioma, including monomorphous oligodendroglioma-like nuclei,

endocrinoid networks of thin capillaries, and frequent microcalcifications⁷⁸⁻⁷⁹. However, due to the higher frequency of FGFR3 fusions in GBM compared to lower-grade glioma, as well as the limited long-term follow-up data for ostensibly low-grade but FGFR3-positive cases, guidelines have recommended cautious evaluation of clinical features and close monitoring of the course of the disease when a histologically low-grade tumor is found to harbor an FGFR3 fusion⁷⁹. One retrospective analysis reported association between higher FGFR3 expression and longer OS in a combined cohort of patients with grade II-IV glioma⁸⁰.

FINDING SUMMARY

FGFR3 (Fibroblast growth factor receptor 3) encodes a receptor tyrosine kinase that typically promotes cell cycle progression and angiogenesis via activation of downstream signaling pathways, including RAS-MAPK and AKT; gain of function mutations in FGFRs have been reported in several cancer types⁸¹⁻⁸³. FGFR3 fusions that retain the kinase domain (exons 11-17) have been shown to be activating and oncogenic^{54,71}. Additionally, fusions that disrupt a binding site of the regulatory microRNA miR-99a in the FGFR3 3' UTR have been demonstrated to increase FGFR3 expression⁶⁹. Rearrangements that include the N-terminal portion of FGFR3 (exons 1-17), such as observed here, are predicted to be activating and oncogenic.

GENE

CDK4

ALTERATION

amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

CDK4 amplification or activation may predict sensitivity to CDK4/6 inhibitors such as abemaciclib, palbociclib, and ribociclib⁸⁴⁻⁸⁷. Clinical benefit has been reported for limited tumor types including patients with

CDK4-amplified liposarcoma and sarcoma in response to treatment with abemaciclib⁸⁸, palbociclib^{84,89}, and ribociclib⁹⁰.

FREQUENCY & PROGNOSIS

Across TCGA and MKSCC studies, CDK4 amplification has been reported in 4.0-9.4% of glioma cases and 14% of glioblastoma multiforme cases (cBioPortal, Sep 2021)⁹¹⁻⁹⁵. A study has reported amplification of the 12q14-15 region, where CDK4 and MDM2 reside, in 5% (2/42) of glioblastomas⁹⁶. Amplification of CDK4 and corresponding increased CDK4 protein expression has been reported to be associated with a poorer patient outcome in anaplastic astrocytoma and

glioblastoma⁹⁷⁻¹⁰⁰.

FINDING SUMMARY

CDK4 encodes the cyclin-dependent kinase 4, which regulates the cell cycle, senescence, and apoptosis¹⁰¹. CDK4 and its functional homolog CDK6 are activated by D-type cyclins and promote cell cycle progression by inactivating the tumor suppressor Rb¹⁰²⁻¹⁰³. Amplification of the chromosomal region that includes CDK4 has been reported in multiple cancer types, including lung cancer, glioblastoma, and liposarcoma, and has been associated with overexpression of CDK4 protein^{84,104-110}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

GENOMIC FINDINGS

GENE

MDM2

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

MDM2 antagonists disrupt the MDM2-p53 interaction, thereby stabilizing p53¹¹¹. Preclinical studies have suggested that the amplification of MDM2, in the absence of concurrent TP53 mutations, may increase sensitivity to these agents¹¹²⁻¹¹³. Preliminary Phase 1 studies of the MDM2-p53 antagonist alrizomadlin (APG-115) reported a PR in a patient with liposarcoma harboring an MDM2 amplification and wildtype for TP53 and SD in 21%–38% (6/28 and 5/13, respectively) of patients in genomically unselected solid tumors¹¹⁴⁻¹¹⁵. A Phase 2 trial of alrizomadlin in combination with pembrolizumab reported a PR in 1 of 3 patients with malignant peripheral nerve sheath tumor that had failed standard therapy, as well as PRs in patients with multiple

types of solid tumors that had failed immunotherapy, including 1 out of 14 patients with non-small cell lung cancer; 1 out of 5 patients with urothelial carcinoma; and 2 out of 5, 1 out of 5, and 1 out of 11 patients with mucosal, uveal, and cutaneous melanoma, respectively¹¹⁶. Phase 1b studies of the MDM2 inhibitor idasanutlin for refractory AML in combination with cytarabine or venetoclax reported anti-leukemic response rates of 33% (25/75) and 37% (11/30), respectively¹¹⁷⁻¹¹⁸; clinical benefit (58% ORR, 7/12) with idasanutlin monotherapy has been reported for patients with polycythemia vera¹¹⁹. The dual MDM2/MDM4 inhibitor ALRN-6924 led to an ORR of 27% (4/15) for patients with TP53 wildtype peripheral T-cell lymphoma in a Phase 2 study¹²⁰; responses have also been observed in TP53 wildtype AML, MDS, Merkel cell carcinoma, colorectal cancer, and liposarcoma¹²¹⁻¹²².

FREQUENCY & PROGNOSIS

In the Glioblastoma Multiforme (GBM) TCGA dataset, amplification of MDM2 has been found in 8% of cases⁹³. A study has reported amplification of the 12q14-15 region, where MDM2 and CDK4 reside, in 5% (2/42) of GBMs⁹⁶. Amplification of

MDM2 has been associated with poor survival in patients with glioblastoma^{96,123}.

FINDING SUMMARY

MDM2 encodes an E3 ubiquitin protein ligase, which mediates the ubiquitination and subsequent degradation of p53, Rb1, and other proteins¹²⁴⁻¹²⁶. MDM2 acts to prevent the activity of the tumor suppressor p53; therefore, overexpression or amplification of MDM2 may be oncogenic¹²⁷⁻¹²⁸. Overexpression or amplification of MDM2 is frequent in cancer¹²⁹. Although two retrospective clinical studies suggest that MDM2 amplification may predict a short time-to-treatment failure on anti-PD-1/PD-L1 immune checkpoint inhibitors, with 4/5 patients with MDM2 amplification¹³⁰ and 2/3 patients with MDM2 or MDM4 amplification¹³¹ experiencing tumor hyperprogression, amplification of MDM2 or MDM4 was not associated with shorter progression-free survival (PFS) in a retrospective analysis of non-small cell lung cancer (NSCLC) outcomes with immune checkpoint inhibitors (hazard ratio of 1.4, p=0.44)¹³². The latter study reported PFS of >2 months for 5/8 patients with MDM2/MDM4 amplification¹³².

GENE

QKI

ALTERATION
Y204*

TRANSCRIPT ID
NM_006775

CODING SEQUENCE EFFECT
612C>G

VARIANT ALLELE FREQUENCY (% VAF)
15.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies approved or in clinical trials that directly address genomic alterations in QKI.

FREQUENCY & PROGNOSIS

Recurrent deletions of QKI have been reported in astrocytomas and glioblastomas¹³³⁻¹³⁵. Decreased expression of quaking has also been reported in colon and gastric cancer, largely due to QKI promoter hypermethylation; in one study, forced

expression of quaking in colon cancer cells inhibited proliferation and tumorigenesis¹³⁶⁻¹³⁷. However, another study reported that QKI acts as an oncogene in breast cancer by repressing the expression of the tumor suppressor FOXO1¹³⁸.

FINDING SUMMARY

QKI encodes quaking, an RNA-binding protein that plays roles in RNA metabolism and signal transduction and is necessary for myelination and embryonic development¹³⁹⁻¹⁴⁰. QKI acts as a negative regulator of cell-cycle progression¹⁴¹.

ORDERED TEST # ORD-1262948-01

GENOMIC FINDINGS

GENE

TERT

ALTERATION

promoter -124C>T

TRANSCRIPT ID

NM_198253

CODING SEQUENCE EFFECT

-124C>T

VARIANT ALLELE FREQUENCY (% VAF)

21.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapeutic options for targeting tumors with TERT mutations are limited, although a variety of approaches have been investigated, including immunotherapies using TERT as a tumor-associated antigen and antisense oligonucleotide- or peptide-based therapies. TERT peptide vaccines showed limited anticancer efficacy in clinical trials¹⁴²; however, in one preclinical study, the combination of a TERT peptide vaccine and anti-CTLA-4 therapy suppressed tumor growth¹⁴³. A Phase 2 study of the TERT inhibitor imetelstat for patients with advanced non-small cell lung cancer reported no improvement in PFS or OS¹⁴⁴.

FREQUENCY & PROGNOSIS

TERT promoter mutations have been reported in 51-59% of gliomas¹⁴⁵⁻¹⁴⁶, most frequently in glioblastoma (GBM, 54-84%), gliosarcoma (81%), oligodendroglioma (78%), and historically in oligoastrocytomas (25-31%) but less frequently in lower grade astrocytomas (10-18%) and in only 1% of ependymomas¹⁴⁵⁻¹⁴⁹. In patients with glioblastoma (GBM), the prevalence of TERT promoter mutation is lower in pediatric primary GBM (11%) and adult secondary GBM (28%) compared with adult primary GBM (58-83%)^{145,147}. One study detected TERT promoter mutations in 78% (181/232) of IDH-wildtype GBM samples analyzed¹⁵⁰. TERT promoter mutation has been shown to be significantly associated with increased TERT gene expression in astrocytoma, oligodendroglioma, and GBM¹⁵¹. TERT promoter mutations significantly associate with poor prognosis in patients with GBM, although this correlation may be due to the association with primary GBM as opposed to IDH-positive secondary GBM^{145,147,151-152}. In the context of IDH-wildtype glioma, TERT mutations are associated with reduced OS, whereas in IDH-mutated, 1p/19q co-deleted oligodendroglioma, TERT mutations are associated with improved OS (NCCN CNS Cancers Guidelines, v5.2020).

FINDING SUMMARY

Telomerase reverse transcriptase (TERT, or hTERT) is a catalytic subunit of the telomerase complex, which is required to maintain appropriate chromosomal length¹⁵³. Activation of TERT is a hallmark of cancer, being detected in up to 80-90% of malignancies and absent in quiescent cells¹⁵⁴⁻¹⁵⁶. Mutations within the promoter region of TERT that confer enhanced TERT promoter activity have been reported in two hotspots, located at -124 bp and -146 bp upstream of the transcriptional start site (also termed C228T and C250T, respectively)¹⁵⁷⁻¹⁵⁹, as well as tandem mutations at positions -124/-125 bp and -138/-139 bp¹⁵⁷.

POTENTIAL DIAGNOSTIC IMPLICATIONS

TERT mutations are associated with 1p/19q co-deletion in oligodendrogliomas, and are highly recurrent in IDH/ATRX-wildtype glioblastoma (GBM) (NCCN CNS Cancers Guidelines, v5.2020). Co-occurring TERT mutation, IDH mutation, and 1p/19q co-deletion is indicative of oligodendroglioma, whereas IDH mutation in the absence of TERT mutation is suggestive of astrocytoma (NCCN CNS Cancers Guidelines, v5.2020).

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Erdaftinib

Assay findings association

FGFR3

FGFR3-TACC3 fusion

AREAS OF THERAPEUTIC USE

Erdaftinib is a pan-fibroblast growth factor receptor (FGFR) inhibitor. It is FDA approved for the treatment of patients with advanced or metastatic urothelial carcinoma who have FGFR2 or FGFR3 alterations and have progressed after prior chemotherapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of strong clinical evidence, FGFR3 fusions^{48,65,160-161} and activating mutations^{47-48,160} may confer sensitivity to erdaftinib.

SUPPORTING DATA

In a Phase 1 study of erdaftinib for patients with advanced solid tumors, 1 patient with glioblastoma harboring an FGFR3 fusion achieved a PR⁴⁸. Another Phase 1 study reported that 2 patients with FGFR3 fusion-positive glioblastoma treated with erdaftinib

experienced clinical benefit, including a 115-day SD and a minor response, respectively⁶⁵. Erdaftinib has been primarily studied for the treatment of FGFR-altered urothelial carcinoma. A Phase 2 study evaluating erdaftinib for the treatment of patients with metastatic or unresectable urothelial carcinoma (mUC) previously treated with chemotherapy and harboring FGFR2/3 fusions or FGFR3 activating mutations reported an ORR of 40% (40/99, 3 CR), and a DCR of 80% (79/99)⁴⁷. A Phase 1 trial of erdaftinib reported clinical responses in for patients with various FGFR2- or FGFR3-altered solid tumors^{48,65,162-163}, including cholangiocarcinoma (27% ORR, 3/11), NSCLC (5% ORR, 1/21), breast (9% ORR, 3/34), and ovarian (9% ORR, 1/11), while other cancers including endometrial carcinoma and glioblastoma showed a low ORR (2%, 1/58)¹⁶⁴. Following progression on multiple other lines of therapy, a patient with metastatic FGFR2-fusion-positive NSCLC treated with erdaftinib exhibited an 11-month PR¹⁶².

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
CDK4
RATIONALE
CDK4 amplification may predict sensitivity to

CDK4/6 inhibitors.

ALTERATION
amplification

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04594005
PHASE 1/2

CDK4/6 Tumor, Abemaciclib, Paclitaxel

TARGETS
CDK4, CDK6

LOCATIONS: Seoul (Korea, Republic of)

NCT03099174
PHASE 1

This Study in Patients With Different Types of Cancer (Solid Tumours) Aims to Find a Safe Dose of Xentuzumab in Combination With Abemaciclib With or Without Hormonal Therapies. The Study Also Tests How Effective These Medicines Are in Patients With Lung and Breast Cancer.

TARGETS
CDK4, CDK6, IGF-1, IGF-2, Aromatase, ER

LOCATIONS: Seoul (Korea, Republic of), Goyang (Korea, Republic of), Aichi, Nagoya (Japan), Kanagawa, Isehara (Japan), Tokyo, Chuo-ku (Japan), Tokyo, Koto-ku (Japan), Chiba, Kashiwa (Japan), Helsinki (Finland), Tampere (Finland), Turku (Finland)

NCT04074785
PHASE NULL

Pilot Study of Abemaciclib With Bevacizumab in Recurrent Glioblastoma Patients With Loss of CDKN2A/B or Gain or Amplification of CDK4/6

TARGETS
VEGFA, CDK4, CDK6

LOCATIONS: Texas

NCT04391595
PHASE NULL

LY3214996 Plus Abemaciclib in Recurrent Glioblastoma Patients

TARGETS
CDK4, CDK6, ERK1, ERK2

LOCATIONS: Arizona

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

CLINICAL TRIALS
NCT03834740
PHASE NULL

PhO/2 Ribociclib & Everolimus

TARGETS
CDK6, CDK4, mTOR

LOCATIONS: Arizona

NCT02933736
PHASE NULL

Ribociclib (LEE011) in Preoperative Glioma and Meningioma Patients

TARGETS
CDK6, CDK4

LOCATIONS: Arizona

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT03994796
PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS
TRKB, ALK, TRKC, ROS1, TRKA, CDK4, CDK6, PI3K, mTOR

LOCATIONS: Alaska, Washington

NCT03158389
PHASE 1/2

NCT Neuro Master Match - N²M² (NOA-20)

TARGETS
ALK, RET, CDK4, CDK6, mTOR, MDM2, PD-L1, SMO

LOCATIONS: Berlin (Germany), Dresden (Germany), Regensburg (Germany), Bochum (Germany), Frankfurt am Main (Germany), Essen (Germany), Mainz (Germany), Heidelberg (Germany), Cologne (Germany), Mannheim (Germany)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

CLINICAL TRIALS
GENE
FGFR3
RATIONALE

FGFR inhibitors may be relevant in tumors with alterations that activate FGFR3.

ALTERATION
FGFR3-TACC3 fusion
NCT04083976
PHASE 2

A Study of Erdafitinib in Participants With Advanced Solid Tumors and Fibroblast Growth Factor Receptor (FGFR) Gene Alterations

TARGETS
FGFRs

LOCATIONS: Taipei (Taiwan), Taoyuan City (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Tainan (Taiwan), Kaohsiung City (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Shanghai (China), Nanjing (China)

NCT03797326
PHASE 2

Efficacy and Safety of Pembrolizumab (MK-3475) Plus Lenvatinib (E7080/MK-7902) in Previously Treated Participants With Select Solid Tumors (MK-7902-005/E7080-G000-224/LEAP-005)

TARGETS
PD-1, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Songpago (Korea, Republic of), Bangkok (Thailand), Nedlands (Australia), Kazan (Russian Federation), Herston (Australia), Arkhangelsk (Russian Federation), Moscow (Russian Federation)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT04189445
PHASE 2

Futibatinib in Patients With Specific FGFR Aberrations

TARGETS
FGFRs

LOCATIONS: Seul (Korea, Republic of), Sapporo-shi (Japan), London (United Kingdom), California, Arizona, Wisconsin, Texas

NCT03564691
PHASE 1

Study of MK-4830 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-4830-001)

TARGETS
ITL4, FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Seoul (Korea, Republic of), Tokyo (Japan), Haifa (Israel), Petah Tikva (Israel), Ramat Gan (Israel), Tel Aviv (Israel), Warszawa (Poland), Gdansk (Poland), Heraklion (Greece), Washington

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

CLINICAL TRIALS
NCT04008797
PHASE 1

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

TARGETS
CBP, Beta-catenin, FGFRs, RET,
PDGFRA, VEGFRs, KIT

LOCATIONS: Osakasayama (Japan), Chuo-Ku (Japan), Kashiwa (Japan)

NCT03547037
PHASE 1

A Study to Evaluate the Safety, Pharmacokinetics, Pharmacodynamics, and Immunogenicity of JNJ-63723283, an Anti-Programmed Cell Death (PD)-1 Monoclonal Antibody, as Monotherapy or in Combination With Erdafitinib in Japanese Participants With Advanced Solid Cancers

TARGETS
PD-1, FGFRs

LOCATIONS: Chuo-Ku (Japan), Kashiwa (Japan)

NCT04424966
PHASE NULL

Infigratinib in Recurrent Glioblastoma Patients

TARGETS
FGFR3, FGFR1, FGFR2

LOCATIONS: Arizona

NCT04042116
PHASE 1/2

A Study to Evaluate Lucitanib in Combination With Nivolumab in Patients With a Solid Tumor

TARGETS
FGFRs, VEGFRs, PD-1

LOCATIONS: Innsbruck (Austria), Essen (Germany), Bologna (Italy), Naples (Italy), Leuven (Belgium), Brussels (Belgium), Ghent (Belgium), Washington, Barcelona (Spain), Madrid (Spain)

NCT04565275
PHASE 1/2

A Study of ICP-192 in Patients With Advanced Solid Tumors

TARGETS
FGFR2, FGFR1, FGFR3, FGFR4

LOCATIONS: Colorado, Minnesota, Arizona, Florida

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

CLINICAL TRIALS
GENE
MDM2
ALTERATION
amplification
RATIONALE

Inhibitors of the MDM2-p53 interaction are being tested in clinical trials. Overexpression or

amplification of MDM2 may increase sensitivity to these agents, but more data are required.

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS
TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha
LOCATIONS: Zhongzheng Dist. (Taiwan), Taipei City (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Beijing (China), Woolloongabba (Australia), Darlinghurst (Australia), Randwick (Australia), Melbourne (Australia), Haifa (Israel)

NCT03449381
PHASE 1

This Study Aims to Find the Best Dose of BI 907828 in Patients With Different Types of Advanced Cancer (Solid Tumors)

TARGETS
MDM2
LOCATIONS: Tokyo, Chuo-ku (Japan), Ottawa (Canada), Connecticut, New York, Tennessee, Florida

NCT03611868
PHASE 1/2

A Study of APG-115 in Combination With Pembrolizumab in Patients With Metastatic Melanomas or Advanced Solid Tumors

TARGETS
MDM2, PD-1
LOCATIONS: Brisbane (Australia), California, Arizona, Missouri, Arkansas, Pennsylvania, New York, Tennessee, Texas

NCT03158389
PHASE 1/2

NCT Neuro Master Match - N²M² (NOA-20)

TARGETS
ALK, RET, CDK4, CDK6, mTOR, MDM2, PD-L1, SMO
LOCATIONS: Berlin (Germany), Dresden (Germany), Regensburg (Germany), Bochum (Germany), Frankfurt am Main (Germany), Essen (Germany), Mainz (Germany), Heidelberg (Germany), Cologne (Germany), Mannheim (Germany)

NCT03107780
PHASE 1

MDM2 Inhibitor AMG-232 in Treating Patients With Recurrent or Newly Diagnosed Glioblastoma

TARGETS
MDM2
LOCATIONS: California, Michigan, Pennsylvania, Massachusetts, New York, Maryland, North Carolina, Alabama

NCT03725436
PHASE 1

ALRN-6924 and Paclitaxel in Treating Patients With Advanced, Metastatic, or Unresectable Solid Tumors

TARGETS
MDM2, MDM4
LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

BRCA2
C3253Y

MLL2
P2349L

SDHC
A66V

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKKN1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPR2S2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX
About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials
Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2021. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 - Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of HER2 overexpression as 20% in breast cancer. Based on the F1CDx HER2 CDx concordance study, approximately 10% of HER2 amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 5.2.0

The median exon coverage for this sample is 870x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # **ORD-1262948-01**
APPENDIX
References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Martinez R, et al. Oncology (2004) PMID: 15331927
7. Martinez R, et al. J. Cancer Res. Clin. Oncol. (2005) PMID: 15672285
8. Martinez R, et al. Cancer Genet. Cytogenet. (2007) PMID: 17498554
9. Szybka M, et al. Clin. Neuropathol. () PMID: 12908754
10. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
11. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
12. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
13. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
14. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
15. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
16. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
17. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
18. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
19. Cristescu R, et al. Science (2018) PMID: 30309915
20. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
21. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
22. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
24. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
25. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
26. Zhao J, et al. Nat. Med. (2019) PMID: 30742119
27. Thout M, et al. Nature (2020) PMID: 32322066
28. Bouffett E, et al. J. Clin. Oncol. (2016) PMID: 27001570
29. Johanns TM, et al. Cancer Discov (2016) PMID: 27683556
30. Lukas RV, et al. J. Neurooncol. (2018) PMID: 30073642
31. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
32. Patel RR, et al. Pediatr Blood Cancer (2020) PMID: 32386112
33. Johnson A, et al. Oncologist (2017) PMID: 28912153
34. Draaisma K, et al. Acta Neuropathol Commun (2015) PMID: 26699864
35. Wang L, et al. BMC Cancer (2020) PMID: 32164609
36. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
37. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
38. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
39. Rizvi NA, et al. Science (2015) PMID: 25765070
40. Johnson BE, et al. Science (2014) PMID: 24336570
41. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
42. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
43. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
44. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
45. Nature (2012) PMID: 22810696
46. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
47. Lorient Y, et al. N. Engl. J. Med. (2019) PMID: 31340094
48. Tabernero J, et al. J. Clin. Oncol. (2015) PMID: 26324363
49. Karkera JD, et al. Mol. Cancer Ther. (2017) PMID: 28416604
50. Necchi et al., 2018; ESMO Abstract 900P
51. Pal SK, et al. Cancer Discov (2018) PMID: 29848605
52. Pal SK, et al. Cancer (2020) PMID: 32208524
53. Schuler M, et al. Lancet Oncol. (2019) PMID: 31405822
54. Nakanishi Y, et al. Mol. Cancer Ther. (2015) PMID: 25589496
55. Voss MH, et al. Clin. Cancer Res. (2019) PMID: 30745300
56. Papadopoulos KP, et al. Br. J. Cancer (2017) PMID: 28972963
57. Bellmunt J, et al. Br. J. Cancer (2018) PMID: 30220708
58. Palma N, et al. Eur. Urol. (2015) PMID: 25766722
59. Gozgit JM, et al. Mol. Cancer Ther. (2012) PMID: 22238366
60. Liao RG, et al. Cancer Res. (2013) PMID: 23786770
61. Bellmunt et al., 2018; ASCO Abstract 4534
62. Necchi et al., 2019; ASCO GU Abstract 409
63. Siefker-Radtke et al., 2019; ASCO Abstract 4511
64. Voss et al., 2017; ASCO Abstract 2500
65. Di Stefano AL, et al. Clin. Cancer Res. (2015) PMID: 25609060
66. Farouk Sait S, et al. JCO Precis Oncol (2021) PMID: 34250399
67. Nogova L, et al. J. Clin. Oncol. (2017) PMID: 27870574
68. Slosberg ED, et al. Oncotarget (2018) PMID: 29765547
69. Parker BC, et al. J. Clin. Invest. (2013) PMID: 23298836
70. Stransky N, et al. Nat Commun (2014) PMID: 25204415
71. Singh D, et al. Science (2012) PMID: 22837387
72. Wu YM, et al. Cancer Discov (2013) PMID: 23558953
73. Bao ZS, et al. Genome Res. (2014) PMID: 25135958
74. Di Stefano AL, et al. Neuro Oncol (2020) PMID: 32413119
75. Schittenhelm J, et al. Neurooncol Pract (2021) PMID: 33898054
76. Huse JT, et al. Acta Neuropathol (2017) PMID: 27812792
77. Loilome W, et al. J. Neurooncol. (2009) PMID: 19340397
78. Bielle F, et al. Brain Pathol. (2018) PMID: 28976058
79. Acta Neuropathol Commun (2020) PMID: 32085805
80. Wang Z, et al. Oncotarget (2016) PMID: 27829236
81. Powers CJ, et al. Endocr. Relat. Cancer (2000) PMID: 11021964
82. Eswarakumar VP, et al. Cytokine Growth Factor Rev. (2005) PMID: 15863030
83. Wesche J, et al. Biochem. J. (2011) PMID: 21711248
84. Dickson MA, et al. J. Clin. Oncol. (2013) PMID: 23569312
85. Flaherty KT, et al. Clin. Cancer Res. (2012) PMID: 22090362
86. Patnaik A, et al. Cancer Discov (2016) PMID: 27217383
87. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
88. Dickson et al., 2019; ASCO Abstract 11004
89. Dickson MA, et al. JAMA Oncol (2016) PMID: 27124835
90. Peguero et al., 2016; ASCO Abstract 2528
91. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
92. Gao J, et al. Sci Signal (2013) PMID: 23550210
93. Brennan CW, et al. Cell (2013) PMID: 24120142
94. Jonsson P, et al. Clin. Cancer Res. (2019) PMID: 31263031
95. Ceccarelli M, et al. Cell (2016) PMID: 26824661
96. Zheng S, et al. Genes Dev. (2013) PMID: 23796897
97. Kim H, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20080666
98. Ruano Y, et al. Am. J. Clin. Pathol. (2009) PMID: 19141386
99. Fischer U, et al. Mol. Cancer Res. (2008) PMID: 18403636
100. Bäcklund LM, et al. Br. J. Cancer (2005) PMID: 15970925
101. Choi YJ, et al. Oncogene (2014) PMID: 23644662
102. Cell (1995) PMID: 7736585
103. Musgrove EA, et al. Nat. Rev. Cancer (2011) PMID: 21734724
104. Wikman H, et al. Genes Chromosomes Cancer (2005) PMID: 15543620
105. Rao SK, et al. J. Neurooncol. (2010) PMID: 19609742
106. Chung L, et al. Am. J. Surg. Pathol. (2009) PMID: 19574885
107. Ragazzini P, et al. Histol. Histopathol. (2004) PMID: 15024701
108. Dujardin F, et al. Mod. Pathol. (2011) PMID: 21336260
109. Zhang K, et al. Cancer Res. (2013) PMID: 23393200
110. Horvai AE, et al. Mod. Pathol. (2009) PMID: 19734852
111. Cheok CF, et al. Nat Rev Clin Oncol (2011) PMID: 20975744
112. Ohnstad HO, et al. Cancer (2013) PMID: 23165797
113. Gamble LD, et al. Oncogene (2012) PMID: 21725357
114. Zhang et al., 2019; ASCO Abstract 3124
115. Rasco et al., 2019; ASCO Abstract 3126
116. Tolcher et al., 2021; ASCO Abstract 2506
117. Martinelli et al., 2016; EHA21 Abstract S504
118. Daver et al., 2018; ASH Abstract 767
119. Mascarenhas et al., 2019; ASH Abstract 134
120. Shustov et al., 2018; ASH Abstract 1623
121. Sallman et al., 2018; ASH Abstract 4066
122. Meric-Bernstam et al., 2017; ASCO Abstract 2505
123. Fischer U, et al. Int. J. Cancer (2010) PMID: 19839052
124. Sdek P, et al. Mol. Cell (2005) PMID: 16337594
125. Brady M, et al. Mol. Cell. Biol. (2005) PMID: 15632057
126. Li M, et al. Mol. Cell (2004) PMID: 15053880
127. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
128. Cordon-Cardo C, et al. Cancer Res. (1994) PMID: 8306343
129. Beroukhir R, et al. Nature (2010) PMID: 20164920
130. Kato S, et al. Clin. Cancer Res. (2017) PMID: 28351930
131. Singavi et al., 2017; ESMO Abstract 1140PD
132. Rizvi H, et al. J. Clin. Oncol. (2018) PMID: 29337640
133. Ichimura K, et al. Oncogene (2006) PMID: 16205629
134. Mulholland PJ, et al. Cell Cycle (2006) PMID: 16582634
135. Yin D, et al. Mol. Cancer Res. (2009) PMID: 19435819
136. Yang G, et al. Gastroenterology (2010) PMID: 19686745
137. Bian Y, et al. Biochem. Biophys. Res. Commun. (2012) PMID: 22569043
138. Yu F, et al. Oncol. Rep. (2014) PMID: 24398626
139. Ebersole TA, et al. Nat. Genet. (1996) PMID: 8589716
140. Hardy RJ, et al. J. Neurosci. (1996) PMID: 8987822
141. Yang G, et al. Cell Cycle (2011) PMID: 21768773
142. Nat Rev Clin Oncol (2017) PMID: 27245281
143. Duperret EK, et al. Mol Ther (2018) PMID: 29249395
144. Chiappori AA, et al. Ann Oncol (2015) PMID: 25467017
145. Killela PJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23530248
146. Killela PJ, et al. Oncotarget (2014) PMID: 24722048
147. Nonoguchi N, et al. Acta Neuropathol. (2013) PMID: 23955565
148. Liu X, et al. Cell Cycle (2013) PMID: 23603989
149. Koelsche C, et al. Acta Neuropathol. (2013) PMID: 24154961
150. Yan et al. 2020; DOI:10.1200/PO.19.00385
151. Arita H, et al. Acta Neuropathol. (2013) PMID: 23764841
152. Reitman ZJ, et al. Acta Neuropathol. (2013) PMID: 24217890
153. Shay JW, et al. Semin. Cancer Biol. (2011) PMID: 22015685

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1262948-01

APPENDIX

References

- | | | |
|---|---|---|
| <p>154. Shay JW, et al. Eur. J. Cancer (1997) PMID: 9282118</p> <p>155. Kim NW, et al. Science (1994) PMID: 7605428</p> <p>156. Hanahan D, et al. Cell (2000) PMID: 10647931</p> <p>157. Horn S, et al. Science (2013) PMID: 23348503</p> | <p>158. Huang FW, et al. Science (2013) PMID: 23348506</p> <p>159. Vinagre J, et al. Nat Commun (2013) PMID: 23887589</p> <p>160. Siefker-Radtke et al., 2018; ASCO Abstract 4503</p> <p>161. Park et al., 2019; ASCO Abstract 4117</p> | <p>162. Qin A, et al. J Thorac Oncol (2019) PMID: 30267839</p> <p>163. Soria et al., 2016; ASCO Abstract 4074</p> <p>164. Bahleda R, et al. Clin. Cancer Res. (2019) PMID: 31088831</p> |
|---|---|---|

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 05 January 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531