

ACT Onco[®] + Report

PATIENT		
Identifier: 李偉誠		Patient ID: 33832814
Date of Birth: Sep 19, 1975		Gender: Male
Diagnosis: R/O Liposarcoma		
ORDERING PHYSICIAN		
Name: 顏厥全醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11153970A	Collection site: Mediastinum	Type: FFPE tissue
Date received: Dec 30, 2022	Lab ID: AA-23-00001	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
Not detected			

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
KDR Amplification	Cabozantinib, Pazopanib, Sunitinib	-
KIT Amplification	-	Imatinib, Nilotinib, Sunitinib
LYN Amplification	Dasatinib	-
PDGFRA Amplification	Imatinib, Pazopanib, Sorafenib, Sunitinib	-

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACTOnco[®] + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
Not detected		

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr17	TP53	Homozygous deletion	0
Chr8	LYN, NBN	Amplification	6
Chr4	KDR, KIT, PDGFRA	Amplification	7

- Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	1.3 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 64% tumor purity.
- TMB was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 3A		
<i>KIT</i> Amplification	Imatinib, Nilotinib, Sunitinib	resistant
Level 3B		
<i>KDR</i> Amplification	Cabozantinib	sensitive
Level 4		
<i>KDR</i> Amplification	Pazopanib, Sunitinib	sensitive
<i>LYN</i> Amplification	Dasatinib	sensitive
<i>PDGFRA</i> Amplification	Imatinib, Pazopanib, Sorafenib, Sunitinib	sensitive

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
Not detected	

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

KDR Amplification

Biological Impact

KDR (kinase domain receptor), also known as VEGFR2 or Flk-1, is a tyrosine kinase receptor for the vascular endothelial growth factor (VEGF) and involves in angiogenesis pathway^[1]. Binding of VEGF to KDR results in activation of phospholipase C (PLC-gamma) and downstream signaling via protein kinase C (PKC) and RAF/MEK/ERK^[2]. Mutations of KDR are rare in tumors, and alterations of KDR activity typically occur via KDR amplification and subsequent overexpression^{[3][4]}.

Therapeutic and prognostic relevance

To date, there are four VEGF inhibitors (sorafenib, sunitinib, pazopanib, bevacizumab) and one VEGFR2 inhibitor cabozantinib that are FDA-approved for the treatment of cancers^{[5][6][7]}. Notably, a case report showed that an angiosarcoma patient with concurrent KDR and FLT4 amplification developed a progressive disease when treated with sorafenib, but experienced a potent antitumor response and achieved clinically stable disease for 6 months after receiving pazopanib therapy^[8]. Besides, an angiosarcoma patient with upregulated VEGFR2 responded to sunitinib treatment^[9]. VEGFR2 inhibitors like apatinib and vandetanib are in early clinical phase trial^{[10][11][12]}.

KDR amplification and/or mutation has been selected as an inclusion criteria for the trial examining cabozantinib in metastatic castrate resistant prostate cancer (mCRPC) (NCT04631744) and sunitinib in malignant solid tumors (NCT03297606). The increased copy number of KIT or KDR significantly correlated with a worse 5-year breast cancer-specific survival (BCSS) in triple-negative breast cancer (TNBC) patients^[13].

KIT Amplification

Biological Impact

KIT is a proto-oncogene that encodes a type 3 transmembrane receptor tyrosine kinase. Activation of KIT through dimerization and autophosphorylation upon binding by its ligand results in increased intracellular PI3K/AKT/mTOR, MAPK/ERK and JAK/STAT signaling pathways to promote cell proliferation and survival^[14]. KIT activating mutations are frequently found in 80 - 90% of gastrointestinal stromal tumors (GISTs) which distributed over multiple exons with different frequencies (exons 11 (66.1%), exon 9 (13%), exon 13 (1.2%), and exon 17 (0.6%))^{[15][16]}.

Therapeutic and prognostic relevance

The NCCN guidelines for cutaneous melanoma suggested KIT hotspots mutations which located in exon 11 and exon 13 (eg. W557, V559, L576P, K642E) have a high level of sensitivity to KIT inhibitors (imatinib, sunitinib, nilotinib)^{[17][18][19]}. However, KIT exon 17 mutations (eg. D816H) and KIT amplification appeared to be resistant to KIT inhibitors in patients with melanoma.

The efficacies of several U.S. FDA-approved KIT-targeting tyrosine kinase inhibitors (TKIs) such as imatinib, sunitinib, regorafenib, and ripretinib are strongly dependent on the location of the activating KIT mutations^{[20][21][22][23][24][25][26][27][28][29]}. Patients with GIST harboring KIT exon 9 mutations showed intermediate sensitivity to imatinib and had better relapse-free survival and overall survival (OS) compared with patients carrying KIT exon 11 mutations^[21].

Ponatinib and dasatinib yielded a disease control rate and partial control rate of 67% and 32%, respectively, in GIST patients harboring KIT exon 11 mutations (DOI: 10.1200/jco.2015.33.15_suppl.10535, 10.1200/jco.2011.29.15_suppl.10006). Results from a Phase II trial involving melanoma showed 38.5% response rate to nilotinib in patients harboring KIT exon 11 mutations^[30].

ACT Onco[®] + Report

Both KIT and PDGFRA overexpression were associated with high tumor grade, high proliferation index, and poor outcome in patients with the serous type of ovarian carcinoma^[31].

The newly developing agents such as avapritinib (BLU-285) and investigational AZD3229 all showed the potential to be better inhibitors for clinically relevant KIT/PDGFRA mutations in GIST^[32].

KIT mutations have been determined as an inclusion criterion for the trials evaluating dasatinib, avapritinib, sunitinib, nilotinib, ponatinib, regorafenib, and ripretinib efficacies in melanoma, solid tumors, GIST, systemic mastocytosis (AdvSM), and relapsed or refractory myeloid malignancies (NCT00700882, NCT04771520, NCT03465722, NCT02693535, NCT02561988, NCT01028222, NCT01099514, NCT03171389, NCT02272998, NCT02501551, and NCT02571036).

A phase II trial of imatinib in melanoma showed that patients with KIT amplification had lower disease control rate compared with patients carrying KIT mutations (18% amplified vs. 77% mutated)^[19].

LYN Amplification

Biological Impact

The LYN Proto-Oncogene, Src Family Tyrosine Kinase (LYN) gene encodes a non-receptor tyrosine protein kinase of the Src family (SFK)^[33]. LYN plays an important role in the regulation of immune responses, hematopoiesis, signal transduction of growth factors and cytokines and is activated in the cellular response to DNA damage and genotoxic agents^{[34][35][36][37]}. LYN has been described to promote tumor growth, invasion, epithelial to mesenchymal transition (EMT) and ERK signaling in different cancer types^{[38][39][40]}. Amplification of LYN has been identified in prostate cancer, breast cancer and ovarian cancer (cBioPortal).

Therapeutic and prognostic relevance

LYN could be pharmacologically targeted with Src-kinase inhibitors. Results of preclinical studies showed that dasatinib, a dual-specificity tyrosine kinase inhibitor of ABL and the Src-family tyrosine kinases, exerted antitumor activity in LYN-expressing breast cancer cells^[38].

NBN Amplification

Biological Impact

The NBN gene encodes a component of the MRE11-RAD50-NBN (MRN) complex, which involves in DNA double-strand break sensing and repair^[41]. NBN mutation is related to Nijmegen breakage syndrome, increased cancer incidence and ionizing radiation sensitivity^{[41][42]}. NBN mutations have been found in various cancers, including cholangiocarcinoma, hepatocellular carcinoma^[43], prostate cancer^[44], leukemia, lymphoma^[45], and triple-negative breast cancer^[46].

Therapeutic and prognostic relevance

In a phase II trial (ARIEL2), an ovarian cancer patient harboring a NBN germline mutation showed responses to rucaparib treatment^[47]. NBN loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in ovarian cancer^[48]; the trials evaluating rucaparib efficacy in ovarian cancer^[49] or prostate cancer^[50]; the trials evaluating talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795) or lung cancer (NCT03377556), and the trials evaluating niraparib efficacy in pancreatic cancer (NCT03553004) or any malignancy (except prostate) cancer (NCT03207347).

Germline and somatic mutations in homologous recombination genes, including NBN, have been suggested to be prognostic biomarkers for platinum-based treatment response and superior survival in patients with ovarian, fallopian tube, peritoneal carcinomas and pancreatic cancer^{[51][52]}.

ACT Onco[®] + Report

In a retrospective study of localized prostate cancer, NBN gene amplification has been demonstrated to associate with overall tumor genomic instability and lower biochemical relapse-free rate following image-guided radiotherapy (IGRT)^[53].

Another retrospective study showed that amplification of the NBN gene is associated with protein overexpression and mostly correlated with poor prognosis in several cancer types, including ovarian cancer, breast invasive carcinoma, uterine corpus endometrial carcinoma, and sarcoma. Besides, in vivo and in vitro assays demonstrated that amplification of the NBN gene could induce cisplatin and PARP inhibitor resistance in breast and ovarian cancer cells^[54].

PDGFRA Amplification

Biological Impact

The PDGFRA gene encodes for the protein platelet-derived growth factor alpha (PDGFRA). The Ligand binding to the extracellular domain of PDGFRA induces receptor dimerization, enabling autophosphorylation of specific tyrosine residues and subsequently results in the activation of downstream pathways such as RAS-MAPK, PI3K and PLC-γ that are involved in developmental and cellular responses^{[55][56]}. Mutations, insertions, deletions, fusions and genomic amplification of PDGFRA lead to its activation in several tumor types: ~7% of gastrointestinal stromal tumors (GISTs) have PDGFRA activating mutations and these mutations are mutually exclusive from KIT mutations^[57]; activating mutations in PDGFRA have been observed in ~5% of Chinese melanoma patients^[58]; amplification of PDGFRA is the second most frequent receptor tyrosine kinase amplification in glioblastoma (GBM)^{[59][60][61][62][63]}, intimal sarcomas^[64], malignant peripheral nerve sheath tumors^[65], non-small cell lung adenocarcinomas and non-small cell lung squamous cell carcinomas^[66].

Therapeutic and prognostic relevance

A retrospective study showed that either KIT, PDGFRA, or EGFR amplification in glioma at the time of the first diagnosis was associated with an unfavorable overall survival^[67].

In a preclinical study, a PDGFRA-amplified cell line was sensitive to imatinib, sunitinib and sorafenib treatment, demonstrated by disruption of downstream signaling and reduced cell viability in vitro^[68]. Another study also showed that pazopanib could inhibit tumor growth in the PDGFRA-amplified pleomorphic liposarcoma xenograft mouse model^[69].

TP53 Homozygous deletion

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[70]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[71].

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[72].

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[73]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[74].

ACT Onco[®] + Report

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[75][76][77]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[78]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[79][80]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[81].

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Cabozantinib (COMETRIQ)

Cabozantinib is a small molecule inhibitors of multiple tyrosine kinases, including RET, MET, VEGFR-1, -2 and -3, KIT, TRKB, FLT-3, AXL, and TIE-2. Cabozantinib is developed and marketed by Exelixis under the trade names COMETRIQ (capsule) and CABOMETYX (tablet).

- FDA Approval Summary of Cabozantinib (COMETRIQ)

EXAM ^[82] NCT00704730	Thyroid cancer (Approved on 2012/11/29)
	-
	Cabozantinib vs. Placebo [PFS(M): 11.2 vs. 4]

Cabozantinib (CABOMETYX)

Cabozantinib is a small molecule inhibitors of multiple tyrosine kinases, including RET, MET, VEGFR-1, -2 and -3, KIT, TRKB, FLT-3, AXL, and TIE-2. Cabozantinib is developed and marketed by Exelixis under the trade names COMETRIQ (capsule) and CABOMETYX (tablet).

- FDA Approval Summary of Cabozantinib (CABOMETYX)

COSMIC-311 NCT03690388	Differentiated thyroid cancer (dct) (Approved on 2021/09/17)
	-
	Cabozantinib vs. Placebo [PFS(M): 11 vs. 1.9, ORR(%): 18.0 vs. 0]
CHECKMATE-9ER NCT03141177	Renal cell carcinoma (Approved on 2021/01/22)
	-
	Nivolumab + cabozantinib vs. Sunitinib [ORR(%): 55.7 vs. 27.1, PFS(M): 16.6 vs. 8.3, OS(M): NR vs. NR]
CELESTIAL ^[7] NCT01908426	Hepatocellular carcinoma (Approved on 2019/01/14)
	-
	Cabozantinib vs. Placebo [OS(M): 10.2 vs. 8]
CABOSUN ^[83] NCT01835157	Renal cell carcinoma (Approved on 2017/12/09)
	-
	Cabozantinib vs. Sunitinib [PFS(M): 8.6 vs. 5.3]
METEOR ^[84] NCT01865747	Renal cell carcinoma (Approved on 2016/04/25)
	-
	Cabozantinib vs. Everolimus [PFS(M): 7.4 vs. 3.8]

Dasatinib (SPRYCEL)

Dasatinib is an oral Bcr-Abl tyrosine kinase inhibitor (inhibits the "Philadelphia chromosome") and Src family tyrosine kinase inhibitor. Dasatinib is produced by Bristol-Myers Squibb and sold under the trade name SPRYCEL.

- FDA Approval Summary of Dasatinib (SPRYCEL)

DASISION ^[85] NCT00481247	Chronic myeloid leukemia (Approved on 2010/10/28)
	-
	Dasatinib vs. Imatinib [ORR(%): 76.8 vs. 66.2]

ACT Onco[®] + Report

[86] NCT00123474	Chronic myeloid leukemia (Approved on 2007/11/08)
	-
	Dasatinib [ORR(%): 63.0]
[87] NCT00123487	Acute lymphocytic leukemia (Approved on 2006/06/28)
	-
	Dasatinib [ORR(%): 38.0]

Imatinib (GLEEVEC)

Imatinib is an oral, small molecule inhibitor of tyrosine kinase enzymes, namely, the Abelson proto-oncogene (ABL), c-KIT, and platelet-derived growth factor receptor (PDGFR). Imatinib is developed and marketed by Novartis under the trade name GLEEVEC.

- FDA Approval Summary of Imatinib (GLEEVEC)

[88] NCT00022737	Acute lymphocytic leukemia (Approved on 2013/01/25)
	-
	Imatinib [EFS(%): 70]
	Gastrointestinal stromal tumor (Approved on 2012/01/31)
	KIT positive
	Imatinib [RFS(%): 42 (imatinib for 12) 25 (imatinib for 36)]
	Gastrointestinal stromal tumor (Approved on 2009/02/10)
	KIT+
	Imatinib vs. Placebo [RFS(%): 21 vs. 28]
	Myelodysplastic myeloproliferative cancer (Approved on 2006/10/19)
	-
	Imatinib [MCyR(%): 39, CHR(%): 45]
[89]	Acute lymphocytic leukemia (Approved on 2006/10/19)
	Ph+
	Imatinib [MCyR(%): 35, CHR(%): 19]
	Dermatofibrosarcoma protuberans (Approved on 2006/10/19)
	-
	Imatinib [ORR(%): 83.0]
	Systemic mastocytosis (Approved on 2006/10/19)
	-
	Imatinib [CHR(%): 29]
	Chronic eosinophilic leukemia (Approved on 2006/10/19)
	-
	Imatinib [CHR(%): 61]
[90] NCT00471497	Chronic myeloid leukemia (Approved on 2003/05/20)
	Ph+
	Imatinib vs. Nilotinib [MMR(%): 22 vs. 44]
[91] NCT00333840	Chronic myeloid leukemia (Approved on 2003/04/18)
	-
	Imatinib vs. Interferon- α + cytarabine [PFS(%): 81.2 vs. 60.6]
[92] NCT00009906	Gastrointestinal stromal tumor (Approved on 2002/02/01)
	-
	Imatinib [PFS(M): 18.9 (imatinib 400 mg) 23.2 (imatinib 800 mg)]

ACT Onco[®] + Report

Pazopanib (VOTRIENT)

Pazopanib is an oral, small molecule, multi-kinase inhibitor that targets receptor tyrosine kinase including vascular endothelial growth factor receptor-1, -2, -3 (VEGFR-1, -2, -3), platelet-derived growth factor receptor- α , - β (PDGFR- α , - β), c-kit, fibroblast growth factor-1 and -3 (FGFR-1, -3), thereby inhibiting angiogenesis. Pazopanib is developed and marketed by GlaxoSmithKline under the trade name VOTRIENT.

- FDA Approval Summary of Pazopanib (VOTRIENT)

PALETTE ^[93] NCT00753688	Sarcoma (Approved on 2016/04/26)
	-
	Pazopanib vs. Placebo [PFS(M): 4.6 vs. 1.6]
VEG105192 ^[94] NCT00334282	Renal cell carcinoma (Approved on 2009/10/19)
	-
	Pazopanib vs. Placebo [PFS(M): 9.2 vs. 4.2]

Sorafenib (NEXAVAR)

Sorafenib is a small molecule multi-kinase inhibitor that targets multiple kinase families including VEGFR, PDGFRB, and the RAF family kinases. Sorafenib is co-developed and co-marketed by Bayer HealthCare Pharmaceuticals and Onyx Pharmaceuticals under the trade name NEXAVAR.

- FDA Approval Summary of Sorafenib (NEXAVAR)

DECISION ^[95] NCT00984282	Differentiated thyroid carcinoma (Approved on 2013/11/22)
	-
	Sorafenib vs. Placebo [PFS(M): 10.8 vs. 5.8]
SHARP ^[96] NCT00105443	Hepatocellular carcinoma (Approved on 2007/11/16)
	-
	Sorafenib vs. Placebo [OS(M): 10.7 vs. 7.9]
TARGET ^[97] NCT00073307	Renal cell carcinoma (Approved on 2005/12/20)
	-
	Sorafenib vs. Placebo [PFS(D): 167 vs. 84]

Sunitinib (SUTENT)

Sunitinib is an oral, small molecule, multi-kinase inhibitor that targets receptor tyrosine kinase including platelet-derived growth factor receptor- α , - β (PDGFR- α , - β), vascular endothelial growth factor receptors-1, -2, -3 (VEGFR-1, -2, -3), c-kit, Fms-like tyrosine kinase-3 (FLT3), colony stimulating factor receptor type 1 (CSF-1R), and the glial cell-line derived neurotrophic factor receptor (RET), thereby inhibiting angiogenesis. Sunitinib is developed and marketed by Pfizer under the trade name SUTENT.

- FDA Approval Summary of Sunitinib (SUTENT)

^{[98][99][100]} NCT00428597	Pancreatic cancer (Approved on 2011/05/20)
	-
	Sunitinib vs. Placebo [PFS(M): 10.2 vs. 5.4]
^{[101][102]} NCT00083889	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib vs. IFN- α [PFS(W): 47.3 vs. 22]

ACT Onco[®] + Report

[103][104][102] NCT00077974	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib [ORR(%): 34.0]
[104][102] NCT00054886	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib [ORR(%): 36.5]
[105] NCT00075218	Gastrointestinal stromal tumor (Approved on 2006/01/26)
	-
	Sunitinib vs. Placebo [TTP(W): 27.3 vs. 6.4]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
Not Detected							

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ADGRA2	Splice region	-	c.1833+7G>A	NM_032777	-	89.6%	163
ADGRA2	L474W	10	c.1421T>G	NM_032777	-	79.3%	357
BLM	M348I	5	c.1044G>A	NM_000057	COSM1580597	62.2%	1512
CCNB2	V301G	7	c.902T>G	NM_004701	-	18.7%	1174
CTLA4	Y139C	2	c.416A>G	NM_005214	-	6.3%	882
FLT4	G1329S	30	c.3985G>A	NM_182925	-	59.4%	475
IRS1	S685_S686dup	1	c.2054_2059dup	NM_005544	-	76.0%	416
KDR	E1245A	28	c.3734A>C	NM_002253	-	67.7%	3238
MUC16	T9958S	3	c.29872A>T	NM_024690	-	57.2%	580
SPEN	R1475Q	11	c.4424G>A	NM_015001	-	12.7%	1429
TAP2	Splice region	11	c.1932C>T	NM_018833	-	14.7%	1542
USH2A	A3973S	61	c.11917G>T	NM_206933	-	80.6%	1766
XIAP	D367G	6	c.1100A>G	NM_001167	-	92.5%	67

Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Dec 22, 2022
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11153970A
- Collection site: Mediastium
- Examined by: Dr. Chien-Ta Chiang
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 10%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 30%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 860x
- Target Base Coverage at 100x: 94%

RNA test

- Average unique RNA Start Sites per control GSP2: 150

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100x \geq 85\%$ with a mean coverage $\geq 500x$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®]+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

ACT Onco[®] + Report

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師黃靖婷 博士
Ching-Ting Huang Ph.D.
檢字第 016511 號

CT Huang

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC1B1*
SLC1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Not Applicable.

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS



1: Pazopanib; 2: Imatinib, Sunitinib, Pazopanib, Dasatinib; 3: Cabozantinib; 4: Sunitinib, Pazopanib, Cabozantinib; 5: Sunitinib, Pazopanib; 6: Sunitinib; 7: Imatinib, Sunitinib, Pazopanib, Sorafenib, Dasatinib

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 23172303; 2013, J Biochem;153(1):13-9
Vascular endothelial growth factor and its receptor system: physiological functions in angiogenesis and pathological roles in various diseases.
2. PMID: 12778165; 2003, Nat Med;9(6):669-76
The biology of VEGF and its receptors.
3. PMID: 21382095; 2011, Neuropathology;31(6):583-8
Amplification of the PDGFRA, KIT and KDR genes in glioblastoma: a population-based study.
4. PMID: 31604903; 2020, Oncologist;25(1):e39-e47
The Pan-Cancer Landscape of Coamplification of the Tyrosine Kinases KIT, KDR, and PDGFRA.
5. PMID: 23028128; 2012, Cold Spring Harb Perspect Med;2(10):
Anti-VEGF therapies in the clinic.
6. PMID: 27718781; 2016, N Engl J Med;375(23):2246-2254
Adjuvant Sunitinib in High-Risk Renal-Cell Carcinoma after Nephrectomy.
7. PMID: 29972759; 2018, N Engl J Med;379(1):54-63
Cabozantinib in Patients with Advanced and Progressing Hepatocellular Carcinoma.
8. PMID: 27160228; 2016, J Natl Compr Canc Netw;14(5):499-502
Antitumor Response of VEGFR2- and VEGFR3-Amplified Angiosarcoma to Pazopanib.
9. PMID: 25639617; 2015, Breast J;21(2):205-7
Refractory angiosarcoma of the breast with VEGFR2 upregulation successfully treated with sunitinib.
10. PMID: 26578684; 2016, Clin Cancer Res;22(8):1940-50
KDR Amplification Is Associated with VEGF-Induced Activation of the mTOR and Invasion Pathways but does not Predict Clinical Benefit to the VEGFR TKI Vandetanib.
11. PMID: 29855279; 2018, BMC Cancer;18(1):618
Apatinib treatment for KIT- and KDR-amplified angiosarcoma: a case report.
12. PMID: 31786466; 2020, Biomed Pharmacother;122():109587
Efficacy and safety of the VEGFR2 inhibitor Apatinib for metastatic soft tissue sarcoma: Chinese cohort data from NCT03121846.
13. PMID: 22170730; 2012, Genes Chromosomes Cancer;51(4):375-83
Increased gene copy number of KIT and VEGFR2 at 4q12 in primary breast cancer is related to an aggressive phenotype and impaired prognosis.
14. PMID: 17546049; 2007, Oncogene;26(54):7560-8
KIT oncogenic signaling mechanisms in imatinib-resistant gastrointestinal stromal tumor: PI3-kinase/AKT is a crucial survival pathway.
15. PMID: 15365079; 2004, J Clin Oncol;22(18):3813-25
Biology of gastrointestinal stromal tumors.
16. PMID: 22089421; 2011, Nat Rev Cancer;11(12):865-78
Gastrointestinal stromal tumours: origin and molecular oncology.
17. PMID: 21690468; 2011, J Clin Oncol;29(21):2904-9
Phase II, open-label, single-arm trial of imatinib mesylate in patients with metastatic melanoma harboring c-Kit mutation or amplification.
18. PMID: 21642685; 2011, JAMA;305(22):2327-34
KIT as a therapeutic target in metastatic melanoma.
19. PMID: 23775962; 2013, J Clin Oncol;31(26):3182-90

ACT Onco[®] + Report

Imatinib for melanomas harboring mutationally activated or amplified KIT arising on mucosal, acral, and chronically sun-damaged skin.

20. PMID: 12181401; 2002, N Engl J Med;347(7):472-80
Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors.
21. PMID: 18955458; 2008, J Clin Oncol;26(33):5352-9
Primary and secondary kinase genotypes correlate with the biological and clinical activity of sunitinib in imatinib-resistant gastrointestinal stromal tumor.
22. PMID: 19164557; 2009, Proc Natl Acad Sci U S A;106(5):1542-7
KIT kinase mutants show unique mechanisms of drug resistance to imatinib and sunitinib in gastrointestinal stromal tumor patients.
23. PMID: 18235121; 2008, J Clin Oncol;26(4):620-5
Long-term results from a randomized phase II trial of standard- versus higher-dose imatinib mesylate for patients with unresectable or metastatic gastrointestinal stromal tumors expressing KIT.
24. PMID: 16098458; 2005, Eur J Cancer;41(12):1751-7
Outcome of patients with advanced gastro-intestinal stromal tumours crossing over to a daily imatinib dose of 800 mg after progression on 400 mg.
25. PMID: 15451219; 2004, Lancet;364(9440):1127-34
Progression-free survival in gastrointestinal stromal tumours with high-dose imatinib: randomised trial.
26. PMID: 23177515; 2013, Lancet;381(9863):295-302
Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial.
27. PMID: 25641662; 2015, Cancer;121(9):1405-13
Clinical outcomes of patients with advanced gastrointestinal stromal tumors: safety and efficacy in a worldwide treatment-use trial of sunitinib.
28. PMID: 19282169; 2009, Eur J Cancer;45(11):1959-68
Clinical evaluation of continuous daily dosing of sunitinib malate in patients with advanced gastrointestinal stromal tumour after imatinib failure.
29. PMID: 32511981; 2020, Lancet Oncol;21(7):923-934
Ripretinib in patients with advanced gastrointestinal stromal tumours (INVICTUS): a double-blind, randomised, placebo-controlled, phase 3 trial.
30. PMID: 28327988; 2017, Ann Oncol;28(6):1380-1387
Efficacy and safety of nilotinib in patients with KIT-mutated metastatic or inoperable melanoma: final results from the global, single-arm, phase II TEAM trial.
31. PMID: 15583695; 2004, Br J Cancer;91(12):2048-55
Genetic alterations and protein expression of KIT and PDGFRA in serous ovarian carcinoma.
32. PMID: 32350132; 2020, Sci Transl Med;12(541):
Discovery and pharmacological characterization of AZD3229, a potent KIT/PDGFRα inhibitor for treatment of gastrointestinal stromal tumors.
33. PMID: 9202419; 1997, Int J Biochem Cell Biol;29(3):397-400
Lyn, a src-like tyrosine kinase.
34. PMID: 19290919; 2009, Immunol Rev;228(1):23-40
Multiple roles of Lyn kinase in myeloid cell signaling and function.
35. PMID: 11825908; 2002, J Biol Chem;277(16):13732-8
Phosphatidylinositol 3-kinase and Src family kinases are required for phosphorylation and membrane recruitment of Dok-1 in c-Kit signaling.
36. PMID: 11435302; 2001, Blood;98(2):343-50
Lyn is required for normal stem cell factor-induced proliferation and chemotaxis of primary hematopoietic cells.
37. PMID: 10574731; 1999, J Exp Biol;202 Pt 24():3541-53

ACT Onco[®] + Report

Redox control in development and evolution: evidence from colonial hydroids

38. PMID: 20215510; 2010, Cancer Res;70(6):2296-306
LYN is a mediator of epithelial-mesenchymal transition and a target of dasatinib in breast cancer.
39. PMID: 22490227; 2012, Clin Cancer Res;18(10):2850-60
Lyn kinase mediates cell motility and tumor growth in EGFRvIII-expressing head and neck cancer.
40. PMID: 22731636; 2012, Mol Cancer;11():43
Lyn is involved in CD24-induced ERK1/2 activation in colorectal cancer.
41. PMID: 9590181; 1998, Cell;93(3):477-86
The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response.
42. PMID: 9590180; 1998, Cell;93(3):467-76
Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome.
43. PMID: 24349281; 2013, PLoS One;8(12):e82426
Mutation inactivation of Nijmegen breakage syndrome gene (NBS1) in hepatocellular carcinoma and intrahepatic cholangiocarcinoma.
44. PMID: 22864661; 2012, Fam Cancer;11(4):595-600
Identification of a novel NBN truncating mutation in a family with hereditary prostate cancer.
45. PMID: 21923652; 2011, Br J Haematol;155(4):468-76
Promising therapy results for lymphoid malignancies in children with chromosomal breakage syndromes (Ataxia teleangiectasia or Nijmegen-breakage syndrome): a retrospective survey.
46. PMID: 26083025; 2015, PLoS One;10(6):e0130393
Prevalence of Germline Mutations in Genes Engaged in DNA Damage Repair by Homologous Recombination in Patients with Triple-Negative and Hereditary Non-Triple-Negative Breast Cancers.
47. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
48. PMID: 30353044; 2018, Br J Cancer;119(11):1401-1409
Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes.
49. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
50. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
51. PMID: 24240112; 2014, Clin Cancer Res;20(3):764-75
Germline and somatic mutations in homologous recombination genes predict platinum response and survival in ovarian, fallopian tube, and peritoneal carcinomas.
52. PMID: 29961768; 2019, Genet Med;21(1):213-223
Germline cancer susceptibility gene variants, somatic second hits, and survival outcomes in patients with resected pancreatic cancer.
53. PMID: 25415046; 2014, Oncotarget;5(22):11081-90
NBN gain is predictive for adverse outcome following image-guided radiotherapy for localized prostate cancer.
54. PMID: 32226530; 2020, Theranostics;10(9):3939-3951
Copy Number Amplification of DNA Damage Repair Pathways Potentiates Therapeutic Resistance in Cancer.

ACT Onco[®] + Report

55. PMID: 18483217; 2008, Genes Dev;22(10):1276-312
Role of platelet-derived growth factors in physiology and medicine.
56. PMID: 3013421; 1986, Cell;46(2):155-69
The biology of platelet-derived growth factor.
57. PMID: 20023271; 2010, Am J Clin Pathol;133(1):149-55
Detection of c-KIT and PDGFRA gene mutations in gastrointestinal stromal tumors: comparison of DHPLC and DNA sequencing methods using a single population-based cohort.
58. PMID: 24132921; 2013, Clin Cancer Res;19(24):6935-42
Large-scale analysis of PDGFRA mutations in melanomas and evaluation of their sensitivity to tyrosine kinase inhibitors imatinib and crenolanib.
59. PMID: 18772890; 2008, Nature;455(7216):1061-8
Comprehensive genomic characterization defines human glioblastoma genes and core pathways.
60. PMID: 19915670; 2009, PLoS One;4(11):e7752
Glioblastoma subclasses can be defined by activity among signal transduction pathways and associated genomic alterations.
61. PMID: 22323597; 2012, Proc Natl Acad Sci U S A;109(8):3041-6
Intratumoral heterogeneity of receptor tyrosine kinases EGFR and PDGFRA amplification in glioblastoma defines subpopulations with distinct growth factor response.
62. PMID: 20129251; 2010, Cancer Cell;17(1):98-110
Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1.
63. PMID: 20889717; 2010, Genes Dev;24(19):2205-18
PDGFRA gene rearrangements are frequent genetic events in PDGFRA-amplified glioblastomas.
64. PMID: 20685895; 2010, Cancer Res;70(18):7304-14
Coactivated platelet-derived growth factor receptor {alpha} and epidermal growth factor receptor are potential therapeutic targets in intimal sarcoma.
65. PMID: 16357008; 2006, Carcinogenesis;27(3):664-71
Mutation and expression of PDGFRA and KIT in malignant peripheral nerve sheath tumors, and its implications for imatinib sensitivity.
66. PMID: 19755855; 2009, Cancer Biol Ther;8(21):2042-50
Amplification of chromosomal segment 4q12 in non-small cell lung cancer.
67. PMID: 17189383; 2006, Mol Cancer Res;4(12):927-34
Amplification of KIT, PDGFRA, VEGFR2, and EGFR in gliomas.
68. PMID: 19366796; 2009, Cancer Res;69(9):3937-46
Ligand-dependent platelet-derived growth factor receptor (PDGFR)-alpha activation sensitizes rare lung cancer and sarcoma cells to PDGFR kinase inhibitors.
69. PMID: 30060824; 2018, Tissue Cell;53():30-36
Doxorubicin-resistant pleomorphic liposarcoma with PDGFRA gene amplification is targeted and regressed by pazopanib in a patient-derived orthotopic xenograft mouse model.
70. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
71. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
72. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-

ACT Onco[®] + Report

Line Therapy Within 3 Months.

73. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
74. PMID: 25669829; 2015, Ann Oncol;26(5):1012-8
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
75. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
76. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
77. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
78. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
79. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
80. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
81. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
82. PMID: 24002501; 2013, J Clin Oncol;31(29):3639-46
Cabozantinib in progressive medullary thyroid cancer.
83. PMID: 28199818; 2017, J Clin Oncol;35(6):591-597
Cabozantinib Versus Sunitinib As Initial Targeted Therapy for Patients With Metastatic Renal Cell Carcinoma of Poor or Intermediate Risk: The Alliance A031203 CABOSUN Trial.
84. PMID: 26406150; 2015, N Engl J Med;373(19):1814-23
Cabozantinib versus Everolimus in Advanced Renal-Cell Carcinoma.
85. PMID: 20525995; 2010, N Engl J Med;362(24):2260-70
Dasatinib versus imatinib in newly diagnosed chronic-phase chronic myeloid leukemia.
86. PMID: 18541900; 2008, J Clin Oncol;26(19):3204-12
Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia.
87. PMID: 17496201; 2007, Blood;110(7):2309-15
Dasatinib induces rapid hematologic and cytogenetic responses in adult patients with Philadelphia chromosome positive acute lymphoblastic leukemia with resistance or intolerance to imatinib: interim results of a phase 2 study.
88. PMID: 19805687; 2009, J Clin Oncol;27(31):5175-81
Improved early event-free survival with imatinib in Philadelphia chromosome-positive acute lymphoblastic leukemia: a children's oncology group study.
89. PMID: 12200353; 2002, Blood;100(6):1965-71
A phase 2 study of imatinib in patients with relapsed or refractory Philadelphia chromosome-positive acute lymphoid leukemias.
90. PMID: 21856226; 2011, Lancet Oncol;12(9):841-51

ACT Onco[®] + Report

Nilotinib versus imatinib for the treatment of patients with newly diagnosed chronic phase, Philadelphia chromosome-positive, chronic myeloid leukaemia: 24-month minimum follow-up of the phase 3 randomised ENESTnd trial.

91. PMID: 18256322; 2008, Blood;111(8):4022-8
Imatinib pharmacokinetics and its correlation with response and safety in chronic-phase chronic myeloid leukemia: a subanalysis of the IRIS study.
92. PMID: 28196207; 2017, JAMA Oncol;3(7):944-952
Correlation of Long-term Results of Imatinib in Advanced Gastrointestinal Stromal Tumors With Next-Generation Sequencing Results: Analysis of Phase 3 SWOG Intergroup Trial S0033.
93. PMID: 22595799; 2012, Lancet;379(9829):1879-86
Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial.
94. PMID: 20100962; 2010, J Clin Oncol;28(6):1061-8
Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial.
95. PMID: 24768112; 2014, Lancet;384(9940):319-28
Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial.
96. PMID: 18650514; 2008, N Engl J Med;359(4):378-90
Sorafenib in advanced hepatocellular carcinoma.
97. PMID: 17189398; 2006, Clin Cancer Res;12(24):7271-8
Sorafenib for the treatment of advanced renal cell carcinoma.
98. PMID: 27924459; 2016, Target Oncol;11(6):815-824
Patient-Reported Outcomes and Quality of Life with Sunitinib Versus Placebo for Pancreatic Neuroendocrine Tumors: Results From an International Phase III Trial.
99. PMID: 27836885; 2017, Ann Oncol;28(2):339-343
Sunitinib in pancreatic neuroendocrine tumors: updated progression-free survival and final overall survival from a phase III randomized study.
100. PMID: 21306237; 2011, N Engl J Med;364(6):501-13
Sunitinib malate for the treatment of pancreatic neuroendocrine tumors.
101. PMID: 17227905; 2007, Oncologist;12(1):107-13
Food and Drug Administration drug approval summary: Sunitinib malate for the treatment of gastrointestinal stromal tumor and advanced renal cell carcinoma.
102. PMID: 27238653; 2016, Eur Urol;70(6):1006-1015
Early Tumour Shrinkage: A Tool for the Detection of Early Clinical Activity in Metastatic Renal Cell Carcinoma.
103. PMID: 16757724; 2006, JAMA;295(21):2516-24
Sunitinib in patients with metastatic renal cell carcinoma.
104. PMID: 25577718; 2015, Eur Urol;67(5):952-8
Depth of remission is a prognostic factor for survival in patients with metastatic renal cell carcinoma.
105. PMID: 17046465; 2006, Lancet;368(9544):1329-38
Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial.