

REPORT SUMMARY

PATIENT AND SAMPLE INFORMATION2

VARIANT(S) WITH CLINICAL RELEVANCE2

THERAPEUTIC IMPLICATIONS.....3

REPORT DETAILS

VARIANT INTERPRETATION5

US FDA-APPROVED DRUG(S)12

ONGOING CLINICAL TRIALS33

DETAILED TEST RESULTS35

HOTSPOT GENOTYPES.....37

TEST DETAILS39

ACTOnco[®]+ GENE LIST43

DISCLAIMER44

APPENDIX

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS.....46

REFERENCES.....50

PATIENT AND SAMPLE INFORMATION

PATIENT

Name: 呂富美
Gender: Female
Date of Birth: Sep 30, 1964
Patient ID: 19283126
Diagnosis: Cholangiocarcinoma

SPECIMEN

Type: FFPE tissue
Date received: Nov 01, 2021
Collection site: Liver
Specimen ID: S10935424
Lab ID: AA-21-04509
D/ID: NA

ORDERING PHYSICIAN

Name: 陳明晃醫師
Facility: 臺北榮總
Tel: 886-228712121
Address: 臺北市北投區石牌路二段 201 號

VARIANT(S) WITH CLINICAL RELEVANCE

Only variant(s) with clinical significance are listed. See the "DETAILED TEST RESULTS" section for full details.

SINGLE NUCLEOTIDE AND SMALL INDEL VARIANTS

Gene	Amino Acid Change	Coverage	Allele Frequency	COSMIC ID
<i>BRAF</i>	G469A	1631	17.5%	COSM460
<i>IDH1</i>	R132C	692	15.2%	COSM28747
<i>PBRM1</i>	F1337fs	903	37.0%	-
<i>TSC1</i>	Splice donor	928	51.9%	-

COPY NUMBER VARIANTS (CNVs)

Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on **58%** tumor purity.

Amplification (Copy number ≥ 8)

Chr	Gene	Copy Number
ND	ND	ND

Homozygous deletion (Copy number=0)

Chr	Gene
chr9	<i>CDKN2A</i>
chr22	<i>NF2</i>

Heterozygous deletion (Copy number=1)

Chr	Gene
chr9	<i>PTCH1</i>
chr13	<i>BRCA2</i>
chr15	<i>RAD51</i>

ND, Not Detected

TUMOR MUTATIONAL BURDEN (TMB)

22.8 muts/Mb (TMB-High)

Muts/Mb, mutations per megabase

Note:

TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is < 30%.

MICROSATELLITE INSTABILITY (MSI)

Microsatellite stable (MSS)

Variant Analysis:

醫檢師張筑芎 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



Sign Off

醫檢師張筑芎 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 1		
IDH1 R132C	Ivosidenib	sensitive
Level 3B		
BRAF G469A	Selumetinib	sensitive
TSC1 Splice donor	Everolimus, Temsirolimus	sensitive
CDKN2A Homozygous deletion	Abemaciclib, Palbociclib, Ribociclib	sensitive
RAD51 Heterozygous deletion	Niraparib, Rucaparib	sensitive
PTCH1 Heterozygous deletion	Sonidegib, Vismodegib	sensitive
Level 4		
IDH1 R132C	Bevacizumab, Dasatinib, Olaparib, Rucaparib, Talazoparib, Sunitinib	sensitive
BRAF G469A	Trametinib	sensitive
NF2 Homozygous deletion	Everolimus	sensitive
BRCA2 Heterozygous deletion	Olaparib, Rucaparib	sensitive
RAD51 Heterozygous deletion	Olaparib	sensitive
BRAF G469A	Cetuximab, Panitumumab, Vemurafenib	resistant

† Refer to "ONGOING CLINICAL TRIALS" section for detailed trial information.

Note: Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence.

Level	Description
1	FDA-recognized biomarker predictive of response to an FDA approved drug in this indication
2	Standard care biomarker (recommended as standard care by the NCCN or other expert panels) predictive of response to an FDA approved drug in this indication
3	A Biomarkers that predict response or resistance to therapies approved by the FDA or professional societies for a different type of tumor
	B Biomarkers that serve as inclusion criteria for clinical trials
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports or preclinical studies

IMMUNE CHECKPOINT INHIBITORS (ICI) THERAPIES

Approved for PATIENT's Tumor Type

Therapies	Genomic Alterations	Effect	Ongoing Clinical Trials
Pembrolizumab Dostarlimab-gxly	TMB-High (22.8 muts/Mb)	Sensitive	NCT02628067

Approved for OTHER Tumor Types

Therapies	Genomic Alterations	Effect	Ongoing Clinical Trials
Durvalumab Nivolumab Ipilimumab Atezolizumab Avelumab Cemiplimab-rwlc	TMB-High (22.8 muts/Mb)	Sensitive	NCT04589845

TMB, Tumor Mutational Burden; Muts/Mb, mutations per megabase

Genomic markers and alterations that are associated with response to ICI therapies

Positive Biomarker	Negative Biomarker
TMB-H: Yes	EGFR aberration: ND
MSI-H: ND	MDM2/MDM4 amplification: ND
MMR biallelic inactivation: ND	STK11 biallelic inactivation: ND
PBRM1 biallelic inactivation: ND	PTEN biallelic inactivation: ND
SERPINB3/SERPINB4 mutation: ND	B2M biallelic inactivation: ND
	JAK1/2 biallelic inactivation: ND

MMR, mismatch repair; ND, not detected

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

VARIANT INTERPRETATION

Tumor mutational burden (TMB): High (22.8 mutations / Mb)

The patient's tumor harbors 22.8 mutations / Mb and is classified as high tumor mutational burden (TMB). High TMB is a potential biomarker that predicts response to immune checkpoint inhibitors, including anti-CTLA-4^{[1][2][3]} and anti-PD-1^{[3][4]} in melanoma, anti-PD-1 in non-small cell lung cancer (NSCLC)^[5] and colorectal cancer (CRC)^[6], cutaneous squamous cell carcinoma (CSCC)^[7], and anti-PD-L1 therapy in bladder cancer^[8]. Of note, the U.S. FDA has approved tumor mutational burden-high (TMB-H) as a predictive biomarker for pembrolizumab in adult and pediatric patients with unresectable or metastatic solid tumor who have progressed following prior treatment and have no satisfactory alternative treatment options. CRCs with defects in mismatch-repair (MMR) are more susceptible to PD-1 blockade^[6]. High mutation load is associated with shorter overall survival in lung cancer^[9] and breast cancer^[10] patients.

BRAF G469A

Biological Impact

BRAF is a serine/threonine kinase that belongs to the RAF family. The protein plays an essential role in the regulation of mitogen-activated protein kinase (MAPK) cascade, which affects a range of cellular response including cell division, differentiation, and secretion^{[11][12]}. Mutations in the BRAF gene, most commonly the V600 residue, are the most frequently identified oncogenic mutations in melanomas, and have been identified in several types of cancers including non-Hodgkin lymphoma, thyroid cancers, non-small cell lung carcinoma, hairy cell leukemia, glioma, gastrointestinal stromal tumor, and colorectal cancers (CRCs)^{[13][14]}. Of note, in the vast majority of cases, BRAF mutations are non-overlapping with other oncogenic mutations (e.g., NRAS mutations, KIT mutations, etc.) found in melanoma. V600E has been determined to be an activating mutation, which results in enhanced BRAF kinase activity and constitutive activation of downstream MEK/ERK signaling cascade^{[15][16]}.

BRAF G469A mutation occurred at the protein kinase domain of the BRAF protein and has been shown to increase BRAF kinase activity and promote downstream signaling in the MAPK pathway^{[17][13]}.

Therapeutic and prognostic relevance

A retrospective study indicated that similar to other BRAF kinase domain mutation subtypes, BRAF non-V600E mutations (G469A included) also predicts a less benefit of anti-EGFR monoclonal antibody treatment in patients with heavily-pretreated colorectal cancer^[18].

In a Phase II trial (NCI-MATCH), trametinib resulted in stable disease in a patient with lung adenocarcinoma harboring BRAF G469A, who had remained on therapy for 20 months without progression^[19].

The preclinical study demonstrated that compared to trametinib or dabrafenib single treatment, a combined trametinib and dabrafenib treatment enhances and prolongs the ERK inhibition and antiproliferative effect in BRAF G469A-expressing NSCLC cell line^[20].

Meanwhile, the mutation BRAF G469A in metastatic melanoma cell lines has shown weak responsiveness to vemurafenib^[21], and vemurafenib treatment did not show efficacy in patients with advanced solid tumors harboring BRAF G469A (NCT02304809, NCT02091141)^{[22][23]}. Abraxane was shown as a promising therapeutic approach in preclinical assay^[21].

BRAF activating mutations have been determined as an inclusion criterion for the trials evaluating selumetinib efficacies in cancers (NCT01089101, NCT00888134, NCT00866177, and NCT00936221).

IDH1 R132C

Biological Impact

IDH1 encodes the isocitrate dehydrogenase 1 (IDH1), an enzyme that catalyzes the conversion of isocitrate to alpha-ketoglutarate (alpha-KG), a crucial step in the tricarboxylic acid (TCA) cycle. Mutations in IDH1 convert alpha-KG to 2-HG, which is an oncogenic metabolite^[24]. Missense mutations of IDH1/2, including the substitution of the amino acids arginine 132 in IDH1 and arginine 172 or 142 in IDH2, leads to accumulate intracellular D-2-hydroxyglutarate (D-2HG) and alter epigenetic regulation, cancer cell differentiation and metabolism^{[25][26][27][28]}. Mutations in IDH1 and IDH2 have been reported a wide range of cancers, including prostate cancer^[29], chondrosarcoma^[30], glioma^[31], cholangiocarcinoma^[32] and acute myeloid leukemia (AML)^[33].

R132C mutation is located in the active site of the IDH1 protein^[34] and has been characterized to be oncogenic to promote tumor formation and progression by increasing conversion of the alpha-ketoglutarate to onco-metabolite 2-HG^[35].

Therapeutic and prognostic relevance

Ivosidenib, the IDH1 inhibitor has been approved by the U.S. Food and Drug Administration (U.S. FDA) for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) carrying a susceptible IDH1 mutation (R132C, R132H, R132G, R132S, and R132L) as detected by an FDA-approved test. Results of a Phase I clinical study showed that IDH305 and AG-120 (Ivosidenib), inhibitors explicitly targeting the mutated IDH1, demonstrate efficacy in patients AML harboring IDH1 R132 mutations^[36]. In another clinical study (n=63), glioma patients harboring IDH1 R132 mutations showed favorable overall survival when treated with bevacizumab or sunitinib^[37]. Of note, in August, 2021, U.S. FDA has also approved ivosidenib for the treatment of adult patients with locally advanced or metastatic cholangiocarcinoma with IDH1 mutation (R132C, R132H, R132G, R132S, and R132L) based on the results of Study AG120-C-005 (NCT02989857)^[38].

A combination therapy consists of vandetanib, a multi-tyrosine kinase inhibitor, temozolomide, and radiotherapy demonstrated a significant increased PFS and OS in glioblastoma patients harboring IDH1 R132H compared to glioblastoma patients without IDH1 R132H^[39].

Preclinical studies in intrahepatic cholangiocarcinoma, glioma, sarcoma, and AML also demonstrated the sensitivity of R132C/H-mutant tumors to multi-kinase inhibitors such as saracatinib and dasatinib^[40], PARP inhibitors^{[41][42]}, as well as IDH1 inhibitor AGI-5198^{[43][44]}. Small molecule inhibitors of IDH1, which have shown activity in the preclinical settings in IDH1-mutant glioma and leukemia cells, are currently in clinical trials^{[43][27]}.

In glioblastoma patients, IDH1 R132 mutation is an independent prognostic factor for improved overall survival^[45].

PBRM1 F1337fs

Biological Impact

The PBRM1 gene encodes the protein BAF180 tumor suppressor, which is a component of the nucleosome-remodeling complex switching defective/sucrose non-fermenting (SWI/SNF)^[46]. Loss of PBRM1 activity is associated with chromosomal instability^[47]. PBRM1, BAP1 and SETD2 are three frequently altered tumor suppressor genes on chromosome 3p in a region that is deleted in over 90% of clear cell renal cell carcinoma (ccRCC)^{[48][49]}.

F1337fs mutation results in a change in the amino acid sequence beginning at 1337, likely to cause premature truncation of the functional PBRM1 protein (UniProtKB). This mutation is predicted to lead to a loss of PBRM1 protein function, despite not being characterized in the literature.

Therapeutic and prognostic relevance

Biallelic loss of PBRM1 has been shown to correlate with clinical benefit in clear cell renal cell carcinoma (ccRCC)^[50], melanoma, lung cancer, bladder cancer, and head and neck squamous carcinoma (HNSCC) patients^[51] treated with immune checkpoint inhibitors.

Decreased expression of PBRM1 has been shown to predict unfavorable clinical outcome in patients with ccRCC^[52].

TSC1 Splice donor

Biological Impact

The tuberous sclerosis complex 1 (TSC1) gene encodes a tumor suppressor, hamartin, a key negative regulator of the mammalian target of rapamycin (mTOR) pathway^{[53][54]}. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis^{[55][56][57]}, while LOH in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)^[58] and endometrial cancer^[59]. Loss of single TSC1 allele (haploinsufficiency) may provide a growth advantage to bladder epithelial cells, contributing to bladder cancer development^[60]. Both TSC1 and TSC2 mutations cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC), in which individuals develop a variety of benign but often

progressive neoplasms^[61].

TSC1 c.2041+1G>A is a variant located at the splice donor region, which may result in the exon skipping.

Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple neoplasms, such as bladder tumors^[62], gastric, sarcoma, thyroid cancer, and HNSCC^[63]. There were case reports demonstrated the efficacy of sirolimus in malignant uterine perivascular epithelioid cell tumors (PEComa) patients harboring mutations/deletions in TSC1 and TSC2 genes, and temsirolimus in PEComa patients with hyperactivated mTOR pathway. Genomic profiling analysis of GOG248, a Phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer showed that mutations in AKT1, TSC1 and TSC2 may predict clinical benefit from temsirolimus^[64]. Recent studies indicate that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets^[65].

BRCA2 Heterozygous deletion

Biological Impact

The BRCA2 gene encodes a tumor suppressor involved in the homologous recombination pathway for double-strand DNA repair^[66]. BRCA2 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[67]. BRCA2 germline mutations confer an increased lifetime risk of developing breast, ovarian, prostate and pancreatic cancer, limited reports of related gastric cancer, and Fanconi anemia subtype D1-associated risk of brain cancer, medulloblastoma, pharyngeal cancer, chronic lymphocytic leukemia and acute myeloid leukemia^[68]. Somatic mutations in BRCA2 are highest in colorectal, non-small cell lung cancer (NSCLC), and ovarian cancers^[69].

Therapeutic and prognostic relevance

The U.S. FDA has approved olaparib in advanced ovarian cancer under several settings including (1) first-line maintenance treatment for patients with deleterious or suspected deleterious germline or somatic BRCA mutation who are in complete or partial response to first-line platinum-based chemotherapy^[70]; (2) in combination with bevacizumab as first-line maintenance treatment for patients with homologous recombination deficiency (HRD)-positive status^[71]; (3) maintenance treatment for patients with germline BRCA-mutated recurrent ovarian cancer who are in complete or partial response to platinum-based chemotherapy^{[72][73]}; (4) treatment for patients with germline BRCA-mutated advanced ovarian cancer who have been treated with three or more prior lines of chemotherapy^[74]. In addition, olaparib has also been approved in patients with deleterious or suspected deleterious germline BRCA-mutated, HER2-negative metastatic breast cancer who have been treated with chemotherapy in either neoadjuvant, adjuvant, or metastatic setting^[75] and germline BRCA-mutated metastatic pancreatic cancer^[76]. Of note, in May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

(HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[77].

Rucaparib has been approved for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy^[78] and patients with BRCA-mutated epithelial ovarian, fallopian tube, or primary peritoneal cancer, who have been treated with two or more chemotherapies^[79]. In May 2020, the U.S. FDA also approved rucaparib to treat adult patients with a deleterious BRCA mutation-associated metastatic castration-resistant prostate cancer (mCRPC) who have been treated with androgen receptor-directed therapy and a taxane-based chemotherapy (TRITON2, NCT02952534).

The U.S. FDA also approved niraparib for the maintenance treatment of patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in response to platinum-based chemotherapy^{[80][81]} and patients who have been treated with three or more prior lines of chemotherapy and associated with HRD positive status^[82]. In addition, talazoparib for patients with deleterious or suspected deleterious germline BRCA-mutated, HER2 negative locally advanced or metastatic breast cancer^[83].

CDKN2A Homozygous deletion

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein^{[84][85]} whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^[86]. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[87]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[88][89]}.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[90][91]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[92][93][94]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[95][96][97]}. However, CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[91][98][99]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[93]. Administration of combined palbociclib and MEK inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[100].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[101].

NF2 Homozygous deletion

Biological Impact

The neurofibromin (NF2) gene encodes the protein Merlin, a tumor suppressor that functions as a negative regulator of the PI3K/AKT/mTOR pathway^{[102][103][104]}. NF2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[105]. Inactivation germline mutations in the NF2 are associated with the hereditary neurofibromatosis type 2, a disorder characterized by the growth of noncancerous tumors in the nervous system^{[102][106]}. Somatic mutations or deletion of NF2 are frequently observed in human cancers, including 20-50% of pleural mesotheliomas^[107], 6% papillary renal cell carcinoma, 5% pancreas cancer, and 4% melanoma (cbioPortal; June 2015), and less frequently in other cancers^[108].

Therapeutic and prognostic relevance

Genomic alterations with activating effects on the mTOR signaling pathway have been identified to confer sensitivity to everolimus across multiple cancer types^{[109][110][111][63]}. There are at least two case studies indicating the clinical efficacy of everolimus in bladder cancer^[62] and urothelial carcinoma^[112], both harboring NF2 truncating mutations. Preclinical evidence has shown the efficacy of MEK1/2 inhibitor selumetinib in KRAS-mutant thyroid cancer model with NF2 loss^[113].

Analysis of afatinib-plus-cetuximab-resistant biopsy specimens revealed a loss-of-function alteration in genes that modulate mTOR signaling pathway, including NF2 and TSC1^[114].

PTCH1 Heterozygous deletion

Biological Impact

The PTCH1 (protein patched homolog 1) gene encodes a multi-pass transmembrane receptor for sonic hedgehog (shh), a tumor suppressor that acts to repress shh signaling in the absence of ligand^[115]. Inactivation of PTCH1 results in hedgehog ligand-independent activation of SMO, causing a downstream activation of the pathway and lead to the neoplastic growth^{[116][117]}. Recurrent PTCH1 mutations have been reported in sporadic basal cell carcinoma (BCCs) and medulloblastoma^{[118][119][120][121]}. Germline PTCH1 mutations are associated with the nevoid basal cell carcinoma syndrome (NBCCS, Gorlin syndrome), predisposing patients to basal cell carcinoma and medulloblastoma^[119]. PTCH1 is a haploinsufficient tumor suppressor gene with one copy loss may be sufficient to promote tumor development in mice^{[116][122]}.

Therapeutic and prognostic relevance

Vismodegib is a small molecule inhibitor of SMO approved by the FDA for the treatment of patients with basal cell carcinoma. A heavily-pretreated patient with metastatic medulloblastoma harboring loss of heterozygosity and somatic mutation of PTCH1, showed rapid regression of the tumor after treated with vismodegib^[123]. Furthermore, a phase II study demonstrated that vismodegib treatment results in extended progression-free survival (PFS) in patients with loss-of-heterozygosity, SHH-driven medulloblastoma^{[124][125]}. In a phase II trial (MyPathway), 3 advanced solid tumors patients harboring PTCH1 loss-of-function mutations had partial responses to vismodegib treatment^[22].

RAD51 Heterozygous deletion

Biological Impact

The RAD51 gene encodes a recombinase that is crucial for homologous recombination (HR)-mediated repair of double-strand DNA breaks (DSBs) by forming complexes with known tumor suppressors including BRCA1, BRCA2, and PALB2^{[126][127][128]}. RAD51 has been characterized as a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[129]. Overexpression of RAD51 has been observed in many cancer cells, including pancreatic cancer and breast cancer and its hyperexpression is implicated in drug resistance^{[130][131][132][133][134][135][136]}. Germline mutations in RAD51 are associated with increased susceptibility to breast cancer^{[137][138][139][140]}.

Therapeutic and prognostic relevance

RAD51 loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in ovarian cancer^[141]; rucaparib efficacy in solid tumor (NCT04171700); talazoparib efficacy in lung cancer (NCT03377556); niraparib efficacy in pancreatic cancer (NCT03553004) or any malignancy (except prostate cancer) (NCT03207347).

Preclinical studies showed that decreased RAD51 expression could sensitize cells to olaparib-induced tumor cell cytotoxicity^{[142][143]}.

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

US FDA-APPROVED DRUG(S)

Abemaciclib (VERZENIO)

Abemaciclib is a cyclin-dependent kinase 4/6 (CDK4/6) inhibitor. Abemaciclib is developed and marketed by Eli Lilly under the trade name VERZENIO.

FDA Approval Summary of Abemaciclib (VERZENIO)

monarchE NCT03155997	Breast cancer (Approved on 2021/10/12)
	HR-positive, HER2-negative
	Abemaciclib+tamoxifen/aromatase inhibitor vs. Tamoxifen/aromatase inhibitor [IDFS at 36 months(%): 86.1 vs. 79.0]
MONARCH 3^[144] NCT00246621	Breast cancer (Approved on 2018/02/26)
	HR-positive, HER2-negative
	Abemaciclib + anastrozole/letrozole vs. Placebo + anastrozole/letrozole [PFS(M): 28.2 vs. 14.8]
MONARCH 1^[145] NCT02102490	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative
	Abemaciclib [ORR(%): 19.7 vs. 17.4]
MONARCH 2^[99] NCT02107703	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative
	Abemaciclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 16.4 vs. 9.3]

Atezolizumab (TECENTRIQ)

Atezolizumab is a humanized, anti-programmed cell death-ligand 1 (PD-L1) monoclonal antibody of the IgG1 isotype, which can lead to the reactivation of immune cells that might recognize and attack tumor cells. Atezolizumab is developed and marketed by Genentech/Roche under the trade name TECENTRIQ.

FDA Approval Summary of Atezolizumab (TECENTRIQ)

IMpower010 NCT02486718	Non-small cell lung carcinoma (Approved on 2021/10/15)
	PD-L1 TC ≥1%
	Atezolizumab vs. Best supportive care (bsc) [DFS (PD-L1 TC≥1%)(M): not reached vs. 35.3]
IMbrave150 NCT03434379	Hepatocellular carcinoma (Approved on 2020/05/29)
	-
	Atezolizumab plus bevacizumab vs. Sorafenib [PFS(M): 6.8 vs. 4.3, OS(M): NR vs. 13.2]

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

Page 12 of 63

IMpower133^[146] NCT02763579	Small cell lung cancer (Approved on 2019/03/18)
	-
IMpassion130^[147] NCT02425891	Breast cancer (Approved on 2019/03/08)
	PD-L1 Atezolizumab plus nab-paclitaxel vs. Nab-paclitaxel [PFS(M): 7.4 vs. 4.8]
OAK^[148] NCT02008227	Non-small cell lung carcinoma (Approved on 2016/10/18)
	PD-L1 Atezolizumab vs. Docetaxel [OS(M): 13.8 vs. 9.6]
POPLAR^[149] NCT01903993	Non-small cell lung carcinoma (Approved on 2016/10/18)
	PD-L1 Atezolizumab vs. Docetaxel [OS(M): 12.6 vs. 9.7]
Invigor 210^[8] NCT02108652	Urinary bladder urothelial carcinoma (Approved on 2016/05/18)
	PD-L1 Atezolizumab [ORR(%): 14.8]

Avelumab (BAVENCIO)

Avelumab is fully human monoclonal programmed death ligand-1 (PD-L1) antibody, belonging to the group of immune checkpoint blockade cancer therapies. Avelumab is developed and marketed by Merck KGaA and Pfizer under the trade name BAVENCIO.

FDA Approval Summary of Avelumab (BAVENCIO)

JAVELIN Renal 101^[150] NCT02684006	Renal cell carcinoma (Approved on 2019/05/14)
	-
JAVELIN Solid Tumor NCT01772004	Avelumab plus axitinib vs. Sunitinib [ORR(%): 51.4 vs. 25.7, PFS(M): 13.8 vs. 8.4]
	Bladder urothelial carcinoma (Approved on 2017/05/09)
	-
	Avelumab [ORR(13W)(%): 13.6, ORR(6M)(%): 16.1]

JAVELIN Merkel 200^[151] NCT02155647	Merkel cell carcinoma (Approved on 2017/03/23)
	-
	Avelumab [ORR(%): 33.0, DOR(M): 2.8 to 23.3+]

Bevacizumab (AVASTIN)

Bevacizumab is a recombinant humanized monoclonal antibody that blocks angiogenesis by inhibiting VEGF-A. Bevacizumab is developed and marketed by Genentech/Roche under the trade name AVASTIN.

FDA Approval Summary of Bevacizumab (AVASTIN)

IMbrave150 NCT03434379	Hepatocellular carcinoma (Approved on 2020/05/29)
	-
	Atezolizumab plus bevacizumab vs. Sorafenib [PFS(M): 6.8 vs. 4.3, OS(M): NR vs. 13.2]
GOG-0213^[152] NCT00565851	Peritoneal carcinoma, Ovary epithelial cancer, Fallopian tube cancer (Approved on 2016/12/06)
	-
	Bevacizumab + carboplatin + paclitaxel vs. Carboplatin + paclitaxel [OS(M): 42.6 vs. 37.3]
OCEANS^[153] NCT00434642	Peritoneal carcinoma, Ovary epithelial cancer, Fallopian tube cancer (Approved on 2016/12/06)
	-
	Bevacizumab + carboplatin + gemcitabine vs. Carboplatin + gemcitabine [PFS(M): 12.4 vs. 8.4]
AURELIA^[154] NCT00976911	Peritoneal carcinoma, Ovary epithelial cancer, Fallopian tube cancer (Approved on 2014/11/14)
	-
	Bevacizumab + chemotherapy vs. Chemotherapy [PFS(M): 6.8 vs. 3.4]
GOG-0240^[155] NCT00803062	Cervical cancer (Approved on 2014/08/14)
	-
	Bevacizumab + chemotherapy vs. Chemotherapy [OS(M): 16.8 vs. 12.9]
ML18147^[156] NCT00700102	Colorectal cancer (Approved on 2013/01/23)
	-
	Bevacizumab + chemotherapy vs. Chemotherapy [OS(M): 11.2 vs. 9.8]

BO17705^[157] NCT00738530	Renal cell carcinoma (Approved on 2009/07/31)
	-
	Bevacizumab + ifn- α 2a vs. Ifn- α 2a [PFS(M): 10.2 vs. 5.4]
AVF3708g^[158] NCT00345163	Glioblastoma multiforme (Approved on 2009/05/06)
	-
	Bevacizumab + irinotecan vs. Bevacizumab [ORR(%): 25.9]
E4599^[159] NCT00021060	Non-small cell lung carcinoma (Approved on 2006/10/11)
	-
	Bevacizumab + paclitaxel + carboplatin vs. Paclitaxel + carboplatin [OS(M): 12.3 vs. 10.3]
E3200^[160] NCT00025337	Colorectal cancer (Approved on 2006/06/20)
	-
	Bevacizumab + oxaliplatin + fluorouracil + leucovorin vs. Oxaliplatin + fluorouracil + leucovorin [OS(M): 13 vs. 10.8]
AVF2107g^[161] NCT00109070	Colorectal cancer (Approved on 2004/02/26)
	-
	Bevacizumab + irinotecan+5-fu + leucovorin vs. Irinotecan + 5-fu + leucovorin [OS(M): 20.3 vs. 15.6]

Cemiplimab-rwlc (LIBTAYO)

Cemiplimab-rwlc is a recombinant human IgG4 monoclonal antibody that binds to PD-1 and blocks its interaction with PD-L1 and PD-L2. Cemiplimab-rwlc is developed and marketed by Sanofi and Regeneron under the trade name LIBTAYO.

FDA Approval Summary of Cemiplimab-rwlc (LIBTAYO)

Study 1624 NCT03088540	Non-small lung cancer (Approved on 2021/02/22)
	PD-L1 TPS \geq 50%
	Cemiplimab-rwlc vs. Platinum-based chemotherapy [PFS(M): 6.2 vs. 5.6, OS(M): 22.1 vs. 14.3]
Study 1620 NCT03132636	Metastatic basal cell carcinoma (mbcc) (Approved on 2021/02/09)
	-
	Cemiplimab-rwlc [ORR(%): 21.0, DOR(M): NR]

Study 1620 NCT03132636	Locally advanced basal cell carcinoma (labcc) (Approved on 2021/02/09)
	-
	Cemiplimab-rwlc [ORR(%): 29.0, DOR(M): NR]
Study 1423, Study 1540 [7] NCT02383212, NCT02760498	cutaneous squamous cell carcinoma (Approved on 2018/09/28)
	-
	Cemiplimab-rwlc [ORR(%): 47.2]

Dasatinib (SPRYCEL)

Dasatinib is an oral Bcr-Abl tyrosine kinase inhibitor (inhibits the "Philadelphia chromosome") and Src family tyrosine kinase inhibitor. Dasatinib is produced by Bristol-Myers Squibb and sold under the trade name SPRYCEL.

FDA Approval Summary of Dasatinib (SPRYCEL)

DASISION^[162] NCT00481247	Chronic myeloid leukemia (Approved on 2010/10/28)
	-
	Dasatinib vs. Imatinib [ORR(%): 76.8 vs. 66.2]
[163] NCT00123474	Chronic myeloid leukemia (Approved on 2007/11/08)
	-
	Dasatinib [ORR(%): 63.0]
[164] NCT00123487	Acute lymphocytic leukemia (Approved on 2006/06/28)
	-
	Dasatinib [ORR(%): 38.0]

Dostarlimab-gxly (JEMPERLI)

Dostarlimab-gxly is a programmed death receptor-1 (PD-1)-blocking antibody. Dostarlimab-gxly is developed and marketed by GlaxoSmithKline LLC under the trade name JEMPERLI.

FDA Approval Summary of Dostarlimab-gxly (JEMPERLI)

GARNET NCT02715284	Cancer (Approved on 2021/08/17)
	dMMR
	Dostarlimab [ORR(%): 41.6, DoR(M): 34.7]

GARNET (Cohort A) NCT02715284	Endometrial carcinoma (Approved on 2021/04/22)
	dMMR
	Dostarlimab-gxly [ORR(%): 42.3, DOR(M): NR]

Durvalumab (IMFINZI)

Durvalumab is a programmed death ligand-1 (PD-L1)-blocking antibody. Durvalumab is developed and marketed by AstraZeneca under the trade name IMFINZI.

FDA Approval Summary of Durvalumab (IMFINZI)

CASPIAN^[165] NCT03043872	Extensive-stage small cell lung cancer (Approved on 2020/03/27)
	-
	Durvalumab + etoposide + carboplatin or durvalumab + etoposide + cisplatin vs. Etoposide + carboplatin or etoposide + cisplatin [OS(M): 13 vs. 10.3]
PACIFIC^[166] NCT02125461	Non-small cell lung carcinoma (Approved on 2018/02/16)
	-
	Durvalumab vs. Placebo [PFS(M): 16.8 vs. 5.6]
CD-ON-MEDI4736-1108^[167] NCT01693562	Bladder urothelial carcinoma (Approved on 2017/05/01)
	-
	Durvalumab [ORR(All)(%): 17.0, ORR(PD-L1 high)(%): 26.3, ORR (PD-L1 low/negative)(%): 4.1]

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4^[168] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	-
	Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]

BOLERO-2^[169] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
RADIANT-3^[170] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1^[171] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
RECORD-1^[172] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

Ipilimumab (YERVOY)

Ipilimumab is a fully human monoclonal antibody against the cytotoxic T-lymphocyte associated protein 4 (CTLA-4), an immune checkpoint protein receptor, to reactivate the immune responses. Ipilimumab is developed by Medarex and Bristol-Myers Squibb, and marketed by the latter under the trade name YERVOY.

FDA Approval Summary of Ipilimumab (YERVOY)

CHECKMATE-743 NCT02899299	Pleural mesothelioma (Approved on 2020/10/02)
	- Nivolumab + ipilimumab vs. Pemetrexed + cisplatin [OS(M): 18.1 vs. 14.1]
CHECKMATE-9LA NCT03215706	Non-small cell lung carcinoma (Approved on 2020/05/26)
	- Nivolumab + ipilimumab + platinum-doublet chemotherapy vs. Platinum-doublet chemotherapy [OS(M): 14.1 vs. 10.7]
CHECKMATE-227 NCT02477826	Non-small cell lung carcinoma (Approved on 2020/05/15)
	PD-L1 tumor expression ≥ 1% Nivolumab + ipilimumab vs. Platinum-doublet chemotherapy [OS(M): 17.1 vs. 14.9]

CHECKMATE-040 NCT01658878	Hepatocellular carcinoma (Approved on 2020/03/10)
	- Nivolumab + ipilimumab [ORR(%): 33.0]
CHECKMATE-142 ^[173] NCT02060188	Colorectal cancer (Approved on 2018/07/10)
	MSI-H or dMMR Ipilimumab plus nivolumab vs. Nivolumab [ORR(%): 49.0 vs. 32.0]
CHECKMATE-214 ^[174] NCT02231749	Renal cell carcinoma (Approved on 2018/04/16)
	- Nivolumab plus ipilimumab vs. Sunitinib [OS(M): 67.1 vs. 55.5]
EORTC 18071 ^[175] NCT00636168	Melanoma (Approved on 2015/10/28)
	- Ipilimumab vs. Placebo [RFS(M): 26 vs. 17]
MDX010-20 ^[176] NCT00094653	Melanoma (Approved on 2011/03/25)
	- Ipilimumab vs. Peptide vaccine with incomplete freund's adjuvant (gp100) [OS(M): 10 vs. 6]

Ivosidenib (TIBSOVO)

Ivosidenib is an isocitrate dehydrogenase-1 (IDH1) inhibitor. Ivosidenib is developed and marketed by Agios under the trade name TIBSOVO.

FDA Approval Summary of Ivosidenib (TIBSOVO)

Study AG120-C-005 NCT02989857	Cholangiocarcinoma (Approved on 2021/08/25)
	IDH1 Ivosidenib vs. Placebo [PFS(M): 2.7 vs. 1.4]
AG120-C-001 ^[36] NCT02074839	Acute myeloid leukemia (Approved on 2018/07/20)
	IDH1-mutated Ivosidenib [pCR(%): 42.9]

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

FDA Approval Summary of Niraparib (ZEJULA)

QUADRA^[82] NCT02354586	Ovarian cancer (Approved on 2019/10/23)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)
	Niraparib [ORR(%): 24.0, DOR(M): 8.3]
NOVA^[81] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	gBRCA+ CR/PR to platinum-based chemotherapy
	Niraparib vs. Placebo [PFS(M): 21 vs. 5.5]
NOVA^[81] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	gBRCA- CR/PR to platinum-based chemotherapy
	Niraparib vs. Placebo [PFS(M): 9.3 vs. 3.9]

Nivolumab (OPDIVO)

Nivolumab is a programmed death receptor-1 (PD-1)-blocking antibody. Nivolumab is developed and marketed by Bristol-Myers Squibb under the trade name OPDIVO.

FDA Approval Summary of Nivolumab (OPDIVO)

CHECKMATE-274 NCT02632409	Urothelial carcinoma (Approved on 2021/08/19)
	-
	Nivolumab vs. Placebo [DFS(M): 20.8 vs. 10.8, DFS(M): NR vs. 8.4]
CHECKMATE-577 NCT02743494	Gastroesophageal junction adenocarcinoma (Approved on 2021/05/20)
	-
	Nivolumab vs. Placebo every 4 weeks beginning at week 17 for up to one year of treatment [DFS(M): 22.4 vs. 11]

CHECKMATE-649 NCT02872116	Gastroesophageal junction adenocarcinoma, Gastric adenocarcinoma (Approved on 2021/04/16)
	-
	Nivolumab + chemotherapy (xelox or folfox) vs. Chemotherapy (xelox or folfox) [PFS(M): 7.7 vs. 6, OS(M): 14.4 vs. 11.1]
CHECKMATE-9ER NCT03141177	Renal cell carcinoma (Approved on 2021/01/22)
	-
	Nivolumab + cabozantinib vs. Sunitinib [ORR(%): 55.7 vs. 27.1, PFS(M): 16.6 vs. 8.3, OS(M): NR vs. NR]
CHECKMATE-743 NCT02899299	Pleural mesothelioma (Approved on 2020/10/02)
	-
	Nivolumab + ipilimumab vs. Pemetrexed + cisplatin [OS(M): 18.1 vs. 14.1]
CHECKMATE-9LA NCT03215706	Non-small cell lung carcinoma (Approved on 2020/05/26)
	-
	Nivolumab + ipilimumab + platinum-doublet chemotherapy vs. Platinum-doublet chemotherapy [OS(M): 14.1 vs. 10.7]
CHECKMATE-227 NCT02477826	Non-small cell lung carcinoma (Approved on 2020/05/15)
	PD-L1 tumor expression $\geq 1\%$
	Nivolumab + ipilimumab vs. Platinum-doublet chemotherapy [OS(M): 17.1 vs. 14.9]
CheckMate 040 NCT01658878	Hepatocellular carcinoma (Approved on 2020/03/10)
	-
	Nivolumab + ipilimumab [ORR(%): 33.0]
CheckMate 032 NCT01928394	Lung small cell carcinoma (Approved on 2018/08/16)
	-
	Nivolumab [ORR(%): 12.0]
CheckMate 040 NCT01658878	Hepatocellular carcinoma (Approved on 2017/09/22)
	-
	Nivolumab [ORR(%): 14.3]

CheckMate 142 NCT02060188	Colorectal cancer (Approved on 2017/07/31)
	MSI-H or dMMR Nivolumab [ORR(%): 32.0]
CheckMate 275^[177] NCT02387996	Urinary bladder urothelial carcinoma (Approved on 2017/02/02)
	- Nivolumab [ORR(%): 19.6]
CheckMate 141^[178] NCT02105636	Squamous cell carcinoma of the head and neck cancer (Approved on 2016/11/10)
	- Nivolumab vs. Investigator's choice of cetuximab, methotrexate or docetaxel [OS(M): 7.5 vs. 5.1]
CheckMate 039^[179] NCT01592370	Hodgkin's lymphoma (Approved on 2016/05/17)
	- Nivolumab [ORR(%): 66.0]
CheckMate 205^[180] NCT02181738	Hodgkin's lymphoma (Approved on 2016/05/17)
	- Nivolumab [ORR(%): 66.0]
CheckMate 067^[181] NCT01844505	Melanoma (Approved on 2016/01/23)
	- Ipilimumab vs. Placebo [PFS(M): 11.5 vs. 2.9]
CheckMate 066^[182] NCT01721772	Melanoma (Approved on 2015/11/24)
	BRAF V600 wild-type Nivolumab vs. Dacarbazine [OS(M): Not Reached vs. 10.8]
CheckMate 025^[183] NCT01668784	Renal cell carcinoma (Approved on 2015/11/23)
	- Nivolumab vs. Everolimus [OS(M): 25 vs. 19.6]
CheckMate 057^[184] NCT01673867	Non-small cell lung carcinoma (Approved on 2015/10/09)
	- Nivolumab vs. Docetaxel [OS(M): 12.2 vs. 9.4]

CheckMate 017^[185] NCT01642004	Non-small cell lung carcinoma (Approved on 2015/03/04)
	-
	Nivolumab vs. Docetaxel [OS(M): 9.2 vs. 6]
CheckMate 037^[186] NCT01721746	Melanoma (Approved on 2014/12/22)
	-
	Nivolumab vs. Dacarbazine or carboplatin + paclitaxel [ORR(%): 32.0]

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

FDA Approval Summary of Olaparib (LYNPARZA)

PROfound^[77] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	ATMm, BRCA1m, BRCA2m, BARD1m, BRIP1m, CDK12m, CHEK1m, CHEK2m, FANCLm, PALB2m, RAD51Bm, RAD51Cm, RAD51Dm, RAD54Lm
	Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1^[71] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)
	Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
POLO^[76] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	Germline BRCA mutation (deleterious/suspected deleterious)
	Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1^[70] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	Germline or somatic BRCA-mutated (gBRCAm or sBRCAm)
	Olaparib vs. Placebo [PFS(M): NR vs. 13.8]

OlympiAD^[75] NCT02000622	Breast cancer (Approved on 2018/02/06)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative
	Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21^[187] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA+
	Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
Study19^[188] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	-
	Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]
Study 42^[189] NCT01078662	Ovarian cancer (Approved on 2014/12/19)
	Germline BRCA mutation (deleterious/suspected deleterious)
	Olaparib [ORR(%): 34.0, DOR(M): 7.9]

Palbociclib (IBRANCE)

Palbociclib is an oral, cyclin-dependent kinase (CDK) inhibitor specifically targeting CDK4 and CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Palbociclib is developed and marketed by Pfizer under the trade name IBRANCE.

FDA Approval Summary of Palbociclib (IBRANCE)

PALOMA-2^[190] NCT01740427	Breast cancer (Approved on 2017/03/31)
	ER+, HER2-
	Palbociclib + letrozole vs. Placebo + letrozole [PFS(M): 24.8 vs. 14.5]
PALOMA-3^[191] NCT01942135	Breast cancer (Approved on 2016/02/19)
	ER+, HER2-
	Palbociclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 9.5 vs. 4.6]

Pembrolizumab (KEYTRUDA)

Pembrolizumab is a programmed death receptor-1 (PD-1)-blocking antibody. Pembrolizumab is developed and marketed by Merck under the trade name KEYTRUDA.

FDA Approval Summary of Pembrolizumab (KEYTRUDA)

KEYNOTE-826 NCT03635567	Cervical cancer (Approved on 2021/10/13)
	PD-L1 (CPS ≥1) Pembrolizumab+paclitaxel+cisplatin with or without bevacizumab vs. Placebo+paclitaxel+cisplatin with or without bevacizumab [OS (PD-L1, CPS ≥1)(M): Not reached vs. 16.3, PFS(M): 10.4 vs. 8.2]
KEYNOTE-522 NCT03036488	Triple-receptor negative breast cancer (Approved on 2021/07/26)
	- Pembrolizumab + chemotherapy as neoadjuvant treatment vs. Placebo in combination with chemotherapy [pCR(%): 63.0 vs. 56.0, EFS(): 123 vs. 93]
KEYNOTE-775 (Study 309) NCT03517449	Endometrial carcinoma (Approved on 2021/07/22)
	Not microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR) Pembrolizumab + lenvatinib vs. Investigator's choice of doxorubicin or paclitaxel [PFS(M): 6.6 vs. 3.8, OS(M): 17.4 vs. 12]
KEYNOTE-811 NCT03615326	Gastroesophageal junction adenocarcinoma (Approved on 2021/05/05)
	Her2+ Pembrolizumab 200 mg every 3 weeks, in combination with trastuzumab and either fluorouracil plus cisplatin or capecitabine plus oxaliplatin vs. Placebo every 3 weeks, in combination with trastuzumab and either fluorouracil plus cisplatin or capecitabine plus oxaliplatin [ORR(%): 74.0 vs. 52.0, DOR(M): 10.6 vs. 9.5]
KEYNOTE-590 NCT03189719	Esophageal cancer, Gastroesophageal junction adenocarcinoma (Approved on 2021/03/22)
	- Pembrolizumab in combination with cisplatin and fluorouracil vs. Placebo with cisplatin and fluorouracil [PFS(M): 6.3 vs. 5.8, OS(M): 12.4 vs. 9.8]

KEYNOTE-355 NCT02819518	Triple-receptor negative breast cancer (Approved on 2020/11/13)
	PD-L1 (CPS ≥ 10) Pembrolizumab + paclitaxel protein-bound, or paclitaxel, or gemcitabine plus carboplatin vs. Placebo + paclitaxel protein-bound, or paclitaxel, or gemcitabine plus carboplatin [PFS(M): 9.7 vs. 5.6]
KEYNOTE-204 NCT02684292	Hodgkin's lymphoma (Approved on 2020/10/14)
	- Pembrolizumab vs. Brentuximab vedotin [PFS(M): 13.2 vs. 8.3]
KEYNOTE-158 NCT02628067	Cancer (Approved on 2020/06/17)
	TMB-H; ≥ 10 mutations/megabase Pembrolizumab (tmb-h) vs. Pembrolizumab (non-tmb-h) [ORR(%): 29.0 vs. 6.0]
KEYNOTE-146 NCT02501096	Endometrial carcinoma (Approved on 2019/09/17)
	not microsatellite instability high (MSI-H) or mismatch repair deficient (dMMR) Pembrolizumab + lenvatinib [ORR(%): 38.3, DOR(M): NR]
KEYNOTE-426 ^[192] NCT02853331	Renal cell carcinoma (Approved on 2019/04/19)
	- Pembrolizumab + axitinib vs. Sunitinib [ORR(%): 59.3 vs. 35.7, PFS(M): 15.1 vs. 11.1]
KEYNOTE-017 ^[193] NCT02267603	Merkel cell carcinoma (Approved on 2018/12/19)
	- Pembrolizumab [ORR(%): 56.0]
KEYNOTE-224 ^[194] NCT02702414	Hepatocellular carcinoma (Approved on 2018/11/09)
	- Pembrolizumab [ORR(%): 17.0]
KEYNOTE-407 ^[195] NCT02775435	Squamous non-small cell lung carcinoma (Approved on 2018/10/30)
	- Pembrolizumab + carboplatin + paclitaxel/nab-paclitaxel vs. Carboplatin + paclitaxel/nab-paclitaxel [ORR(%): 58.0 vs. 35.0, PFS(M): 6.4 vs. 4.8]

KEYNOTE-189^[195] NCT02578680	Nonsquamous non-small cell lung carcinoma (Approved on 2018/08/20)
	- Pembrolizumab + pemetrexed + platinum vs. Pemetrexed + platinum [PFS(M): 8.8 vs. 4.9, OS(M): NR vs. 11.3]
KEYNOTE-170 NCT02576990	Mediastinal large b-cell lymphoma (Approved on 2018/06/13)
	- Pembrolizumab [ORR(%): 45.0]
KEYNOTE-158 NCT02628067	Cervical cancer (Approved on 2018/06/13)
	- Pembrolizumab [ORR(%): 14.3]
KEYNOTE-059 NCT02335411	Gastric adenocarcinoma, Gastroesophageal junction adenocarcinoma (Approved on 2017/09/22)
	- Pembrolizumab [ORR(%): 13.3]
KEYNOTE-158 NCT02628067	Cancer (Approved on 2017/05/23)
	MSI-H or dMMR Pembrolizumab [ORR(%): 39.6]
KEYNOTE-164 NCT02460198	Cancer (Approved on 2017/05/23)
	MSI-H or dMMR Pembrolizumab [ORR(%): 39.6]
KEYNOTE-028^{[196][197]} NCT02054806	Cancer (Approved on 2017/05/23)
	MSI-H or dMMR Pembrolizumab [ORR(%): 39.6]
KEYNOTE-016^[6] NCT01876511	Cancer (Approved on 2017/05/23)
	MSI-H or dMMR Pembrolizumab [ORR(%): 39.6]
KEYNOTE-012^{[198][199][200][201]} NCT01848834	Cancer (Approved on 2017/05/23)
	MSI-H or dMMR Pembrolizumab [ORR(%): 39.6]

KEYNOTE-052 NCT02335424	Urinary bladder urothelial carcinoma (Approved on 2017/05/18)
	-
KEYNOTE-045 ^[202] NCT02256436	Urinary bladder urothelial carcinoma (Approved on 2017/05/18)
	-
KEYNOTE-087 ^[203] NCT02453594	Hodgkin's lymphoma (Approved on 2017/03/14)
	-
KEYNOTE-024 ^[204] NCT02142738	Non-small cell lung carcinoma (Approved on 2016/10/24)
	PD-L1 expression (TPS >= 50%)
KEYNOTE-012 ^[199] NCT01848834	Head and neck squamous cell carcinoma (Approved on 2016/08/05)
	-
KEYNOTE-006 ^[205] NCT01866319	Melanoma (Approved on 2015/12/18)
	-
KEYNOTE-010 ^[206] NCT01905657	Non-small cell lung carcinoma (Approved on 2015/10/02)
	PD-L1 expression (TPS >= 1%)
KEYNOTE-002 ^[207] NCT01704287	Melanoma (Approved on 2014/09/24)
	-

Ribociclib (KISQALI)

Ribociclib is a cyclin-dependent kinase (CDK) inhibitor specifically targeting cyclin D1/CDK4 and cyclin D3/CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Ribociclib is developed by Novartis and Astex Pharmaceuticals and marketed by Novartis under the trade name KISQALI.

FDA Approval Summary of Ribociclib (KISQALI)

MONALEESA-2^[98] NCT01958021	Breast cancer (Approved on 2017/03/13)
	HR+, HER2-
	Ribociclib vs. Letrozole [PFS(M): NR vs. 14.7]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA+, sBRCA
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
ARIEL3^[78] NCT01968213	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	All HRD tBRCA
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]
ARIEL2^[208] NCT01482715, NCT01891344	Ovarian cancer (Approved on 2016/12/19)
	Germline and/or somatic BRCA mutation
	Rucaparib [ORR(%): 54.0]

Selumetinib (KOSELUGO)

Selumetinib is a kinase inhibitor. Selumetinib is developed and marketed by AstraZeneca under the trade name KOSELUGO.

FDA Approval Summary of Selumetinib (KOSELUGO)

SPRINT NCT01362803	Plexiform neurofibromas (Approved on 2020/04/10)
	Neurofibromatosis type 1
	Selumetinib [ORR(%): 66.0]

Sonidegib (ODOMZO)

Sonidegib is a Hedgehog signaling pathway inhibitor by blocking its key component, smoothened (smo). Sonidegib is developed and marketed by Novartis under the trade name ODOMZO.

FDA Approval Summary of Sonidegib (ODOMZO)

BOLT^[209] NCT01327053	Basal cell carcinoma (Approved on 2015/07/24)
	-
	Sonidegib [ORR(%): 58.0]

Sunitinib (SUTENT)

Sunitinib is an oral, small molecule, multi-kinase inhibitor that targets receptor tyrosine kinase including platelet-derived growth factor receptor- α , - β (PDGFR- α , - β), vascular endothelial growth factor receptors-1, -2, -3 (VEGFR-1, -2, -3), c-kit, Fms-like tyrosine kinase-3 (FLT3), colony stimulating factor receptor type 1 (CSF-1R), and the glial cell-line derived neurotrophic factor receptor (RET), thereby inhibiting angiogenesis. Sunitinib is developed and marketed by Pfizer under the trade name SUTENT.

FDA Approval Summary of Sunitinib (SUTENT)

[210][211][212] NCT00428597	Pancreatic cancer (Approved on 2011/05/20)
	-
	Sunitinib vs. Placebo [PFS(M): 10.2 vs. 5.4]
[213][214][215] NCT00077974	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib [ORR(%): 34.0]

[214][215] NCT00054886	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib [ORR(%): 36.5]
[216][215] NCT00083889	Renal cell carcinoma (Approved on 2007/02/02)
	-
	Sunitinib vs. Ifn- α [PFS(W): 47.3 vs. 22]
[217] NCT00075218	Gastrointestinal stromal tumor (Approved on 2006/01/26)
	-
	Sunitinib vs. Placebo [TTP(W): 27.3 vs. 6.4]

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

FDA Approval Summary of Talazoparib (TALZENNA)

EMBRACA ^[83] NCT01945775	Breast cancer (Approved on 2018/10/16)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

FDA Approval Summary of Temsirolimus (TORISEL)

[218] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	-
	Temsirolimus vs. IFN- α [OS(M): 10.9 vs. 7.3]

Trametinib (MEKINIST)

Trametinib is an anti-cancer inhibitor which targets MEK1 and MEK2. Trametinib is developed and marketed by GlaxoSmithKline (GSK) under the trade name MEKINIST.

FDA Approval Summary of Trametinib (MEKINIST)

BRF117019 ^[219] NCT02034110	Anaplastic thyroid cancer (Approved on 2018/05/04)
	BRAF V600E
	Dabrafenib + trametinib [ORR(%): 61.0]
BRF113928 ^[220] NCT01336634	Non-small cell lung cancer (Approved on 2017/06/22)
	BRAF V600E
	Trametinib + dabrafenib vs. Dabrafenib [ORR(%): 63.0 vs. 27.0, DOR(M): 12.6 vs. 9.9]
COMBI-d ^[221] NCT01584648	Melanoma (Approved on 2014/01/10)
	BRAF V600E/K
	Trametinib + dabrafenib vs. Dabrafenib + placebo [PFS(M): 9.3 vs. 8.8]
METRIC ^[222] NCT01245062	Melanoma (Approved on 2013/05/29)
	BRAF V600E/K
	Trametinib vs. Dacarbazine or paclitaxel [PFS(M): 4.8 vs. 1.5]

Vismodegib (ERIVEDGE)

Vismodegib is a cyclopamine-competitive antagonist and acts as a first-in-class Hedgehog signaling pathway inhibitor by blocking its key component smoothened (smo). Vismodegib is developed by Genentech and marketed by Roche under the trade name ERIVEDGE.

FDA Approval Summary of Vismodegib (ERIVEDGE)

ERIVANCE BCC ^[223] NCT00833417	Basal cell carcinoma (Approved on 2012/01/30)
	-
	Vismodegib [ORR (mBCC)(%): 30.3, ORR (laBCC)(%): 42.9]

d=day; w=week; m=month

ONGOING CLINICAL TRIALS

Clinical trials shown below were selected by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

IMMUNE CHECKPOINT INHIBITORS

Drugs	Pembrolizumab
NCT ID	NCT02628067
Phase	II
Content	In this study, participants with multiple types of advanced (unresectable and/or metastatic) solid tumors who have progressed on standard of care therapy will be treated with pembrolizumab (MK-3475).
Contact	Name: Toll Free Number Phone: 1-888-577-8839 Email: NA
Location	Status: Recruiting Country: Taiwan City: Taipei Name: Merck Sharp & Dohme (I.A.) Corp.

Drugs	Atezolizumab
NCT ID	NCT04589845
Phase	II
Content	TAPISTRY is a Phase II, global, multicenter, open-label, multi-cohort study designed to evaluate the safety and efficacy of targeted therapies or immunotherapy as single agents or in rational, specified combinations in participants with unresectable, locally advanced or metastatic solid tumors determined to harbor specific oncogenic genomic alterations or who are tumor mutational burden (TMB)-high as identified by a validated next-generation sequencing (NGS) assay. Participants with solid tumors will be treated with a drug or drug regimen tailored to their NGS assay results at screening. Participants will be assigned to the appropriate cohort based on their genetic alteration(s). Treatment will be assigned on the basis of relevant oncogenotype, will have cohort-specific

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

Page 33 of 63

	inclusion/exclusion criteria, and, unless otherwise specified, will continue until disease progression, loss of clinical benefit, unacceptable toxicity, participant or physician decision to discontinue, or death, whichever occurs first.
Contact	<p>Name: Reference Study ID Number: BO41932</p> <p>www.roche.com/about_roche/roche_worldwide.htm</p> <p>Phone: 888-662-6728 (U.S. and Canada)</p> <p>Email: Global-Roche-Genentech-Trials@gene.com</p>
Location	<p>Status: Recruiting</p> <p>Country: Taiwan</p> <p>City: Tainan</p> <p>Name: National Cheng Kung University Hospital; Oncology</p> <p>Status: Recruiting</p> <p>Country: Taiwan</p> <p>City: Taipei City</p> <p>Name: Taipei Veterans General Hospital; Department of Oncology</p> <p>Status: Recruiting</p> <p>Country: Taiwan</p> <p>City: Taoyuan County</p> <p>Name: Chang Gung Memorial Hospital-Linkou; Dept of Oncology</p> <p>Status: Recruiting</p> <p>Country: Taiwan</p> <p>City: Zhongzheng Dist.</p> <p>Name: National Taiwan University Hospital; Oncology</p>

DETAILED TEST RESULTS


SINGLE NUCLEOTIDE AND SMALL INDEL VARIANTS

Gene	Chr	Exon	Accession Number	cDNA Change	Amino Acid Change	Coverage	Allele Frequency	COSMIC ID
ABL1	9	11	NM_005157	c.2246C>T	S749L	693	25.7%	-
ABL1	9	11	NM_005157	c.3322_3324delinsTCG	P1108S	436	7.1%	-
ARID1B	6	-	NM_017519	c.4071+5G>C	Splice region	477	5.5%	-
ARID2	12	15	NM_152641	c.4253G>A	G1418E	265	7.9%	-
ATM	11	13	NM_000051	c.1903C>T	H635Y	221	44.8%	-
BCL6	3	8	NM_001130845	c.1742C>T	A581V	463	5.4%	-
BRAF	7	11	NM_004333	c.1406G>C	G469A	1631	17.5%	COSM460
BUB1B	15	18	NM_001211	c.2347G>A	V783M	637	6.3%	-
CD274 (PD-L1)	9	-	NM_014143	c.791-4C>G	Splice region	652	24.4%	-
DPYD	1	22	NM_000110	c.2869G>A	G957S	477	5.0%	-
DTX1	12	2	NM_004416	c.424C>T	L142F	421	6.7%	-
EPHA2	1	6	NM_004431	c.1428G>T	K476N	881	43.8%	-
ERCC3	2	9	NM_000122	c.1411G>A	V471I	1383	27.3%	COSM3042197
FLT4	5	-	NM_182925	c.986-6C>T	Splice region	240	8.3%	COSM7347173
FLT4	5	28	NM_182925	c.3782C>T	T1261I	505	7.5%	-
GATA2	3	3	NM_032638	c.571G>A	A191T	365	5.8%	-
IDH1	2	4	NM_005896	c.394C>T	R132C	692	15.2%	COSM28747
IDH2	15	2	NM_002168	c.136G>A	V46M	401	6.0%	COSM5778128
KIT	4	-	NM_000222	c.1880-8A>G	Splice region	630	52.4%	-
MET	7	2	NM_001127500	c.632T>G	L211W	1840	56.7%	COSM7216189
MSH2	2	11	NM_000251	c.1690A>G	T564A	258	70.9%	COSM7450527
MSH6	2	4	NM_000179	c.1445G>A	R482Q	2871	46.7%	COSM1021268
MUC16	19	3	NM_024690	c.29255C>T	S9752F	592	6.4%	COSM3542212
MUC16	19	-	NM_024690	c.36067+1G>A	Splice donor	437	5.0%	-
MUC6	11	11	NM_005961	c.1334C>T	S445L	206	10.7%	-
NOTCH4	6	6	NM_004557	c.1145C>T	P382L	350	6.0%	-
NTRK3	15	13	NM_001012338	c.1559G>A	G520E	150	14.7%	COSM5609688
PBRM1	3	26	NM_018313	c.4011del	F1337fs	903	37.0%	-
PIK3C2G	12	14	NM_004570	c.1972G>T	A658S	863	45.8%	COSM6244593
POLD1	19	-	NM_001256849	c.1893-7A>G	Splice region	140	56.4%	-
PTPRD	9	42	NM_002839	c.5561T>C	F1854S	530	5.1%	-
RBM10	X	15	NM_005676	c.1692C>T	Splice region	1082	56.4%	-
SYNE1	6	44	NM_182961	c.6513G>A	M2171I	523	5.7%	-
TEK	9	2	NM_000459	c.136G>A	A46T	404	5.7%	-
TEK	9	3	NM_000459	c.368C>T	S123F	409	5.1%	-
TSC1	9	-	NM_000368	c.2041+1G>A	Splice donor	928	51.9%	-
TSC2	16	31	NM_000548	c.3713C>T	A1238V	308	7.5%	-
USH2A	1	41	NM_206933	c.7718G>C	R2573P	896	51.2%	-
WT1	11	2	NM_024426	c.764C>T	S255L	372	6.2%	-

Mutations with clinical relevance are highlighted in red.

COPY NUMBER VARIANTS (CNVs)

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



HOTSPOT GENOTYPES

Listed variants are biomarkers or hotspots that are recommended as standard care by the NCCN or other expert panels and not necessarily FDA-recognized for a particular indication. The genotypes have been manually checked to ensure sufficient coverage for each hotspot of the target gene.

Gene	Variant	Genotype Detected
<i>BRAF</i>	V600X	Not detected
<i>EGFR</i>	A763_Y764insFQEA, E709K, E709_T710delinsD, Exon 19 deletion, Exon 19 insertion, Exon 20 insertion, G719A/C/D/S, L747P, L833V, L858R, L861Q/R, S768I, T790M	Not detected
<i>IDH2</i>	R140Q, R172G/K/M/S	Not detected
<i>KIT</i>	A502_Y503dup, D419del, D579del, D816F/V/Y, D820A/E/G/Y, E554_I571del, E554_K558del, E554_V559del, Exon 11 mutation, F522C, H697Y, I563_L576del, I653T, K550_W557del, K558N, K558_E562del, K558_V559del, K558delinsNP, K642E, M552_W557del, N505I, N564_Y578del, N822H/I/K/Y, P551_M552del, P573_D579del, P577_D579del, P577_W582delinsPYD, P838L, Q556_K558del, T417_D419delinsI, T417_D419delinsRG, T574_Q575insTQLPYD, V530I, V555_L576del, V555_V559del, V559A/C/D/G, V559_V560del, V559del, V560D/G, V560del, V569_L576del, V654A, W557G/R, W557_K558del, Y553N, Y553_K558del, Y570H, Y578C	Not detected
<i>KRAS</i>	A146T/V/P, G12X, G13X, Q61X	Not detected
<i>MET</i>	D1028H/N/Y	Not detected
<i>NRAS</i>	G12X, G13X, Q61X	Not detected
<i>PDGFRA</i>	A633T, C450_K451insMIEWMI, C456_N468del, C456_R481del, D568N, D842I/V, D842_H845del, D842_M844del, D846Y, E311_K312del, G853D, H650Q, H845Y, H845_N848delinsP, I843del, N659K/R/S, N848K, P577S, Q579R, R560_V561insER, R748G, R841K, S566_E571delinsR, S584L, V469A, V536E, V544_L545insAVLVLLVIVISLI, V561A/D, V561_I562insER, V658A, W559_R560del, Y375_K455del, Y555C, Y849C/S	Not detected
<i>PIK3CA</i>	C420R, E542K/V, E545A/D/G/K, H1047X, Q546E/R	Not detected

V600X= any mutation in the valine (V) at amino acid 600 being replaced by a different amino acid.

G12X= any mutation in the glycine (G) at amino acid 12 being replaced by a different amino acid.

G13X= any mutation in the glycine (G) at amino acid 13 being replaced by a different amino acid.

Q61X= any mutation in the glutamine (Q) at amino acid 61 being replaced by a different amino acid.

H1047X= any mutation in the histidine (H) at amino acid 1047 being replaced by a different amino acid.

Gene	Copy Number Detected
<i>CDK4</i>	2
<i>EGFR</i>	2
<i>ERBB2</i>	2
<i>MET</i>	2

Copy number ≥ 8 is considered amplification

Other known alterations that are associated with sensitivity, resistance, and toxicity to therapies.

Gene	Variant	Genotype Detected
AKT1	E17K	Not detected
ALK	C1156Y, D1203N, G1202R, L1152R, S1206Y, T1151_L1152insT	Not detected
BRAF	K601E, L597V/Q/R/S	Not detected
DPYD	D949V, I560S, splice-site mutation	Not detected
EGFR	A750P, C797S/Y, S492R	Not detected
ERBB2	V659E	Not detected
ESR1	D538G, E380Q, L469V, L536H/P/Q/R, S432L, S463P, V422del, V534E, Y537C/N/S	Not detected
FGFR3	G370C, G380R, K650E/N/R/M/T/Q, R248C, S249C, S371C, Y373C	Not detected
IDH1	R132C/G/H/L/Q/S	R132C
MAP2K1	D67N, E203K, F53L, K57E/N, P124S, Q56P, Q56_V60del, R47Q, R49L, S222D	Not detected
PTEN	R130*/fs/G/L/P/Q	Not detected
TPMT	A154T, Y240C	Not detected

Gene	Copy Number Detected
FGFR1	2
MDM2	2
MDM4	2

Copy number ≥ 8 is considered amplification

TEST DETAILS

ABOUT ACTOnco®+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (INDELs) (≤ 15 nucleotides) and large-scale genomic alterations like copy number variations (CNVs).

See ACTOnco®+ Gene List' Section for details of gene sequenced.

DATABASE USED

- Reference genome: human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210208)
- ACT Genomics in-house database

NEXT-GENERATION SEQUENCING (NGS) METHODS

Extracted genomic DNA was amplified using four pools of primer pairs targeting coding exons of analyzed genes. Amplicons were ligated with barcoded adaptors. Quality and quantity of amplified library were determined using the fragment analyzer (AATI) and Qubit (Invitrogen). Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system (Thermo Fisher Scientific) according to the Ion PI Hi-Q Chef Kit protocol (Thermo Fisher Scientific) or Ion 540 Kit-Chef protocol (Thermo Fisher Scientific). Sequencing was performed on the Ion Proton or Ion S5 sequencer (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite (version 5.10). Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (INDELs) were identified using the Torrent Variant Caller plug-in (version 5.10). The coverage was down-sampled to 4000. VEP (Variant Effect Predictor) (version 100) was used to annotate every variant using databases from Clinvar (version 20210208), COSMIC v.92 and Genome Aggregation database r2.1.1. Variants with coverage ≥ 25 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained.

This test provides uniform coverage of the targeted regions, enabling target base coverage at $100x \geq 85\%$ with a mean coverage $\geq 500x$.

Variants reported in Genome Aggregation database r2.1.1 with $> 1\%$ minor allele frequency (MAF) were

considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number variations (CNVs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations from samples in ACT Genomics in-house database.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as “TMB-High”, “TMB-Low” or “Cannot Be Determined”. TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as “Cannot Be Determined” if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

STANDARD OPERATING PROCEDURES (SOPS)

Standard operating procedures (SOPs) are shown below:

- AG2-QP-15 Specimen Management Procedure
- AG3-QP16-03 SOP of Cancer Cell DNA and RNA Extraction
- AG3-QP16-07 SOP of Nucleic Acid Extraction with QIAasympyphony SP
- AG3-QP16-08 SOP of FFPE Nucleic Acid Extraction
- AG3-QP16-10 SOP of HE Staining
- AG3-QP16-13 SOP of Library Construction and Preparation
- AG3-QP16-17 SOP of DNA Quantification with Qubit Fluorometer
- AG3-QP16-20 SOP of CE-Fragment Analysis
- AG3-QP16-22 SOP of Variant Calling
- AG3-QP16-24 SOP of Ion Torrent System Sequencing Reaction
- AG3-QP16-26 SOP of Ion Chef Preparation

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

- AG3-QP16-35 SOP of Variant Annotation
- AG3-QP16-96 SOP of Manual Inspection for SNVIndel Variant
- AG3-QP16-95 SOP of Manual Inspection for Copy Number Variant
- AG3-QP40-08 (02) Standard protocol for variant interpretation, curation and classification
- AG3-QP16-41 SOP of The user manual for clinical report system (CRS)

LIMITATIONS

This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.

NOTES

We do not exclude the possibility that pathogenic variants may not be reported by one or more of the tools and the parameters used.


PATHOLOGY EVALUATION

- H&E-stained section No.: S10935424
- Collection site: Liver
- Examined by: Dr. Yeh-Han Wang
- Estimated neoplastic nuclei (whole sample): The percentage of viable tumor cells in total cells in the whole slide (%): 25%
The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 70%
The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
Additional comment: NA
- Manual macrodissection: Performed on the highlighted region



The outline highlights the area of malignant neoplasm annotated by a pathologist.

SPECIMEN PHOTO(S)



- Collection date: Oct 2020
- Facility retrieved: 臺北榮總

RUN QC

- Panel: ACTOnco®+
- Mean Depth: 792x
- Target Base Coverage at 100x: 93%

ACTOnco® + GENE LIST

ABCB1*	AURKB	CBL	CDKN2B	E2F3	FAT1	GRIN2A	JAK2	MED12	NOTCH4	PMS1	RAD51D	SLCO1B3*	TNFRSF14
ABCC2*	AXIN1	CCNA1	CDKN2C	EGFR	FBXW7	GSK3B	JAK3	MEF2B	NPM1	PMS2	RAD52	SMAD2	TNFSF11
ABCG2*	AXIN2	CCNA2	CEBPA*	EP300	FCGR2B	GSTP1*	JUN*	MEN1	NQO1*	POLB	RAD54L	SMAD3	TOP1
ABL1	AXL	CCNB1	CHEK1	EPCAM	FGF1*	GSTT1*	KAT6A	MET	NRAS	POLD1	RAF1	SMAD4	TP53
ABL2	B2M	CCNB2	CHEK2	EPHA2	FGF10	HGF	KDM5A	MITF	NSD1	POLE	RARA	SMARCA4	TPMT*
ADAMTS1	BAP1	CCNB3	CIC	EPHA3	FGF14	HIF1A	KDM5C	MLH1	NTRK1	PPARG	RB1	SMARCB1	TSC1
ADAMTS13	BARD1	CCND1	CREBBP	EPHA5	FGF19*	HIST1H1C*	KDM6A	MPL	NTRK2	PPP2R1A	RBM10	SMO	TSC2
ADAMTS15	BCL10	CCND2	CRKL	EPHA7	FGF23	HIST1H1E*	KDR	MRE11	NTRK3	PRDM1	RECQL4	SOC1*	TSHR
ADAMTS16	BCL2*	CCND3	CRLF2	EPHB1	FGF3	HNF1A	KEAP1	MSH2	PAK3	PRKAR1A	REL	SOX2*	TYMS
ADAMTS18	BCL2L1	CCNE1	CSF1R	ERBB2	FGF4*	HR	KIT	MSH6	PALB2	PRKCA	RET	SOX9	U2AF1
ADAMTS6	BCL2L2*	CCNE2	CTCF	ERBB3	FGF6	HRAS*	KMT2A	MTHFR*	PARP1	PRKCB	RHOA	SPEN	UBE2A*
ADAMTS9	BCL6	CCNH	CTLA4	ERBB4	FGFR1	HSP90AA1	KMT2C	MTOR	PAX5	PRKCG	RICTOR	SPOP	UBE2K
ADAMTSL1	BCL9	CD19	CTNNA1	ERCC1	FGFR2	HSP90AB1	KMT2D	MUC16	PAX8	PRKCI	RNF43	SRC	UBR5
ADGRA2	BCOR	CD274	CTNNB1	ERCC2	FGFR3	HSPA4	KRAS	MUC4	PBRM1	PRKCQ	ROS1	STAG2	UGT1A1*
ADH1C*	BIRC2	CD58	CUL3	ERCC3	FGFR4	HSPA5	LCK	MUC6	PDCD1	PRKDC	RPPH1	STAT3	USH2A
AKT1	BIRC3	CD70*	CYLD	ERCC4	FH	IDH1	LIG1	MUTYH	PDCD1LG2	PRKN	RPTOR	STK11	VDR*
AKT2	BLM	CD79A	CYP1A1*	ERCC5	FLCN	IDH2	LIG3	MYC	PDGFRA	PSMB8	RUNX1	SUFU	VEGFA
AKT3	BMPR1A	CD79B	CYP2B6*	ERG	FLT1	IFNL3*	LMO1	MYCL	PDGFRB	PSMB9	RUNX1T1	SYK	VEGFB
ALDH1A1*	BRAF	CDC73	CYP2C19*	ESR1	FLT3	IGF1	LRP1B	MYCN	PDIA3	PSME1	RXRA	SYNE1	VHL
ALK	BRCA1	CDH1	CYP2C8*	ESR2	FLT4	IGF1R	LYN	MYD88	PGF	PSME2	SDHA	TAF1	WT1
AMER1	BRCA2	CDK1	CYP2D6	ETV1	FOXL2*	IGF2	MALT1	NAT2*	PHOX2B*	PSME3	SDHB	TAP1	XIAP
APC	BRD4	CDK12	CYP2E1*	ETV4	FOXP1	IKBKB	MAP2K1	NBN	PIK3C2B	PTCH1	SDHC	TAP2	XPO1
AR	BRIP1	CDK2	CYP3A4*	EZH2	FRG1	IKBKE	MAP2K2	NEFH	PIK3C2G	PTEN	SDHD	TAPBP	XRCC2
ARAF	BTG1*	CDK4	CYP3A5*	FAM46C	FUBP1	IKZF1	MAP2K4	NF1	PIK3C3	PTGS2	SERPINB3	TBX3	ZNF217
ARID1A	BTG2*	CDK5	DAXX	FANCA	GATA1	IL6	MAP3K1	NF2	PIK3CA	PTPN11	SERPINB4	TEK	
ARID1B	BTK	CDK6	DCUN1D1	FANCC	GATA2	IL7R	MAP3K7	NFE2L2	PIK3CB	PTPRD	SETD2	TERT	
ARID2	BUB1B	CDK7	DDR2	FANCD2	GATA3	INPP4B	MAPK1	NFKB1	PIK3CD	PTPRT	SF3B1	TET1	
ASXL1	CALR	CDK8	DICER1	FANCE	GNA11	INSR	MAPK3	NFKBIA	PIK3CG	RAC1	SGK1	TET2	
ATM	CANX	CDK9	DNMT3A	FANCF	GNA13	IRF4	MAX	NKX2-1*	PIK3R1	RAD50	SH2D1A*	TGFBR2	
ATR	CARD11	CDKN1A	DOT1L	FANCG	GNAQ	IRS1	MCL1	NOTCH1	PIK3R2	RAD51	SLC19A1*	TMSB4X*	
ATRX	CASP8	CDKN1B	DPYD	FANCL	GNAS	IRS2*	MDM2	NOTCH2	PIK3R3	RAD51B	SLC22A2*	TNF	
AURKA	CBFB	CDKN2A	DTX1	FAS	GREM1	JAK1	MDM4	NOTCH3	PIM1	RAD51C	SLCO1B1*	TNFAIP3	

*Analysis of copy number alteration not available.

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

DISCLAIMER

Legal Statement

This test was developed by ACT Genomics and its performing characteristics were determined by ACT Genomics. This test result is to be used for clinical consultative purposes only and is not intended as a substitute for a clinical guidance of your doctor or another qualified medical practitioner. It should not be regarded as investigational or used for research.

The detection of genomic alterations does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; the detection of no genomic alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Treatment Decisions are the Responsibility of the Physician

Decisions on clinical care and treatment should be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, including physical examinations, information from other diagnostics tests and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test, or the information contained in this report.

In terms of consulting a different treating physician, the patient must file an application and fulfill the listed criteria for ACT Genomics to provide the patient's report to the assigned physician. The report may not be copied or reproduced except in its totality.

Genetic Alterations and Drugs Not Presented in Ranked Order

In this report, neither any biomarker alteration nor any drug associated with a potential clinical benefit (or potential lack of clinical benefit), are ranked in order of potential or predicted efficacy.

Level of Evidence Provided

Drugs with a potential clinical benefit (or potential lack of clinical benefit) are evaluated for level of published evidence with at least one clinical efficacy case report or preclinical study. We endeavor to keep the information in the report up to date. However, customers must be aware that scientific understanding and technologies change over time, and we make no warranty as to the accuracy, suitability or currency of information provided in this report at any time.

No Guarantee of Clinical Benefit

This report makes no promises or guarantees about the effectiveness of a particular drug or any treatment procedure in any disease or in any patient. This report also makes no promises or guarantees that a drug without an association of reportable genomic alteration will, in fact, provide no clinical benefit.

Liability

ACT Genomics is not affiliated with any medical facility or medical practitioner. We provide information for informational purposes only, therefore, ACT Genomics and their employees cannot be held responsible for any direct, indirect, special, incidental or consequential damages that may arise from the use of information provided in the report.

免責聲明

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。


證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。


責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。


SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS



1: Everolimus, Temsirolimus; 2: Trametinib, Selumetinib




1: Abemaciclib, Palbociclib, Ribociclib



1: Olaparib, Niraparib, Rucaparib, Talazoparib; 2: Dasatinib

Hedgehog Signalling



1: Sonidegib, Vismodegib

REFERENCES

1. PMID: 26559592; 2015, N Engl J Med;373(20):1984
Genetic Basis for Clinical Response to CTLA-4 Blockade in Melanoma.
2. PMID: 26359337; 2015, Science;350(6257):207-211
Genomic correlates of response to CTLA-4 blockade in metastatic melanoma.
3. PMID: 28251903; 2017, Sci Transl Med;9(379):
Integrated molecular analysis of tumor biopsies on sequential CTLA-4 and PD-1 blockade reveals markers of response and resistance.
4. PMID: 26997480; 2016, Cell;165(1):35-44
Genomic and Transcriptomic Features of Response to Anti-PD-1 Therapy in Metastatic Melanoma.
5. PMID: 25765070; 2015, Science;348(6230):124-8
Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer.
6. PMID: 26028255; 2015, N Engl J Med;372(26):2509-20
PD-1 Blockade in Tumors with Mismatch-Repair Deficiency.
7. PMID: 29863979; 2018, N Engl J Med;379(4):341-351
PD-1 Blockade with Cemiplimab in Advanced Cutaneous Squamous-Cell Carcinoma.
8. PMID: 26952546; 2016, Lancet;387(10031):1909-20
Atezolizumab in patients with locally advanced and metastatic urothelial carcinoma who have progressed following treatment with platinum-based chemotherapy: a single-arm, multicentre, phase 2 trial.
9. PMID: 27009843; 2016, Oncotarget;7(16):22857-64
Analysis of ultra-deep targeted sequencing reveals mutation burden is associated with gender and clinical outcome in lung adenocarcinoma.
10. PMID: 24839032; 2014, Breast Cancer Res Treat;146(1):211-20
Somatic mutation load of estrogen receptor-positive breast tumors predicts overall survival: an analysis of genome sequence data.
11. PMID: 15520807; 2004, Nat Rev Mol Cell Biol;5(11):875-85
The RAF proteins take centre stage.
12. PMID: 24737949; 2014, J Carcinog;13():1
BRAF and beyond: Tailoring strategies for the individual melanoma patient.
13. PMID: 12068308; 2002, Nature;417(6892):949-54
Mutations of the BRAF gene in human cancer.
14. PMID: 24071849; 2013, Nat Genet;45(10):1113-20
The Cancer Genome Atlas Pan-Cancer analysis project.
15. PMID: 20179705; 2010, Nature;464(7287):427-30
RAF inhibitors transactivate RAF dimers and ERK signalling in cells with wild-type BRAF.
16. PMID: 21388974; 2011, Mol Cancer Ther;10(3):385-94
BRAFV600E: implications for carcinogenesis and molecular therapy.

17. PMID: 19010912; 2008, Cancer Res;68(22):9375-83
Genetic predictors of MEK dependence in non-small cell lung cancer.
18. PMID: 28972961; 2017, Br J Cancer;117(10):1450-1458
Clinical significance of BRAF non-V600E mutations on the therapeutic effects of anti-EGFR monoclonal antibody treatment in patients with pretreated metastatic colorectal cancer: the Biomarker Research for anti-EGFR monoclonal Antibodies by Comprehensive Cancer genomics (BREAC) study.
19. PMID: 31924734; 2020, Clin Cancer Res;26(8):1812-1819
Trametinib Activity in Patients with Solid Tumors and Lymphomas Harboring BRAF Non-V600 Mutations or Fusions: Results from NCI-MATCH (EAY131).
20. PMID: 28947956; 2017, Oncotarget;8(36):60094-60108
Non-V600 BRAF mutations recurrently found in lung cancer predict sensitivity to the combination of Trametinib and Dabrafenib.
21. PMID: 26070258; 2015, Cancer Chemother Pharmacol;76(2):433-8
Metastatic melanoma cells with BRAF G469A mutation: nab-paclitaxel better than vemurafenib?
22. PMID: 29320312; 2018, J Clin Oncol;36(6):536-542
Targeted Therapy for Advanced Solid Tumors on the Basis of Molecular Profiles: Results From MyPathway, an Open-Label, Phase IIa Multiple Basket Study.
23. PMID: 31959346; 2020, Ann Oncol;31(2):289-294
Vemurafenib in non-small-cell lung cancer patients with BRAF^{V600} and BRAF^{nonV600} mutations.
24. PMID: 23999441; 2013, J Clin Invest;123(9):3672-7
Isocitrate dehydrogenase mutations in leukemia.
25. PMID: 21130701; 2010, Cancer Cell;18(6):553-67
Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation.
26. PMID: 22343901; 2012, Nature;483(7390):474-8
IDH mutation impairs histone demethylation and results in a block to cell differentiation.
27. PMID: 23558173; 2013, Science;340(6132):622-6
Targeted inhibition of mutant IDH2 in leukemia cells induces cellular differentiation.
28. PMID: 27956631; 2016, Proc Natl Acad Sci U S A;113(52):15084-15089
The IDH2 R172K mutation associated with angioimmunoblastic T-cell lymphoma produces 2HG in T cells and impacts lymphoid development.
29. PMID: 19378339; 2009, Int J Cancer;125(2):353-5
Mutational analysis of IDH1 codon 132 in glioblastomas and other common cancers.
30. PMID: 21598255; 2011, J Pathol;224(3):334-43
IDH1 and IDH2 mutations are frequent events in central chondrosarcoma and central and periosteal chondromas but not in other mesenchymal tumours.
31. PMID: 19246647; 2009, Am J Pathol;174(4):1149-53
IDH1 mutations are early events in the development of astrocytomas and oligodendrogliomas.
32. PMID: 22180306; 2012, Oncologist;17(1):72-9
Frequent mutation of isocitrate dehydrogenase (IDH)1 and IDH2 in cholangiocarcinoma identified through broad-

based tumor genotyping.

33. PMID: 19657110; 2009, N Engl J Med;361(11):1058-66
Recurring mutations found by sequencing an acute myeloid leukemia genome.
34. PMID: 19228619; 2009, N Engl J Med;360(8):765-73
IDH1 and IDH2 mutations in gliomas.
35. PMID: 19935646; 2009, Nature;462(7274):739-44
Cancer-associated IDH1 mutations produce 2-hydroxyglutarate.
36. PMID: 29860938; 2018, N Engl J Med;378(25):2386-2398
Durable Remissions with Ivosidenib in IDH1-Mutated Relapsed or Refractory AML.
37. PMID: 22199315; 2011, Anticancer Res;31(12):4457-63
Correlation between IDH1 gene mutation status and survival of patients treated for recurrent glioma.
38. PMID: 32416072; 2020, Lancet Oncol;21(6):796-807
Ivosidenib in IDH1-mutant, chemotherapy-refractory cholangiocarcinoma (ClarIDHy): a multicentre, randomised, double-blind, placebo-controlled, phase 3 study.
39. PMID: 25910950; 2015, Clin Cancer Res;21(16):3610-8
A Multicenter, Phase II, Randomized, Noncomparative Clinical Trial of Radiation and Temozolomide with or without Vandetanib in Newly Diagnosed Glioblastoma Patients.
40. PMID: 27231123; 2016, Cancer Discov;6(7):727-39
Isocitrate Dehydrogenase Mutations Confer Dasatinib Hypersensitivity and SRC Dependence in Intrahepatic Cholangiocarcinoma.
41. PMID: 28148839; 2017, Sci Transl Med;9(375):
2-Hydroxyglutarate produced by neomorphic IDH mutations suppresses homologous recombination and induces PARP inhibitor sensitivity.
42. PMID: 29339439; 2018, Clin Cancer Res;24(7):1705-1715
<i>IDH1/2</i> Mutations Sensitize Acute Myeloid Leukemia to PARP Inhibition and This Is Reversed by IDH1/2-Mutant Inhibitors.
43. PMID: 23558169; 2013, Science;340(6132):626-30
An inhibitor of mutant IDH1 delays growth and promotes differentiation of glioma cells.
44. PMID: 28232670; 2017, Leukemia;31(10):2020-2028
Pan-mutant-IDH1 inhibitor BAY1436032 is highly effective against human IDH1 mutant acute myeloid leukemia in vivo.
45. PMID: 20127344; 2010, Acta Neuropathol;119(4):487-94
The prognostic IDH1(R132) mutation is associated with reduced NADP+-dependent IDH activity in glioblastoma.
46. PMID: 19234488; 2009, Oncogene;28(14):1653-68
The SWI/SNF complex and cancer.
47. PMID: 24613357; 2014, Cell Rep;6(6):973-981
BAF180 promotes cohesion and prevents genome instability and aneuploidy.
48. PMID: 23867514; 2013, Cancer J;19(4):324-32
PBRM1 and BAP1 as novel targets for renal cell carcinoma.

49. PMID: 10897333; 2000, Mol Pathol;53(3):137-44
Role of chromosome 3p12-p21 tumour suppressor genes in clear cell renal cell carcinoma: analysis of VHL dependent and VHL independent pathways of tumorigenesis.
50. PMID: 29301960; 2018, Science;359(6377):801-806
Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma.
51. PMID: 30150660; 2018, Nat Genet;50(9):1271-1281
Genomic correlates of response to immune checkpoint blockade in microsatellite-stable solid tumors.
52. PMID: 26003625; 2015, Urol Oncol;33(8):340.e9-16
Decreased PBRM1 expression predicts unfavorable prognosis in patients with clear cell renal cell carcinoma.
53. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35
mTOR: from growth signal integration to cancer, diabetes and ageing.
54. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
55. PMID: 9242607; 1997, Science;277(5327):805-8
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
56. PMID: 8269512; 1993, Cell;75(7):1305-15
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
57. PMID: 1303246; 1992, Nat Genet;2(1):37-41
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
58. PMID: 18538015; 2008, BMC Cancer;8():163
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
59. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
60. PMID: 20610279; 2010, Urol Oncol;28(4):409-28
Bladder cancer or bladder cancers? Genetically distinct malignant conditions of the urothelium.
61. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56
The tuberous sclerosis complex.
62. PMID: 22923433; 2012, Science;338(6104):221
Genome sequencing identifies a basis for everolimus sensitivity.
63. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
64. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
65. PMID: 26412398; 2015, Sci Rep;5():14534
PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.

66. PMID: 11239455; 2001, Mol Cell;7(2):263-72
BRCA2 is required for homology-directed repair of chromosomal breaks.
67. PMID: 17597348; 2007, Ann Surg Oncol;14(9):2510-8
Heterogenic loss of the wild-type BRCA allele in human breast tumorigenesis.
68. PMID: 22193408; 2011, Nat Rev Cancer;12(1):68-78
BRCA1 and BRCA2: different roles in a common pathway of genome protection.
69. PMID: 27283171; 2016, J Natl Compr Canc Netw;14(6):795-806
The Relevance of Hereditary Cancer Risks to Precision Oncology: What Should Providers Consider When Conducting Tumor Genomic Profiling?
70. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
71. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
72. PMID: 28884698; 2017, Lancet Oncol;18(9):e510
Correction to Lancet Oncol 2017; 18: 1274-84.
73. PMID: 22452356; 2012, N Engl J Med;366(15):1382-92
Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer.
74. PMID: 26187614; 2015, Clin Cancer Res;21(19):4257-61
FDA Approval Summary: Olaparib Monotherapy in Patients with Deleterious Germline BRCA-Mutated Advanced Ovarian Cancer Treated with Three or More Lines of Chemotherapy.
75. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
76. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline *BRCA*-Mutated Metastatic Pancreatic Cancer.
77. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
78. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
79. PMID: 28882436; 2017, Gynecol Oncol;147(2):267-275
Antitumor activity and safety of the PARP inhibitor rucaparib in patients with high-grade ovarian carcinoma and a germline or somatic BRCA1 or BRCA2 mutation: Integrated analysis of data from Study 10 and ARIEL2.
80. PMID: 31562799; 2019, N Engl J Med;381(25):2391-2402
Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
81. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
82. PMID: 30948273; 2019, Lancet Oncol;20(5):636-648
Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial.

83. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
84. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
85. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
86. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
87. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.
88. PMID: 7550353; 1995, Nat Genet;11(2):210-2
Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.
89. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
90. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
91. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
92. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
93. PMID: 27217383; 2016, Cancer Discov;6(7):740-53
Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.
94. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
95. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
96. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
97. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.

98. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
99. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
100. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
101. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
102. PMID: 25893302; 2016, Oncogene;35(5):537-48
Role of Merlin/NF2 inactivation in tumor biology.
103. PMID: 19451229; 2009, Mol Cell Biol;29(15):4235-49
Loss of the tumor suppressor gene NF2, encoding merlin, constitutively activates integrin-dependent mTORC1 signaling.
104. PMID: 19451225; 2009, Mol Cell Biol;29(15):4250-61
NF2/merlin is a novel negative regulator of mTOR complex 1, and activation of mTORC1 is associated with meningioma and schwannoma growth.
105. PMID: 17655741; 2007, Brain Pathol;17(4):371-6
Role of NF2 haploinsufficiency in NF2-associated polyneuropathy.
106. PMID: 19545378; 2009, Orphanet J Rare Dis;4():16
Neurofibromatosis type 2 (NF2): a clinical and molecular review.
107. PMID: 21642991; 2011, Nat Genet;43(7):668-72
The nuclear deubiquitinase BAP1 is commonly inactivated by somatic mutations and 3p21.1 losses in malignant pleural mesothelioma.
108. PMID: 24393766; 2014, Oncotarget;5(1):67-77
NF2/merlin in hereditary neurofibromatosis 2 versus cancer: biologic mechanisms and clinical associations.
109. PMID: 27091708; 2016, J Clin Oncol;34(18):2115-24
Molecular Alterations and Everolimus Efficacy in Human Epidermal Growth Factor Receptor 2-Overexpressing Metastatic Breast Cancers: Combined Exploratory Biomarker Analysis From BOLERO-1 and BOLERO-3.
110. PMID: 26503204; 2016, J Clin Oncol;34(5):419-26
Correlative Analysis of Genetic Alterations and Everolimus Benefit in Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Advanced Breast Cancer: Results From BOLERO-2.
111. PMID: 24833916; 2014, Breast Cancer (Dove Med Press);6():43-57
Use of mTOR inhibitors in the treatment of breast cancer: an evaluation of factors that influence patient outcomes.
112. PMID: 25630452; 2015, Eur Urol;67(6):1195-1196
Exceptional Response on Addition of Everolimus to Taxane in Urothelial Carcinoma Bearing an NF2 Mutation.
113. PMID: 26359368; 2015, Cancer Discov;5(11):1178-93
NF2 Loss Promotes Oncogenic RAS-Induced Thyroid Cancers via YAP-Dependent Transactivation of RAS Proteins and Sensitizes Them to MEK Inhibition.

114. PMID: 24813888; 2014, Cell Rep;7(4):999-1008
Acquired resistance of EGFR-mutant lung adenocarcinomas to afatinib plus cetuximab is associated with activation of mTORC1.
115. PMID: 8906794; 1996, Nature;384(6605):176-9
Biochemical evidence that patched is the Hedgehog receptor.
116. PMID: 12016144; 2002, Carcinogenesis;23(5):727-33
Unbalanced overexpression of the mutant allele in murine Patched mutants.
117. PMID: 11130178; 2000, Cell Mol Life Sci;57(12):1720-31
Hedgehog signalling in cancer.
118. PMID: 8782823; 1996, Nat Genet;14(1):78-81
The role of the human homologue of Drosophila patched in sporadic basal cell carcinomas.
119. PMID: 8658145; 1996, Science;272(5268):1668-71
Human homolog of patched, a candidate gene for the basal cell nevus syndrome.
120. PMID: 9422511; 1998, Nature;391(6662):90-2
Activating Smoothened mutations in sporadic basal-cell carcinoma.
121. PMID: 22832583; 2012, Nature;488(7409):100-5
Dissecting the genomic complexity underlying medulloblastoma.
122. PMID: 10738305; 2000, Genes Chromosomes Cancer;28(1):77-81
Evidence that haploinsufficiency of Ptch leads to medulloblastoma in mice.
123. PMID: 19726761; 2009, N Engl J Med;361(12):1173-8
Treatment of medulloblastoma with hedgehog pathway inhibitor GDC-0449.
124. PMID: 26169613; 2015, J Clin Oncol;33(24):2646-54
Vismodegib Exerts Targeted Efficacy Against Recurrent Sonic Hedgehog-Subgroup Medulloblastoma: Results From Phase II Pediatric Brain Tumor Consortium Studies PBTC-025B and PBTC-032.
125. PMID: 25595896; 2015, Oncotarget;6(2):1190-201
Temozolomide resistance in glioblastoma occurs by miRNA-9-targeted PTCH1, independent of sonic hedgehog level.
126. PMID: 20930833; 2010, Nature;467(7316):667-8
DNA repair: A protein giant in its entirety.
127. PMID: 20729858; 2010, Nat Struct Mol Biol;17(10):1263-5
The breast cancer tumor suppressor BRCA2 promotes the specific targeting of RAD51 to single-stranded DNA.
128. PMID: 20729832; 2010, Nature;467(7316):678-83
Purified human BRCA2 stimulates RAD51-mediated recombination.
129. PMID: 22305526; 2012, Am J Hum Genet;90(2):301-7
RAD51 haploinsufficiency causes congenital mirror movements in humans.
130. PMID: 18243065; 2008, DNA Repair (Amst);7(5):686-93
The consequences of Rad51 overexpression for normal and tumor cells.
131. PMID: 24811120; 2014, Oncotarget;5(10):3261-72
Rad51 supports triple negative breast cancer metastasis.

132. PMID: 26317153; 2015, Cell Cycle;14(19):3190-202
High levels of RAD51 perturb DNA replication elongation and cause unscheduled origin firing due to impaired CHK1 activation.
133. PMID: 21807066; 2011, Biochim Biophys Acta;1816(2):209-18
RAD51 as a potential biomarker and therapeutic target for pancreatic cancer.
134. PMID: 10851081; 2000, Oncogene;19(23):2791-5
DNA repair and recombination factor Rad51 is over-expressed in human pancreatic adenocarcinoma.
135. PMID: 24741789; 2014, Rev Med Chir Soc Med Nat Iasi;118(1):133-40
Rad51 overexpression and resistance to genotoxic agents. A study in the fission yeast Schizosaccharomyces pombe.
136. PMID: 18618591; 2009, Mol Carcinog;48(2):105-9
Rad51 overexpression rescues radiation resistance in BRCA2-defective cancer cells.
137. PMID: 10807537; 2000, J Hum Genet;45(3):133-7
Identification of Rad51 alteration in patients with bilateral breast cancer.
138. PMID: 26108708; 2015, Sci Rep;5():11588
RAD51 135G>C substitution increases breast cancer risk in an ethnic-specific manner: a meta-analysis on 21,236 cases and 19,407 controls.
139. PMID: 11248061; 2001, Proc Natl Acad Sci U S A;98(6):3232-6
A single nucleotide polymorphism in the RAD51 gene modifies cancer risk in BRCA2 but not BRCA1 carriers.
140. PMID: 17999359; 2007, Am J Hum Genet;81(6):1186-200
RAD51 135G-->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies.
141. PMID: 30353044; 2018, Br J Cancer;119(11):1401-1409
Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes.
142. PMID: 24577941; 2014, Mol Cancer Ther;13(5):1170-80
The use of Olaparib (AZD2281) potentiates SN-38 cytotoxicity in colon cancer cells by indirect inhibition of Rad51-mediated repair of DNA double-strand breaks.
143. PMID: 28759753; 2017, Biomed Pharmacother;94():165-168
Inhibition of Rad51 sensitizes breast cancer cells with wild-type PTEN to olaparib.
144. PMID: 28968163; 2017, J Clin Oncol;35(32):3638-3646
MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer.
145. PMID: 28533223; 2017, Clin Cancer Res;23(17):5218-5224
MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR+/HER2- Metastatic Breast Cancer.
146. PMID: 30280641; 2018, N Engl J Med;379(23):2220-2229
First-Line Atezolizumab plus Chemotherapy in Extensive-Stage Small-Cell Lung Cancer.
147. PMID: 30345906; 2018, N Engl J Med;379(22):2108-2121
Atezolizumab and Nab-Paclitaxel in Advanced Triple-Negative Breast Cancer.
148. PMID: 27979383; 2017, Lancet;389(10066):255-265
Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial.

149. PMID: 26970723; 2016, Lancet;387(10030):1837-46
Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial.
150. PMID: 30779531; 2019, N Engl J Med;380(12):1103-1115
Avelumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma.
151. PMID: 27592805; 2016, Lancet Oncol;17(10):1374-1385
Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial.
152. PMID: 28438073; 2018, Cogn Emot;32(7):1499-1513
The meaning profiles of anxiety and depression: similarities and differences in two age groups.
153. PMID: 22529265; 2012, J Clin Oncol;30(17):2039-45
OCEANS: a randomized, double-blind, placebo-controlled phase III trial of chemotherapy with or without bevacizumab in patients with platinum-sensitive recurrent epithelial ovarian, primary peritoneal, or fallopian tube cancer.
154. PMID: 24637997; 2014, J Clin Oncol;32(13):1302-8
Bevacizumab combined with chemotherapy for platinum-resistant recurrent ovarian cancer: The AURELIA open-label randomized phase III trial.
155. PMID: 24552320; 2014, N Engl J Med;370(8):734-43
Improved survival with bevacizumab in advanced cervical cancer.
156. PMID: 23168366; 2013, Lancet Oncol;14(1):29-37
Continuation of bevacizumab after first progression in metastatic colorectal cancer (ML18147): a randomised phase 3 trial.
157. PMID: 18156031; 2007, Lancet;370(9605):2103-11
Bevacizumab plus interferon alfa-2a for treatment of metastatic renal cell carcinoma: a randomised, double-blind phase III trial.
158. PMID: 19720927; 2009, J Clin Oncol;27(28):4733-40
Bevacizumab alone and in combination with irinotecan in recurrent glioblastoma.
159. PMID: 17167137; 2006, N Engl J Med;355(24):2542-50
Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer.
160. PMID: 17442997; 2007, J Clin Oncol;25(12):1539-44
Bevacizumab in combination with oxaliplatin, fluorouracil, and leucovorin (FOLFOX4) for previously treated metastatic colorectal cancer: results from the Eastern Cooperative Oncology Group Study E3200.
161. PMID: 15175435; 2004, N Engl J Med;350(23):2335-42
Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer.
162. PMID: 20525995; 2010, N Engl J Med;362(24):2260-70
Dasatinib versus imatinib in newly diagnosed chronic-phase chronic myeloid leukemia.
163. PMID: 18541900; 2008, J Clin Oncol;26(19):3204-12
Intermittent target inhibition with dasatinib 100 mg once daily preserves efficacy and improves tolerability in imatinib-resistant and -intolerant chronic-phase chronic myeloid leukemia.
164. PMID: 17496201; 2007, Blood;110(7):2309-15
Dasatinib induces rapid hematologic and cytogenetic responses in adult patients with Philadelphia chromosome positive acute lymphoblastic leukemia with resistance or intolerance to imatinib: interim results of a phase 2 study.

165. PMID: 31590988; 2019, Lancet;394(10212):1929-1939
Durvalumab plus platinum-etoposide versus platinum-etoposide in first-line treatment of extensive-stage small-cell lung cancer (CASPIAN): a randomised, controlled, open-label, phase 3 trial.
166. PMID: 28885881; 2017, N Engl J Med;377(20):1919-1929
Durvalumab after Chemoradiotherapy in Stage III Non-Small-Cell Lung Cancer.
167. PMID: 27269937; 2016, J Clin Oncol;34(26):3119-25
Safety and Efficacy of Durvalumab (MEDI4736), an Anti-Programmed Cell Death Ligand-1 Immune Checkpoint Inhibitor, in Patients With Advanced Urothelial Bladder Cancer.
168. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
169. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
170. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
171. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
172. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
173. PMID: 28734759; 2017, Lancet Oncol;18(9):1182-1191
Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study.
174. PMID: 29562145; 2018, N Engl J Med;378(14):1277-1290
Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma.
175. PMID: 25840693; 2015, Lancet Oncol;16(5):522-30
Adjuvant ipilimumab versus placebo after complete resection of high-risk stage III melanoma (EORTC 18071): a randomised, double-blind, phase 3 trial.
176. PMID: 20525992; 2010, N Engl J Med;363(8):711-23
Improved survival with ipilimumab in patients with metastatic melanoma.
177. PMID: 28131785; 2017, Lancet Oncol;18(3):312-322
Nivolumab in metastatic urothelial carcinoma after platinum therapy (CheckMate 275): a multicentre, single-arm, phase 2 trial.
178. PMID: 27718784; 2016, N Engl J Med;375(19):1856-1867
Nivolumab for Recurrent Squamous-Cell Carcinoma of the Head and Neck.
179. PMID: 25482239; 2015, N Engl J Med;372(4):311-9
PD-1 blockade with nivolumab in relapsed or refractory Hodgkin's lymphoma.
180. PMID: 27451390; 2016, Lancet Oncol;17(9):1283-94
Nivolumab for classical Hodgkin's lymphoma after failure of both autologous stem-cell transplantation and brentuximab vedotin: a multicentre, multicohort, single-arm phase 2 trial.

181. PMID: 26027431; 2015, N Engl J Med;373(1):23-34
Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma.
182. PMID: 25399552; 2015, N Engl J Med;372(4):320-30
Nivolumab in previously untreated melanoma without BRAF mutation.
183. PMID: 26406148; 2015, N Engl J Med;373(19):1803-13
Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma.
184. PMID: 26412456; 2015, N Engl J Med;373(17):1627-39
Nivolumab versus Docetaxel in Advanced Nonsquamous Non-Small-Cell Lung Cancer.
185. PMID: 26028407; 2015, N Engl J Med;373(2):123-35
Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer.
186. PMID: 25795410; 2015, Lancet Oncol;16(4):375-84
Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial.
187. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
188. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
189. PMID: 25366685; 2015, J Clin Oncol;33(3):244-50
Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation.
190. PMID: 27959613; 2016, N Engl J Med;375(20):1925-1936
Palbociclib and Letrozole in Advanced Breast Cancer.
191. PMID: 26030518; 2015, N Engl J Med;373(3):209-19
Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer.
192. PMID: 30779529; 2019, N Engl J Med;380(12):1116-1127
Pembrolizumab plus Axitinib versus Sunitinib for Advanced Renal-Cell Carcinoma.
193. PMID: 27093365; 2016, N Engl J Med;374(26):2542-52
PD-1 Blockade with Pembrolizumab in Advanced Merkel-Cell Carcinoma.
194. PMID: 29875066; 2018, Lancet Oncol;19(7):940-952
Pembrolizumab in patients with advanced hepatocellular carcinoma previously treated with sorafenib (KEYNOTE-224): a non-randomised, open-label phase 2 trial.
195. PMID: 29658856; 2018, N Engl J Med;378(22):2078-2092
Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell Lung Cancer.
196. PMID: 28291584; 2017, Lancet Oncol;18(5):623-630
Clinical safety and activity of pembrolizumab in patients with malignant pleural mesothelioma (KEYNOTE-028): preliminary results from a non-randomised, open-label, phase 1b trial.
197. PMID: 28489510; 2017, J Clin Oncol;35(22):2535-2541
Safety and Antitumor Activity of Pembrolizumab in Advanced Programmed Death Ligand 1-Positive Endometrial Cancer: Results From the KEYNOTE-028 Study.

198. PMID: 27138582; 2016, J Clin Oncol;34(21):2460-7
Pembrolizumab in Patients With Advanced Triple-Negative Breast Cancer: Phase Ib KEYNOTE-012 Study.
199. PMID: 28081914; 2017, Lancet Oncol;18(2):212-220
Safety and activity of pembrolizumab in patients with locally advanced or metastatic urothelial cancer (KEYNOTE-012): a non-randomised, open-label, phase 1b study.
200. PMID: 27247226; 2016, Lancet Oncol;17(7):956-965
Safety and clinical activity of pembrolizumab for treatment of recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-012): an open-label, multicentre, phase 1b trial.
201. PMID: 27157491; 2016, Lancet Oncol;17(6):717-726
Pembrolizumab for patients with PD-L1-positive advanced gastric cancer (KEYNOTE-012): a multicentre, open-label, phase 1b trial.
202. PMID: 28212060; 2017, N Engl J Med;376(11):1015-1026
Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma.
203. PMID: 28441111; 2017, J Clin Oncol;35(19):2125-2132
Phase II Study of the Efficacy and Safety of Pembrolizumab for Relapsed/Refractory Classic Hodgkin Lymphoma.
204. PMID: 27718847; 2016, N Engl J Med;375(19):1823-1833
Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer.
205. PMID: 25891173; 2015, N Engl J Med;372(26):2521-32
Pembrolizumab versus Ipilimumab in Advanced Melanoma.
206. PMID: 26712084; 2016, Lancet;387(10027):1540-50
Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial.
207. PMID: 26115796; 2015, Lancet Oncol;16(8):908-18
Pembrolizumab versus investigator-choice chemotherapy for ipilimumab-refractory melanoma (KEYNOTE-002): a randomised, controlled, phase 2 trial.
208. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
209. PMID: 25981810; 2015, Lancet Oncol;16(6):716-28
Treatment with two different doses of sonidegib in patients with locally advanced or metastatic basal cell carcinoma (BOLT): a multicentre, randomised, double-blind phase 2 trial.
210. PMID: 27924459; 2016, Target Oncol;11(6):815-824
Patient-Reported Outcomes and Quality of Life with Sunitinib Versus Placebo for Pancreatic Neuroendocrine Tumors: Results From an International Phase III Trial.
211. PMID: 27836885; 2017, Ann Oncol;28(2):339-343
Sunitinib in pancreatic neuroendocrine tumors: updated progression-free survival and final overall survival from a phase III randomized study.
212. PMID: 21306237; 2011, N Engl J Med;364(6):501-13
Sunitinib malate for the treatment of pancreatic neuroendocrine tumors.
213. PMID: 16757724; 2006, JAMA;295(21):2516-24
Sunitinib in patients with metastatic renal cell carcinoma.

214. PMID: 25577718; 2015, Eur Urol;67(5):952-8
Depth of remission is a prognostic factor for survival in patients with metastatic renal cell carcinoma.
215. PMID: 27238653; 2016, Eur Urol;70(6):1006-1015
Early Tumour Shrinkage: A Tool for the Detection of Early Clinical Activity in Metastatic Renal Cell Carcinoma.
216. PMID: 17227905; 2007, Oncologist;12(1):107-13
Food and Drug Administration drug approval summary: Sunitinib malate for the treatment of gastrointestinal stromal tumor and advanced renal cell carcinoma.
217. PMID: 17046465; 2006, Lancet;368(9544):1329-38
Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial.
218. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsirrolimus, interferon alfa, or both for advanced renal-cell carcinoma.
219. PMID: 29072975; 2018, J Clin Oncol;36(1):7-13
Dabrafenib and Trametinib Treatment in Patients With Locally Advanced or Metastatic BRAF V600-Mutant Anaplastic Thyroid Cancer.
220. PMID: 27080216; 2016, Lancet Oncol;17(5):642-50
Dabrafenib in patients with BRAF(V600E)-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, phase 2 trial.
221. PMID: 25265492; 2014, N Engl J Med;371(20):1877-88
Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma.
222. PMID: 22663011; 2012, N Engl J Med;367(2):107-14
Improved survival with MEK inhibition in BRAF-mutated melanoma.
223. PMID: 22670903; 2012, N Engl J Med;366(23):2171-9
Efficacy and safety of vismodegib in advanced basal-cell carcinoma.

11110171301

ACTFusion™ Report

PATIENT	SPECIMEN	ORDERING PHYSICIAN
Name: 呂富美	Type: FFPE tissue	Name: 陳明晃醫師
Gender: Female	Date received: Nov 01, 2021	Facility: 臺北榮總
Date of Birth: Sep 30, 1964	Collection site: Liver	Tel: 886-228712121
Patient ID: 19283126	Specimen ID: S10935424	Address: 臺北市北投區石牌路二段 201 號
Diagnosis: Cholangiocarcinoma	Lab ID: AA-21-04509	
	D/ID: NA	

ABOUT ACTFusion™

The test is a next-generation sequencing (NGS) based in vitro diagnostic assay to detect fusion transcripts of 13 genes, including *ALK*, *BRAF*, *EGFR*, *FGFR1*, *FGFR2*, *FGFR3*, *MET*, *NRG1*, *NTRK1*, *NTRK2*, *NTRK3*, *RET*, and *ROS1*.

VARIANT(S) WITH CLINICAL RELEVANCE

FUSION RESULTS

No fusion gene detected in this sample.

Variant Analysis:

醫檢師張筑芎 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



Sign Off

醫檢師張筑芎 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

Not Applicable.

VARIANT INTERPRETATION

Not Applicable.

US FDA-APPROVED DRUG(S)

Not Applicable.

ONGOING CLINICAL TRIAL(S)

Clinical trials shown below were selected by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACTFusion™ GENE LIST

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1
NTRK1	NTRK2	NTRK3	RET	ROS1	-	-	-

TEST DETAILS

SPECIMEN RECEIVED



- H&E-stained section No.: S10935424
- Collection date: Oct 2020
- Collection site: Liver
- Facility retrieved: 臺北榮總
- Examined by: Dr. Yeh-Han Wang
- Estimated neoplastic nuclei (whole sample): The percentage of viable tumor cells in total cells in the whole slide (%): 25%
The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 70%

The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%

The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%

Additional comment: NA

- Manual macrodissection: Performed on the highlighted region

The outline highlights the area of malignant neoplasm annotated by a pathologist.

NEXT-GENERATION SEQUENCING (NGS) METHODS

The extracted RNA was reverse-transcribed and subjected to library construction. The quality and quantity of the amplified library was determined using the fragment analyzer (AATI) and Qubit (Invitrogen). Sequencing was performed on the Ion 540™ Chip/ Ion 550™ Chip / Ion P1™ Chip and Ion GeneStudio™ S5 Prime System / Ion Proton™ System (Life Technologies). All assays were performed in accordance with ACT Genomics testing SOPs.

Data processing and statistical analysis for the identification of relevant fusions was performed using in-house fusion calling pipeline with default parameter setting. The four internal controls for the purpose of monitoring the overall sequencing quality of the sample were built into the assay, including CHMP2A, RABA7A, GPI, and VCP. Amplification of these genes using gene specific primers was performed, and the sequencing results were applied to the analysis pipeline to assess RNA quality. The inability of the software to detect these genes was considered a run failure. To ensure optimal sequencing quality for variant analysis, all samples had to meet the following sample quality control (QC) criteria: 1) Average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) ≥ 10 (default), and 2) Total reads after sequencing $\geq 500,000$ (recommended).

Samples passed the sample QC would be subjected to the fusion analysis pipeline for fusion transcript calling. Briefly, the analysis pipeline aligned sequenced reads to a reference genome, identified regions that map to noncontiguous regions of the genome, and applied filters to exclude probable false-positive events and annotate previously characterized fusion events. A minimum of 5 reads with 3 unique sequencing start sites that cross the breakpoints was set as the cutoff value to indicate strong evidence of fusions. RNA fusions would need to be in frame in order to generate productive transcripts. In addition, databases with details for documented fusions were used to authenticate the fusion sequence identified. Known fusions were queried using Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In summary, samples with detectable fusions had to meet the following criteria: 1) Number of unique start sites (SS) for the GSP2 ≥ 3 . 2) Number of supporting reads spanning the fusion junction ≥ 5 . 3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$. 4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

Quiver Gene Fusion Database version 5.1.18

LIMITATIONS

This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

STANDARD OPERATING PROCEDURES (SOPs)

Standard operating procedures (SOPs) are shown below:

- AG2-QP-15 Specimen Management Procedure
- AG3-QP16-08 SOP of FFPE Nucleic Acid Extraction
- AG3-QP16-10 SOP of HE Staining
- AG3-QP16-17 SOP of DNA Quantification with Qubit Fluorometer
- AG3-QP16-20 SOP of CE-Fragment Analysis
- AG3-QP16-24 SOP of Ion Torrent System Sequencing Reaction
- AG3-QP16-26 SOP of Ion Chef Preparation
- AG3-QP40-08 (02) Standard protocol for variant interpretation, curation and classification
- AG3-QP16-94 (01) SOP of ACTFusion v3 Library Construction and Preparation
- AG3-QP16-36(02) SOP of Fusion Gene Detection
- AG3-QP16-41 SOP of The user manual for clinical report system (CRS)

RUN QC

- Panel: ACTFusion™
- Total reads: 1338564
- Average unique RNA Start Sites per control GSP2: 21

DISCLAIMER**Legal Statement**

This test was developed by ACT Genomics and its performing characteristics were determined by ACT Genomics. This test result is to be used for clinical consultative purposes only and is not intended as a substitute for a clinical guidance of your doctor or another qualified medical practitioner. It should not be regarded as investigational or used for research.

The detection of genomic alterations does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; the detection of no genomic alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Treatment Decisions are the Responsibility of the Physician

Decisions on clinical care and treatment should be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, including physical examinations, information from other diagnostics tests and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test, or the information contained in this report.

In terms of consulting a different treating physician, the patient must file an application and fulfill the listed criteria for ACT Genomics to provide the patient's report to the assigned physician. The report may not be copied or reproduced except in its totality.

Genetic Alterations and Drugs Not Presented in Ranked Order

In this report, neither any biomarker alteration nor any drug associated with a potential clinical benefit (or potential lack of clinical benefit), are ranked in order of potential or predicted efficacy.

Level of Evidence Provided

Drugs with a potential clinical benefit (or potential lack of clinical benefit) are evaluated for level of published evidence with at least one clinical efficacy case report or preclinical study. We endeavor to keep the information in the report up to date. However, customers must be aware that scientific understanding and technologies change over time, and we make no warranty as to the accuracy, suitability or currency of information provided in this report at any time.

No Guarantee of Clinical Benefit

This report makes no promises or guarantees about the effectiveness of a particular drug or any treatment procedure in any disease or in any patient. This report also makes no promises or guarantees that a drug without an association of reportable genomic alteration will, in fact, provide no clinical benefit.

Liability

ACT Genomics is not affiliated with any medical facility or medical practitioner. We provide information for informational purposes only, therefore, ACT Genomics and their employees cannot be held responsible for any direct, indirect, special, incidental or consequential damages that may arise from the use of information provided in the report.

免責聲明

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

REFERENCES

Not Applicable.