



## Sample Information

**Patient Name:** 柯宏美  
**Gender:** Female  
**ID No.:** E220273168  
**History No.:** 39774839  
**Age:** 60

**Ordering Doctor:** DOC3109L 邱昭華  
**Ordering REQ.:** C233PBA  
**Signing in Date:** 2021/03/17

**Path No.:** S110-98403  
**MP No.:** F21024  
**Assay:** Oncomine Focus Assay  
**Sample Type:** FFPE  
**Block No.:** S110-75569C  
**Percentage of tumor cells:** 40%  
**Note:**

## Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variant Details	2
Biomarker Descriptions	3
Relevant Therapy Summary	4
Relevant Therapy Details	9
Clinical Trials Summary	20
Alert Details	29

**Report Highlights**  
2 Relevant Biomarkers  
9 Therapies Available  
116 Clinical Trials

## Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	Not detected	NTRK1	Not detected
BRAF	Not detected	NTRK2	Not detected
EGFR	<b>EGFR p.(L858R) c.2573T&gt;G, EGFR p.(T790M) c.2369C&gt;T</b>	NTRK3	Not detected
ERBB2	Not detected	RET	Not detected
KRAS	Not detected	ROS1	Not detected
MET	Not detected		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>EGFR p.(L858R) c.2573T&gt;G</i> epidermal growth factor receptor Allele Frequency: 12.29%	<b>bevacizumab* + erlotinib</b> <sup>2</sup> <b>erlotinib + ramucirumab</b> <sup>1,2</sup> <b>osimertinib</b> <sup>1,2</sup> afatinib + cetuximab atezolizumab + bevacizumab + chemotherapy bevacizumab + gefitinib gefitinib + chemotherapy	None	100
IA	<i>EGFR p.(T790M) c.2369C&gt;T</i> epidermal growth factor receptor Allele Frequency: 8.46%	<b>osimertinib</b> <sup>1,2</sup>	None	72

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

**Tier Reference:** Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

\* Includes biosimilars

 **Alerts informed by public data sources:**  Contraindicated,  Resistance

*EGFR p.(T790M) c.2369C>T*  **gefitinib**<sup>2</sup>  
 afatinib, dacomitinib, erlotinib, gefitinib

Public data sources included in alerts: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

## Variant Details

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
EGFR	p.(T790M)	c.2369C>T	COSM6240	chr7:55249071	8.46%	NM_005228.4	missense	1998
EGFR	p.(L858R)	c.2573T>G	COSM6224	chr7:55259515	12.29%	NM_005228.4	missense	1994
JAK1	p.(=)	c.2199A>G	.	chr1:65310489	99.70%	NM_002227.3	synonymous	1986
ALK	p.(D1529E)	c.4587C>G	.	chr2:29416366	99.90%	NM_004304.4	missense	1999
ALK	p.(I1461V)	c.4381A>G	.	chr2:29416572	99.85%	NM_004304.4	missense	2000
ALK	p.(=)	c.3375C>A	.	chr2:29445458	99.90%	NM_004304.4	synonymous	1993
FGFR3	p.(=)	c.1953G>A	.	chr4:1807894	99.70%	NM_000142.4	synonymous	1992
PDGFRA	p.(=)	c.939T>G	.	chr4:55133726	45.94%	NM_006206.5	synonymous	1996
PDGFRA	p.(=)	c.1701A>G	.	chr4:55141055	99.80%	NM_006206.5	synonymous	1999
PDGFRA	p.(=)	c.2472C>T	.	chr4:55152040	45.44%	NM_006206.5	synonymous	1996
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.35%	NM_213647.2	missense	2000
EGFR	p.(=)	c.2361G>A	.	chr7:55249063	47.44%	NM_005228.4	synonymous	1996
RET	p.(=)	c.2307G>T	.	chr10:43613843	99.80%	NM_020975.4	synonymous	1996

## Biomarker Descriptions

### EGFR (epidermal growth factor receptor)

**Background:** The EGFR gene encodes the epidermal growth factor receptor (EGFR) tyrosine kinase, a member of the ERBB/human epidermal growth factor receptor (HER) family. In addition to EGFR/ERBB1/HER1, other members of the ERBB/HER family include ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4<sup>1</sup>. EGFR ligand induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways. Activation of these pathways promote cell proliferation, differentiation, and survival<sup>2,3</sup>.

**Alterations and prevalence:** Recurrent somatic mutations in the tyrosine kinase domain (TKD) of EGFR are observed in approximately 10-20% of lung adenocarcinoma, and at higher frequencies in never-smoker, female, and Asian populations<sup>4,5,6,7</sup>. The most common mutations occur near the ATP-binding pocket of the TKD and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21<sup>8</sup>. These mutations constitutively activate EGFR resulting in downstream signaling, and represent 80% of the EGFR mutations observed in lung cancer. A second group of less prevalent activating mutations include E709K, G719X, S768I, L861Q, and short in-frame insertion mutations in exon 20<sup>9,10,11,12</sup>. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations<sup>13</sup>. In contrast, a different set of recurrent activating EGFR mutations in the extracellular domain include R108K, A289V and G598V and are primarily observed in glioblastoma<sup>8,14</sup>. Amplification of EGFR is observed in several cancer types including 30% of glioblastoma, 12% of esophageal cancer, 10% of head and neck cancer, 5% of bladder cancer, and 5% of lung squamous cell carcinoma<sup>5,6,7,14,15</sup>. Deletion of exons 2-7, encoding the extracellular domain of EGFR (EGFRvIII), results in overexpression of a ligand-independent constitutively active protein and is observed in approximately 30% of glioblastoma<sup>16,17,18</sup>.

**Potential relevance:** Approved first-generation EGFR tyrosine kinase inhibitors (TKIs) include erlotinib<sup>19</sup> (2004) and gefitinib<sup>20</sup> (2015), which block the activation of downstream signaling by reversible interaction with the ATP-binding site. Although initially approved for advanced lung cancer, the discovery that drug sensitivity was associated with exon 19 and exon 21 activating mutations allowed first-generation TKIs to become subsequently approved for front-line therapy in lung cancer tumors containing exon 19 or exon 21 activating mutations. Second-generation TKIs afatinib<sup>21</sup> (2013) and dacomitinib<sup>22</sup> (2018) bind EGFR and other ERBB/HER gene family members irreversibly and were subsequently approved. First- and second-generation TKIs afatinib, dacomitinib, erlotinib, and gefitinib are recommended for the treatment NSCLC harboring EGFR exon 19 insertions, exon 19 deletions, point mutations L861Q, L858R, S768I, and codon 719 mutations, whereas most EGFR exon 20 insertions, except p.A763\_Y764insFQEA, confer resistance to the same therapies<sup>23,24,25,26</sup>. In lung cancer containing EGFR exon 19 or 21 activating mutations, treatment with TKIs is eventually associated with the emergence of drug resistance<sup>27</sup>. The primary resistance mutation that emerges following treatment with first-generation TKI is T790M, accounting for 50-60% of resistant cases<sup>8</sup>. Third generation TKIs were developed to maintain sensitivity in the presence of T790M. Osimertinib<sup>28</sup> (2015) is an irreversible inhibitor indicated for metastatic EGFR T790M positive lung cancer and for the first-line treatment of metastatic NSCLC containing EGFR exon 19 deletions or exon 21 L858R mutations. Like first-generation TKIs, treatment with osimertinib is associated with acquired resistance. In this case, resistance is associated with the C797S mutation, and occurs in 22-44% of cases<sup>27</sup>. The T790M and C797S mutations may be each selected following sequential treatment with a first-generation TKI followed by a third-generation TKI or vice versa<sup>29</sup>. T790M and C797S can occur in either cis or trans allelic orientation<sup>29</sup>. If C797S is observed following progression after treatment with a third-generation TKI in the first-line setting, sensitivity may be retained to first-generation TKIs<sup>29</sup>. If C797S co-occurs in trans with T790M following sequential treatment with first- and third-generation TKIs, patients may exhibit sensitivity to combination first- and third-generation TKIs, but resistance to third-generation TKIs alone<sup>29,30</sup>. However, C797S occurring in cis conformation with T790M, confers resistance to first- and third-generation TKIs<sup>29</sup>. Fourth-generation TKIs are in development to overcome acquired C797S and T790M resistance mutations after osimertinib treatment. EGFR targeting antibodies including cetuximab (2004), panitumumab (2006), and necitumumab (2016) are under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations. The bispecific antibody, JNJ-61186372<sup>31</sup>, targeting EGFR and MET, and the TKI mobocertinib<sup>32</sup>, each received a breakthrough designation from the FDA (2020) for NSCLC tumors harboring EGFR exon 20 insertion mutations. The Oncoprex immunogene therapy CNVN-202<sup>33</sup> in combination with osimertinib received a fast track designation from the FDA (2020) for NSCLC tumors harboring EGFR mutations that progressed on osimertinib alone. BDTX-189<sup>34</sup> was granted a fast track designation (2020) for the treatment of solid tumors harboring an EGFR exon 20 insertion mutation.

## Relevant Therapy Summary

● In this cancer type   
 ○ In other cancer type   
 ● In this cancer type and other cancer types   
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (III)
erlotinib + ramucirumab	●	●	●	●	✕
bevacizumab + erlotinib	✕	●	●	●	✕
afatinib + cetuximab	✕	●	✕	✕	✕
bevacizumab (Allergan) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Fujifilm Kyowa Kirin Biologics) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Samsung Bioepis) + erlotinib	✕	✕	●	✕	✕
atezolizumab + bevacizumab + carboplatin + paclitaxel	✕	✕	✕	●	✕
bevacizumab + gefitinib	✕	✕	✕	●	✕
gefitinib + carboplatin + pemetrexed	✕	✕	✕	●	✕
apatinib + EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (IV)
bevacizumab + osimertinib, osimertinib	✕	✕	✕	✕	● (IV)
EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (IV)
icotinib hydrochloride	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, radiation therapy	✕	✕	✕	✕	● (IV)
amivantamab, lazertinib, osimertinib	✕	✕	✕	✕	● (III)
atezolizumab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (III)
durvalumab, chemotherapy	✕	✕	✕	✕	● (III)
EGFR tyrosine kinase inhibitor, chemotherapy	✕	✕	✕	✕	● (III)
osimertinib, bevacizumab	✕	✕	✕	✕	● (III)
osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
D-0316, icotinib hydrochloride	✕	✕	✕	✕	● (II/III)
almonertinib, radiation therapy	✕	✕	✕	✕	● (II)
anlotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (II)
anlotinib hydrochloride, icotinib hydrochloride	✕	✕	✕	✕	● (II)
anlotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (II)
atezolizumab, bevacizumab	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
atezolizumab, chemotherapy	✕	✕	✕	✕	● (II)
avitinib, AZD-3759	✕	✕	✕	✕	● (II)
bevacizumab, atezolizumab	✕	✕	✕	✕	● (II)
bevacizumab, atezolizumab, chemotherapy	✕	✕	✕	✕	● (II)
bevacizumab, osimertinib	✕	✕	✕	✕	● (II)
bintrafusp alfa, chemoradiation therapy, durvalumab	✕	✕	✕	✕	● (II)
chemotherapy, atezolizumab, bevacizumab	✕	✕	✕	✕	● (II)
chemotherapy, durvalumab	✕	✕	✕	✕	● (II)
crizotinib + chemotherapy	✕	✕	✕	✕	● (II)
durvalumab, tremelimumab, chemotherapy	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor + chemotherapy, EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, apatinib	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, radiation therapy	✕	✕	✕	✕	● (II)
famitinib, almonertinib	✕	✕	✕	✕	● (II)
nivolumab, ipilimumab	✕	✕	✕	✕	● (II)
olaparib, durvalumab	✕	✕	✕	✕	● (II)
osimertinib, radiation therapy	✕	✕	✕	✕	● (II)
osimertinib, ramucirumab	✕	✕	✕	✕	● (II)
pembrolizumab, chemotherapy	✕	✕	✕	✕	● (II)
ramucirumab, chemotherapy, cytokine	✕	✕	✕	✕	● (II)
SH-1028	✕	✕	✕	✕	● (II)
tyrosine kinase inhibitors, radiation therapy	✕	✕	✕	✕	● (II)
ASK120067	✕	✕	✕	✕	● (I/II)
CBT-502, anlotinib hydrochloride	✕	✕	✕	✕	● (I/II)
DZD-9008	✕	✕	✕	✕	● (I/II)
EMB01	✕	✕	✕	✕	● (I/II)
KP-673	✕	✕	✕	✕	● (I/II)
alisertib, osimertinib	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
alisertib, sapanisertib, osimertinib	✕	✕	✕	✕	● (I)
amivantamab, lazertinib	✕	✕	✕	✕	● (I)
BBP-398	✕	✕	✕	✕	● (I)
BCA101	✕	✕	✕	✕	● (I)
C-005	✕	✕	✕	✕	● (I)
CK-101	✕	✕	✕	✕	● (I)
etrumadenant, zimberelimab, chemotherapy	✕	✕	✕	✕	● (I)
genolimzumab, fruquintinib	✕	✕	✕	✕	● (I)
lazertinib, amivantamab	✕	✕	✕	✕	● (I)
nazartinib, trametinib	✕	✕	✕	✕	● (I)
neratinib, palbociclib, everolimus, trametinib	✕	✕	✕	✕	● (I)
niraparib, osimertinib	✕	✕	✕	✕	● (I)
osimertinib, ipilimumab	✕	✕	✕	✕	● (I)
osimertinib, necitumumab	✕	✕	✕	✕	● (I)
pirotinib	✕	✕	✕	✕	● (I)
telaglenastat, sapanisertib	✕	✕	✕	✕	● (I)
telisotuzumab vedotin, osimertinib	✕	✕	✕	✕	● (I)
TNO-155, nazartinib	✕	✕	✕	✕	● (I)
TQB 3804	✕	✕	✕	✕	● (I)
TY-9591	✕	✕	✕	✕	● (I)
U3-1402	✕	✕	✕	✕	● (I)

### EGFR p.(T790M) c.2369C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (II)
anlotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (IV)
apatinib + EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (IV)
icotinib hydrochloride, radiation therapy	✕	✕	✕	✕	● (IV)
durvalumab, chemotherapy	✕	✕	✕	✕	● (III)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(T790M) c.2369C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
EGFR tyrosine kinase inhibitor, chemotherapy	✕	✕	✕	✕	● (III)
icotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (III)
osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
sintilimab, bevacizumab (Innovent Biologics), chemotherapy	✕	✕	✕	✕	● (III)
toripalimab, chemotherapy	✕	✕	✕	✕	● (III)
abivertinib	✕	✕	✕	✕	● (II)
anlotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (II)
apatinib, chemotherapy	✕	✕	✕	✕	● (II)
atezolizumab, bevacizumab	✕	✕	✕	✕	● (II)
avitinib	✕	✕	✕	✕	● (II)
bevacizumab, osimertinib	✕	✕	✕	✕	● (II)
D-0316	✕	✕	✕	✕	● (II)
durvalumab, tremelimumab, chemotherapy	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor + chemotherapy, EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, apatinib	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, radiation therapy	✕	✕	✕	✕	● (II)
famitinib, almonertinib	✕	✕	✕	✕	● (II)
icotinib hydrochloride	✕	✕	✕	✕	● (II)
KN046	✕	✕	✕	✕	● (II)
nivolumab, ipilimumab	✕	✕	✕	✕	● (II)
olaparib, durvalumab	✕	✕	✕	✕	● (II)
osimertinib, bevacizumab	✕	✕	✕	✕	● (II)
osimertinib, radiation therapy	✕	✕	✕	✕	● (II)
ramucirumab, chemotherapy, cytokine	✕	✕	✕	✕	● (II)
SH-1028	✕	✕	✕	✕	● (II)
toripalimab, anlotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (II)
tyrosine kinase inhibitors, radiation therapy	✕	✕	✕	✕	● (II)
ASK120067	✕	✕	✕	✕	● (I/II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

 In this cancer type    
  In other cancer type    
  In this cancer type and other cancer types    
  No evidence

### EGFR p.(T790M) c.2369C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
DZD-9008	×	×	×	×	● (I/II)
EMB01	×	×	×	×	● (I/II)
KP-673	×	×	×	×	● (I/II)
alisertib, osimertinib	×	×	×	×	● (I)
alisertib, sapanisertib, osimertinib	×	×	×	×	● (I)
amivantamab	×	×	×	×	● (I)
BBP-398	×	×	×	×	● (I)
BCA101	×	×	×	×	● (I)
BEBT-109	×	×	×	×	● (I)
C-005	×	×	×	×	● (I)
CK-101	×	×	×	×	● (I)
lazertinib, amivantamab	×	×	×	×	● (I)
nazartinib, trametinib	×	×	×	×	● (I)
neratinib, palbociclib, everolimus, trametinib	×	×	×	×	● (I)
osimertinib, necitumumab	×	×	×	×	● (I)
Palcitoclast, osimertinib	×	×	×	×	● (I)
pirotenib	×	×	×	×	● (I)
SPH3348, osimertinib	×	×	×	×	● (I)
telisotuzumab vedotin, osimertinib	×	×	×	×	● (I)
TQB 3804	×	×	×	×	● (I)
TQB3456	×	×	×	×	● (I)
TY-9591	×	×	×	×	● (I)
U3-1402	×	×	×	×	● (I)
YK-029A	×	×	×	×	● (I)
YZJ-0318	×	×	×	×	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2020-12-16. For the most up-to-date information, search [www.fda.gov](http://www.fda.gov).

#### EGFR p.(L858R) c.2573T>G

#### ● erlotinib + ramucirumab

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2020-07-06

**Variant class:** EGFR L858R mutation

##### Indications and usage:

CYRAMZA® is a human vascular endothelial growth factor receptor 2 (VEGFR2) antagonist indicated:

- as a single agent or in combination with paclitaxel, for treatment of advanced or metastatic gastric or gastro-esophageal junction adenocarcinoma with disease progression on or after prior fluoropyrimidine- or platinum-containing chemotherapy.
- in combination with erlotinib, for first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) mutations.
- in combination with docetaxel, for treatment of metastatic non-small cell lung cancer with disease progression on or after platinum-based chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving CYRAMZA®.
- in combination with FOLFIRI, for the treatment of metastatic colorectal cancer with disease progression on or after prior therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine.
- as a single agent, for the treatment of hepatocellular carcinoma in patients who have an alpha fetoprotein of  $\geq 400$  ng/mL and have been treated with sorafenib.

##### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/125477s037lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125477s037lbl.pdf)

#### ● osimertinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2020-12-18

**Variant class:** EGFR L858R mutation

##### Indications and usage:

TAGRISSO® is a kinase inhibitor indicated for:

- as adjuvant therapy after tumor resection in adult patients with non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the first-line treatment of adult patients with metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the treatment of adult patients with metastatic EGFR T790M mutation positive NSCLC, as detected by an FDA-approved test, whose disease has progressed on or after EGFR TKI therapy.

##### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/208065s021lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208065s021lbl.pdf)

**EGFR p.(T790M) c.2369C>T****● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Label as of:** 2020-12-18**Variant class:** EGFR T790M mutation**Indications and usage:**

TAGRISSO® is a kinase inhibitor indicated for:

- as adjuvant therapy after tumor resection in adult patients with non-small cell lung cancer (NSCLC) whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the first-line treatment of adult patients with metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test.
- the treatment of adult patients with metastatic EGFR T790M mutation positive NSCLC, as detected by an FDA-approved test, whose disease has progressed on or after EGFR TKI therapy.

**Reference:**[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/208065s021lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208065s021lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2020-12-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### EGFR p.(L858R) c.2573T>G

#### osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

#### afatinib + cetuximab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Progression (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

#### bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

#### erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Other recommended intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

**EGFR p.(L858R) c.2573T>G (continued)****● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR L858R mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]**● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Leptomeningeal Metastases, Spine Metastases (Line of therapy not specified)

**Reference:** NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 3.2020]**EGFR p.(T790M) c.2369C>T****● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR T790M mutation**NCCN Recommendation category:** 1**Population segment (Line of therapy):**

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression, Asymptomatic, Symptomatic (Subsequent therapy)

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]**● osimertinib****Cancer type:** Non-Small Cell Lung Cancer**Variant class:** EGFR T790M mutation**NCCN Recommendation category:** 2A**Population segment (Line of therapy):**

- Brain Metastases, Leptomeningeal Metastases, Spine Metastases (Line of therapy not specified)

**Reference:** NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 3.2020]

## EGFR p.(T790M) c.2369C>T (continued)

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Leptomeningeal Metastases, Spine Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 3.2020]

## Current EMA Information

- ☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

EMA information is current as of 2020-12-16. For the most up-to-date information, search [www.ema.europa.eu/ema](http://www.ema.europa.eu/ema).

### EGFR p.(L858R) c.2573T>G

#### ● bevacizumab (Allergan) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-11-03

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/mvasi-epar-product-information_en.pdf)

#### ● bevacizumab (Fujifilm Kyowa Kirin Biologics) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-11-16

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/equidacent-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/equidacent-epar-product-information_en.pdf)

#### ● bevacizumab (Samsung Bioepis) + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-12-09

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/aybintio-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/aybintio-epar-product-information_en.pdf)

#### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-03-11

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/avastin-epar-product-information_en.pdf)

#### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-07-02

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/cyramza-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/cyramza-epar-product-information_en.pdf)

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-10-16

Variant class: EGFR L858R mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information_en.pdf)

## EGFR p.(T790M) c.2369C>T

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-10-16

Variant class: EGFR T790M mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tagrisso-epar-product-information_en.pdf)

## Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2020-12-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### EGFR p.(L858R) c.2573T>G

#### ● atezolizumab + bevacizumab + carboplatin + paclitaxel

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR L858R mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Non-squamous Cell; Metastatic (First-line therapy); ESMO-MCBS v1.1 score: 3
- Metastatic (Second-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFRi sensitizing mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Advanced (First-line therapy); ESMO-MCBS v1.1 score: 4

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

#### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]



**EGFR p.(L858R) c.2573T>G (continued)****● bevacizumab + gefitinib**

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

**● erlotinib + ramucirumab**

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

**● gefitinib + carboplatin + pemetrexed**

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

**● bevacizumab + erlotinib**

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Stage IV (First-line therapy); ESMO-MCBS v1.1 score: 3

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Advanced (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● bevacizumab + erlotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(L858R) c.2573T>G (continued)

### ● bevacizumab + gefitinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● erlotinib + ramucirumab

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

### ● gefitinib + carboplatin + pemetrexed

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR activating mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Stage IV (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## EGFR p.(T790M) c.2369C>T

### ● osimertinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: EGFR T790M mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Stage IV (Second-line therapy); ESMO-MCBS v1.1 score: 4

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

## Clinical Trials Summary

### EGFR p.(L858R) c.2573T>G + EGFR p.(T790M) c.2369C>T

NCT ID	Title	Phase
NCT04552613	Efficacy of Targeted Therapy Combined Chemotherapy in Advanced EGFR Positive NSCLC Patients With Concurrent Driver Gene Mutations	III
NCT04035486	A Phase III, Open-label, Randomized Study of Osimertinib With or Without Platinum Plus Pemetrexed Chemo, as First-line Treatment in Patients With Epidermal Growth Factor Receptor (EGFR) Mutation Positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer (FLAURA2)	III
NCT04029350	A Multi-center, One-arm, Phase II Trial of Anlotinib Combined With Osimertinib as the Second-line Treatment in Stage IIIB-IV NSCLC With Confirmed EGFRm and T790M.	II
NCT04426825	A Single Arm, Phase II Study of Atezolizumab (MPDL3280A, Anti-PD-L1 Antibody) in Combination With Bevacizumab in Patients With EGFR Mutation Positive Stage IIIB/IV Non-Squamous Non-Small Cell Lung Cancer Pretreated With Epidermal Growth Factor Receptor Tyrosine-Kinase Inhibitors	II
NCT03994393	A Phase II Trial of Durvalumab (MEDI4736) and Tremelimumab With Chemotherapy in Metastatic EGFR Mutant Non-squamous Non-small Cell Lung Cancer (NSCLC) Following Progression on EGFR Tyrosine Kinase Inhibitors (TKIs)	II
No NCT ID	A Phase II Study Of Osimertinib For Untreated CNS Metastasis, EGFR T790M-Positive Non-Small Cell Lung Cancer(LOGIK1603/ WJOG9116L).	II
NCT02811354	Phase II Study of AZD9291 in Patients With Advanced Stage Non-small Cell Lung Cancer Following Prior EGFR TKI Therapy With EGFR and T790M Mutations Detected in Plasma Circulating Tumor DNA (PLASMA)	II
NCT02824952	Neo-adjuvant Trial With AZD9291 in EGFRm+ Stage IIIA/B NSCLC - a Phase II Open-label Study	II
NCT03823807	A Multicenter, Open-label, Phase II Study to Evaluate the Safety and Efficacy of SH-1028 in Locally Advanced or Metastatic NSCLC	II
NCT03502850	A Phase I/II Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumour Activity of ASK120067 in Patients With Locally Advanced or Metastatic T790M Mutation-positive Non-Small Cell Lung Cancer Who Have Progressed Following Prior Therapy With an Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Agent	I/II
NCT03516214	An Open-label, Multicenter, Phase I Dose-escalation Trial of EGF816 and Trametinib in Patients With Non-small Cell Lung Cancer and Acquired EGFR p.T790M-positive Resistance to 1st or 2nd Generation EGFR TKI Therapy	I
NCT02496663	A Phase I Trial of AZD9291 and Necitumumab in EGFR-Mutant Non-small Cell Lung Cancer After Progression on a Previous EGFR Tyrosine Kinase Inhibitor	I
NCT03618043	A Phase I, Open-label Study to Assess the Safety and Tolerability of Ascending Doses of SH-1028 Tablets in Patients With Advanced Solid Cancer	I
NCT02099058	A Multicenter, Phase I/Ib, Open-Label, Dose-Escalation Study of ABBV-399, an Antibody Drug Conjugate, in Subjects With Advanced Solid Tumors	I
NCT04204473	Phase I, Open-label, Single-arm Study to Evaluate the Safety, Tolerance, Pharmacokinetics and Preliminary Efficacy of TY-9591 Tablets in Advanced NSCLC Patients With Epidermal Growth Factor Receptor( EGFR) Positive Mutation	I
NCT03260491	A Multicenter, Open-Label Phase I Study of U3-1402 in Subjects With Metastatic or Unresectable Non-small Cell Lung Cancer.	I
NCT04233021	A Phase II, Multi-centre Study, to Evaluate the Efficacy and Safety of Osimertinib Treatment for Patients With EGFR-mutated Non-small Cell Lung Cancer (NSCLC) With Brain or Leptomeningeal Metastases	II

## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G

NCT ID	Title	Phase
No NCT ID	The Efficacy and Safety of Osimertinib Combined with Bevacizumab in the Treatment of SD Patients with Non-Squamous Cell Lung Cancer	IV
NCT01665417	Randomized, Open Label, Positive Controlled, Multicenter Trial to Evaluate Icotinib as First-line and Maintenance Treatment in EGFR Mutated Patients With Lung Adenocarcinoma	IV
NCT02103257	Sequential Icotinib Plus Chemotherapy Versus Icotinib Alone as First-line Treatment in Stage IIIB/IV Lung Adenocarcinoma: a Randomized, Open-label, Multicenter Study	IV
NCT04487080	A Phase III, Randomized Study of Amivantamab and Lazertinib Combination Therapy Versus Osimertinib Versus Lazertinib as First-Line Treatment in Patients With EGFR-Mutated Locally Advanced or Metastatic Non-Small Cell Lung Cancer.	III
NCT04058704	A Multi-center, Prospective Study to Determine the Efficiency of Icotinib Combined With Radiation Therapy Early Intervention or Late Intervention For NSCLC Patients With Brain Metastases and EGFR(Epidermal Growth Factor Receptor) Mutation	III
NCT03521154	A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter, International Study of Osimertinib as Maintenance Therapy in Patients With Locally Advanced, Unresectable EGFR Mutation-positive Non-Small Cell Lung Cancer (Stage III) Whose Disease Has Not Progressed Following Definitive Platinum-based Chemoradiation Therapy (LAURA)	III
NCT04181060	Randomized Phase III Study of Combination AZD9291 (Osimertinib) and Bevacizumab Versus AZD9291 (Osimertinib) Alone as First-Line Treatment for Patients With Metastatic EGFR-Mutant Non-Small Cell Lung Cancer (NSCLC)	III
NCT04206072	A Phase II/III, Open-Label, Randomised Study to Assess the Safety and Efficacy of D-0316 Versus Icotinib as First Line Treatment in Patients With EGFR Sensitising Mutation, Locally Advanced or Metastatic NSCLC	II/III
NCT04636593	Almonertinib With Concurrent Radiotherapy in The Treatment of Unresectable, Stage ? Non-small-cell Lung Cancer Harboring EGFR Mutations: A Phase ? Cohort Study	II
NCT03736837	A Multi-center, One-arm Clinical Study of Anlotinib Combined With Icotinib as the First-line Treatment in Patients With EGFR Mutation-positive Advanced NSCLC. The Trial Aims to Evaluate the Efficacy and Safety of This Treatment.	II
No NCT ID	Phase II Study of Platinum-Based Doublet Chemotherapy Plus Atezolizumab, In Completely Resected, P-Stage II-IIIA NSCLC Patients Harboring EGFR Mutation. (WJOG11719L Investigator-Initiated Clinical Trial)	II
NCT04099836	Single Arm Phase II Trial of Atezolizumab and Bevacizumab in Epidermal Growth Factor Receptor (EGFR) Mutant Non-Small Cell Lung Cancer in Patients With Progressive Disease After Receiving Osimertinib (TOP 1901).	II
No NCT ID	Low-dose Decitabine plus Crizotinib in EGFR-Mutation-Positive Advanced Non-Small Cell Lung Cancer Patients :A Open-Label,Single-Arm Phase II Study	II
NCT03349203	Icotinib as Neoadjuvant and Adjuvant Therapy in EGFR-mutant Stage IIIB or Oligometastasis Non-small Cell Lung Cancer: a Single Arm, Phase II Clinical Study	II
NCT03396185	Icotinib as Consolidation Therapy After Synchronous or Sequential Chemoradiotherapy in Stage IIIA-IIIB Non-small Cell Lung Cancer With EGFR Sensitive Mutation: A Single Center, Single Arm, Open Label and Prospective Clinical Study	II
NCT03749213	Icotinib as Neoadjuvant Therapy in EGFR-mutant Stage IIIA-N2 Non-small Cell Lung Cancer: a Single Arm, Phase II Clinical Study	II
NCT02726568	A Phase II Study to Determine the Efficacy and Safety of High Dose Icotinib Combined With Stereotatic Radiosurgery for NSCLC Patients Harboring EGFR Mutation With Brain Metastases	II

## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G (continued)

NCT ID	Title	Phase
No NCT ID	A Phase II trial of Osimertinib for Elderly patients with Advanced or Postoperative Recurrent Non-Small-Cell Lung Cancer (SPIRAL-0 Study)	II
NCT03433469	A Phase II Study to Evaluate Neoadjuvant Osimertinib Therapy in Patients with Surgically Resectable, EGFR-Mutant Non-Small Cell Lung Cancer	II
NCT03969823	Whole Genomic Landscape of EGFR Mutation-Positive Advanced Non-Small Cell Lung Cancer Treated With First-Line Osimertinib (WARRIOR)	II
NCT03497767	A Randomised Phase II Trial of Osimertinib With or Without Stereotactic Radiosurgery for EGFR Mutated Non-Small Cell Lung Cancer (NSCLC) With Brain Metastases	II
NCT03769103	Open Label, Multicenter, Phase II Study of Patients With Treatment Naive Metastatic Epidermal Growth Factor Receptor (EGFR) Mutation-Positive Non-Small Cell Lung Cancer (NSCLC) With Brain Metastases Randomized to Stereotactic Radiosurgery (SRS) and Osimertinib or Osimertinib Alone	II
NCT03909334	An Open-Label Randomized Phase II Study of Combining Osimertinib With and Without Ramucirumab in Tyrosine Kinase Inhibitor (TKI)-naïve Epidermal Growth Factor Receptor (EGFR)-Mutant Locally Advanced or Metastatic NSCLC	II
NCT03242915	Phase II Multi-center Study of Pembrolizumab in Combination With Platinum-based Doublet Chemotherapy in NSCLC (Non-small Cell Lung Cancer) Patients With Targetable Genetic Alterations in Their Tumor Previously Treated With Appropriate Targeted Agents With Progressive Disease	II
NCT03446417	A Phase 1/2 Open Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of ZN-e4 (KP-673) in Patients With Advanced Non-Small Cell Lung Cancer with Activating Epidermal Growth Factor Receptor (EGFR) Mutations	I/II
NCT04085315	A Phase I/Ib Study of Alisertib in Combination With Osimertinib in Metastatic EGFR-mutant Lung Cancer	I
NCT04479306	A Ph Ib Study of Osimertinib + Alisertib or Sapanisertib for Osimertinib-Resistant EGFR Mutant Non-Small Cell Lung Cancer (NSCLC) (Crossover Study)	I
NCT02609776	A Phase I, First-in-Human, Open-Label, Dose Escalation Study of JNJ-61186372, a Human Bispecific EGFR and cMet Antibody, in Subjects With Advanced Non-Small Cell Lung Cancer	I
No NCT ID	Phase I Clinical Study of C-005 Tablet In The Treatment Of Advanced Non-Small Cell Lung Cancer	I
No NCT ID	Phase I Clinical Study of Safety, Tolerability, Pharmacokinetics and Initial Efficacy of RX518 in Patients with Advanced Non-small Cell Lung Cancer	I
No NCT ID	The Clinical Observational Study of Neoadjuvant Target Therapy in Stage IIIB and IV Non-small Cell Lung Cancer Patients	I
NCT03976856	A Phase Ib Clinical Study With Extension Phase to Evaluate Safety and Efficacy of Genolimzumab (GB226) in Combination With Fruquintinib in the Treatment of Relapsed or Metastatic NSCLC Patients	I
No NCT ID	Study Of Immunologic Factor In Re-Biopsy Specimen, Peritumoral BALF, And The Peripheral Blood For Predicting Response To Osimertinib In NSCLC Patients	I
NCT01996098	A Multicenter, Randomized, Phase III Trial of Chemotherapy Followed by 6-month or 12-month Icotinib Versus Chemotherapy as Adjuvant Therapy in Stage IIA-IIIA Non-small Cell Lung Cancer Harboring Epidermal Growth Factor Receptor Mutation	III
NCT03574402	An Open-label, Multi-center, Phase II Umbrella Study to Assess Efficacy of Targeted Therapy or Immunotherapy Directed by Next Generation Sequencing (NGS) in Chinese Patients With Advanced NSCLC (TRUMP)	II

## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G (continued)

NCT ID	Title	Phase
NCT04042558	A Multicentre Phase II, Open-label, Non-randomized Study Evaluating Platinum-Pemetrexed-Atezolizumab (+/- Bevacizumab) for Patients With Stage IIIB/IV Non-squamous Non-small Cell Lung Cancer With EGFR Mutations, ALK Rearrangement or ROS1 Fusion Progressing After Targeted Therapies	II
NCT03840902	A Multicenter, Double Blind, Randomized, Controlled Study of M7824 With Concurrent Chemoradiation Followed by M7824 Versus Concurrent Chemoradiation Plus Placebo Followed by Durvalumab in Participants With Unresectable Stage III Non-small Cell Lung Cancer	II
NCT03944772	A Biomarker-directed Phase II Platform Study in Patients With Advanced Non-Small Lung Cancer Whose Disease Has Progressed on First-Line Osimertinib Therapy	II
NCT02264210	A Randomized, Phase II Trial of Icotinib Versus Observation as Adjuvant Treatment in Stage IB Non-Small Cell Lung Cancer Harboring Activating Epidermal Growth Factor Receptor Mutation	II
NCT02820116	An Open-label, Multicenter, Single-arm, Phase II Clinical Study of Icotinib for IIIA - IIIB NSCLC Patients with Epidermal Growth Factor Receptor Mutation	II
NCT03983928	A Phase Ib, Open-label, Single Center, Non-randomized Study for Safety and Efficacy of TQB2450 Combined With Anlotinib in Subjects With Advanced Mutation Positive Non-Small Cell Lung Cancer	I/II
NCT03846310	A Phase I/Ib Study to Evaluate the Safety and Tolerability of Immunotherapy Combinations in Participants With Lung Cancer	I
NCT04141644	A Phase Ib Study to Evaluate the Safety and Efficacy of Osimertinib in Combination With Ipilimumab in Patients With EGFR Mutated Non-Small-Cell Lung Cancer Tumors	I
No NCT ID	The Continuous Evaluation of EGFR Mutation in EGFR-mutation Positive Lung Cancer Patients During EGFR TKI Treatment.	IV
NCT02404675	High Dose Icotinib in Advanced Non-small Cell Lung Cancer With EGFR 21 Exon Mutation (INCREASE): a Randomized, Open-label Study	IV
NCT03991403	Study of Atezolizumab in Combination With Carboplatin + Paclitaxel + Bevacizumab vs With Pemetrexed + Cisplatin or Carboplatin With Stage IV NON-SQUAMOUS NON-SMALL CELL LUNG CANCER With EGFR(+) or ALK(+)	III
NCT03786692	TH-138: Phase II Randomized Trial of Carboplatin + Pemetrexed + Bevacizumab, With or Without Atezolizumab in Stage IV Non-squamous NSCLC Patients Who Harbor a Sensitizing EGFR Mutation or Have Never Smoked	II
NCT02044328	Icotinib as an Adjuvant Therapy for Patients With Stage IIA-IIIA Adenocarcinoma With EGFR Mutation: a Prospective, Exploratory Study	II
NCT03804580	First-Line Treatment With Osimertinib In EGFR-Mutated Non-Small Cell Lung Cancer, Coupled To Extensive Translational Studies	II
NCT04410796	A Phase 2 Randomized Study of Osimertinib Versus Osimertinib Plus Chemotherapy for Patients With Metastatic EGFR-Mutant Lung Cancers That Have Detectable EGFR-Mutant cfDNA in Plasma After Initiation of Osimertinib	II
NCT03667820	Phase II Trial of Osimertinib in Combination With Stereotactic Ablative Radiation (SABR) in EGFR Mutant Advanced Non-Small Cell Lung Cancer (NSCLC)	II
NCT03891615	Phase I Study of Niraparib in Combination With Osimertinib in EGFR-Mutated Advanced Lung Cancer	I
NCT04250545	A Phase I Trial of MLN0128 (Sapanisertib) and CB-839 HCl (Telaglenastat) in Advanced NSCLC Patients	I



## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G (continued)

NCT ID	Title	Phase
NCT03114319	An Open-label, Multi-center, Phase I, Dose Finding Study of Oral TNO155 in Adult Patients With Advanced Solid Tumors	I
No NCT ID	Apatinib Combined With EGFR-TKI For Patients With EGFR Mutation Who Failed EGFR-TKI: A Prospective Study	IV
No NCT ID	Clinical Study Of Combined Action Of Icotinib And Brain Radiotherapy On EGFR-Mutated Non-Small-Cell Lung Cancer Patients With Brain Metastasis	IV
NCT03800134	A Phase III, Double-blind, Placebo-controlled, Multi-center International Study of Neoadjuvant/Adjuvant Durvalumab for the Treatment of Patients With Resectable Stages II and III Non-small Cell Lung Cancer (AEGEAN)	III
NCT03992885	Combination Therapy With Icotinib, Pemetrexed and Platinum in Patients With Metastatic Non-squamous Non-small Cell Lung Cancer With EGFR Mutations Who Did Not Progress After Pemetrexed in Combination With Platinum-based Chemotherapy:a Single-arm, Open, Multicenter Clinical Study.	III
NCT04619563	A Single-arm Exploratory Clinical Study of Anlotinib Hydrochloride Combined With Docetaxel in EGFR Mutations Advanced Non Small Cell Lung Cancer Patients Who Have Progressed After Targeted Therapy and Chemotherapy	II
No NCT ID	Osimertinib Combined Bevacizumab in Untreated Epidermal Growth Factor Receptor Mutated Non-small-cell Lung Cancer Patients with Malignant Pleural And/Or Pericardial Effusion -phase II Trial	II
No NCT ID	Randomized Controlled Trial for EGFR-TKIs Plus S-1 or EGFR-TKIs as the First-Line Therapy for Patients with Advanced Non-small Cell Lung Cancer Harboring EGFR Mutations	II
No NCT ID	Single arm, Exploratory Study for Apatinib mesylate Combined with EGFR-TKI in Patients with EGFR Mutation-positive Advanced Non-squamous Non-small-cell Lung Cancer	II
No NCT ID	EGFR-TKI Combined With Stereotactic Body Radiation Therapy Versus TKI alone for Stage IV Oncogene-Driven Non-Small Cell Lung Cancer.	II
NCT03904823	An Open, Single-arm, Multi-center, Phase II Clinical Trial of Famitinib Combined With Epidermal Growth Factor Receptor (EGFR) Inhibitor HS-10296 in Patients With Advanced EGFR-mutant Non-Small Cell Lung Cancer (NSCLC)	II
NCT02960607	A Phase II Study of High-dose Icotinib in Previously Treated Non-small Cell Lung Cancer Patients With Epidermal Growth Factor Receptor Mutation	II
NCT03091491	Randomised Phase II Study of Nivolumab Versus Nivolumab and Ipilimumab Combination in EGFR Mutant Non-small Cell Lung Cancer	II
NCT04538378	Phase II Trial of Olaparib (LYNPARZA) Plus Durvalumab (IMFINZI) in EGFR-Mutated Adenocarcinomas That Transform to Small Cell Lung Cancer (SCLC) and Other Neuroendocrine Tumors.	II
NCT03460275	Osimertinib as First-line Therapy for Patients With EGFR Mutation-positive Locally Advanced or Metastatic Non-squamous Non-Small Cell Lung Cancer(NSCLC), a Single-Arm, Open-Label, Prospective, Multicenter, Phase II Clinical Trial	II
NCT04425681	Phase II Study of Osimertinib With Bevacizumab for Leptomeningeal Metastasis From EGFR-mutation Non-Small Cell Lung Cancer	II
No NCT ID	Phase II Trial Of Docetaxel Plus Ramucirumab Combination Therapy In Patients With Advanced EGFR Gene Mutation Positive Advanced Stage Non-Squamous Cell Non small Cell Lung Cancer	II
No NCT ID	Clinical Study of Combined Action of the First Generation of TKIs and Brain Radiotherapy on EGFR-Mutated Non-Small-Cell Lung Cancer Patients with Brain Metastasis	II



## Clinical Trials Summary (continued)

### EGFR p.(L858R) c.2573T>G (continued)

NCT ID	Title	Phase
NCT03974022	A Phase I/II, Open-Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumor Efficacy of DZD9008 in Patients With Advanced Non-Small Cell Lung Cancer (NSCLC) With EGFR or HER2 Mutation	I/II
NCT03797391	First-in-human, Phase I/II, Multicenter, Open-Label Study of EMB-01 in Patients With Advanced/ Metastatic Solid Tumors	I/II
NCT04528836	A Phase I/IB First-in-Human Study of the SHP2 Inhibitor BBP-398 (Formerly Known as IACS-15509) in Patients With Advanced Solid Tumors	I
No NCT ID	Phase I Study of DZD9008 in EGFR or HER2 Mutant NSCLC Chinese Patients	I
NCT04077463	An Open-label Phase 1/1b Study to Evaluate the Safety and Pharmacokinetics of JNJ-73841937 (Lazertinib), a Third Generation EGFR-TKI, as Monotherapy or in Combinations With JNJ-61186372, a Human Bispecific EGFR and cMet Antibody in Participants With Advanced Non-Small Cell Lung Cancer	I
No NCT ID	Pharmacokinetic and dose finding study of osimertinib in patients with impaired renal function and low body weight	I
NCT03535363	Phase I Trial of Osimertinib With Stereotactic Radiosurgery (SRS) in Patients With Brain Metastases From EGFR Positive Non-Small-Cell Lung Cancer (NSCLC)	I
No NCT ID	Phase I Clinical Study With Advanced Solid Tumors KBP-5209 Treatment	I
NCT03346811	Efficiency of Icotinib in Plasma ctDNA EGFR Mutation-positive Patients Diagnosed With Lung Cancer:a Single Arm,Multi-center,Open-label Study	II
NCT03065387	Phase I Study of the Pan-ERBB Inhibitor Neratinib Given in Combination With Everolimus, Palbociclib, or Trametinib in Advanced Cancer Subjects With EGFR Mutation/Amplification, HER2 Mutation/ Amplification, or HER3/4 Mutation or KRAS Mutation	I
NCT04128085	A Phase I, Open-label, Multicenter, Dose Escalation and Expansion Study to Evaluate the Tolerance and Pharmacokinetics of TQB3804 in Subjects With Advanced Malignant Tumors	I
NCT04429542	First-in-Human, Phase I/Ib, Open-label, Multicenter Study of Bifunctional EGFR/TGFβ Fusion Protein BCA101 Alone and in Combination With Pembrolizumab in Patients With EGFR-Driven Advanced Solid Tumors	I

### EGFR p.(T790M) c.2369C>T

NCT ID	Title	Phase
No NCT ID	Clinical Study Anlotinib Combined with Osimertinib in the Treatment of Advanced Non-Squamous Non-Small Cell Lung Cancer with 1st Generation and 2nd Generation EGFR-TKI Resistance with T790M Mutation.	IV
NCT03300115	Single-arm,Multi-center,Phase II Clinical Trial of the Efficacy and Safety of AC0010 in the Treatment of EGFR T790M Mutation-positive Patients With Advanded NSCLC	II
NCT03574402	An Open-label, Multi-center, Phase II Umbrella Study to Assess Efficacy of Targeted Therapy or Immunotherapy Directed by Next Generation Sequencing (NGS) in Chinese Patients With Advanced NSCLC (TRUMP)	II
NCT03861156	A Phase II Study to Assess the Safety and Efficacy of D0316 in Patients With Locally Advanced/ Metastatic Non Small Cell Lung Cancer Whose Tumors Are Epidermal Growth Factor Receptor Mutation Positive	II

## Clinical Trials Summary (continued)

### EGFR p.(T790M) c.2369C>T (continued)

NCT ID	Title	Phase
No NCT ID	An Evaluation of Tumor Response to Osimertinib by Early FDG-PET Finding In Patients with T790M Positive EGFR Mutated Non-Small Cell Lung Cancer	II
No NCT ID	Phase II Study of Osimertinib in Patients with Non-Small Cell Lung Cancer Who Acquired Resistance to Afatinib (ALCSG-04)	II
NCT03433469	A Phase II Study to Evaluate Neoadjuvant Osimertinib Therapy in Patients with Surgically Resectable, EGFR-Mutant Non-Small Cell Lung Cancer	II
NCT03497767	A Randomised Phase II Trial of Osimertinib With or Without Stereotactic Radiosurgery for EGFR Mutated Non-Small Cell Lung Cancer (NSCLC) With Brain Metastases	II
NCT04316351	Efficacy and Safety of Toripalimab (JS001) Combined With Pemetrexed and Anlotinib for Patients With T790M Positive Non-Small Cell Lung Cancer After Osimertinib Resistance: a Phase II, Multi-center, Single Arm Study	II
NCT03446417	A Phase 1/2 Open Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics, and Anti-tumor Activity of ZN-e4 (KP-673) in Patients With Advanced Non-Small Cell Lung Cancer with Activating Epidermal Growth Factor Receptor (EGFR) Mutations	I/II
NCT04085315	A Phase I/Ib Study of Alisertib in Combination With Osimertinib in Metastatic EGFR-mutant Lung Cancer	I
NCT04479306	A Ph Ib Study of Osimertinib + Alisertib or Sapanisertib for Osimertinib-Resistant EGFR Mutant Non-Small Cell Lung Cancer (NSCLC) (Crossover Study)	I
No NCT ID	Evaluation of Phase I Clinical Trial of BEBT-109 in Patients with Advanced Non-small Cell Lung Cancer	I
No NCT ID	Phase I Clinical Study of C-005 Tablet In The Treatment Of Advanced Non-Small Cell Lung Cancer	I
No NCT ID	Phase I Clinical Study of Safety, Tolerability, Pharmacokinetics and Initial Efficacy of RX518 in Patients with Advanced Non-small Cell Lung Cancer	I
NCT03754244	A Phase I Study of TQ-B3456 on Tolerance and Pharmacokinetics	I
No NCT ID	Evaluation Of The Phase I Clinical Trial Of Safety, Tolerability, Pharmacokinetic Characteristics, And Preliminary Efficacy Of YK-029A Tablets In Patients With Advanced Non-Small Cell Lung Cancer (NSCLC)	I
No NCT ID	A Multicenter, Open and Dose Escalation Phase I Study of YZJ-0318 Maleate Tablets in Patients with Advanced Non-Small Cell Lung Cancer with Positive Epidermal Growth Factor Receptor T790M Mutation.	I
NCT03802240	A Randomized, Double-blind, Multi-center, Phase III Clinical Study Assessing the Efficacy and Safety of Sintilimab ± IBI305 Combined With Pemetrexed and Cisplatin in Patients With EGFR-mutant Locally Advanced or Metastatic Non-squamous Non-small Cell Lung Cancer Who Have Failed Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor (EGFR-TKI) Treatment (ORIENT-31)	III
NCT03924050	A Randomized, Double-Blind, Placebo-Controlled, Multicenter, Phase III Study of Pemetrexed + Platinum Chemotherapy With or Without Toripalimab (JS001) in Advanced Non-small Cell Lung Cancer (NSCLC) Participants With TKI-resistant EGFR-mutated Tumors	III
NCT03376737	A Single-Arm Phase II Clinical Trial of Apatinib as the Maintenance Therapy in Advanced Lung Adenocarcinoma	II
NCT03838848	A Phase II Study to Evaluate the Efficacy, Safety, and Tolerability of KN046 in Patients With Advanced Non-small Cell Lung Cancer	II
NCT03804580	First-Line Treatment With Osimertinib In EGFR-Mutated Non-Small Cell Lung Cancer, Coupled To Extensive Translational Studies	II

## Clinical Trials Summary (continued)

### EGFR p.(T790M) c.2369C>T (continued)

NCT ID	Title	Phase
NCT02609776	A Phase I, First-in-Human, Open-Label, Dose Escalation Study of JNJ-61186372, a Human Bispecific EGFR and cMet Antibody, in Subjects With Advanced Non-Small Cell Lung Cancer	I
NCT04001777	A Phase Ib Study of Safety and Efficacy of APG-1252 in Combination With Osimertinib (AZD9291) in EGFR TKI Resistant NSCLC Patients	I
No NCT ID	Phase I Clinical Trial of SPH3348 Single-agent Combined with Oxitinib in Patients with Advanced Solid Tumors with Abnormal c-Met in China.	I
No NCT ID	Apatinib Combined With EGFR-TKI For Patients With EGFR Mutation Who Failed EGFR-TKI: A Prospective Study	IV
No NCT ID	Clinical Study Of Combined Action Of Icotinib And Brain Radiotherapy On EGFR-Mutated Non-Small-Cell Lung Cancer Patients With Brain Metastasis	IV
NCT03800134	A Phase III, Double-blind, Placebo-controlled, Multi-center International Study of Neoadjuvant/Adjuvant Durvalumab for the Treatment of Patients With Resectable Stages II and III Non-small Cell Lung Cancer (AEGEAN)	III
NCT03992885	Combination Therapy With Icotinib, Pemetrexed and Platinum in Patients With Metastatic Non-squamous Non-small Cell Lung Cancer With EGFR Mutations Who Did Not Progress After Pemetrexed in Combination With Platinum-based Chemotherapy:a Single-arm, Open, Multicenter Clinical Study.	III
NCT04619563	A Single-arm Exploratory Clinical Study of Anlotinib Hydrochloride Combined With Docetaxel in EGFR Mutations Advanced Non Small Cell Lung Cancer Patients Who Have Progressed After Targeted Therapy and Chemotherapy	II
No NCT ID	Osimertinib Combined Bevacizumab in Untreated Epidermal Growth Factor Receptor Mutated Non-small-cell Lung Cancer Patients with Malignant Pleural And/Or Pericardial Effusion -phase II Trial	II
No NCT ID	Randomized Controlled Trial for EGFR-TKIs Plus S-1 or EGFR-TKIs as the First-Line Therapy for Patients with Advanced Non-small Cell Lung Cancer Harboring EGFR Mutations	II
No NCT ID	Single arm, Exploratory Study for Apatinib mesylate Combined with EGFR-TKI in Patients with EGFR Mutation-positive Advanced Non-squamous Non-small-cell Lung Cancer	II
No NCT ID	EGFR-TKI Combined With Stereotactic Body Radiation Therapy Versus TKI alone for Stage IV Oncogene-Driven Non-Small Cell Lung Cancer.	II
NCT03904823	An Open, Single-arm, Multi-center, Phase II Clinical Trial of Famitinib Combined With Epidermal Growth Factor Receptor (EGFR) Inhibitor HS-10296 in Patients With Advanced EGFR-mutant Non-Small Cell Lung Cancer (NSCLC)	II
NCT02960607	A Phase II Study of High-dose Icotinib in Previously Treated Non-small Cell Lung Cancer Patients With Epidermal Growth Factor Receptor Mutation	II
NCT03091491	Randomised Phase II Study of Nivolumab Versus Nivolumab and Ipilimumab Combination in EGFR Mutant Non-small Cell Lung Cancer	II
NCT04538378	Phase II Trial of Olaparib (LYNPARZA) Plus Durvalumab (IMFINZI) in EGFR-Mutated Adenocarcinomas That Transform to Small Cell Lung Cancer (SCLC) and Other Neuroendocrine Tumors.	II
NCT03460275	Osimertinib as First-line Therapy for Patients With EGFR Mutation-positive Locally Advanced or Metastatic Non-squamous Non-Small Cell Lung Cancer(NSCLC), a Single-Arm, Open-Label, Prospective, Multicenter, Phase II Clinical Trial	II
NCT04425681	Phase II Study of Osimertinib With Bevacizumab for Leptomeningeal Metastasis From EGFR-mutation Non-Small Cell Lung Cancer	II




## Clinical Trials Summary (continued)

### EGFR p.(T790M) c.2369C>T (continued)

NCT ID	Title	Phase
No NCT ID	Phase II Trial Of Docetaxel Plus Ramucirumab Combination Therapy In Patients With Advanced EGFR Gene Mutation Positive Advanced Stage Non-Squamous Cell Non small Cell Lung Cancer	II
No NCT ID	Clinical Study of Combined Action of the First Generation of TKIs and Brain Radiotherapy on EGFR-Mutated Non-Small-Cell Lung Cancer Patients with Brain Metastasis	II
NCT03974022	A Phase I/II, Open-Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumor Efficacy of DZD9008 in Patients With Advanced Non-Small Cell Lung Cancer (NSCLC) With EGFR or HER2 Mutation	I/II
NCT03797391	First-in-human, Phase I/II, Multicenter, Open-Label Study of EMB-01 in Patients With Advanced/ Metastatic Solid Tumors	I/II
NCT04528836	A Phase I/IB First-in-Human Study of the SHP2 Inhibitor BBP-398 (Formerly Known as IACS-15509) in Patients With Advanced Solid Tumors	I
No NCT ID	Phase I Study of DZD9008 in EGFR or HER2 Mutant NSCLC Chinese Patients	I
NCT04077463	An Open-label Phase 1/1b Study to Evaluate the Safety and Pharmacokinetics of JNJ-73841937 (Lazertinib), a Third Generation EGFR-TKI, as Monotherapy or in Combinations With JNJ-61186372, a Human Bispecific EGFR and cMet Antibody in Participants With Advanced Non-Small Cell Lung Cancer	I
No NCT ID	Study Of Immunologic Factor In Re-Biopsy Specimen, Peritumoral BALF, And The Peripheral Blood For Predicting Response To Osimertinib In NSCLC Patients	I
No NCT ID	Pharmacokinetic and dose finding study of osimertinib in patients with impaired renal function and low body weight	I
NCT03535363	Phase I Trial of Osimertinib With Stereotactic Radiosurgery (SRS) in Patients With Brain Metastases From EGFR Positive Non-Small-Cell Lung Cancer (NSCLC)	I
No NCT ID	Phase I Clinical Study With Advanced Solid Tumors KBP-5209 Treatment	I
NCT03065387	Phase I Study of the Pan-ERBB Inhibitor Neratinib Given in Combination With Everolimus, Palbociclib, or Trametinib in Advanced Cancer Subjects With EGFR Mutation/Amplification, HER2 Mutation/ Amplification, or HER3/4 Mutation or KRAS Mutation	I
NCT04128085	A Phase I, Open-label, Multicenter, Dose Escalation and Expansion Study to Evaluate the Tolerance and Pharmacokinetics of TQB3804 in Subjects With Advanced Malignant Tumors	I
NCT04429542	First-in-Human, Phase I/Ib, Open-label, Multicenter Study of Bifunctional EGFR/TGFβ Fusion Protein BCA101 Alone and in Combination With Pembrolizumab in Patients With EGFR-Driven Advanced Solid Tumors	I

## Alerts Informed By Public Data Sources

### Current NCCN Information

 Contraindicated
  Not recommended
  Resistance

NCCN information is current as of 2020-12-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### EGFR p.(L858R) c.2573T>G

##### alectinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

##### brigatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

##### ceritinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

##### crizotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

## EGFR p.(L858R) c.2573T>G (continued)

### – lorlatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFRi sensitizing mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Crizotinib, ceritinib, alectinib, brigatinib, or lorlatinib are not recommended as subsequent therapy for patients with sensitizing EGFR mutations who relapse on EGFR TKI therapy."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### – atezolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### – nivolumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### – pembrolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

## EGFR p.(T790M) c.2369C>T

### atezolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### nivolumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### pembrolizumab

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "subsequent therapy with pembrolizumab, nivolumab, or atezolizumab is not recommended in patients with EGFR mutations or ALK fusions."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### afatinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib, or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### dacomitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib, or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

## EGFR p.(T790M) c.2369C>T (continued)

### erlotinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib, or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

### gefitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR T790M mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "The most common known mechanism is the acquisition of T790M (which is a secondary mutation in EGFR), which renders the kinase resistant to erlotinib, gefitinib, dacomitinib, or afatinib."

**Reference:** NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]

## Current EMA Information

 Contraindicated     Not recommended     Resistance

EMA information is current as of 2020-12-16. For the most up-to-date information, search [www.ema.europa.eu/ema](http://www.ema.europa.eu/ema).

## EGFR p.(T790M) c.2369C>T

### gefitinib

**Cancer type:** Non-Small Cell Lung Cancer

**Label as of:** 2019-05-28

**Variant class:** EGFR T790M mutation

**Reference:**

[https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/iressa-epar-product-information_en.pdf)



## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985 Sep 6;229(4717):974-6. PMID: 2992089
2. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
3. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
4. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
5. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
9. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
10. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
11. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
12. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
13. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
14. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
15. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
16. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469
17. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene*. 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
18. Gan et al. The EGFRvIII variant in glioblastoma multiforme. *J Clin Neurosci.* 2009 Jun;16(6):748-54. PMID: 19324552
19. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)
20. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/206995s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/206995s003lbl.pdf)
21. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2019/201292s015lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/201292s015lbl.pdf)
22. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/211288s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/211288s000lbl.pdf)
23. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2021]
24. Naidoo et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer*. 2015 Sep 15;121(18):3212-3220. PMID: 26096453
25. Vyse et al. *Signal Transduct Target Ther.* 2019;4:5. PMID: 30854234
26. Yi et al. A comparison of epidermal growth factor receptor mutation testing methods in different tissue types in non-small cell lung cancer. *Int J Mol Med.* 2014 Aug;34(2):464-74. PMID: 24891042
27. Madic et al. EGFR C797S, EGFR T790M and EGFR sensitizing mutations in non-small cell lung cancer revealed by six-color crystal digital PCR. *Oncotarget*. 2018 Dec 21;9(100):37393-37406. PMID: 30647840
28. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/208065s021lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/208065s021lbl.pdf)
29. Niederst et al. The Allelic Context of the C797S Mutation Acquired upon Treatment with Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin. Cancer Res.* 2015 Sep 1;21(17):3924-33. PMID: 25964297
30. Wang et al. Lung Adenocarcinoma Harboring EGFR T790M and In Trans C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol.* 2017 Nov;12(11):1723-1727. PMID: 28662863

## References (continued)

31. <https://www.jnj.com/janssen-announces-u-s-fda-breakthrough-therapy-designation-granted-for-jnj-6372-for-the-treatment-of-non-small-cell-lung-cancer>
32. <https://www.takeda.com/newsroom/newsreleases/2020/takeda-announces-u.s.-fda-breakthrough-therapy-designation-for-mobocertinib-tak-788-for-the-treatment-of-nsclc-patients-with-egfr-exon-20-insertion-mutations/>
33. <https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>
34. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>