

REPORT SUMMARY

PATIENT AND SAMPLE INFORMATION	2
VARIANT(S) WITH CLINICAL RELEVANCE	2
THERAPEUTIC IMPLICATIONS.....	3

REPORT DETAILS

VARIANT INTERPRETATION	4
US FDA-APPROVED DRUG(S)	6
ONGOING CLINICAL TRIALS	7
DETAILED TEST RESULTS	8
HOTSPOT GENOTYPES.....	10
TEST DETAILS	12
ACTOnco®+ GENE LIST	16
DISCLAIMER	17

APPENDIX

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS.....	19
REFERENCES.....	20

PATIENT AND SAMPLE INFORMATION

PATIENT

Name: 莊智惠
Gender: Female
Date of Birth: Oct 05, 1943
Patient ID: 32558966
Diagnosis: Metastatic adenoid cystic carcinoma

SPECIMEN

Type: FFPE tissue
Date received: Dec 30, 2021
Collection site: Lung
Specimen ID: S11039576
Lab ID: AA-21-06573
D/ID: NA

ORDERING PHYSICIAN

Name: 周德盈醫師
Facility: 臺北榮總
Tel: 886-228712121
Address: 臺北市北投區石牌路二段 201 號

VARIANT(S) WITH CLINICAL RELEVANCE

Only variant(s) with clinical significance are listed. See the "DETAILED TEST RESULTS" section for full details.

SINGLE NUCLEOTIDE AND SMALL INDEL VARIANTS

Gene	Amino Acid Change	Coverage	Allele Frequency	COSMIC ID
DICER1	G1708*	325	50.2%	-
FUBP1	Y541*	1435	45.0%	COSM1716368
KDM6A	L617fs	609	45.8%	-
KDM6A	R612*	1851	27.8%	COSM28784

COPY NUMBER VARIANTS (CNVS)

Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on **30%** tumor purity.

Amplification (Copy number ≥ 8)

Chr	Gene	Copy Number
ND	ND	ND

Homozygous deletion (Copy number=0)

Chr	Gene
ND	ND

Heterozygous deletion (Copy number=1)

Chr	Gene
ND	ND

ND, Not Detected

TUMOR MUTATIONAL BURDEN (TMB)

< 1 muts/Mb

Muts/Mb, mutations per megabase

MICROSATELLITE INSTABILITY (MSI)

Microsatellite stable (MSS)

Note:

TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is < 30%.

Variant Analysis:

醫檢師黃靖婷 博士
Ching-Ting Huang Ph.D.
檢字第 016511 號

CT Huang

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

No genomic alterations predicted to confer sensitivity or lack of benefit to targeted therapy approved in this tumor type.

IMMUNE CHECKPOINT INHIBITORS (ICI) THERAPIES

Genomic markers and alterations that are associated with response to ICI therapies

Positive Biomarker	Negative Biomarker
TMB-H: ND	EGFR aberration: ND
MSI-H: ND	MDM2/MDM4 amplification: ND
MMR biallelic inactivation: ND	STK11 biallelic inactivation: ND
PBRM1 biallelic inactivation: ND	PTEN biallelic inactivation: ND
SERPINB3/SERPINB4 mutation: ND	B2M biallelic inactivation: ND
	JAK1/2 biallelic inactivation: ND

MMR, mismatch repair; ND, not detected

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

VARIANT INTERPRETATION

DICER1 G1708*

Biological Impact

DICER1 gene encodes an endoribonuclease that functions in the RNA interference (RNAi) pathway to cleave long double-stranded RNA (dsRNA) molecules into short dsRNA molecules including microRNA (miRNA) and small interfering RNA (siRNA)^{[1][2]}. Germline loss-of-function mutations of DICER1 has been linked with familial pleuropulmonary blastoma (PPB), cystic nephroma (CN), and ovarian Sertoli-Leydig cell tumor^{[3][4][5]}. DICER1 has been implicated as a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[6][7]}.

G1708* mutation results in a premature truncation of the DICER1 protein at amino acid 1708 (UniProtKB). This mutation is predicted to lead to a loss of DICER1 function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

Relative low expression of DICER1 has been reported to associate with poor survival in hepatocellular carcinoma^[8], lung cancer^[9], breast cancer^{[10][11]}, and endometrial adenocarcinoma^[12].

FUBP1 Y541*

Biological Impact

The far upstream element-binding protein 1 (FUBP1) is a DNA- and RNA-binding protein which regulates c-Myc expression and RNA stabilization^[13]. Although overexpression of FUBP1 can stimulate MYC expression^[14], it has also been shown that FUBP1 protein participates in a complex with PUF60 that negatively regulates MYC expression^[15]. The missense, nonsense, and silent mutations, whole gene deletions, and frameshift deletions and insertions are frequently observed in cancers of the central nervous system such as oligodendroglioma^{[16][17]}, and low-grade glioma^[18].

In particular, inactivating mutations in CIC and FUBP1 genes, which mapped in chromosome 19q and 1p, respectively, have been frequently observed in 1p/19q-co-deleted oligodendroglioma^[17] as well as 1p/19q-co-deleted oligoastrocytoma^[19].

Y541* mutation results in a premature truncation of the FUBP1 protein at amino acid 541 (UniProtKB). This mutation is predicted to lead to a loss of FUBP1 function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

Loss of FUBP1 expression has been associated with unfavorable progression-free survival and overall survival in oligodendroglioma^[20].

KDM6A L617fs, R612*

Biological Impact

KDM6A (lysine-specific demethylase 6A) gene encodes a histone demethylase for histone 3 lysine 27 (H3K27) which mediates tissue-specific expression of various genes and is mostly involved in embryogenesis, developmental processes and the cell cycle^[21]. Germline deletions and point mutations of KDM6A are associated with Kabuki syndrome, which is characterized by unique facial appearance, skeletal anomalies, and growth retardation^{[22][23]}. Somatic inactivating mutations of KDM6A have been reported in bladder cancer^{[24][25]}, multiple myeloma^[26], T-cell acute lymphoblastic leukemia^{[27][28]}, renal cell carcinoma^[29], pancreatic cancer^[30], and prostate cancer^[31].

L617fs mutation results in a change in the amino acid sequence beginning at 617, likely to cause premature truncation of the functional KDM6A protein (UniProtKB). This mutation is predicted to lead to a loss of KDM6A protein function, despite not being characterized in the literature. R612* mutation results in a premature truncation of the KDM6A protein at amino acid 612 (UniProtKB). This mutation is predicted to lead to a loss of KDM6A function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

Low expression of KDM6A has been reported to associate with poorer survival in esophageal squamous cell carcinoma and B cell lymphoma^{[32][33]}. Defective KDM6A (mutations or deletions) was associated with shorter survival in squamous-like pancreatic cancer^[34].

US FDA-APPROVED DRUG(S)

No genomic alterations predicted to confer sensitivity or lack of benefit to targeted therapy approved.

ONGOING CLINICAL TRIALS

Clinical trials shown below were selected by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

DETAILED TEST RESULTS


SINGLE NUCLEOTIDE AND SMALL INDEL VARIANTS

Gene	Chr	Exon	Accession Number	cDNA Change	Amino Acid Change	Coverage	Allele Frequency	COSMIC ID
ADAMTS16	5	13	NM_139056	c.1964A>G	H655R	712	47.3%	-
ARID2	12	15	NM_152641	c.4390C>T	R1464C	1740	50.6%	COSM2067866
BIRC3	11	6	NM_001165	c.1156G>A	V386M	512	53.1%	-
CARD11	7	7	NM_032415	c.941A>C	E314A	993	51.8%	-
CCNB3	X	10	NM_033031	c.3839C>T	T1280I	762	50.4%	-
DICER1	14	24	NM_177438	c.5122G>T	G1708*	325	50.2%	-
FAT1	4	22	NM_005245	c.11998G>A	E4000K	1789	46.4%	-
FUBP1	1	17	NM_003902	c.1623C>G	Y541*	1435	45.0%	COSM1716368
GSTP1	11	6	NM_000852	c.370C>G	P124A	345	48.7%	-
KDM6A	X	16	NM_021140	c.1849_1850del	L617fs	609	45.8%	-
KDM6A	X	16	NM_021140	c.1834C>T	R612*	1851	27.8%	COSM28784
MUC16	19	12	NM_024690	c.36585C>A	S12195R	674	44.5%	-
MUC6	11	24	NM_005961	c.3176G>A	R1059H	176	50.6%	-
PDGFRB	5	7	NM_002609	c.1108C>T	R370C	1158	50.7%	COSM6743163
RAD54L	1	16	NM_003579	c.1748T>C	I583T	1105	49.6%	-

Mutations with clinical relevance are highlighted in red.

COPY NUMBER VARIANTS (CNVS)

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



HOTSPOT GENOTYPES

Listed variants are biomarkers or hotspots that are recommended as standard care by the NCCN or other expert panels and not necessarily FDA-recognized for a particular indication. The genotypes have been manually checked to ensure sufficient coverage for each hotspot of the target gene.

Gene	Variant	Genotype Detected
<i>BRAF</i>	V600X	Not detected
<i>EGFR</i>	A763_Y764insFQEA, E709K, E709_T710delinsD, Exon 19 deletion, Exon 19 insertion, Exon 20 insertion, G719A/C/D/S, L747P, L833V, L858R, L861Q/R, S768I, T790M	Not detected
<i>IDH2</i>	R140Q, R172G/K/M/S	Not detected
<i>KIT</i>	A502_Y503dup, D419del, D579del, D816F/V/Y, D820A/E/G/Y, E554_I571del, E554_K558del, E554_V559del, Exon 11 mutation, F522C, H697Y, I563_L576del, I653T, K550_W557del, K558N, K558_E562del, K558_V559del, K558delinsNP, K642E, M552_W557del, N505I, N564_Y578del, N822H/I/K/Y, P551_M552del, P573_D579del, P577_D579del, P577_W582delinsPYD, P838L, Q556_K558del, T417_D419delinsI, T417_D419delinsRG, T574_Q575insTQLPYD, V530I, V555_L576del, V555_V559del, V559A/C/D/G, V559_V560del, V559del, V560D/G, V560del, V569_L576del, V654A, W557G/R, W557_K558del, Y553N, Y553_K558del, Y570H, Y578C	Not detected
<i>KRAS</i>	A146T/V/P, G12X, G13X, Q61X	Not detected
<i>MET</i>	D1028H/N/Y	Not detected
<i>NRAS</i>	G12X, G13X, Q61X	Not detected
<i>PDGFRA</i>	A633T, C450_K451insMIEWMI, C456_N468del, C456_R481del, D568N, D842I/V, D842_H845del, D842_M844del, D846Y, E311_K312del, G853D, H650Q, H845Y, H845_N848delinsP, I843del, N659K/R/S, N848K, P577S, Q579R, R560_V561insER, R748G, R841K, S566_E571delinsR, S584L, V469A, V536E, V544_L545insAVLVLLVIVISLI, V561A/D, V561_I562insER, V658A, W559_R560del, Y375_K455del, Y555C, Y849C/S	Not detected
<i>PIK3CA</i>	C420R, E542K/V, E545A/D/G/K, H1047X, Q546E/R	Not detected

V600X= any mutation in the valine (V) at amino acid 600 being replaced by a different amino acid.

G12X= any mutation in the glycine (G) at amino acid 12 being replaced by a different amino acid.

G13X= any mutation in the glycine (G) at amino acid 13 being replaced by a different amino acid.

Q61X= any mutation in the glutamine (Q) at amino acid 61 being replaced by a different amino acid.

H1047X= any mutation in the histidine (H) at amino acid 1047 being replaced by a different amino acid.

Gene	Copy Number Detected
<i>CDK4</i>	2
<i>EGFR</i>	2
<i>ERBB2</i>	2
<i>MET</i>	2

Copy number ≥ 8 is considered amplification

Other known alterations that are associated with sensitivity, resistance, and toxicity to therapies.

Gene	Variant	Genotype Detected
<i>AKT1</i>	E17K	Not detected
<i>ALK</i>	C1156Y, D1203N, G1202R, L1152R, S1206Y, T1151_L1152insT	Not detected
<i>BRAF</i>	K601E, L597V/Q/R/S	Not detected
<i>DPYD</i>	D949V, I560S, splice-site mutation	Not detected
<i>EGFR</i>	A750P, C797S/Y, S492R	Not detected
<i>ERBB2</i>	V659E	Not detected
<i>ESR1</i>	D538G, E380Q, L469V, L536H/P/Q/R, S432L, S463P, V422del, V534E, Y537C/N/S	Not detected
<i>FGFR3</i>	G370C, G380R, K650E/N/R/M/T/Q, R248C, S249C, S371C, Y373C	Not detected
<i>IDH1</i>	R132C/G/H/L/Q/S	Not detected
<i>MAP2K1</i>	D67N, E203K, F53L, K57E/N, P124S, Q56P, Q56_V60del, R47Q, R49L, S222D	Not detected
<i>PTEN</i>	R130*/fs/G/L/P/Q	Not detected
<i>TPMT</i>	A154T, Y240C	Not detected

Gene	Copy Number Detected
<i>FGFR1</i>	5
<i>MDM2</i>	2
<i>MDM4</i>	2

Copy number ≥ 8 is considered amplification

TEST DETAILS

ABOUT ACTOnco®+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (INDELs) (≤ 15 nucleotides) and large-scale genomic alterations like copy number variations (CNVs).

See ACTOnco®+ Gene List' Section for details of gene sequenced.

DATABASE USED

- Reference genome: human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210208)
- ACT Genomics in-house database

NEXT-GENERATION SEQUENCING (NGS) METHODS

Extracted genomic DNA was amplified using four pools of primer pairs targeting coding exons of analyzed genes. Amplicons were ligated with barcoded adaptors. Quality and quantity of amplified library were determined using the fragment analyzer (AATI) and Qubit (Invitrogen). Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system (Thermo Fisher Scientific) according to the Ion PI Hi-Q Chef Kit protocol (Thermo Fisher Scientific) or Ion 540 Kit-Chef protocol (Thermo Fisher Scientific). Sequencing was performed on the Ion Proton or Ion S5 sequencer (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite (version 5.10). Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (INDELs) were identified using the Torrent Variant Caller plug-in (version 5.10). The coverage was down-sampled to 4000. VEP (Variant Effect Predictor) (version 100) was used to annotate every variant using databases from Clinvar (version 20210208), COSMIC v.92 and Genome Aggregation database r2.1.1. Variants with coverage ≥ 25 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained.

This test provides uniform coverage of the targeted regions, enabling target base coverage at $100x \geq 85\%$ with a mean coverage $\geq 500x$.

Variants reported in Genome Aggregation database r2.1.1 with $> 1\%$ minor allele frequency (MAF) were

considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number variations (CNVs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations from samples in ACT Genomics in-house database.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as “TMB-High”, “TMB-Low” or “Cannot Be Determined”. TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as “Cannot Be Determined” if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

STANDARD OPERATING PROCEDURES (SOPS)

Standard operating procedures (SOPs) are shown below:

- AG2-QP-15 Specimen Management Procedure
- AG3-QP16-03 SOP of Cancer Cell DNA and RNA Extraction
- AG3-QP16-07 SOP of Nucleic Acid Extraction with QIAasympyphony SP
- AG3-QP16-08 SOP of FFPE Nucleic Acid Extraction
- AG3-QP16-10 SOP of HE Staining
- AG3-QP16-13 SOP of Library Construction and Preparation
- AG3-QP16-17 SOP of DNA Quantification with Qubit Fluorometer
- AG3-QP16-20 SOP of CE-Fragment Analysis
- AG3-QP16-22 SOP of Variant Calling
- AG3-QP16-24 SOP of Ion Torrent System Sequencing Reaction
- AG3-QP16-26 SOP of Ion Chef Preparation

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

- AG3-QP16-35 SOP of Variant Annotation
- AG3-QP16-96 SOP of Manual Inspection for SNVIndel Variant
- AG3-QP16-95 SOP of Manual Inspection for Copy Number Variant
- AG3-QP40-08 (02) Standard protocol for variant interpretation, curation and classification
- AG3-QP16-41 SOP of The user manual for clinical report system (CRS)

LIMITATIONS

This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.

NOTES

We do not exclude the possibility that pathogenic variants may not be reported by one or more of the tools and the parameters used.

PATHOLOGY EVALUATION

- H&E-stained section No.: S11039576
- Collection site: Lung
- Examined by: Dr. Yeh-Han Wang
- Estimated neoplastic nuclei (whole sample): The percentage of viable tumor cells in total cells in the whole slide (%): 70%
The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 70%
The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
Additional comment: NA
- Manual macrodissection: Not performed



The outline highlights the area of malignant neoplasm annotated by a pathologist.

SPECIMEN PHOTO(S)



- Collection date: Dec 2021
- Facility retrieved: 臺北榮總

RUN QC

- Panel: ACTOnco[®]+
- Mean Depth: 808x
- Target Base Coverage at 100x: 94%

ACTOnco® + GENE LIST

ABCB1*	AURKB	CBL	CDKN2B	E2F3	FAT1	GRIN2A	JAK2	MED12	NOTCH4	PMS1	RAD51D	SLCO1B3*	TNFRSF14
ABCC2*	AXIN1	CCNA1	CDKN2C	EGFR	FBXW7	GSK3B	JAK3	MEF2B	NPM1	PMS2	RAD52	SMAD2	TNFSF11
ABCG2*	AXIN2	CCNA2	CEBPA*	EP300	FCGR2B	GSTP1*	JUN*	MEN1	NQO1*	POLB	RAD54L	SMAD3	TOP1
ABL1	AXL	CCNB1	CHEK1	EPCAM	FGF1*	GSTT1*	KAT6A	MET	NRAS	POLD1	RAF1	SMAD4	TP53
ABL2	B2M	CCNB2	CHEK2	EPHA2	FGF10	HGF	KDM5A	MITF	NSD1	POLE	RARA	SMARCA4	TPMT*
ADAMTS1	BAP1	CCNB3	CIC	EPHA3	FGF14	HIF1A	KDM5C	MLH1	NTRK1	PPARG	RB1	SMARCB1	TSC1
ADAMTS13	BARD1	CCND1	CREBBP	EPHA5	FGF19*	HIST1H1C*	KDM6A	MPL	NTRK2	PPP2R1A	RBM10	SMO	TSC2
ADAMTS15	BCL10	CCND2	CRKL	EPHA7	FGF23	HIST1H1E*	KDR	MRE11	NTRK3	PRDM1	RECQL4	SOC1*	TSHR
ADAMTS16	BCL2*	CCND3	CRLF2	EPHB1	FGF3	HNF1A	KEAP1	MSH2	PAK3	PRKAR1A	REL	SOX2*	TYMS
ADAMTS18	BCL2L1	CCNE1	CSF1R	ERBB2	FGF4*	HR	KIT	MSH6	PALB2	PRKCA	RET	SOX9	U2AF1
ADAMTS6	BCL2L2*	CCNE2	CTCF	ERBB3	FGF6	HRAS*	KMT2A	MTHFR*	PARP1	PRKCB	RHOA	SPEN	UBE2A*
ADAMTS9	BCL6	CCNH	CTLA4	ERBB4	FGFR1	HSP90AA1	KMT2C	MTOR	PAX5	PRKCG	RICTOR	SPOP	UBE2K
ADAMTSL1	BCL9	CD19	CTNNA1	ERCC1	FGFR2	HSP90AB1	KMT2D	MUC16	PAX8	PRKCI	RNF43	SRC	UBR5
ADGRA2	BCOR	CD274	CTNNB1	ERCC2	FGFR3	HSPA4	KRAS	MUC4	PBRM1	PRKCQ	ROS1	STAG2	UGT1A1*
ADH1C*	BIRC2	CD58	CUL3	ERCC3	FGFR4	HSPA5	LCK	MUC6	PDCD1	PRKDC	RPPH1	STAT3	USH2A
AKT1	BIRC3	CD70*	CYLD	ERCC4	FH	IDH1	LIG1	MUTYH	PDCD1LG2	PRKN	RPTOR	STK11	VDR*
AKT2	BLM	CD79A	CYP1A1*	ERCC5	FLCN	IDH2	LIG3	MYC	PDGFRA	PSMB8	RUNX1	SUFU	VEGFA
AKT3	BMPR1A	CD79B	CYP2B6*	ERG	FLT1	IFNL3*	LMO1	MYCL	PDGFRB	PSMB9	RUNX1T1	SYK	VEGFB
ALDH1A1*	BRAF	CDC73	CYP2C19*	ESR1	FLT3	IGF1	LRP1B	MYCN	PDIA3	PSME1	RXRA	SYNE1	VHL
ALK	BRCA1	CDH1	CYP2C8*	ESR2	FLT4	IGF1R	LYN	MYD88	PGF	PSME2	SDHA	TAF1	WT1
AMER1	BRCA2	CDK1	CYP2D6	ETV1	FOXL2*	IGF2	MALT1	NAT2*	PHOX2B*	PSME3	SDHB	TAP1	XIAP
APC	BRD4	CDK12	CYP2E1*	ETV4	FOXP1	IKBKB	MAP2K1	NBN	PIK3C2B	PTCH1	SDHC	TAP2	XPO1
AR	BRIP1	CDK2	CYP3A4*	EZH2	FRG1	IKBKE	MAP2K2	NEFH	PIK3C2G	PTEN	SDHD	TAPBP	XRCC2
ARAF	BTG1*	CDK4	CYP3A5*	FAM46C	FUBP1	IKZF1	MAP2K4	NF1	PIK3C3	PTGS2	SERPINB3	TBX3	ZNF217
ARID1A	BTG2*	CDK5	DAXX	FANCA	GATA1	IL6	MAP3K1	NF2	PIK3CA	PTPN11	SERPINB4	TEK	
ARID1B	BTK	CDK6	DCUN1D1	FANCC	GATA2	IL7R	MAP3K7	NFE2L2	PIK3CB	PTPRD	SETD2	TERT	
ARID2	BUB1B	CDK7	DDR2	FANCD2	GATA3	INPP4B	MAPK1	NFKB1	PIK3CD	PTPRT	SF3B1	TET1	
ASXL1	CALR	CDK8	DICER1	FANCE	GNA11	INSR	MAPK3	NFKBIA	PIK3CG	RAC1	SGK1	TET2	
ATM	CANX	CDK9	DNMT3A	FANCF	GNA13	IRF4	MAX	NKX2-1*	PIK3R1	RAD50	SH2D1A*	TGFBR2	
ATR	CARD11	CDKN1A	DOT1L	FANCG	GNAQ	IRS1	MCL1	NOTCH1	PIK3R2	RAD51	SLC19A1*	TMSB4X*	
ATRX	CASP8	CDKN1B	DPYD	FANCL	GNAS	IRS2*	MDM2	NOTCH2	PIK3R3	RAD51B	SLC22A2*	TNF	
AURKA	CBFB	CDKN2A	DTX1	FAS	GREM1	JAK1	MDM4	NOTCH3	PIM1	RAD51C	SLCO1B1*	TNFAIP3	

*Analysis of copy number alteration not available.

行動基因僅提供技術檢測服務及檢測報告，檢測結果之臨床解釋及相關醫療處置，請諮詢專業醫師。報告結果僅對此試驗件有效。

行動基因臨床分子醫學實驗室 台北市內湖區新湖二路 345 號 3F

Email: service@actgenomics.com T: +886-2-2795-3660 | F: +886-2-2795-5016

AG4-QP4001-02(05)

DISCLAIMER

Legal Statement

This test was developed by ACT Genomics and its performing characteristics were determined by ACT Genomics. This test result is to be used for clinical consultative purposes only and is not intended as a substitute for a clinical guidance of your doctor or another qualified medical practitioner. It should not be regarded as investigational or used for research.

The detection of genomic alterations does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; the detection of no genomic alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Treatment Decisions are the Responsibility of the Physician

Decisions on clinical care and treatment should be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, including physical examinations, information from other diagnostics tests and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test, or the information contained in this report.

In terms of consulting a different treating physician, the patient must file an application and fulfill the listed criteria for ACT Genomics to provide the patient's report to the assigned physician. The report may not be copied or reproduced except in its totality.

Genetic Alterations and Drugs Not Presented in Ranked Order

In this report, neither any biomarker alteration nor any drug associated with a potential clinical benefit (or potential lack of clinical benefit), are ranked in order of potential or predicted efficacy.

Level of Evidence Provided

Drugs with a potential clinical benefit (or potential lack of clinical benefit) are evaluated for level of published evidence with at least one clinical efficacy case report or preclinical study. We endeavor to keep the information in the report up to date. However, customers must be aware that scientific understanding and technologies change over time, and we make no warranty as to the accuracy, suitability or currency of information provided in this report at any time.

No Guarantee of Clinical Benefit

This report makes no promises or guarantees about the effectiveness of a particular drug or any treatment procedure in any disease or in any patient. This report also makes no promises or guarantees that a drug without an association of reportable genomic alteration will, in fact, provide no clinical benefit.

Liability

ACT Genomics is not affiliated with any medical facility or medical practitioner. We provide information for informational purposes only, therefore, ACT Genomics and their employees cannot be held responsible for any direct, indirect, special, incidental or consequential damages that may arise from the use of information provided in the report.

免責聲明

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS

Not Applicable.

REFERENCES

1. PMID: 25176334; 2014, Nat Rev Cancer;14(10):662-72
DICER1: mutations, microRNAs and mechanisms.
2. PMID: 28473628; 2017, Biochem J;474(10):1603-1618
Molecular mechanisms of Dicer: endonuclease and enzymatic activity.
3. PMID: 21266384; 2011, J Med Genet;48(4):273-8
DICER1 syndrome: clarifying the diagnosis, clinical features and management implications of a pleiotropic tumour predisposition syndrome.
4. PMID: 19556464; 2009, Science;325(5943):965
DICER1 mutations in familial pleuropulmonary blastoma.
5. PMID: 24761742; 1993, ;():
DICER1-Related Disorders
6. PMID: 19903759; 2009, Genes Dev;23(23):2700-4
Dicer1 functions as a haploinsufficient tumor suppressor.
7. PMID: 20019750; 2010, Cell Death Differ;17(4):633-41
Monoallelic but not biallelic loss of Dicer1 promotes tumorigenesis in vivo.
8. PMID: 27313724; 2016, Oncol Lett;11(6):3961-3966
Prognostic significance of Dicer expression in hepatocellular carcinoma.
9. PMID: 15723655; 2005, Cancer Sci;96(2):111-5
Reduced expression of Dicer associated with poor prognosis in lung cancer patients.
10. PMID: 19672267; 2009, Br J Cancer;101(4):673-83
Prognostic value of Dicer expression in human breast cancers and association with the mesenchymal phenotype.
11. PMID: 24386264; 2013, PLoS One;8(12):e83724
Prognostic significance of deregulated dicer expression in breast cancer.
12. PMID: 21425145; 2011, Cancer;117(7):1446-53
DICER1 expression and outcomes in endometrioid endometrial adenocarcinoma.
13. PMID: 22926519; 2013, Oncogene;32(24):2907-16
Far upstream element binding protein 1: a commander of transcription, translation and beyond.
14. PMID: 8125259; 1994, Genes Dev;8(4):465-80
A sequence-specific, single-strand binding protein activates the far upstream element of c-myc and defines a new DNA-binding motif.
15. PMID: 20422426; 2011, J Cancer Res Clin Oncol;137(2):321-30
Potential role of cyclooxygenase-2 on the regulation of the drug efflux transporter ABCG2 in breast cancer cell lines.
16. PMID: 24117486; 2014, Neuropathol Appl Neurobiol;40(2):205-16
Loss of FUBP1 expression in gliomas predicts FUBP1 mutation and is associated with oligodendroglial differentiation, IDH1 mutation and 1p/19q loss of heterozygosity.
17. PMID: 21817013; 2011, Science;333(6048):1453-5
Mutations in CIC and FUBP1 contribute to human oligodendroglioma.

18. PMID: 29617667; 2018, Cell Rep;23(1):282-296.e4
Somatic Mutational Landscape of Splicing Factor Genes and Their Functional Consequences across 33 Cancer Types.
19. PMID: 22588899; 2012, Acta Neuropathol;123(6):853-60
CIC and FUBP1 mutations in oligodendrogliomas, oligoastrocytomas and astrocytomas.
20. PMID: 24030748; 2014, Mod Pathol;27(3):332-42
Loss of CIC and FUBP1 expressions are potential markers of shorter time to recurrence in oligodendroglial tumors.
21. PMID: 24561908; 2014, Epigenetics;9(5):658-68
The H3K27me3 demethylase UTX in normal development and disease.
22. PMID: 22197486; 2012, Am J Hum Genet;90(1):119-24
Deletion of KDM6A, a histone demethylase interacting with MLL2, in three patients with Kabuki syndrome.
23. PMID: 23076834; 2013, Hum Mutat;34(1):108-10
KDM6A point mutations cause Kabuki syndrome.
24. PMID: 25225064; 2014, Clin Cancer Res;20(18):4935-48
Concurrent alterations in TERT, KDM6A, and the BRCA pathway in bladder cancer.
25. PMID: 25092538; 2015, Eur Urol;67(2):198-201
Genomic predictors of survival in patients with high-grade urothelial carcinoma of the bladder.
26. PMID: 27235425; 2016, Clin Cancer Res;22(23):5783-5794
The Spectrum and Clinical Impact of Epigenetic Modifier Mutations in Myeloma.
27. PMID: 22377896; 2012, Leukemia;26(8):1881-3
Sequencing histone-modifying enzymes identifies UTX mutations in acute lymphoblastic leukemia.
28. PMID: 25320243; 2015, Blood;125(1):13-21
The H3K27me3 demethylase UTX is a gender-specific tumor suppressor in T-cell acute lymphoblastic leukemia.
29. PMID: 21248752; 2011, Nature;469(7331):539-42
Exome sequencing identifies frequent mutation of the SWI/SNF complex gene PBRM1 in renal carcinoma.
30. PMID: 25719666; 2015, Nature;518(7540):495-501
Whole genomes redefine the mutational landscape of pancreatic cancer.
31. PMID: 22722839; 2012, Nature;487(7406):239-43
The mutational landscape of lethal castration-resistant prostate cancer.
32. PMID: 29351209; 2018, Int J Mol Sci;19(1):
The Prognostic Significance of Histone Demethylase UTX in Esophageal Squamous Cell Carcinoma.
33. PMID: 30006524; 2018, Nat Commun;9(1):2720
UTX is an escape from X-inactivation tumor-suppressor in B cell lymphoma.
34. PMID: 29533787; 2018, Cancer Cell;33(3):512-526.e8
Loss of KDM6A Activates Super-Enhancers to Induce Gender-Specific Squamous-like Pancreatic Cancer and Confers Sensitivity to BET Inhibitors.

215

ACTFusion™ Report

PATIENT		
Name: 莊智惠		Patient ID: 32558966
Date of Birth: Oct 05, 1943		Gender: Female
Diagnosis: Metastatic adenoid cystic carcinoma		
ORDERING PHYSICIAN		
Name: 周德盈醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11039576	Collection site: Lung	Date received: Dec 30, 2021
Lab ID: AA-21-06573	Type: FFPE tissue	D/ID: NA

ABOUT ACTFusion™

The test is a next-generation sequencing (NGS) based in vitro diagnostic assay to detect fusion transcripts of 13 genes, including *ALK*, *BRAF*, *EGFR*, *FGFR1*, *FGFR2*, *FGFR3*, *MET*, *NRG1*, *NTRK1*, *NTRK2*, *NTRK3*, *RET*, and *ROS1*.

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample.	

ACTFusion™ Report

THERAPEUTIC IMPLICATION

Not Applicable.

VARIANT INTERPRETATION

Not Applicable.

US FDA-APPROVED DRUG(S)

Not Applicable.

ACTFusion™ Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACTFusion™ Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Dec 2021
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11039576
- Collection site: Lung
- Examined by: Dr. Yeh-Han Wang
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 70%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 70%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
 5. Additional comment: NA
- Manual macrodissection: Not performed
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTFusion™
- Total reads: 230169
- Average unique RNA Start Sites per control GSP2: 107

LIMITATIONS

This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

ACTFusion™ Report

NEXT-GENERATION SEQUENCING (NGS) METHODS

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX.

STANDARD OPERATING PROCEDURES (SOPs)

- AG2-QP-15 Specimen Management Procedure
- AG3-QP16-08 SOP of FFPE Nucleic Acid Extraction
- AG3-QP16-10 SOP of HE Staining
- AG3-QP16-17 SOP of DNA Quantification with Qubit Fluorometer
- AG3-QP16-20 SOP of CE-Fragment Analysis
- AG3-QP16-24 SOP of Ion Torrent System Sequencing Reaction
- AG3-QP16-26 SOP of Ion Chef Preparation
- AG3-QP40-08 (02) Standard protocol for variant interpretation, curation and classification
- AG3-QP16-94 (01) SOP of ACTFusion v3 Library Construction and Preparation
- AG3-QP16-36(02) SOP of Fusion Gene Detection
- AG3-QP16-41 SOP of The user manual for clinical report system (CRS)

DATABASES USED

- Quiver Gene Fusion Database version 5.1.18

GENE LIST

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1
NTRK1	NTRK2	NTRK3	RET	ROS1			

Variant Analysis:

醫檢師黃靖婷 博士
Ching-Ting Huang Ph.D.
檢字第 016511 號

CT Huang

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

ACTFusion™ Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACTFusion™ Report

REFERENCE

Not Applicable.