

ACT Onco[®] + Report

PATIENT		
Name: 黃淳涵		Patient ID: 46791633
Date of Birth: Apr 05, 2000		Gender: Female
Diagnosis: Hepatocellular carcinoma		
ORDERING PHYSICIAN		
Name: 陳三奇醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S0908269D		Type: FFPE tissue
Collection site: Liver		
Date received: Sep 06, 2022	Lab ID: AA-22-05182	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
Not detected			

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
ERBB3 R1127H	Lapatinib	-
MCL1 Amplification	Regorafenib, Sorafenib	-

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACTOnco[®] + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
AXIN1	V683fs	50.8%
ERBB3	R1127H	63.8%

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr16	AXIN1, PALB2, TSC2	Heterozygous deletion	1
Chr4	FBXW7	Heterozygous deletion	1
Chr1	MCL1, NTRK1	Amplification	6

- Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	< 1 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 57% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is < 30%.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 4		
ERBB3 R1127H	Lapatinib	sensitive
MCL1 Amplification	Regorafenib, Sorafenib	sensitive

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
Not detected	

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

AXIN1 V683fs, Heterozygous deletion

Biological Impact

AXIN1 gene encodes a scaffold protein that is a component of the beta-catenin destruction complex. In the complex, AXIN1 provide scaffolding for the APC, GSK-3 beta and beta-catenin, which enables the degradation of beta-catenin in the absence of WNT ligand binding and negatively regulates WNT signaling pathway^{[1][2]}. Somatic mutations and deletions of AXIN1 have been reported in hepatocellular carcinoma^[3], medulloblastoma^{[4][5]}, gastrointestinal cancer^[6], and colorectal cancer^[7].

V683fs mutation results in a change in the amino acid sequence beginning at 683, likely to cause premature truncation of the functional AXIN1 protein (UniProtKB). This mutation is predicted to lead to a loss of AXIN1 protein function, despite not being characterized in the literature. Loss of the second wild-type allele resulted in the biallelic inactivation of the gene.

Therapeutic and prognostic relevance

A retrospective study (n = 124) has demonstrated that reduced AXIN expression in tumor specimen was associated with poor survival in patients with esophageal squamous cell carcinoma (ESCC)^[8].

ERBB3 R1127H

Biological Impact

The ERBB3 (also known as HER3) gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases^[9]. HER3 lacks or has little intrinsic tyrosine kinase activity. Upon binding of its ligand, neu differentiation factor (NDF), HER3 forms a heterodimer with ErbB2^[10] and subsequently activates various mitogenic signaling cascades, including the PI3K/AKT/mTOR, STAT and RAS/RAF/MAPK^{[11][12][13]}. Aberrant expression or alterations of the ErbB family play crucial roles in the development and progression of cancer^[14]. Enhanced expression of HER2 has been observed in a broad spectrum of human cancers, including gastric, bladder, uterine, colorectal, and breast cancers^[15]. HER3 is the preferred heterodimeric partner for EGFR in melanoma and pancreatic carcinoma^{[16][17]}, while in breast cancer, HER3 preferably heterodimerizes with HER2 and plays a critical role in HER2-mediated tumorigenesis.

ERBB3 R1127H locates within the cytoplasmic domain of the HER3 protein (UniProtKB). However, this variant has not been characterized; therefore, its biological impact on the ERBB3 protein remains unknown.

Therapeutic and prognostic relevance

Currently, there are no FDA-approved anti-HER3 therapies for patients with solid tumors. A variety of strategies targeting HER3 including pan HER approach, abrogating its dimerization partners' kinase activity using small molecule inhibitors (e.g. lapatinib, erlotinib, gefitinib, afatinib, and neratinib) or direct targeting of its extracellular domain (e.g. including AV-203 (Abstract nr 2509, AACR 2012)) are under investigation^{[18][19]}.

Preclinical data indicated that HER3 expression level is a predictive biomarker of pertuzumab (an anti-HER2 antibody) efficacy in HER2 low-expressing pancreatic cancer^[20].

ERBB3 mutation has been selected as an inclusion criteria for the trial examining afatinib in urothelial tract carcinoma, non-small cell lung carcinoma (NSCLC) and malignant solid tumor (NCT02780687, NCT01523587, NCT03810872)^{[21][22]}.

One patient with intrahepatic cholangiocarcinoma carrying an ERBB3 R1127H mutation responded to lapatinib, an ERBB inhibitor treatment, had 20.5 and 24.8 months of PFS, and OS, respectively^[23].

ACT Onco[®] + Report

FBXW7 Heterozygous deletion

Biological Impact

The F-box/WD repeat-containing protein 7 (FBXW7) gene encodes a protein that belongs to the SCF (SKP1-CUL1-F-box protein) E3 ligase complex. FBXW7 is recognized as a tumor suppressor which is involved in the negative regulation of oncogenes such as c-Myc^{[24][25]}, c-Jun^[26], cyclin E^[27], Notch family members^{[28][29]}, Aurora-A^[30], mTOR^[31], KLF5^[32], and MCL-1^[33]. Inactivating FBXW7 mutation or copy number loss may result in the accumulation of oncoproteins and therefore lead to malignant transformation^[34]. FBXW7 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[32][33][35]}.

Therapeutic and prognostic relevance

Clinical efficacy of mTOR inhibitors was seen in patients harboring aberrations in the FBXW7 gene (one patient with refractory fibrolamellar hepatocellular carcinoma, and one patient with lung adenocarcinoma)^{[36][37]}. Moreover, in vitro assay also suggested that loss or inactivation of FBXW7 may confer sensitivity to mTOR inhibitor^[31].

Preclinical studies suggested that mutations or loss of FBXW7 were associated with regorafenib and oxaliplatin resistance in CRC cell lines and gefitinib resistance in lung cancer cells^{[38][39][40][41]}.

Retrospective studies have indicated that a relatively low expression level of FBXW7 is an independent prognostic marker of poor survival for patients with hepatocellular carcinoma, lung adenocarcinoma and squamous cell carcinoma^{[42][40]}.

MCL1 Amplification

Biological Impact

The myeloid cell leukemia 1 (MCL1) gene encodes a member of the BCL2 pro-survival family^[43]. MCL1 is highly regulated by various oncogenic signaling pathways, including the mitogen-activated protein kinase (MAPK) pathway^[44], the mTOR pathway^[45], and the phosphatidylinositol-3 kinase (PI3K) pathway^[46]. Oncogenic roles for MCL1 have been previously suggested by the report of increased rates of lymphoma in transgenic mice^[47]. Somatic amplification of MCL1 may be a common mechanism in cancer cells to increase cell survival^[48]. MCL1 overexpression was observed from a retrospective analysis of parotid gland carcinomas, including adenoid cystic carcinoma^[49].

Therapeutic and prognostic relevance

Therapies targeting MCL1 and other BCL2 family members with the pan-BCL2 family inhibitors are currently under investigation^[50]. A case report has demonstrated clinical efficacy of sorafenib, when combined with vorinostat, in a metastatic triple-negative breast cancer (TNBC) patient with MCL1-amplified tumor^[51]. Several in vitro studies also showed that sorafenib induces cell death via inhibition of MCL1 expression in multiple cancer types including, hepatocellular carcinoma (HCC), lung cancer, breast cancer, cholangiocarcinoma, endometrial cancer and chronic lymphocytic leukemia^{[52][53][54][55][56][57]}. Preclinical studies have also demonstrated the efficacy of regorafenib in reducing MCL1 expression in human colorectal cancer (CRC) cell lines^{[58][59]}. In vivo models of colon cancer showed that MCL1 expression is inhibited by targeting of the mTOR pathway using everolimus, promoting increased tumor cell killing of cancers with KRAS or BRAF mutations^[60].

ACT Onco[®] + Report

NTRK1 Amplification

Biological Impact

The NTRK1 gene encodes the TRKA (tropomyosin receptor kinase) receptor which plays an important role in the development and function of the nervous system. Gene fusions of NTRK1 lead to constitutive activation of MAP-kinase, PI3-kinase, and PLC-γ pathways, and represent the main molecular alterations with known oncogenic and transforming potential in various malignancies, including soft tissue sarcoma, non-small cell lung cancer (NSCLC), glioblastoma multiforme (GBM), thyroid carcinoma, and pilocytic astrocytomas^{[61][62]}. A pan-cancer study (n=1250) demonstrated that 2.2% of the metastatic cancer patients harbored NTRK amplification and NTRK protein overexpression was observed in 14.8% of NTRK-amplified tumors (doi.org/10.23838/pfm.2017.00142).

Therapeutic and prognostic relevance

Patients with NTRK1 amplification had only limited benefit from larotrectinib treatment according to the few clinical studies. One of them had a partial response with larotrectinib of short duration (3.7 months)^[63], and the other one with metastatic NTRK1-amplified (copy number=8) esophageal carcinoma showed clinical efficacy for six weeks, and then a progressive disease of new lesions were observed^[64].

PALB2 Heterozygous deletion

Biological Impact

The partner and localizer of BRCA2 (PALB2) gene encodes a protein that plays a critical role in homologous recombination repair (HRR) through its ability to interact with BRCA2 in nuclear foci, promoting its localization and stability in key nuclear structures^[65]. The Fanconi anemia complementation group (FANC) which includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM and FANCN (also called PALB2) are involved in the repair of DNA double-strand breaks (DSBs) by homologous recombination (HR)^{[66][67][68]}. PALB2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological function^[69]. Biallelic germline loss-of-function mutations in PALB2 cause Fanconi anemia, whereas monoallelic loss-of-function mutations are associated with an increased risk of breast cancer and pancreatic cancer^[70]. Fanconi Anemia is an autosomal recessive disease characterized by hematological abnormalities, bone marrow failure, limb deformities, skin hyperpigmentation, and susceptibility to hematologic and solid malignancies, such as acute myeloid leukemia and head and neck carcinoma^{[71][72]}.

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[73].

NCCN guideline for pancreatic cancer recommended that rucaparib as maintenance therapy following prior platinum-based therapy for patients with pancreatic adenocarcinoma harboring PALB2 mutations^[74].

In a Phase I trial, a pancreatic cancer patient harboring a PALB2 mutation had a partial response to talazoparib treatment^[75]. In a preclinical study, talazoparib demonstrated anti-tumor activity on a Wilms tumor cell line with a PALB2 truncating mutation in mice xenograft model^[76]. PALB2 loss of function mutation has been determined as an inclusion criterion for the trial evaluating rucaparib efficacy in ovarian cancer^[77] or prostate cancer^[78]; talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795) or lung cancer (NCT03377556); niraparib efficacy in pancreatic cancer (NCT03553004), prostate cancer (NCT02854436), or any malignancy (except prostate) cancer (NCT03207347).

ACT Onco[®] + Report

A case report demonstrated an exceptional response to mitomycin C and cisplatin treatment in a gemcitabine-resistant pancreatic cancer patient with biallelic inactivation of PALB2^[79].

TSC2 Heterozygous deletion

Biological Impact

The tuberous sclerosis complex 2 (TSC2) gene encodes a protein called tuberin, which interact with a protein called hamartin (encoded by the TSC1 gene). This hamartin-tuberin tumor suppressor complex plays a critical role in growth control as a negative regulator of the mammalian target of rapamycin (mTOR) pathway^{[80][81]}. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis complex^{[82][83][84]}, while the loss of heterozygosity (LOH) in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)^[85] and endometrial cancer^[86]. TSC2 deletion, splicing-mutant, and inactivating mutations such as A1141T, G305V, S1514X, and R1032X, has been identified in TSC2-null hepatocellular carcinoma (HCC) cell lines, patient-derived xenograft, and primary tumors. Mutations in the TSC1 and TSC2 genes cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC)^[87].

Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple cancer types, such as bladder cancer, gastric cancer, sarcoma, thyroid cancer, hepatocellular carcinoma (HCC) as well as head and neck squamous cell carcinoma (HNSCC)^{[88][89][90]}. Results from one Phase II study of advanced endometrial cancer showed that mutations in AKT1, TSC1, and TSC2 might predict sensitivity to temsirolimus^[91]. Recent studies indicated that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets^[92].

Everolimus has been approved by the U.S. FDA for Tuberous Sclerosis Complex (TSC)-associated renal angiomyolipoma and Tuberous Sclerosis Complex (TSC)-associated subependymal giant cell astrocytoma (SEGA). This approval is based on the results from EXIST-1, EXIST-2, and Study 2485 trials (NCT00789828, NCT00790400, and NCT00411619).

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4 ^[93] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
BOLERO-2 ^[94] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
EXIST-2 NCT00790400	Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
RADIANT-3 ^[95] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1 ^[96] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
RECORD-1 ^[97] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

Lapatinib (TYKERB)

Lapatinib is a dual tyrosine kinase inhibitor that interrupts the HER2/neu and epidermal growth factor receptor (EGFR) pathways. Lapatinib is developed and marketed by GlaxoSmithKline under the trade name TYKERB.

- FDA Approval Summary of Lapatinib (TYKERB)

^[98] NCT00073528	Her2-receptor positive breast cancer (Approved on 2010/01/29)
	ER+/HER2+ Lapatinib + letrozole vs. Letrozole [PFS(W): 35.4 vs. 13]
^[99] NCT00078572	Her2-receptor positive breast cancer (Approved on 2007/03/13)
	HER2+ Lapatinib + capecitabine vs. Capecitabine [TTP(W): 27.1 vs. 18.6]

ACT Onco[®] + Report

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (ZEJULA)

PRIMA NCT02655016	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29)
	- Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
QUADRA ^[100] NCT02354586	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2019/10/23)
	HRD+ Niraparib [ORR(%): 24.0, DOR(M): 8.3]
NOVA ^[101] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	- Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (LYNPARZA)

OlympiA NCT02032823	Her2-negative high-risk early breast cancer (Approved on 2022/03/11)
	HER2-/gBRCA mutation Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):]
PROfound ^[73] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	HRR genes mutation Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1 ^[102] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD+ Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
POLO ^[103] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	gBRCA mutation Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1 ^[104] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	gBRCA mutation or sBRCA mutation Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
OlympiAD ^[105] NCT02000622	Breast cancer (Approved on 2018/02/06)
	HER2-/gBRCA mutation Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21 ^[106] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA mutation Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]

ACT Onco[®] + Report

Study19 ^[107] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	-
	Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]
Study 42 ^[108] NCT01078662	Ovarian cancer (Approved on 2014/12/19)
	gBRCA mutation
	Olaparib [ORR(%): 34.0, DOR(M): 7.9]

Regorafenib (STIVARGA)

Regorafenib is a multi-kinase inhibitor which targets angiogenic, stromal and oncogenic receptor tyrosine kinases (RTKs). Regorafenib is developed and marketed by Bayer HealthCare Pharmaceuticals under the trade name STIVARGA.

- FDA Approval Summary of Regorafenib (STIVARGA)

RESORCE ^[109] NCT01774344	Hepatocellular carcinoma, Hepatocellular carcinoma (Approved on 2017/04/27)
	-
	Bsc vs. Placebo [OS(M): 10.6 vs. 7.8]
GRID ^[110] NCT01271712	Gastrointestinal stromal tumor (Approved on 2013/02/25)
	-
	Regorafenib vs. Placebo [PFS(M): 4.8 vs. 0.9]
CORRECT ^[111] NCT01103323	Colorectal cancer (Approved on 2012/09/27)
	-
	Regorafenib vs. Placebo [OS(M): 6.4 vs. 5]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA mutation or sBRCA mutation
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
ARIEL3 ^[77] NCT01968213	Ovarian cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	-
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]

ACT Onco[®] + Report

Sorafenib (NEXAVAR)

Sorafenib is a small molecule multi-kinase inhibitor that targets multiple kinase families including VEGFR, PDGFRB, and the RAF family kinases. Sorafenib is co-developed and co-marketed by Bayer HealthCare Pharmaceuticals and Onyx Pharmaceuticals under the trade name NEXAVAR.

- FDA Approval Summary of Sorafenib (NEXAVAR)

DECISION ^[112] NCT00984282	Differentiated thyroid carcinoma (Approved on 2013/11/22)
	-
	Sorafenib vs. Placebo [PFS(M): 10.8 vs. 5.8]
SHARP ^[113] NCT00105443	Hepatocellular carcinoma (Approved on 2007/11/16)
	-
	Sorafenib vs. Placebo [OS(M): 10.7 vs. 7.9]
TARGET ^[114] NCT00073307	Renal cell carcinoma (Approved on 2005/12/20)
	-
	Sorafenib vs. Placebo [PFS(D): 167 vs. 84]

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

EMBRACA ^[115] NCT01945775	Breast cancer (Approved on 2018/10/16)
	HER2-/gBRCA mutation
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

[116] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	-
	Temsirolimus vs. Ifn-α [OS(M): 10.9 vs. 7.3]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
AXIN1	V683fs	8	c.2048del	NM_003502	-	50.8%	593
ERBB3	R1127H	27	c.3380G>A	NM_001982	-	63.8%	1190

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
DPYD	I843T	20	c.2528T>C	NM_000110	-	22.4%	997
FAT1	E1141D	3	c.3423G>C	NM_005245	COSM6476272	74.8%	1033
KEAP1	G9R	2	c.25G>A	NM_203500	COSM2812662	48.4%	1001
KMT2C	R380L	8	c.1139G>T	NM_170606	COSM225885	5.3%	3991
MUC16	D12307N	16	c.36919G>A	NM_024690	-	50.0%	596
MUC16	S9527G	3	c.28579A>G	NM_024690	-	48.7%	1303
NKX2-1	G322S	2	c.964G>A	NM_003317	-	29.6%	81
RUNX1	Q308H	6	c.924G>T	NM_001001890	-	51.1%	824
SMARCB1	Splice region	8	c.1116G>A	NM_003073	COSM1003	52.3%	974
TSC2	A1141T	30	c.3421G>A	NM_000548	-	71.5%	200
TSC2	I485F	15	c.1453A>T	NM_000548	-	25.2%	630


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco® + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Mar 20, 2020
- Facility retrieved: 臺中榮總
- H&E-stained section No.: S0908269D
- Collection site: Liver
- Examined by: Dr. Chien-Ta Chiang
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 70%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 70%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
 5. Additional comment: NA
- Manual macrodissection: Not performed
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco®+

DNA test

- Mean Depth: 877x
- Target Base Coverage at 100x: 94%

RNA test

- Average unique RNA Start Sites per control GSP2: 35

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

ACT Onco[®] + Report

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師張筑芃 博士
Chu-Yuan Chang Ph.D.
檢字第 020115 號



Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號



ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTS11	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC01B1*
SLC01B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION


Gene	Therapies	Possible effect
<i>FBXW7</i>	Everolimus, Temsirolimus	sensitive
<i>TSC2</i>	Everolimus, Temsirolimus	sensitive
<i>PALB2</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>FBXW7</i>	Gefitinib, Regorafenib	resistant

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Everolimus, Temsirolimus; 2: Sorafenib

ACT Onco[®] + Report



1: Olaparib, Niraparib, Rucaparib, Talazoparib

Receptor Tyrosine Kinase/Growth Factor Signalling



1: Lapatinib; 2: Regorafenib; 3: Sorafenib, Regorafenib; 4: Lapatinib

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 9734785; 1998, Genes Cells;3(6):395-403
Axin, an inhibitor of the Wnt signalling pathway, interacts with beta-catenin, GSK-3beta and APC and reduces the beta-catenin level.
2. PMID: 23169527; 2013, Cold Spring Harb Perspect Biol;5(1):a007898
The β -catenin destruction complex.
3. PMID: 12101426; 2002, Oncogene;21(31):4863-71
Mutational spectrum of beta-catenin, AXIN1, and AXIN2 in hepatocellular carcinomas and hepatoblastomas.
4. PMID: 12555076; 2003, Oncogene;22(4):632-6
AXIN1 mutations but not deletions in cerebellar medulloblastomas.
5. PMID: 11585731; 2001, Cancer Res;61(19):7039-43
Deletions of AXIN1, a component of the WNT/wingless pathway, in sporadic medulloblastomas.
6. PMID: 25236910; 2014, Cancer Lett;355(1):1-8
AXIN1 and AXIN2 variants in gastrointestinal cancers.
7. PMID: 14566817; 2003, Int J Cancer;107(5):696-9
Detection of point mutations of the Axin1 gene in colorectal cancers.
8. PMID: 19582507; 2009, Ann Surg Oncol;16(9):2486-93
Reduced axin protein expression is associated with a poor prognosis in patients with squamous cell carcinoma of esophagus.
9. PMID: 19536107; 2009, Nat Rev Cancer;9(7):463-75
Novel anticancer targets: revisiting ERBB2 and discovering ERBB3.
10. PMID: 9130710; 1997, EMBO J;16(7):1647-55
ErbB-2, the preferred heterodimerization partner of all ErbB receptors, is a mediator of lateral signaling.
11. PMID: 22785351; 2012, Nat Rev Cancer;12(8):553-63
The ERBB network: at last, cancer therapy meets systems biology.
12. PMID: 7515147; 1994, Mol Cell Biol;14(6):3550-8
ErbB3 is involved in activation of phosphatidylinositol 3-kinase by epidermal growth factor.
13. PMID: 12853564; 2003, Proc Natl Acad Sci U S A;100(15):8933-8
The ErbB2/ErbB3 heterodimer functions as an oncogenic unit: ErbB2 requires ErbB3 to drive breast tumor cell proliferation.
14. PMID: 19208461; 2009, Curr Opin Cell Biol;21(2):177-84
ErbB receptors and signaling pathways in cancer.
15. PMID: 23680147; 2013, Cancer Cell;23(5):603-17
Oncogenic ERBB3 mutations in human cancers.
16. PMID: 18698037; 2008, Clin Cancer Res;14(16):5188-97
HER3 is a determinant for poor prognosis in melanoma.
17. PMID: 20647770; 2010, Cancer Biol Ther;10(6):555-63
ErbB3 expression promotes tumorigenesis in pancreatic adenocarcinoma.
18. PMID: 30057690; 2018, Oncol Rev;12(1):355
HER3 signaling and targeted therapy in cancer.
19. PMID: 30109175; 2018, Acta Pharm Sin B;8(4):503-510
Understanding the biology of HER3 receptor as a therapeutic target in human cancer.

ACT Onco[®] + Report

20. PMID: 25216528; 2014, Oncotarget;5(16):7138-48
HER3 as biomarker and therapeutic target in pancreatic cancer: new insights in pertuzumab therapy in preclinical models.
21. PMID: 27044931; 2016, J Clin Oncol;34(18):2165-71
Afatinib Activity in Platinum-Refractory Metastatic Urothelial Carcinoma in Patients With ERBB Alterations.
22. PMID: 29902295; 2018, JAMA Oncol;4(9):1189-1197
Association of ERBB Mutations With Clinical Outcomes of Afatinib- or Erlotinib-Treated Patients With Lung Squamous Cell Carcinoma: Secondary Analysis of the LUX-Lung 8 Randomized Clinical Trial.
23. PMID: 32323774; 2020, Oncol Rep;43(4):1089-1102
Targeted deep sequencing contributes to guiding personalized targeted therapy for advanced biliary tract cancer patients with non-radical resection: A real-world study.
24. PMID: 15498494; 2004, Curr Biol;14(20):1852-7
A nucleolar isoform of the Fbw7 ubiquitin ligase regulates c-Myc and cell size.
25. PMID: 15103331; 2004, EMBO J;23(10):2116-25
Phosphorylation-dependent degradation of c-Myc is mediated by the F-box protein Fbw7.
26. PMID: 16023596; 2005, Cancer Cell;8(1):25-33
The v-Jun point mutation allows c-Jun to escape GSK3-dependent recognition and destruction by the Fbw7 ubiquitin ligase.
27. PMID: 11533444; 2001, Science;294(5540):173-7
Phosphorylation-dependent ubiquitination of cyclin E by the SCFFbw7 ubiquitin ligase.
28. PMID: 11461910; 2001, J Biol Chem;276(38):35847-53
The Notch intracellular domain is ubiquitinated and negatively regulated by the mammalian Sel-10 homolog.
29. PMID: 11425854; 2001, J Biol Chem;276(37):34371-8
Functional interaction between SEL-10, an F-box protein, and the nuclear form of activated Notch1 receptor.
30. PMID: 16863506; 2006, Cancer Sci;97(8):729-36
Fbxw7 contributes to tumor suppression by targeting multiple proteins for ubiquitin-dependent degradation.
31. PMID: 18787170; 2008, Science;321(5895):1499-502
FBXW7 targets mTOR for degradation and cooperates with PTEN in tumor suppression.
32. PMID: 20484041; 2010, Cancer Res;70(11):4728-38
The Fbw7 tumor suppressor targets KLF5 for ubiquitin-mediated degradation and suppresses breast cell proliferation.
33. PMID: 21368833; 2011, Nature;471(7336):104-9
SCF(FBW7) regulates cellular apoptosis by targeting MCL1 for ubiquitylation and destruction.
34. PMID: 18094723; 2008, Nat Rev Cancer;8(2):83-93
FBW7 ubiquitin ligase: a tumour suppressor at the crossroads of cell division, growth and differentiation.
35. PMID: 23032637; 2012, Cancer Inform;11():157-71
Haploinsufficiency of Tumor Suppressor Genes is Driven by the Cumulative Effect of microRNAs, microRNA Binding Site Polymorphisms and microRNA Polymorphisms: An In silico Approach.
36. PMID: 24586741; 2014, PLoS One;9(2):e89388
FBXW7 mutations in patients with advanced cancers: clinical and molecular characteristics and outcomes with mTOR inhibitors.
37. PMID: 24360397; 2014, Lung Cancer;83(2):300-1
Temozolimide therapy in a patient with lung adenocarcinoma harboring an FBXW7 mutation.
38. PMID: 27399335; 2017, Oncogene;36(6):787-796
FBW7 mutations mediate resistance of colorectal cancer to targeted therapies by blocking Mcl-1 degradation.

ACT Onco[®] + Report

39. PMID: 25860929; 2015, Oncotarget;6(11):9240-56
FBXW7-mutated colorectal cancer cells exhibit aberrant expression of phosphorylated-p53 at Serine-15.
40. PMID: 29633504; 2018, Mol Oncol;12(6):883-895
FBXW7 deletion contributes to lung tumor development and confers resistance to gefitinib therapy.
41. PMID: 28522751; 2017, Cancer Res;77(13):3527-3539
Targeting FBW7 as a Strategy to Overcome Resistance to Targeted Therapy in Non-Small Cell Lung Cancer.
42. PMID: 24884509; 2014, Mol Cancer;13():110
Fbxw7 is an independent prognostic marker and induces apoptosis and growth arrest by regulating YAP abundance in hepatocellular carcinoma.
43. PMID: 18955968; 2008, Oncogene;27(50):6398-406
Bcl-2 family proteins and cancer.
44. PMID: 18676833; 2008, Cancer Res;68(15):6109-17
Down-regulation of myeloid cell leukemia-1 through inhibiting Erk/Pin 1 pathway by sorafenib facilitates chemosensitization in breast cancer.
45. PMID: 18664580; 2008, Proc Natl Acad Sci U S A;105(31):10853-8
mTORC1 promotes survival through translational control of Mcl-1.
46. PMID: 11314001; 2001, Oncogene;20(6):677-85
The involvement of PI 3-K/Akt-dependent up-regulation of Mcl-1 in the prevention of apoptosis of Hep3B cells by interleukin-6.
47. PMID: 11389033; 2001, Blood;97(12):3902-9
MCL1 transgenic mice exhibit a high incidence of B-cell lymphoma manifested as a spectrum of histologic subtypes.
48. PMID: 20164920; 2010, Nature;463(7283):899-905
The landscape of somatic copy-number alteration across human cancers.
49. PMID: 21734342; 2011, Dis Markers;30(5):229-33
Mcl-1 expression is up-regulated in malignancies of the parotid gland.
50. PMID: 26045609; 2015, Blood;126(3):363-72
Inhibition of Mcl-1 with the pan-Bcl-2 family inhibitor (-)BI97D6 overcomes ABT-737 resistance in acute myeloid leukemia.
51. PMID: 27293397; 2016, Case Rep Oncol;9(1):112-8
A Combination of Targeted Therapy with Chemotherapy Backbone Induces Response in a Treatment-Resistant Triple-Negative MCL1-Amplified Metastatic Breast Cancer Patient.
52. PMID: 17178882; 2006, Cancer Res;66(24):11851-8
Sorafenib blocks the RAF/MEK/ERK pathway, inhibits tumor angiogenesis, and induces tumor cell apoptosis in hepatocellular carcinoma model PLC/PRF/5.
53. PMID: 23392173; 2013, Cell Death Dis;4():e485
Mcl-1-dependent activation of Beclin 1 mediates autophagic cell death induced by sorafenib and SC-59 in hepatocellular carcinoma cells.
54. PMID: 16007148; 2005, Oncogene;24(46):6861-9
The role of Mcl-1 downregulation in the proapoptotic activity of the multikinase inhibitor BAY 43-9006.
55. PMID: 19821497; 2009, Hepatology;50(6):1861-70
Sorafenib inhibits signal transducer and activator of transcription-3 signaling in cholangiocarcinoma cells by activating the phosphatase shatterproof 2.
56. PMID: 20071162; 2010, Eur J Cancer;46(4):836-50
The multikinase inhibitor Sorafenib induces apoptosis and sensitises endometrial cancer cells to TRAIL by different mechanisms.
57. PMID: 21979753; 2012, Mol Med;18():19-28
Sorafenib-induced apoptosis of chronic lymphocytic leukemia cells is associated with downregulation of RAF and myeloid cell leukemia

ACT Onco[®] + Report

sequence 1 (Mcl-1).

58. PMID: 24763611; 2014, Clin Cancer Res;20(13):3472-84
Regorafenib inhibits colorectal tumor growth through PUMA-mediated apoptosis.
59. PMID: 26561209; 2015, Cancer Biol Ther;16(12):1710-9
Regorafenib with a fluoropyrimidine for metastatic colorectal cancer after progression on multiple 5-FU-containing combination therapies and regorafenib monotherapy.
60. PMID: 24163374; 2014, Cancer Discov;4(1):42-52
mTOR inhibition specifically sensitizes colorectal cancers with KRAS or BRAF mutations to BCL-2/BCL-XL inhibition by suppressing MCL-1.
61. PMID: 24162815; 2013, Nat Med;19(11):1469-1472
Oncogenic and drug-sensitive NTRK1 rearrangements in lung cancer.
62. PMID: 27843590; 2016, ESMO Open;1(2):e000023
NTRK gene fusions as novel targets of cancer therapy across multiple tumour types.
63. PMID: 30624546; 2019, Ann Oncol;30(2):325-331
Larotrectinib in adult patients with solid tumours: a multi-centre, open-label, phase I dose-escalation study.
64. PMID: 32323889; 2020, Oncologist;25(6):e881-e886
Antitumor Activity of Larotrectinib in Esophageal Carcinoma with NTRK Gene Amplification.
65. PMID: 16793542; 2006, Mol Cell;22(6):719-29
Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2.
66. PMID: 23325218; 2013, Nature;493(7432):356-63
Fanconi anaemia and the repair of Watson and Crick DNA crosslinks.
67. PMID: 15905196; 2005, Carcinogenesis;26(10):1731-40
The Fanconi anemia group A protein modulates homologous repair of DNA double-strand breaks in mammalian cells.
68. PMID: 27037238; 2016, EMBO J;35(9):909-23
Interplay between Fanconi anemia and homologous recombination pathways in genome integrity.
69. PMID: 24153426; 2013, Nat Commun;4():2578
Heterozygous mutations in PALB2 cause DNA replication and damage response defects.
70. PMID: 20858716; 2010, Cancer Res;70(19):7353-9
PALB2/FANCN: recombining cancer and Fanconi anemia.
71. PMID: 25754594; 2015, Hum Mutat;36(5):562-8
Loss-of-Function FANCL Mutations Associate with Severe Fanconi Anemia Overlapping the VACTERL Association.
72. PMID: 28678401; 2017, Cancer;123(20):3943-3954
Assessing the spectrum of germline variation in Fanconi anemia genes among patients with head and neck carcinoma before age 50.
73. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
74. PMID: 33970687; 2021, J Clin Oncol;39(22):2497-2505
Phase II Study of Maintenance Rucaparib in Patients With Platinum-Sensitive Advanced Pancreatic Cancer and a Pathogenic Germline or Somatic Variant in BRCA1, BRCA2, or PALB2.
75. PMID: 28242752; 2017, Cancer Discov;7(6):620-629
Phase I, Dose-Escalation, Two-Part Trial of the PARP Inhibitor Talazoparib in Patients with Advanced Germline BRCA1/2 Mutations and Selected Sporadic Cancers.
76. PMID: 25263539; 2015, Pediatr Blood Cancer;62(1):91-8

ACT Onco[®] + Report

Initial testing (stage 1) of the PARP inhibitor BMN 673 by the pediatric preclinical testing program: PALB2 mutation predicts exceptional in vivo response to BMN 673.

77. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
78. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
79. PMID: 21135251; 2011, Mol Cancer Ther;10(1):3-8
Personalizing cancer treatment in the age of global genomic analyses: PALB2 gene mutations and the response to DNA damaging agents in pancreatic cancer.
80. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35
mTOR: from growth signal integration to cancer, diabetes and ageing.
81. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
82. PMID: 9242607; 1997, Science;277(5327):805-8
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
83. PMID: 8269512; 1993, Cell;75(7):1305-15
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
84. PMID: 1303246; 1992, Nat Genet;2(1):37-41
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
85. PMID: 18538015; 2008, BMC Cancer;8():163
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
86. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
87. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56
The tuberous sclerosis complex.
88. PMID: 22923433; 2012, Science;338(6104):221
Genome sequencing identifies a basis for everolimus sensitivity.
89. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
90. PMID: 25724664; 2015, Mol Cancer Ther;14(5):1224-35
Loss of Tuberous Sclerosis Complex 2 (TSC2) Is Frequent in Hepatocellular Carcinoma and Predicts Response to mTORC1 Inhibitor Everolimus.
91. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
92. PMID: 26412398; 2015, Sci Rep;5():14534
PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.
93. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.

ACT Onco[®] + Report

94. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
95. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
96. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
97. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
98. PMID: 20156908; 2010, Oncologist;15(2):122-9
Lapatinib plus letrozole as first-line therapy for HER-2+ hormone receptor-positive metastatic breast cancer.
99. PMID: 17192538; 2006, N Engl J Med;355(26):2733-43
Lapatinib plus capecitabine for HER2-positive advanced breast cancer.
100. PMID: 30948273; 2019, Lancet Oncol;20(5):636-648
Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial.
101. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
102. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
103. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
104. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
105. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
106. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
107. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
108. PMID: 25366685; 2015, J Clin Oncol;33(3):244-50
Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation.
109. PMID: 27932229; 2017, Lancet;389(10064):56-66
Regorafenib for patients with hepatocellular carcinoma who progressed on sorafenib treatment (RESORCE): a randomised, double-blind, placebo-controlled, phase 3 trial.
110. PMID: 23177515; 2013, Lancet;381(9863):295-302
Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial.
111. PMID: 23177514; 2013, Lancet;381(9863):303-12
Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial.

ACT Onco[®] + Report

112. PMID: 24768112; 2014, Lancet;384(9940):319-28
Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial.
113. PMID: 18650514; 2008, N Engl J Med;359(4):378-90
Sorafenib in advanced hepatocellular carcinoma.
114. PMID: 17189398; 2006, Clin Cancer Res;12(24):7271-8
Sorafenib for the treatment of advanced renal cell carcinoma.
115. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
116. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsirrolimus, interferon alfa, or both for advanced renal-cell carcinoma.