

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Colon adenocarcinoma (CRC)	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Lung
	NAME Kuo, Pi-Liang		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S111-75443 B
	DATE OF BIRTH 20 February 1955		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Male		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 11 February 2022
	MEDICAL RECORD # 44789686		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 13 April 2022

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 4 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

KRAS wildtype
NRAS wildtype
APC W593*
MYC amplification
LYN amplification
RAD21 amplification
RBM10 R726*
TP53 R273H

3 Disease relevant genes with no reportable alterations: **BRAF, KRAS, NRAS**

Report Highlights

- Targeted therapies with **NCCN categories of evidence** in this tumor type: Cetuximab (p. 9), Panitumumab (p. 9)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 10)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 4 Muts/Mb

GENOMIC FINDINGS

KRAS - wildtype

0 Trials

NRAS - wildtype

0 Trials

APC - W593*

4 Trials see p. 10

MYC - amplification

4 Trials see p. 11

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)		THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
Cetuximab	2A	none
Panitumumab	2A	
Cetuximab	2A	none
Panitumumab	2A	
none		none
none		none

☐ NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

LYN - amplification	p. 6	RBM10 - R726*	p. 7
RAD21 - amplification	p. 7	TP53 - R273H	p. 8

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵. For patients with chemotherapy-refractory microsatellite-stable (MSS) metastatic colorectal cancer (CRC), a Phase 3 trial reported no OS advantage from the combination of the PD-L1 inhibitor atezolizumab plus cobimetinib relative to regorafenib (8.9 vs. 8.5 months, HR=1.00);

atezolizumab monotherapy similarly did not prolong OS (7.1 vs. 8.5 months, HR=1.19)⁶. For patients with MSS CRC, a Phase 2 study combining ipilimumab and nivolumab reported an overall DCR of 25% (10/40)⁷. Two Phase 1 studies for patients with MSS CRC treated with regorafenib and nivolumab reported PFSs of 7.9 months⁸ and 5.7 months⁹, and a patient with MSS CRC refractory to chemotherapy treated with the PD-1 inhibitor sintilimab and regorafenib reported a CR¹⁰.

— Nontargeted Approaches —

MSI has not been found to be a predictive biomarker for combination chemotherapy regimens, including FOLFOX¹¹⁻¹² and FOLFIRI¹³⁻¹⁴. Patients with MSS CRC are more likely to benefit from postsurgical fluorouracil (FU)-based adjuvant therapy¹⁵⁻¹⁶ but less likely to benefit from irinotecan chemotherapy¹⁷.

FREQUENCY & PROGNOSIS

MSS colorectal cancers (CRCs) make up 70-85% of CRC cases^{3,18-22}. MSS colorectal cancers are

molecularly heterogeneous, driven by diverse mechanisms such as extensive DNA methylation, oncogenic mutations in KRAS or BRAF, or chromosomal instability²². Multiple studies have shown that MSS CRCs have a worse prognosis than MSI-high tumors^{18,23-29}.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor²⁰. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2^{20,30-31}. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers^{19,32-33}. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{19-20,31,33}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

4 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1³⁴⁻³⁶, anti-PD-1 therapies³⁴⁻³⁷, and combination nivolumab and ipilimumab³⁸⁻⁴³. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors^{34-37,44}. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment for patients with 9 types of advanced tumors³⁴. Analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with chemotherapy⁴⁵ or those with lower TMB treated with PD-1 or PD-L1-targeting agents³⁵. However, the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors found significant improvement in ORR for patients with TMB ≥ 10 Muts/Mb (based on this assay or others) compared to those with TMB < 10 Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials^{37,44}. Together, these studies suggest that patients with TMB ≥ 10 Muts/Mb may derive clinical benefit from PD-1 or PD-L1

inhibitors. In CRC specifically, a retrospective analysis of immune checkpoint inhibitor efficacy reported significantly improved OS for patients with tumors harboring TMB ≥ 9.8 Muts/Mb compared with those with tumors with TMB < 9.8 Muts/Mb (~ equivalency < 12 Muts/Mb as measured by this assay)³⁴. Another retrospective study reported that a TMB ≥ 12 Muts/Mb cutoff identifies $> 99\%$ of MSI-High CRC cases but only 3% of MSS cases, indicating the utility of this cutoff for identification of patients with CRC likely to benefit from treatment with immune checkpoint inhibitors⁴⁶.

FREQUENCY & PROGNOSIS

Elevated tumor mutational burden (TMB) has been reported in 8-25% of colorectal cancer (CRC) samples^{21,47-48}. Multiple studies have reported that up to 90% of hypermutated CRC cases exhibit high levels of microsatellite instability (MSI-H) and mismatch repair deficiency (MMR-D)^{21,47}. Increased TMB is significantly associated with MSI-H and MMR-D, with studies reporting that 100% of MSI-H CRCs harbor elevated TMB and conversely that 100% of tumors with low TMB harbor intact MMR⁴⁷. A subset of CRCs that harbor increased TMB but not MSI-H are driven by mutations in POLE, which leads to an "ultramutated" phenotype with especially high TMB^{21,47}. Tumors with increased TMB harbor BRAF V600E mutations more frequently than those with low TMB^{21,47}, whereas TMB-low tumors more frequently harbor mutations in TP53 and APC²¹. The prognostic value of tumor mutational burden (TMB) in colorectal cancer (CRC) is context- and therapy-dependent. A study of tissue TMB (tTMB) in 145 CRC samples showed longer OS in TMB-high samples compared with TMB-low ones⁴⁹. Similarly, for patients with metastatic CRC treated with first-line chemotherapy combined with bevacizumab or

cetuximab, high tissue TMB (tTMB-H) was associated with longer OS⁵⁰. For patients treated with adjuvant chemotherapy, tTMB-H was associated with better 5-year relapse-free survival⁵¹. However, for patients with EGFR/BRAF-inhibitor-treated, BRAF-mutated microsatellite stable (MSS) metastatic CRC, intermediate tTMB was associated with significantly poorer PFS and OS compared with TMB-low status; patients with primary resistance to EGFR/BRAF blockage had higher TMB than those sensitive to these therapies⁵². In a study for 61 patients with metastatic, MSS CRC treated with best standard of care, plasma TMB scores ≥ 28 Muts/Mb (approximately 14 Muts/Mb as measured by this assay) were associated with reduced OS compared with plasma TMB scores < 28 Muts/Mb (3.0 vs. 5.3 months, HR=0.76, $p=0.007$), whereas tTMB was not found to be prognostic in this population⁵³.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁵⁴⁻⁵⁵ and cigarette smoke in lung cancer⁵⁶⁻⁵⁷, treatment with temozolomide-based chemotherapy in glioma⁵⁸⁻⁵⁹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{21,60-63}, and microsatellite instability (MSI)^{21,60,63}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{34,44,46}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

GENOMIC FINDINGS

GENE

KRAS

ALTERATION

wildtype

with clinical benefit of treatment with EGFR-targeting antibodies cetuximab⁶⁴⁻⁶⁷ or panitumumab⁶⁸⁻⁷⁰ for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Guidelines v3.2021).

that KRAS wild-type status is associated with decreased metastasis, better clinicopathological features, and longer survival of patients with CRC^{73-76,80-81}.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated

FREQUENCY & PROGNOSIS

Approximately 50-65% of colorectal cancers (CRCs) have been reported to lack KRAS mutations⁷¹⁻⁷⁹. Numerous studies have reported

FINDING SUMMARY

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor formation⁸²⁻⁸³. No alterations in KRAS were identified in this case.

GENE

NRAS

ALTERATION

wildtype

targeting antibodies cetuximab⁶⁴⁻⁶⁷ or panitumumab⁶⁸⁻⁷⁰ for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Guidelines v3.2021).

of patients with CRC.

FINDING SUMMARY

NRAS encodes a member of the RAS family of small GTPases that mediate transduction of growth signals. Activation of RAS signaling causes cell growth, differentiation, and survival by activating the RAF-MAPK-ERK, PI3K, and other pathways⁸². No alterations in NRAS were identified in this case.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated with clinical benefit of treatment with EGFR-

FREQUENCY & PROGNOSIS

The majority of colorectal cancers (CRCs) (91-98%) have been reported to lack NRAS mutations^{21,79,84-89}. NRAS wild-type status has been reported to be associated with decreased frequency of metastasis⁷⁹ and longer survival⁸⁹⁻⁹⁰

GENE

APC

ALTERATION

W593*

TRANSCRIPT ID

NM_000038

CODING SEQUENCE EFFECT

1778G>A

VARIANT ALLELE FREQUENCY (% VAF)

36.9%

In a Phase 1 trial of the CBP/beta-catenin antagonist E7386, 1 patient with APC-mutated small bowel adenocarcinoma achieved a PR with tumor shrinkage of ~69% and response duration of 165 days⁹⁴; preclinical data support sensitivity of APC-deficient gastric or colorectal cancer models to E7386⁹⁵⁻⁹⁶.

cell division and adhesion. APC interacts with beta-catenin and controls signaling in the WNT pathway, which regulates embryonic development and cell differentiation¹⁰⁰. Alterations such as seen here may disrupt APC function or expression¹⁰¹⁻¹⁰⁵.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved drugs targeting APC inactivation in cancer. Loss of APC function leads to accumulation of beta-catenin and upregulation of WNT pathway transcription programs⁹¹, and potential therapeutic approaches to target this pathway include CBP/beta-catenin antagonists, which interfere with the ability of beta-catenin to interact with transcriptional co-activator CBP⁹²⁻⁹³.

FREQUENCY & PROGNOSIS

APC alterations have been found in 77% of tumors in the Colorectal Adenocarcinoma TCGA dataset²¹. Inactivation of APC leads to activation of the Wnt/beta-catenin pathway, which is thought to play a role in the adenoma-carcinoma transition in some cancers, including colorectal cancer (CRC)⁹⁷. The prognostic significance of APC mutations in sporadic CRC remains unclear⁹⁸. Solid tumors with WNT/beta-catenin pathway alterations, as seen here, were observed to have significantly less T-cell inflammation in one study⁹⁹.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the APC variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with familial adenomatous polyposis (ClinVar, Sep 2021)¹⁰⁶. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in APC are found in more than 90% of patients with familial adenomatous polyposis (FAP)¹⁰⁷⁻¹⁰⁹. The prevalence for FAP in the general population is estimated to be 1:8,300 from birth¹¹⁰, and in the appropriate clinical context germline testing of APC is recommended.

FINDING SUMMARY

APC (adenomatous polyposis coli) encodes a tumor suppressor with critical roles in regulating

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

GENOMIC FINDINGS

GENE

MYC

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no available therapies that directly target MYC. However, preclinical data indicate that MYC overexpression may predict sensitivity to investigational agents targeting CDK1¹¹¹⁻¹¹², CDK2¹¹³, Aurora kinase A¹¹⁴⁻¹²¹, Aurora kinase B¹²²⁻¹²⁵, glutaminase¹²⁶⁻¹²⁹, or BET bromodomain-containing proteins¹³⁰⁻¹³³, as well as agents targeting both HDAC and PI3K¹³⁴⁻¹³⁶. A Phase 2 study reported a PFS benefit associated with a combination of the Aurora A kinase inhibitor alisertib and paclitaxel as second-line therapy for patients with MYC-overexpressed small cell lung cancer but not for patients without MYC overexpression¹³⁷. A patient with MYC-amplified

invasive ductal breast carcinoma experienced a PR to an Aurora kinase inhibitor¹³⁸. The glutaminase inhibitor CB-839, in combination with either everolimus or cabozantinib, has demonstrated encouraging efficacy in Phase 1 and 2 studies enrolling patients with pretreated advanced renal cell carcinoma¹³⁹⁻¹⁴⁰.

— Nontargeted Approaches —

MYC amplification has also been suggested to predict response to chemotherapy in patients with breast cancer in some studies¹⁴¹⁻¹⁴². Preclinical evidence suggests that colon cancer cells with MYC amplification may be more sensitive to 5-fluorouracil and paclitaxel¹⁴³⁻¹⁴⁴.

FREQUENCY & PROGNOSIS

Mutation and amplification of MYC have been observed in up to 2% and 5% of patients with colorectal cancer, respectively (cBioPortal, COSMIC, Sep 2021)¹⁴⁵⁻¹⁴⁷. Overexpression of MYC in colorectal carcinoma has been reported to occur in the absence of MYC amplification¹⁴⁸⁻¹⁵⁰. MYC is

a target gene of beta-catenin and may be upregulated by aberrant WNT signaling in colorectal cancer¹⁵¹. MYC overexpression has been reported in 62-91% of colorectal carcinomas studied^{148,150,152-153}. MYC protein overexpression was reported to be a favorable prognostic biomarker in patients with colorectal cancer, and patients with low-level MYC amplification have been found to have significantly longer survival¹⁵³⁻¹⁵⁴. However, MYC amplification and high expression have been associated with metastatic and aggressive colorectal tumors^{150,155}.

FINDING SUMMARY

MYC (c-MYC) encodes a transcription factor that regulates many genes related to cell cycle regulation and cell growth. It is an oncogene and may be activated in as many as 20% of cancers¹⁵⁶. MYC dysregulation (amplification, overexpression, translocation) has been identified in a number of different cancer types¹⁵⁷. MYC amplification has been significantly linked with increased mRNA and protein levels and results in the dysregulation of a large number of target genes^{156,158-159}.

GENE

LYN

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Dasatinib is a kinase inhibitor that targets the BCR-ABL fusion protein, SRC family kinases including LYN (specifically at low nanomolar concentration)¹⁶⁰⁻¹⁶¹, and other kinases, and has been approved to treat chronic myelocytic leukemia (CML) and acute lymphoblastic leukemia (ALL). A pediatric patient with relapsed B-cell acute lymphoblastic leukemia and an NCOR1-LYN

fusion achieved complete remission after 2 weeks of treatment with dasatinib¹⁶². Similarly, a preclinical study showed that treatment with dasatinib significantly increased survival in a xenograft model of leukemic blast cells harboring NCOR1-LYN¹⁶³. In preclinical studies of LYN-expressing breast and prostate cancer, dasatinib has been reported to inhibit cell migration and invasion^{160,164}. However, amplification or other genomic alterations in LYN in solid tumors, and their potential predictive value for sensitivity of these tumors to dasatinib and other kinase inhibitors, remains poorly understood.

FREQUENCY & PROGNOSIS

LYN alterations are rare in solid tumors¹⁶⁵⁻¹⁶⁶. However, LYN amplification has been reported more frequently, including in ovarian (3.1%),

melanoma (2.3%), prostate (2.2%), breast (1.9%), and endometrial (1.6%) cancers¹⁶⁵⁻¹⁶⁶. LYN expression and activation have also been reported in several types of solid tumors, including glioblastoma¹⁶⁷, prostate cancer¹⁶⁸, head and neck squamous cell carcinoma (HNSCC)¹⁶⁹, Ewing sarcoma¹⁷⁰, and breast cancer¹⁶⁴. High LYN expression was associated with lower survival rates for patients with breast, colorectal, and renal cancers^{164,171-172}.

FINDING SUMMARY

LYN encodes a SRC family intracellular membrane-associated tyrosine protein kinase. LYN is expressed predominantly in hematopoietic cells and conveys signals from the B-cell receptor (BCR) and other receptors to activate the PI3K, STAT, and other signaling pathways¹⁷³⁻¹⁷⁴.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

GENOMIC FINDINGS
GENE

RAD21

ALTERATION
 amplification

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no therapies to target alterations in this gene.

FREQUENCY & PROGNOSIS

RAD21 amplifications have been reported in solid

 tumors, including breast cancers (7%), melanoma (5.4%), and prostate (2.4%) cancers¹⁶⁵. RAD21 overexpression has been correlated with poor prognosis in endometrial cancer¹⁷⁵, breast cancer¹⁷⁶⁻¹⁷⁷, Ewing sarcoma¹⁷⁸, and colorectal cancer (CRC), especially in KRAS-mutant CRC¹⁷⁹.

FINDING SUMMARY

 RAD21 encodes a protein involved in DNA double-strand break repair and sister chromatid cohesion as a part of the cohesin complex¹⁸⁰⁻¹⁸³. In preclinical studies, downregulation of RAD21 or other cohesin components leads to loss of expression from amplified genes, as well as amplifications themselves upon cell passaging¹⁸⁴,

 but also leads to an increase in deletions, insertions, and other rearrangements¹⁸⁵. High RAD21 expression has also been associated with increased genomic instability¹⁸⁶. Cohesin complex also organizes chromatin domains and regulates gene expression¹⁸⁷⁻¹⁸⁸. Both overexpression and reduction of expression of RAD21 has been reported to alter gene expression¹⁸⁹. RAD21 amplification has been correlated with increased expression in breast^{176,186,190} and endometrial¹⁷⁵ cancers. Other RAD21 alterations, including truncating and point mutations, have been reported in the context of cancer, but the majority have not been characterized.

GENE

RBM10

ALTERATION
 R726*

TRANSCRIPT ID
 NM_005676

CODING SEQUENCE EFFECT
 2176C>T

VARIANT ALLELE FREQUENCY (% VAF)
 40.6%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies approved or

clinical trials that directly address genomic alterations in RBM10.

FREQUENCY & PROGNOSIS

 Mutations in RBM10 have been identified in numerous cancers at low frequencies, including non-small cell lung cancer (NSCLC) (7.4%), bladder cancer (5.4%), and thyroid cancer (4.6%)¹⁶⁵⁻¹⁶⁶. Several reports have analyzed the prognostic relevance of RBM10 mutations and expression levels in cancer, but results have been mixed. RBM10 mutations were associated with shorter OS in lung adenocarcinomas¹⁹¹⁻¹⁹², but with enhanced survival in pancreatic ductal adenocarcinomas compared with RBM10 wildtype tumors¹⁹³. Low RBM10 expression was associated with poor prognosis and shorter OS in patients

 with hepatocellular carcinoma (HCC) and pancreatic ductal cancer¹⁹⁴⁻¹⁹⁵, while in lung adenocarcinoma, it was associated with a better prognosis and longer OS¹⁹⁶.

FINDING SUMMARY

 RBM10 encodes RNA binding motif protein 10, a nuclear RNA-binding protein involved in the regulation of alternative splicing¹⁹⁷⁻¹⁹⁸. Germline mutations in RBM10 cause TARP syndrome, an X-linked recessive disorder characterized by development of micrognathia, glossoptosis, and cleft palate¹⁹⁹⁻²⁰⁰.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

R273H

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

818G>A

VARIANT ALLELE FREQUENCY (% VAF)

44.0%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁰¹⁻²⁰⁴, or p53 gene therapy and immunotherapeutics such as SGT-53²⁰⁵⁻²⁰⁹ and ALT-801²¹⁰. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²¹¹. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²¹². A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²¹³. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²¹⁴. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel²¹⁵. A Phase 1

trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations²¹⁶. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²¹⁷. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁰⁹. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutated p53 such as APR-246²¹⁸⁻²²⁰. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR²²¹. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies²²²⁻²²³; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies²²⁴⁻²²⁵. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

TP53 mutations have been reported in up to 60% of colorectal cancer cases^{21,226-231}. A study reported p53 expression in 49% of analyzed colorectal cancer cases²³². TP53 mutation has not been consistently demonstrated to be a significant independent prognostic marker in the context of CRC²³³.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²³⁴. Alterations such

as seen here may disrupt TP53 function or expression²³⁵⁻²³⁹.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Sep 2021)¹⁰⁶. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²⁴⁰⁻²⁴², including sarcomas²⁴³⁻²⁴⁴. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²⁴⁵ to 1:20,000²⁴⁴. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30²⁴⁶. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁴⁷⁻²⁵². CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²⁴⁷⁻²⁴⁸. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease²⁵³. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{251,254-255}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

ORDERED TEST # ORD-1342720-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Cetuximab

Assay findings association
KRAS
wildtype

NRAS
wildtype

AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including cetuximab, have been shown to have significant clinical activity for patients with CRC^{64-67,256-257}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of cetuximab in metastatic CRC (NCCN Colon Cancer Guidelines v3.2021).

SUPPORTING DATA

Cetuximab has been shown to improve OS, PFS, and response rate for patients with KRAS-wild-type CRC, both as first-line combination therapy with FOLFIRI or FOLFOX₄^{64-65,257} and as monotherapy or combination therapy with irinotecan for chemotherapy-refractory

patients^{66-67,256}. A prospective study of first-line cetuximab for patients with KRAS/NRAS/BRAF mutation-negative metastatic CRC resulted in limited efficacy, with 10.5% (2/19) of participants experiencing PRs and 57.9% (11/19) experiencing SDs²⁵⁸. The Phase 2 AVETUX trial of cetuximab combined with avelumab and mFOLFOX6 for patients with RAS- and BRAF-wild-type metastatic CRC resulted in an ORR of 79.5% (6 CR and 25 PRs, n=39) and a DCR of 92.3%²⁵⁹. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)²⁶⁰. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)²⁶¹.

Panitumumab

Assay findings association
KRAS
wildtype

NRAS
wildtype

AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including panitumumab, have been shown to have significant clinical activity for patients with CRC^{68,260,262}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of panitumumab in metastatic CRC (NCCN Colon Cancer Guidelines v3.2021).

SUPPORTING DATA

Panitumumab has been shown to improve OS, PFS, and ORR for patients with KRAS wild-type CRC, both as first-line combination therapy with FOLFOX₄⁶⁸ and as

monotherapy for chemotherapy-refractory patients^{260,262}. An open-label, randomized Phase 2 trial reported that for patients with unresectable RAS-wild-type colorectal adenocarcinoma treated with first-line panitumumab plus FOLFOX₄, maintenance with a combination of panitumumab plus fluorouracil and leucovorin was superior to panitumumab monotherapy (10-month PFS 59% vs. 49%)²⁶³. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)²⁶⁰. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)²⁶¹.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
APC
ALTERATION
W593*

RATIONALE

Based on preclinical and limited clinical data, APC inactivation may be associated with sensitivity to

CBP/beta-catenin interaction inhibitors.

NCT03833700
PHASE 1

A Study of E7386 in Participants With Advanced Solid Tumor Including Colorectal Cancer (CRC)

TARGETS
CBP, Beta-catenin

LOCATIONS: Fukuoka (Japan), Nagaizumi-cho (Japan), Chuo Ku (Japan), Kashiwa (Japan)

NCT05091346
PHASE 1/2

A Study of E7386 in Combination With Pembrolizumab in Previously Treated Participants With Selected Solid Tumors

TARGETS
CBP, Beta-catenin, PD-1

LOCATIONS: Tokyo (Japan)

NCT04008797
PHASE 1

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

TARGETS
CBP, Beta-catenin, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Osakasayama (Japan), Chuo-Ku (Japan), Kashiwa (Japan)

NCT03264664
PHASE 1

Study of E7386 in Participants With Selected Advanced Neoplasms

TARGETS
CBP, Beta-catenin

LOCATIONS: Glasgow (United Kingdom), Manchester (United Kingdom), Sutton (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

CLINICAL TRIALS

GENE

MYC

ALTERATION

amplification

RATIONALE

MYC overexpression may predict sensitivity to inhibition of CDKs, especially CDK1 and CDK2, of Aurora kinases, including Aurora kinase A and B,

and of BET domain proteins, which are reported to downregulate MYC expression and MYC-dependent transcriptional programs.

NCT04983810
PHASE 1/2

A Study to Investigate Fadraciclub (CYC065), in Subjects With Advanced Solid Tumors and Lymphoma

TARGETS
CDK2, CDK9

LOCATIONS: Seoul (Korea, Republic of), Barcelona (Spain), California, Texas

NCT03220347
PHASE 1

A Study to Assess the Safety, Tolerability, Pharmacokinetics and Preliminary Efficacy of CC-90010 in Subjects With Advanced Solid Tumors and Relapsed/Refractory Non-Hodgkin's Lymphomas

TARGETS
BRD2, BRD3, BRD4, BRDT

LOCATIONS: Chikusa-ku (Japan), Koto-ku (Japan), Kashiwa (Japan), Madrid (Spain)

NCT04555837
PHASE 1/2

Alisertib and Pembrolizumab for the Treatment of Patients With Rb-deficient Head and Neck Squamous Cell Cancer

TARGETS
Aurora kinase A, PD-1

LOCATIONS: Texas

NCT01434316
PHASE 1

Veliparib and Dinaciclib in Treating Patients With Advanced Solid Tumors

TARGETS
PARP, CDK1, CDK9, CDK5, CDK2

LOCATIONS: Massachusetts

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ARFRP1
amplification

BCL2L1
amplification

CBL
amplification

FGF19
amplification

KMT2A (MLL)
amplification

NBN
amplification

ZNF217
amplification

ASXL1
amplification

BCL2L2
P33A

CCND1
amplification

FGF3
amplification

MRE11A
amplification

SDHD
amplification

ATM
C353S and amplification

BCORL1
T1111M

CHEK1
amplification

FGF4
amplification

MSH2
I169V

SRC
amplification

AURKA
amplification

C11ORF30 (EMSY)
amplification

EED
amplification

GNAS
amplification

MST1R
R504C

TSC2
A1141T

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKKN1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKB1A	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV1
ETV4	ETVS	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT
KMT2A (MLL)	MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA
RAF1	RARA	RET	ROS1	RSPO2	SDC4	SLC34A2	TERC*	TERT**
TPRSS2								

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX
About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.


ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table
 Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 - Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 291656669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 6.1.0

The median exon coverage for this sample is 1,011x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX
References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Ciardiello et al., 2018; ESMO Abstract LBA-004
7. Parikh et al., 2021; DOI: 10.1038/s43018-021-00269-7
8. Fukuoka S, et al. J. Clin. Oncol. (2020) PMID: 32343640
9. Kim et al., 2020; DOI: 10.1016/j.jannonc.2020.04.073
10. Zhang Y, et al. BMC Gastroenterol (2021) PMID: 34688262
11. Sinicrope FA, et al. J. Clin. Oncol. (2013) PMID: 24019539
12. Gavin PG, et al. Clin. Cancer Res. (2012) PMID: 23045248
13. Bertagnolli MM, et al. J. Clin. Oncol. (2009) PMID: 19273709
14. Van Cutsem E, et al. J. Clin. Oncol. (2009) PMID: 19451425
15. Ribic CM, et al. N. Engl. J. Med. (2003) PMID: 12867608
16. Sargent DJ, et al. J. Clin. Oncol. (2010) PMID: 20498393
17. Fallik D, et al. Cancer Res. (2003) PMID: 14522894
18. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
19. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
20. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
21. Nature (2012) PMID: 22810696
22. Histopathology (2007) PMID: 17204026
23. Samowitz WS, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11535541
24. Elsaleh H, et al. Clin Colorectal Cancer (2001) PMID: 12445368
25. Brueckl WM, et al. Anticancer Res. () PMID: 12820457
26. Guidoboni M, et al. Am. J. Pathol. (2001) PMID: 11438476
27. Gryfe R, et al. N. Engl. J. Med. (2000) PMID: 10631274
28. Sinicrope FA, et al. Gastroenterology (2006) PMID: 16952542
29. Laghi L, et al. Dig Dis (2012) PMID: 22722556
30. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
31. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
32. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
33. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
34. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
35. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
36. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
37. Cristescu R, et al. Science (2018) PMID: 30309915
38. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
39. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
40. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
41. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
42. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
43. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
44. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32199526
45. Legrand et al., 2018; ASCO Abstract 12000
46. Fabrizio DA, et al. J Gastrointest Oncol (2018) PMID: 30151257
47. Stadler ZK, et al. J. Clin. Oncol. (2016) PMID: 27022117
48. Shao C, et al. JAMA Netw Open (2020) PMID: 33119110
49. Schwartz et al., 2018; ASCO Abstract 572
50. Innocenti F, et al. J Clin Oncol (2019) PMID: 30865548
51. Lee DW, et al. Clin Cancer Res (2019) PMID: 31285374
52. Randon G, et al. Eur J Cancer (2022) PMID: 34933155
53. Chen EX, et al. JAMA Oncol (2020) PMID: 32379280
54. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
55. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
56. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
57. Rizvi NA, et al. Science (2015) PMID: 25765070
58. Johnson BE, et al. Science (2014) PMID: 24336570
59. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
60. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
61. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
62. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
63. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
64. Van Cutsem E, et al. J. Clin. Oncol. (2011) PMID: 21502544
65. Bokemeyer C, et al. Ann. Oncol. (2011) PMID: 21228335
66. Karapetis CS, et al. N. Engl. J. Med. (2008) PMID: 18946061
67. De Roock W, et al. Ann. Oncol. (2008) PMID: 17998284
68. Douillard JY, et al. Ann. Oncol. (2014) PMID: 24718886
69. Douillard JY, et al. N. Engl. J. Med. (2013) PMID: 24024839
70. Amado RG, et al. J. Clin. Oncol. (2008) PMID: 18316791
71. Lièvre A, et al. Cancer Res. (2006) PMID: 16618717
72. De Roock W, et al. Lancet Oncol. (2011) PMID: 21163703
73. Chen J, et al. BMC Cancer (2014) PMID: 25367198
74. Li W, et al. BMC Cancer (2015) PMID: 25929517
75. Hu J, et al. Medicine (Baltimore) (2016) PMID: 27977612
76. Zekri J, et al. Genet. Mol. Res. (2017) PMID: 28218784
77. Staudacher JJ, et al. Clin Transl Gastroenterol (2017) PMID: 29048416
78. Wang Y, et al. Virchows Arch. (2018) PMID: 29705968
79. Guo F, et al. Sci Rep (2018) PMID: 29666387
80. Mármol I, et al. Int J Mol Sci (2017) PMID: 28106826
81. Kwak MS, et al. Medicine (Baltimore) (2017) PMID: 28858102
82. Pylyayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) PMID: 21993244
83. Kahn S, et al. Anticancer Res. () PMID: 3310850
84. Pentheroudakis G, et al. BMC Cancer (2013) PMID: 23374602
85. Vaughn CP, et al. Genes Chromosomes Cancer (2011) PMID: 21305640
86. Janku F, et al. Target Oncol (2013) PMID: 23400451
87. De Roock W, et al. Lancet Oncol. (2010) PMID: 20619739
88. Irahara N, et al. Diagn. Mol. Pathol. (2010) PMID: 20736745
89. Schirripa M, et al. Int. J. Cancer (2015) PMID: 24806288
90. Cercek A, et al. Clin. Cancer Res. (2017) PMID: 28446505
91. Zhan T, et al. Oncogene (2017) PMID: 27617575
92. Jung YS, et al. Exp Mol Med (2020) PMID: 32037398
93. Krishnamurthy N, et al. Cancer Treat Rev (2018) PMID: 29169144
94. Kawazoe et al., 2021; ESMO Abstract 473P
95. Yamada K, et al. Cancer Res (2021) PMID: 33408116
96. Kanda Y, et al. Biochem Biophys Res Commun (2022) PMID: 34837838
97. Fu Y, et al. Int. J. Cancer (2011) PMID: 21455986
98. Quyn AJ, et al. Surgeon (2008) PMID: 19110823
99. Luke JJ, et al. Clin Cancer Res (2019) PMID: 30635339
100. Logan CY, et al. Annu. Rev. Cell Dev. Biol. (2004) PMID: 15473860
101. Eklof Spink K, et al. EMBO J. (2001) PMID: 11707392
102. Liu J, et al. J. Mol. Biol. (2006) PMID: 16753179
103. Dikovskaya D, et al. J. Cell. Sci. (2010) PMID: 20144988
104. Murphy SJ, et al. Dig. Dis. Sci. (2007) PMID: 17410430
105. Aretz S, et al. Hum. Mutat. (2004) PMID: 15459959
106. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
107. Kerr SE, et al. J Mol Diagn (2013) PMID: 23159591
108. Annu Rev Pathol (2011) PMID: 21090969
109. Kastritis E, et al. Int. J. Cancer (2009) PMID: 18844223
110. Half E, et al. Orphanet J Rare Dis (2009) PMID: 19822006
111. Horiuchi D, et al. J. Exp. Med. (2012) PMID: 22430491
112. Goga A, et al. Nat. Med. (2007) PMID: 17589519
113. Molenaar JJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19525400
114. Dammert MA, et al. Nat Commun (2019) PMID: 31375684
115. Mollaoglu G, et al. Cancer Cell (2017) PMID: 28089889
116. Cardnell RJ, et al. Oncotarget (2017) PMID: 29088717
117. Wang L, et al. Mol Oncol (2017) PMID: 28417568
118. Takahashi Y, et al. Ann. Oncol. (2015) PMID: 25632068
119. Li Y, et al. Thyroid (2018) PMID: 30226440
120. Mahadevan D, et al. PLoS ONE (2014) PMID: 24893165
121. Park SI, et al. Target Oncol (2019) PMID: 31429028
122. Helfrich BA, et al. Mol. Cancer Ther. (2016) PMID: 27496133
123. Hook KE, et al. Mol. Cancer Ther. (2012) PMID: 22222631
124. Yang D, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20643922
125. He J, et al. Anticancer Drugs (2019) PMID: 30540594
126. Shroff EH, et al. Proc. Natl. Acad. Sci. U.S.A. (2015) PMID: 25964345
127. Effenberger M, et al. Oncotarget (2017) PMID: 29156762
128. Qu X, et al. Biochem. Biophys. Res. Commun. (2018) PMID: 30103944
129. Xiang Y, et al. J. Clin. Invest. (2015) PMID: 25915584
130. Delmore JE, et al. Cell (2011) PMID: 21889194
131. Bandopadhyay P, et al. Clin. Cancer Res. (2014) PMID: 24297863
132. Lovén J, et al. Cell (2013) PMID: 23582323
133. Otto C, et al. Neoplasia (2019) PMID: 31734632
134. Dong LH, et al. J Hematol Oncol (2013) PMID: 23866964
135. Pei Y, et al. Cancer Cell (2016) PMID: 26977882
136. Fu XH, et al. Acta Pharmacol. Sin. (2019) PMID: 30224636
137. Owonikoko TK, et al. J Thorac Oncol (2020) PMID: 31655296
138. Ganesan P, et al. Mol. Cancer Ther. (2014) PMID: 25253784
139. Tannir et al., 2018; ASCO GU Abstract 603
140. Motzer et al., 2019; ESMO Abstract LBA54
141. Pereira CB, et al. PLoS ONE (2013) PMID: 23555992
142. Yasojima H, et al. Eur. J. Cancer (2011) PMID: 21741827
143. Arango D, et al. Cancer Res. (2001) PMID: 11406570
144. Bottone MG, et al. Exp. Cell Res. (2003) PMID: 14516787
145. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
146. Gao J, et al. Sci Signal (2013) PMID: 23550210
147. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
148. Erisman MD, et al. Mol. Cell. Biol. (1985) PMID: 3837853
149. Erisman MD, et al. Oncogene (1988) PMID: 3283655
150. Wang J, et al. Chin. Med. Sci. J. (1994) PMID: 8086630

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1342720-01

APPENDIX **References**

151. Herbst A, et al. BMC Genomics (2014) PMID: 24467841
152. Huang MY, et al. Cancer Biomark (2013) PMID: 24240588
153. Toon CW, et al. PLoS ONE (2014) PMID: 24503701
154. Augenlicht LH, et al. Cancer Res. (1997) PMID: 9135021
155. Monnat M, et al. Int. J. Cancer (1987) PMID: 3040597
156. Dang CV, et al. Semin. Cancer Biol. (2006) PMID: 16904903
157. Nesbit CE, et al. Oncogene (1999) PMID: 10378696
158. Blancato J, et al. Br. J. Cancer (2004) PMID: 15083194
159. Fromont G, et al. Hum. Pathol. (2013) PMID: 23574779
160. Nam S, et al. Cancer Res. (2005) PMID: 16230377
161. Williams NK, et al. J. Biol. Chem. (2009) PMID: 18984583
162. Dai HP, et al. Front Oncol (2020) PMID: 32266142
163. Tomii T, et al. Leukemia (2021) PMID: 33199837
164. Choi YL, et al. Cancer Res. (2010) PMID: 20215510
165. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
166. Nguyen B, et al. Cell (2022) PMID: 35120664
167. Stettner MR, et al. Cancer Res. (2005) PMID: 15994925
168. Goldenberg-Furmanov M, et al. Cancer Res. (2004) PMID: 14871838
169. Wheeler SE, et al. Clin. Cancer Res. (2012) PMID: 22490227
170. Guan H, et al. Mol. Cancer Ther. (2008) PMID: 18644993
171. Huang TH, et al. Cancer Cell (2013) PMID: 23764002
172. Roseweir AK, et al. BMC Cancer (2016) PMID: 26984511
173. Xu Y, et al. Immunity (2005) PMID: 15664155
174. Cell Commun. Signal (2012) PMID: 22805580
175. Supernat A, et al. Oncol Lett (2012) PMID: 23205091
176. Xu H, et al. Breast Cancer Res. (2011) PMID: 21255398
177. Sharaf R, et al. Genome Med (2022) PMID: 35227290
178. Su XA, et al. Genes Dev (2021) PMID: 33766983
179. Deb S, et al. Br. J. Cancer (2014) PMID: 24548858
180. Xu H, et al. Nat. Rev. Cancer (2011) PMID: 21326324
181. Hill VK, et al. Biochim. Biophys. Acta (2016) PMID: 27207471
182. Solomon DA, et al. BMB Rep (2014) PMID: 24856830
183. Bauerschmidt C, et al. Nucleic Acids Res. (2010) PMID: 19906707
184. Yun J, et al. Nucleic Acids Res. (2016) PMID: 26420833
185. Gelot C, et al. Nucleus (2016) PMID: 27326661
186. Yan M, et al. Breast Cancer Res. (2012) PMID: 22537934
187. Sofueva S, et al. EMBO J. (2013) PMID: 24185899
188. Deng Z, et al. EMBO J. (2012) PMID: 23010778
189. Yun J, et al. EMBO Rep. (2016) PMID: 27466323
190. Mahmood SF, et al. Carcinogenesis (2014) PMID: 24148822
191. Yin LL, et al. Oncol Lett (2018) PMID: 30405763
192. Li Z, et al. Cell Cycle (2020) PMID: 33064970
193. Witkiewicz AK, et al. Nat Commun (2015) PMID: 25855536
194. Zhao Z, et al. Eur Rev Med Pharmacol Sci (2020) PMID: 32572914
195. Xiao W, et al. Am J Cancer Res (2021) PMID: 33520366
196. Sun X, et al. Int J Oncol (2019) PMID: 30483773
197. Wang Y, et al. EMBO Mol Med (2013) PMID: 24000153
198. Inoue A, et al. FEBS Lett. (2014) PMID: 24530524
199. Johnston JJ, et al. Am. J. Hum. Genet. (2010) PMID: 20451169
200. Gripp KW, et al. Am. J. Med. Genet. A (2011) PMID: 21910224
201. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
202. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
203. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
204. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
205. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
206. Xu L, et al. Mol. Med. (2001) PMID: 11713371
207. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
208. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
209. Pirolo KF, et al. Mol. Ther. (2016) PMID: 27357628
210. Hajdenberg et al., 2012; ASCO Abstract e15010
211. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
212. Moore et al., 2019; ASCO Abstract 5513
213. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
214. Oza et al., 2015; ASCO Abstract 5506
215. Lee J, et al. Cancer Discov (2019) PMID: 31315834
216. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
217. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
218. Lehmann S, et al. J. Clin. Oncol. (2012) PMID: 22965953
219. Mohell N, et al. Cell Death Dis (2015) PMID: 26086967
220. Franssón Á, et al. J Ovarian Res (2016) PMID: 27179933
221. Gourley et al., 2016; ASCO Abstract 5571
222. Kwok M, et al. Blood (2016) PMID: 26563132
223. Boudny M, et al. Haematologica (2019) PMID: 30975914
224. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
225. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
226. Goh HS, et al. Cancer Res. (1995) PMID: 7585578
227. Berg M, et al. PLoS ONE (2010) PMID: 21103049
228. Han SW, et al. PLoS ONE (2013) PMID: 23700467
229. Peeters M, et al. Clin. Cancer Res. (2013) PMID: 23325582
230. Malhotra P, et al. Tumour Biol. (2013) PMID: 23526092
231. Di Bartolomeo M, et al. Target Oncol (2014) PMID: 23821376
232. Wangefjord S, et al. Diagn Pathol (2013) PMID: 23337059
233. Russo A, et al. J. Clin. Oncol. (2005) PMID: 16172461
234. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
235. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
236. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
237. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
238. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
239. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
240. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
241. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100
242. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
243. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
244. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
245. Lalloo F, et al. Lancet (2003) PMID: 12672316
246. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713
247. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
248. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
249. Xie M, et al. Nat. Med. (2014) PMID: 25326804
250. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
251. Severson EA, et al. Blood (2018) PMID: 29678827
252. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
253. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
254. Chabon JJ, et al. Nature (2020) PMID: 32269342
255. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
256. Cunningham D, et al. N. Engl. J. Med. (2004) PMID: 15269313
257. Jonker DJ, et al. N. Engl. J. Med. (2007) PMID: 18003960
258. Moiseyenko VM, et al. Clin Drug Investig (2018) PMID: 29470838
259. Stein et al., 2020; ASCO GI Abstract 96
260. Price TJ, et al. Lancet Oncol. (2014) PMID: 24739896
261. Sakai D, et al. Eur J Cancer (2020) PMID: 32526634
262. Van Cutsem E, et al. J. Clin. Oncol. (2007) PMID: 17470858
263. Pietrantonio F, et al. JAMA Oncol (2019) PMID: 31268481

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531