



Sample Information

Patient Name: 黃欣捷  
Gender: Female  
ID No.: F227152734  
History No.: 45589466  
Age: 34  
  
Ordering Doctor: DOC6258D 林益庭  
Ordering REQ.: OCNKSSG  
Signing in Date: 2023/07/18

Path No.: M112-00186  
MP No.: MY23046  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/07/13

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	16 Therapies Available
Relevant Therapy Summary	3	0 Clinical Trials
Relevant Therapy Details	4	
Alert Details	14	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	None detected	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	FLT3 ITD mutation	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>FLT3 ITD mutation</b> fms related receptor tyrosine kinase 3 Allele Frequency: 47.96%	<b>gilteritinib</b> <sup>1,2</sup> <b>midostaurin + chemotherapy</b> <sup>1,2</sup> allogeneic stem cells azacitidine cytarabine cytarabine + fludarabine + idarubicin + filgrastim daunorubicin decitabine gemtuzumab ozogamicin + chemotherapy gilteritinib + chemotherapy idarubicin midostaurin mitoxantrone sorafenib sorafenib + chemotherapy venetoclax + chemotherapy	None	0

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
FLT3	p.(Y597_E598insDDF REY)	c.1793_1794insTGAT TTCAGAGAATATGA	COSM5879511	chr13:28608262	47.96%	NM_004119.3	nonframeshift Insertion	1981
CEBPA	p.(H195_P196dup)	c.589_590insACCCG C		chr19:33792731	34.27%	NM_004364.4	nonframeshift Insertion	1036

## Biomarker Descriptions

### FLT3 (fms related receptor tyrosine kinase 3)

**Background:** The FLT3 gene encodes the fms related tyrosine kinase 3, a tyrosine kinase receptor that is a member of the class III receptor tyrosine kinase family that also includes PDGFR, FMS, and KIT<sup>1</sup>. FLT3 is highly expressed in hematopoietic progenitor cells<sup>2</sup>. Genomic alterations in FLT3 activate downstream oncogenic pathways including PI3K/AKT/mTOR and RAS/RAF/MEK/ERK pathways which promote cellular proliferation, survival, and inhibition of differentiation<sup>1</sup>.

**Alterations and prevalence:** Somatic mutations occur in approximately 30% of acute myeloid leukemia (AML), 7-10% of melanoma, and up to 8% of uterine cancer<sup>3,4,5,6</sup>. The most common activating FLT3 mutations are internal tandem duplications (ITD) that range from 3 to 400 base pairs in length within exons 14 and 15 in the juxtamembrane (JM) domain<sup>7</sup>. The second most frequent mutations are point mutations in exon 20 within the tyrosine kinase domain (TKD)<sup>8</sup>. FLT3 is amplified in up to 8% of colorectal cancer, 3% of stomach cancer, and is commonly overexpressed in AML<sup>5,6,9</sup>.

**Potential relevance:** FLT3 rearrangements are recognized by the World Health Organization (WHO) as one of the possible molecular abnormality requirements that define myeloid/lymphoid neoplasms with eosinophilia and tyrosine kinase gene fusions<sup>10</sup>. The presence of FLT3-ITD confers poor prognosis in myelodysplastic syndrome (MDS)<sup>11</sup>. Concurrent expression of FLT-ITD with mutant or wild-type NPM1 (when lacking adverse risk genetic lesions) confers intermediate risk in AML<sup>12,13</sup>. FLT3 TKD mutation at D835 confers poor prognosis in MDS<sup>11</sup>. Midostaurin<sup>14</sup> (2017) and gilteritinib<sup>15</sup> (2018) are kinase inhibitors approved for AML patients with FLT3-ITD and TKD mutations including D835 and I836 mutations. The FDA granted fast track designations in 2017 to crenolanib<sup>16</sup> and in 2022 to tuspentinib (HM43239)<sup>17</sup> for FLT3 mutation-positive relapsed or refractory AML. In 2018 the FDA granted breakthrough therapy designation to quizartinib<sup>18</sup> for AML with FLT3-ITD. A phase II trial testing crenolanib in 34 patients with FLT3-ITD and TKD mutated relapsed/refractory AML, reported that FLT3 inhibitor naïve patients demonstrated a longer overall survival (OS) and event free

## Biomarker Descriptions (continued)

survival (EFS) in comparison to previously treated patients (median OS: 55 weeks vs 13 weeks; median EFS: 13 weeks vs 7 weeks)<sup>19</sup>. Another phase II trial of crenolanib with chemotherapy in newly diagnosed FLT3 mutated AML reported complete remission in 24/29 (83%) patients<sup>20</sup>. Several multi-targeted tyrosine kinase inhibitors such as sorafenib (2005), sunitinib (2006), cabozantinib (2012), and ponatinib (2012) are FDA approved and include FLT3 as a target. Sorafenib is recommended in combination with chemotherapy in FLT3-ITD mutated AML<sup>12</sup>.

## Relevant Therapy Summary

● In this cancer type    
 ○ In other cancer type    
 ① In this cancer type and other cancer types    
 ✕ No evidence

FLT3 ITD mutation					
Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
gilteritinib	●	●	●	●	✕
midostaurin + cytarabine + daunorubicin	●	✕	●	●	✕
Allogeneic hematopoietic stem cell transplantation	✕	●	✕	✕	✕
azacitidine	✕	●	✕	✕	✕
cytarabine	✕	●	✕	✕	✕
cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
daunorubicin	✕	●	✕	✕	✕
decitabine	✕	●	✕	✕	✕
gemtuzumab ozogamicin + cytarabine + daunorubicin	✕	●	✕	✕	✕
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
gemtuzumab ozogamicin + daunorubicin	✕	●	✕	✕	✕
gilteritinib + azacitidine	✕	●	✕	✕	✕
idarubicin	✕	●	✕	✕	✕
midostaurin	✕	●	✕	✕	✕
midostaurin + cytarabine	✕	●	✕	✕	✕
midostaurin + daunorubicin	✕	●	✕	✕	✕
mitoxantrone	✕	●	✕	✕	✕
sorafenib	✕	●	✕	✕	✕
sorafenib + azacitidine	✕	●	✕	✕	✕
sorafenib + decitabine	✕	●	✕	✕	✕
venetoclax + azacitidine	✕	●	✕	✕	✕
venetoclax + cytarabine	✕	●	✕	✕	✕
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕

## Relevant Therapy Summary (continued)

☒ In this cancer type   ☐ In other cancer type   ☒ In this cancer type and other cancer types   ☒ No evidence

### FLT3 ITD mutation (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + decitabine	×	●	×	×	×

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type   ☐ In other cancer type   ☒ In this cancer type and other cancer types

FDA information is current as of 2023-05-17. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### FLT3 ITD mutation

#### ● gilteritinib

**Cancer type:** Acute Myeloid Leukemia

**Label as of:** 2022-01-12

**Variant class:** FLT3 ITD mutation

**Indications and usage:**

XOSPATA® is a kinase inhibitor indicated for the treatment of adult patients who have relapsed or refractory acute myeloid leukemia (AML) with a FLT3 mutation as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/211349s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/211349s003lbl.pdf)

#### ● midostaurin + cytarabine + daunorubicin

**Cancer type:** Acute Myeloid Leukemia

**Label as of:** 2021-11-15

**Variant class:** FLT3 ITD mutation

**Indications and usage:**

RYDAPT® is a kinase inhibitor indicated for the treatment of adult patients with:

- Newly diagnosed acute myeloid leukemia (AML) that is FLT3 mutation-positive as detected by an FDA-approved test, in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation.

Limitations of Use: RYDAPT® is not indicated as a single-agent induction therapy for the treatment of patients with AML.

- Aggressive systemic mastocytosis (ASM), systemic mastocytosis with associated hematological neoplasm (SM-AHN), or mast cell leukemia (MCL).

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/207997s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/207997s008lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-05-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

### FLT3 ITD mutation

#### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

#### ● daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

#### ● gilteritinib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Residual (Recurrence therapy)
- (Induction therapy, Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: CD33 positive, NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)
- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● gemtuzumab ozogamicin + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: CD33 positive, NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)
- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● midostaurin + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● midostaurin + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Residual (Recurrence therapy)
- (Induction therapy, Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● sorafenib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● sorafenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]



## FLT3 ITD mutation (continued)

### ● sorafenib + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention
- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● gilteritinib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● gilteritinib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● midostaurin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)
- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

Other criteria: NPM1 wild type

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● sorafenib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● sorafenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## FLT3 ITD mutation (continued)

### ● sorafenib + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

### ● venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

## Current EMA Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

EMA information is current as of 2023-05-17. For the most up-to-date information, search [www.ema.europa.eu/ema](https://www.ema.europa.eu/ema).

### FLT3 ITD mutation

#### ☒ gilteritinib

Cancer type: Acute Myeloid Leukemia

Label as of: 2021-09-08

Variant class: FLT3 ITD mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/xospata-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/xospata-epar-product-information_en.pdf)

#### ☒ midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Label as of: 2023-05-11

Variant class: FLT3 ITD mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/rydapt-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/rydapt-epar-product-information_en.pdf)

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

ESMO information is current as of 2023-05-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### FLT3 ITD mutation

#### ☒ gilteritinib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

#### ☒ midostaurin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 ITD mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Alerts Informed By Public Data Sources

### Current FDA Information

☒ Contraindicated    ☒ Not recommended    ☒ Resistance    ☒ Breakthrough    ☒ Fast Track

FDA information is current as of 2023-05-17. For the most up-to-date information, search [www.fda.gov](http://www.fda.gov).

### FLT3 ITD mutation

#### ☒ crenolanib

Cancer type: Acute Myeloid Leukemia

Variant class: FLT3 mutation

Supporting Statement:

The FDA has granted Fast Track Designation to the benzimidazole type I kinase inhibitor, crenolanib, for:

- FLT3 mutation-positive relapsed or refractory acute myeloid leukemia (AML)
- PDGFRA D842V mutated unresectable or metastatic gastrointestinal stromal tumors (GIST)

Reference:

<https://www.globenewswire.com/news-release/2017/12/01/1216122/0/en/Arog-Pharmaceuticals-Receives-FDA-Fast-Track-Designation-for-Crenolanib-in-Relapsed-or-Refractory-FLT3-Positive-AML.html>

## FLT3 ITD mutation (continued)

### **A** tuspetinib

**Cancer type:** Acute Myeloid Leukemia

**Variant class:** FLT3 mutation

**Supporting Statement:**

The FDA has granted Fast Track Designation to tuspetinib (HM43239), a myeloid kinase inhibitor, for relapsed or refractory (R/R) acute myeloid leukemia (AML) with FLT3 mutation.

**Reference:**

<https://www.aptose.com/news-media/press-releases/detail/230/aptose-receives-fast-track-designation-for-hm43239-in>

## References

1. Grafone et al. An overview on the role of FLT3-tyrosine kinase receptor in acute myeloid leukemia: biology and treatment. *Oncol Rev.* 2012 Mar 5;6(1):e8. PMID: 25992210
2. Short et al. Emerging treatment paradigms with FLT3 inhibitors in acute myeloid leukemia. *Ther Adv Hematol.* 2019; 10: 2040620719827310. PMID: 30800259
3. Small. FLT3 mutations: biology and treatment. *Hematology Am Soc Hematol Educ Program.* 2006:178-84. PMID: 17124058
4. Cancer Genome Atlas Research Network. Genomic and epigenomic landscapes of adult de novo acute myeloid leukemia. *N Engl J Med.* 2013 May 30;368(22):2059-74. doi: 10.1056/NEJMoa1301689. Epub 2013 May 1. PMID: 23634996
5. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
6. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
7. Nakao et al. Internal tandem duplication of the *flt3* gene found in acute myeloid leukemia. *Leukemia.* 1996 Dec;10(12):1911-8. PMID: 8946930
8. Yamamoto et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood.* 2001 Apr 15;97(8):2434-9. PMID: 11290608
9. Carow et al. Expression of the hematopoietic growth factor receptor FLT3 (STK-1/Flk2) in human leukemias. *Blood.* 1996 Feb 1;87(3):1089-96. PMID: 8562934
10. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia.* 2022 Jul;36(7):1703-1719. PMID: 35732831
11. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
12. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]
13. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
14. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/207997s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/207997s008lbl.pdf)
15. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/211349s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/211349s003lbl.pdf)
16. <https://www.globenewswire.com/news-release/2017/12/01/1216122/0/en/Arog-Pharmaceuticals-Receives-FDA-Fast-Track-Designation-for-Crenolanib-in-Relapsed-or-Refractory-FLT3-Positive-AML.html>
17. <https://www.aptose.com/news-media/press-releases/detail/230/aptose-receives-fast-track-designation-for-hm43239-in>
18. [https://www.daiichisankyo.com/media\\_investors/media\\_relations/press\\_releases/detail/006896.html](https://www.daiichisankyo.com/media_investors/media_relations/press_releases/detail/006896.html)
19. Jasleen et al. Results of a Phase II Study of Crenolanib in Relapsed/Refractory Acute Myeloid Leukemia Patients (Pts) with Activating FLT3 Mutations. *Blood.* 124:389
20. Bartus et al. Crenolanib in combination with standard chemotherapy in newly diagnosed FLT3-mutated AML patients. *ASH 2017. Abstract #566*