



Sample Information

Patient Name: 許丕煌  
Gender: Male  
ID No.: W100071546  
History No.: 42673063  
Age: 56  
  
Ordering Doctor: DOC1697J 蔡淳光  
Ordering REQ.: 0CERPVI  
Signing in Date: 2023/01/12

Path No.: M112-00003  
MP No.: MY23001  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/01/06

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	4 Therapies Available
Relevant Therapy Summary	4	0 Clinical Trials
Relevant Therapy Details	4	
Prognostic Details	8	
Diagnostic Details	9	

Relevant Acute Myeloid Leukemia Variants

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	None detected	MYH11	<b>CBFB-MYH11 fusion</b>
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>CBFB-MYH11 fusion</b> core-binding factor subunit beta - myosin heavy chain 11  <b>Prognostic significance:</b> ELN 2017: Favorable <b>Diagnostic significance:</b> Acute Myeloid Leukemia	cytarabine + daunorubicin cytarabine + idarubicin cytarabine + mitoxantrone gemtuzumab ozogamicin + chemotherapy	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

### Prevalent cancer biomarkers without relevant evidence based on included data sources

*KRAS p.(G12D) c.35G>A, KRAS p.(G12V) c.35G>T, NRAS p.(Q61H) c.183A>T*

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
NRAS	p.(Q61H)	c.183A>T	COSM585	chr1:115256528	21.08%	NM_002524.5	missense	1988
KRAS	p.(G12V)	c.35G>T	COSM520	chr12:25398284	9.16%	NM_033360.4	missense	1997
KRAS	p.(G12D)	c.35G>A	COSM521	chr12:25398284	12.82%	NM_033360.4	missense	1997

### Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
CBFB-MYH11	CBFB-MYH11.C4M33	chr16:67100701 - chr16:15814908	498
CBFB-MYH11	CBFB-MYH11.C5M33	chr16:67116211 - chr16:15814908	19473

## Biomarker Descriptions

### CBFB (core-binding factor subunit beta)

**Background:** The CBFB gene encodes the core-binding factor subunit beta, a member of the PEBP2/CBF transcription factor family<sup>1</sup>. CBFB is capable of heterodimerization with the RUNX protein family (RUNX1, RUNX2, and RUNX3) which results in the formation of the core binding factor (CBF) complex, a transcription factor complex responsible for the regulation of many critical functions in hematopoiesis and osteogenesis<sup>2,3,4</sup>. Although possessing no DNA-binding activity, CBFB has been observed to enhance stability and transcriptional activity of RUNX proteins, thereby exhibiting a critical role in RUNX mediated transcriptional regulation<sup>3,4</sup>. In cancer, mutations in CBFB have been implicated in decreased protein stability and loss of function, supporting a tumor suppressor role for CBFB<sup>4</sup>.

**Alterations and prevalence:** Somatic mutations in CBFB are observed in 2% of diffuse large B-cell lymphoma, breast invasive carcinoma, and uterine corpus endometrial carcinoma<sup>5</sup>. Biallelic deletions in CBFB are found in 2% of ovarian serous cystadenocarcinoma, prostate adenocarcinoma, and breast invasive carcinoma<sup>5</sup>. Translocations including inv(16) and t(16;16) have been observed to be recurrent in de novo AML, occurring in 7-10% of patients, and have been associated with the AML M4 with bone marrow eosinophilia (M4Eo) subtype<sup>6</sup>. Translocations often result in CBFB-MYH11 fusion, which can exist as one of multiple transcripts, depending on the exons fused<sup>6</sup>.

## Biomarker Descriptions (continued)

**Potential relevance:** Currently, no therapies are approved for CBFB aberrations. Translocations, including inv(16) and t(16;16) which result in CBFB-MYH11 fusion, are diagnostic markers for acute myeloid leukemia and are associated with favorable prognosis<sup>7</sup>.

### KRAS (KRAS proto-oncogene, GTPase)

**Background:** The KRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes NRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>8,9,10</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. KRAS mutations are observed in up to 10-20% of uterine cancer, 30-35% of lung adenocarcinoma and colorectal cancer, and about 60% of pancreatic cancer<sup>5</sup>. The majority of KRAS mutations consist of point mutations occurring at G12, G13, and Q61<sup>5,11,12</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>13,14</sup>.

**Potential relevance:** The KRAS inhibitor, sotorasib<sup>15</sup>, is approved (2021) for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC). The FDA has granted breakthrough therapy designation (2021) to the small molecule inhibitor, adagrasib, for KRAS G12C positive in non-small cell lung cancer following prior systemic therapy<sup>16</sup>. The small molecular inhibitor, RO-5126766, was also granted breakthrough designation (2021) alone for KRAS G12V mutant non-small cell lung cancer or in combination with defactinib, for KRAS mutant endometrial carcinoma and KRAS G12V mutant non-small cell lung cancer<sup>17</sup>. The PLK1 inhibitor, onvansertib<sup>18</sup>, was granted fast track designation (2020) in combination with bevacizumab and FOLFIRI for second-line treatment of patients with KRAS-mutated metastatic colorectal cancer (mCRC). Additionally, the SHP2 inhibitor, BBP-398<sup>19</sup> was granted fast track designation (2022) in combination with sotorasib for previously treated patients with KRAS G12C-mutated metastatic NSCLC. The EGFR antagonists, cetuximab<sup>20</sup> and panitumumab<sup>21</sup>, are contraindicated for treatment of colorectal cancer patients with KRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>14</sup>. Additionally, KRAS mutations are associated with poor prognosis in NSCLC<sup>22</sup>.

### MYH11 (myosin heavy chain 11)

**Background:** MYH11 encodes myosin heavy-chain 11 which is a key contractile protein involved in smooth muscle movement, protein transport, and cell-cell interaction<sup>23</sup>. MYH11 belongs to the myosin family of motor proteins that functions to convert chemical energy into mechanical energy through ATP hydrolysis<sup>24,25</sup>. MYH11 consists of two pairs of light chains and two heavy chains produced from the MYH11 gene<sup>1</sup>. MYH11 is frequently translocated in acute myeloid leukemia (AML) where the C-terminal of MYH11 protein that contains a coiled-coil smooth muscle myosin heavy chain (SMMHC) fuses with the core binding factor  $\beta$  (CBF $\beta$ ) gene<sup>26,27</sup>. The CBFB-MYH11 fusion dimerizes with RUNX1, a key transcription factor in myelodysplastic malignancies, and inhibits RUNX1's function in hematopoiesis<sup>28</sup>.

**Alterations and prevalence:** Somatic missense, nonsense, and splice-site mutations of MYH11 are observed in 13% of melanoma, 10% of uterine, 6% of stomach as well as bladder cancers, and 2% of AML cases<sup>13</sup>. The inv(16)(p13.1;q22) or t(16;16)(p13.1;q22) translocation results in the CBFB-MYH11 fusion and is observed in over 5% of AML cases<sup>27</sup>.

**Potential relevance:** The inv(16)(p13.1;q22)/t(16;16)(p13.1;q22), that results in CBFB-MYH11 fusion, is a diagnostic entity for AML, as defined by the World Health Organization (WHO)<sup>7,29</sup>. The CBFB-MYH11 fusion is also associated with favorable risk stratification in AML<sup>7</sup>.

### NRAS (NRAS proto-oncogene, GTPase)

**Background:** The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>8,9,10</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers<sup>5,30</sup>. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61<sup>5,31</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>13,14</sup>.

**Potential relevance:** Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab<sup>20</sup> and panitumumab<sup>21</sup>, are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>14</sup>. The FDA has granted fast track designation to the pan-RAF inhibitor, KIN-2787<sup>32</sup>, for the treatment of NRAS mutation positive metastatic or unresectable melanoma. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome<sup>33</sup> as well as melanoma<sup>34</sup>. In a phase III clinical trial in patients with

## Biomarker Descriptions (continued)

advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively<sup>35</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### CBFB-MYH11 fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
gemtuzumab ozogamicin + cytarabine + daunorubicin	×	●	×	●	×
cytarabine + daunorubicin	×	●	×	×	×
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×

## Relevant Therapy Details

### Current NCCN Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### CBFB-MYH11 fusion

#### ● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 1

Population segment (Line of therapy):  
☒ (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**CBFB-MYH11 fusion (continued)****● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

**● gemtuzumab ozogamicin + cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

## CBFB-MYH11 fusion (continued)

### ● gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

### ● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

## Current ESMO Information

- ☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

ESMO information is current as of 2022-11-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### CBFB-MYH11 fusion

#### ☒ gemtuzumab ozogamicin + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- Core Binding Factor (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Prognostic Details

### Current NCCN Information

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### CBFB-MYH11 fusion

### Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

NCCN Recommendation category: 2A

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

### Current ESMO Information

ESMO information is current as of 2022-11-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

#### CBFB-MYH11 fusion

### Prognostic significance: ELN 2017: Favorable

Cancer type: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]



## Diagnostic Details

### Current NCCN Information

NCCN information is current as of 2022-11-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

#### CBFB-MYH11 fusion

##### Diagnostic significance: Acute Myeloid Leukemia

Variant class: inv(16) or t(16;16)

NCCN Recommendation category: 2A

##### Diagnostic notes:

- WHO 2016 classification defined as  $\geq 20\%$  blasts in blood or bone marrow, or in appropriate clinical setting with  $< 20\%$  blasts and recurrent cytogenetic abnormalities.

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]

##### Diagnostic significance: Acute Myeloid Leukemia

Variant class: inv(16)

NCCN Recommendation category: 2A

##### Diagnostic notes:

- Karyotypes t(8;21), inv16, and t(15;17) are considered to be Acute Myeloid Leukemia even if the marrow blast count is less than 20%

Reference: NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]

### Current ESMO Information

ESMO information is current as of 2022-11-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

#### CBFB-MYH11 fusion

##### Diagnostic significance: Acute Myeloid Leukemia

Variant class: MYH11-CBFB fusion [t(16;16)(p13;q22)]

##### Diagnostic notes:

- Core Binding Factor-Acute Myeloid Leukemia

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

## References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Link et al. Core binding factor at the crossroads: determining the fate of the HSC. *J Cell Physiol.* 2010 Jan;222(1):50-6. PMID: 19813271
3. Qin et al. Cbfb regulates bone development by stabilizing Runx family proteins. *J Bone Miner Res.* 2015 Apr;30(4):706-14. PMID: 25262822
4. Malik et al. The transcription factor CBFβ suppresses breast cancer through orchestrating translation and transcription. *Nat Commun.* 2019 May 6;10(1):2071. PMID: 31061501
5. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
6. Lesser et al. Tables of power for the F-test for comparing two exponential survival distributions. *J Chronic Dis.* 1981;34(11):533-44. PMID: 17287858
7. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2022]
8. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer.* 2011 Oct 13;11(11):761-74. PMID: 21993244
9. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
10. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
11. Román et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. *Mol Cancer.* 2018 Feb 19;17(1):33. doi: 10.1186/s12943-018-0789-x. PMID: 29455666
12. Dinu et al. Prognostic significance of KRAS gene mutations in colorectal cancer—preliminary study. *J Med Life.* 2014 Oct-Dec;7(4):581-7. PMID: 25713627
13. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
14. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
15. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/214665s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214665s000lbl.pdf)
16. <https://ir.mirati.com/press-releases/press-release-details/2021/Mirati-Therapeutics-Adagrasib-Receives-Breakthrough-Therapy-Designation-from-U.S.-Food-and-Drug-Administration-for-Patients-with-Advanced-Non-Small-Cell-Lung-Cancer-Harboring-the-KRAS-G12C-Mutation/default.aspx>
17. <https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-receives-breakthrough-therapy-designation-vs>
18. <https://cardiffoncology.investorroom.com/2020-05-28-Cardiff-Oncology-Announces-Fast-Track-Designation-Granted-by-the-FDA-to-Onvansertib-for-Second-Line-Treatment-of-KRAS-Mutated-Colorectal-Cancer>
19. <https://bridgebio.com/news/bridgebio-pharma-announces-first-lung-cancer-patient-dosed-in-phase-1-2-trial-and-us-fda-fast-track-designation-for-shp2-inhibitor-bbp-398-in-combination-with-amgens-lumakras-sotorasib/>
20. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)
21. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125147s210lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf)
22. Slebos et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *N. Engl. J. Med.* 1990 Aug 30;323(9):561-5. PMID: 2199829
23. Matsuoka et al. Human smooth muscle myosin heavy chain gene mapped to chromosomal region 16q12. *Am. J. Med. Genet.* 1993 Apr 1;46(1):61-7. PMID: 7684189
24. Kwartler et al. Overexpression of smooth muscle myosin heavy chain leads to activation of the unfolded protein response and autophagic turnover of thick filament-associated proteins in vascular smooth muscle cells. *J. Biol. Chem.* 2014 May 16;289(20):14075-88. PMID: 24711452
25. Halstead et al. An unusual transduction pathway in human tonic smooth muscle myosin. *Biophys. J.* 2007 Nov 15;93(10):3555-66. PMID: 17704147
26. Huang et al. Genetic approaches to identify pathological limitations in aortic smooth muscle contraction. *PLoS ONE.* 2018;13(3):e0193769. PMID: 29494672
27. Durst et al. The inv(16) fusion protein associates with corepressors via a smooth muscle myosin heavy-chain domain. *Mol. Cell. Biol.* 2003 Jan;23(2):607-19. PMID: 12509458

## References (continued)

28. Shigesada et al. Mechanism of leukemogenesis by the inv(16) chimeric gene CBFB/PEBP2B-MHY11. *Oncogene*. 2004 May 24;23(24):4297-307. PMID: 15156186
29. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016 May 19;127(20):2391-405. PMID: 27069254
30. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE*. 2011;6(7):e22769. PMID: 21829508
31. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res*. 2013 May 1;19(9):2584-91. PMID: 23515407
32. <https://investors.kinnate.com/news-releases/news-release-details/kinnate-biopharma-inc-receives-fast-track-designation-us-food>
33. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
34. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol*. 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
35. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol*. 2017 Apr;18(4):435-445. PMID: 28284557