



Sample Information

Patient Name: 陳明成  
Gender: Male  
ID No.: F123159119  
History No.: 27874482  
Age: 55

Ordering Doctor: DOC1686E 陳玟均  
Ordering REQ.: 0CMCSRS  
Signing in Date: 2023/06/16

Path No.: M112-00152  
MP No.: MY23033  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/06/13

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	12 Therapies Available
Relevant Therapy Summary	3	0 Clinical Trials
Relevant Therapy Details	4	

Relevant Myelodysplastic Syndrome Variants

Gene	Finding	Gene	Finding
ASXL1	None detected	NPM1	None detected
BCOR	None detected	NRAS	<b>NRAS p.(G12D) c.35G&gt;A</b>
CBL	None detected	NUP214	None detected
CREBBP	None detected	RUNX1	<b>RUNX1 p.(R107C) c.319C&gt;T</b>
ETV6	None detected	SF3B1	None detected
EZH2	None detected	SRSF2	None detected
FLT3	None detected	STAG2	None detected
GATA2	None detected	TP53	None detected
IDH2	None detected	U2AF1	None detected
KIT	None detected	WT1	None detected
KMT2A	None detected	ZRSR2	None detected
MECOM	None detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	<i>RUNX1</i> p.(R107C) c.319C>T RUNX family transcription factor 1 Allele Frequency: 22.90%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine liposomal cytarabine-daunorubicin CPX-351 venetoclax + chemotherapy	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

*NRAS* p.(G12D) c.35G>A

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
NRAS	p.(G12D)	c.35G>A	COSM564	chr1:115258747	20.94%	NM_002524.5	missense	1996
RUNX1	p.(R107C)	c.319C>T	COSM24736	chr21:36259172	22.90%	NM_001754.4	missense	2000
TET2	p.(R814C)	c.2440C>T	.	chr4:106157539	48.01%	NM_001127208.2	missense	1131
EZH2	p.(K498*)	c.1492A>T	.	chr7:148513789	43.57%	NM_004456.5	nonsense	1999
ETV6	p.(I370T)	c.1109T>C	.	chr12:12037478	23.40%	NM_001987.5	missense	2000
NF1	p.(K2456=)	c.7368A>G	.	chr17:29677247	27.16%	NM_001042492.3	synonymous	1999
SRSF2	p.(R61=)	c.183C>T	.	chr17:74733060	22.70%	NM_003016.4	synonymous	1996

Biomarker Descriptions

NRAS (NRAS proto-oncogene, GTPase)

**Background:** The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>1,2,3</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers<sup>4,5</sup>. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61<sup>4,6</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>7,8</sup>.

**Potential relevance:** Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab<sup>9</sup> and panitumumab<sup>10</sup>, are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons

## Biomarker Descriptions (continued)

59 and 61), and exon 4 (codons 117 and 146)<sup>8</sup>. The FDA has granted fast track designation to the pan-RAF inhibitor, KIN-2787<sup>11</sup>, for the treatment of NRAS mutation positive metastatic or unresectable melanoma. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome<sup>12</sup> as well as melanoma<sup>13</sup>. In a phase III clinical trial in patients with advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively<sup>14</sup>.

### RUNX1 (RUNX family transcription factor 1)

**Background:** The RUNX1 gene encodes the runt-related transcription factor (RUNX) 1, part of the RUNX family of transcription factors which also includes RUNX2 and RUNX3<sup>15</sup>. All RUNX proteins share several conserved regions with similar functionality including a highly conserved N-terminal 'runt' domain responsible for binding DNA, a C-terminal region composed of an activation domain, inhibitory domain, protein interacting motifs, and a nuclear matrix targeting signal<sup>16</sup>. Each of these proteins are capable of interacting with core binding factor beta (CBFβ) to form the core binding factor (CBF) complex. Consequently, RUNX1, RUNX2, and RUNX3 are collectively known as core binding factor alpha (CBFα) since they can each function as the alpha subunit of CBF. Specifically, CBFβ binds to the 'runt' domain of RUNX1 leading to RUNX1 stabilization and increased affinity of the CBF complex for promoters involved in hematopoietic differentiation and cell cycle regulation<sup>17,18</sup>. RUNX1 is frequently mutated in various hematological malignancies<sup>18</sup>. Germline mutations in RUNX1 result in a rare autosomal dominant condition known as familial platelet disorder, with predisposition to acute myeloid leukemia (FPD/AML)<sup>19,20</sup>. Somatic mutations and chromosomal translocations in RUNX1 are often observed in myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), and chronic myelomonocytic leukemia (CMML)<sup>18</sup>.

**Alterations and prevalence:** RUNX1 is frequently rearranged in hematological malignancies with over 50 different observed translocations<sup>21</sup>. The most recurrent translocation, t(12;21)(q34;q11), results in ETV6-RUNX1 fusion and is observed in 20-25% of childhood ALL<sup>22,23,24</sup>. This translocation is also observed in adult ALL at a lower frequency (2%)<sup>23,24</sup>. Another recurrent translocation, t(8;21)(q22;q22), results in RUNX1-RUNX1T1 fusion and is observed in 5-10% of AML<sup>25</sup>. The RUNX1-RUNX1T1 fusion, consists of the RHD domain of RUNX1 and the majority of RUNX1T1, which promotes oncogenesis by altering transcriptional regulation of RUNX1 target genes<sup>18,25</sup>. Somatic mutations in RUNX1 include missense, nonsense, and frameshift mutations resulting in loss of function or dominant negative effects<sup>18</sup>. RUNX1 mutations are reported in approximately 10% of de novo AML as well as 10-15% of MDS<sup>4,12,18,26</sup>.

**Potential relevance:** AML with RUNX1-RUNX1T1 fusions is considered a distinct molecular subtype by the World Health Organization (WHO)<sup>26,27</sup>. Translocations involving RUNX1, specifically t(8;21)(q22;q22)/RUNX1-RUNX1T1 in AML and t(12;21)(q34;q11)/ETV6-RUNX1 in ALL, are associated with favorable risk<sup>26,28</sup>. On the other hand, mutations in RUNX1 confer poor prognosis in AML, MDS, and systemic mastocytosis (SM)<sup>12,26,29</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### RUNX1 p.(R107C) c.319C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✗	○	✗	✗	✗
azacitidine	✗	○	✗	✗	✗
cytarabine	✗	○	✗	✗	✗
cytarabine + daunorubicin	✗	○	✗	✗	✗
cytarabine + daunorubicin + etoposide	✗	○	✗	✗	✗
cytarabine + etoposide + idarubicin	✗	○	✗	✗	✗
cytarabine + fludarabine + idarubicin + filgrastim	✗	○	✗	✗	✗
cytarabine + idarubicin	✗	○	✗	✗	✗

## Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### RUNX1 p.(R107C) c.319C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cytarabine + mitoxantrone	×	○	×	×	×
decitabine	×	○	×	×	×
liposomal cytarabine-daunorubicin CPX-351	×	○	×	×	×
venetoclax + azacitidine	×	○	×	×	×
venetoclax + cytarabine	×	○	×	×	×
venetoclax + decitabine	×	○	×	×	×

## Relevant Therapy Details

### Current NCCN Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-03-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/international\\_adaptations.aspx](http://www.nccn.org/global/international_adaptations.aspx).

### RUNX1 p.(R107C) c.319C>T

#### ☐ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

#### ☐ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**RUNX1 p.(R107C) c.319C>T (continued)****○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**RUNX1 p.(R107C) c.319C>T (continued)****○ cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ liposomal cytarabine-daunorubicin CPX-351**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**RUNX1 p.(R107C) c.319C>T (continued)****○ venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**○ cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]

**RUNX1 p.(R107C) c.319C>T (continued)****○ cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]



## References

1. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer*. 2011 Oct 13;11(11):761-74. PMID: 21993244
2. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
3. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
4. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
5. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE*. 2011;6(7):e22769. PMID: 21829508
6. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res.* 2013 May 1;19(9):2584-91. PMID: 23515407
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
8. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
9. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)
10. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125147s210lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf)
11. <https://investors.kinnate.com/news-releases/news-release-details/kinnate-biopharma-inc-receives-fast-track-designation-us-food>
12. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
13. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol.* 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
14. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2017 Apr;18(4):435-445. PMID: 28284557
15. de et al. Runx transcription factors in the development and function of the definitive hematopoietic system. *Blood.* 2017 Apr 13;129(15):2061-2069. PMID: 28179276
16. Chuang et al. RUNX family: Regulation and diversification of roles through interacting proteins. *Int. J. Cancer.* 2013 Mar 15;132(6):1260-71. PMID: 23180629
17. Jung et al. Prognostic factor analysis in core-binding factor-positive acute myeloid leukemia. *Anticancer Res.* 2014 Feb;34(2):1037-45. PMID: 24511052
18. Sood et al. Role of RUNX1 in hematological malignancies. *Blood.* 2017 Apr 13;129(15):2070-2082. PMID: 28179279
19. Béri-Dexheimer et al. Clinical phenotype of germline RUNX1 haploinsufficiency: from point mutations to large genomic deletions. *Eur. J. Hum. Genet.* 2008 Aug;16(8):1014-8. PMID: 18478040
20. Hayashi et al. Myeloid neoplasms with germ line RUNX1 mutation. *Int. J. Hematol.* 2017 Aug;106(2):183-188. PMID: 28534116
21. De et al. RUNX1 translocations and fusion genes in malignant hemopathies. *Future Oncol.* 2011 Jan;7(1):77-91. PMID: 21174539
22. De et al. ETV6 fusion genes in hematological malignancies: a review. *Leuk. Res.* 2012 Aug;36(8):945-61. PMID: 22578774
23. Pui et al. Acute lymphoblastic leukemia. *N. Engl. J. Med.* 2004 Apr 8;350(15):1535-48. PMID: 15071128
24. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
25. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D920-4. PMID: 23161685
26. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2022]
27. Arber et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood.* 2016 May 19;127(20):2391-405. PMID: 27069254
28. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 1.2022]
29. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]