

**ABOUT THE TEST** FoundationOne® Heme is a comprehensive genomic profiling test designed to identify genomic alterations within hundreds of cancer-related genes in hematologic malignancies and sarcomas.

<b>PATIENT</b>	<b>DISEASE</b> Soft tissue perivascular epithelioid cell tumor (PEComa)	<b>PHYSICIAN</b>	<b>ORDERING PHYSICIAN</b> Yeh, Yi-Chen	<b>SPECIMEN</b>	<b>SPECIMEN SITE</b> Vagina
	<b>NAME</b> Wang, Fu Ting		<b>MEDICAL FACILITY</b> Taipei Veterans General Hospital		<b>SPECIMEN ID</b> S112-66474 N (PF23038)
	<b>DATE OF BIRTH</b> 18 April 1996		<b>ADDITIONAL RECIPIENT</b> None		<b>SPECIMEN TYPE</b> Slide Deck
	<b>SEX</b> Female		<b>MEDICAL FACILITY ID</b> 205872		<b>DATE OF COLLECTION</b> 07 April 2023
	<b>MEDICAL RECORD #</b> 49390887		<b>PATHOLOGIST</b> Not Provided		<b>SPECIMEN RECEIVED</b> 17 April 2023

## Biomarker Findings

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 0 Muts/Mb

## Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

**TFE3** NONO-TFE3 fusion

## Report Highlights

- Variants with **diagnostic implications** that may indicate a specific cancer type: **TFE3** NONO-TFE3 fusion (p. [3](#))
- Targeted therapies with potential clinical benefit **approved in another tumor type**: **Sunitinib** (p. [4](#))
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. [5](#))

### BIOMARKER FINDINGS

**Microsatellite status** - MS-Stable

**Tumor Mutational Burden** - 0 Muts/Mb

### GENOMIC FINDINGS

**TFE3** - NONO-TFE3 fusion

**10 Trials** see p. [5](#)

### THERAPY AND CLINICAL TRIAL IMPLICATIONS

**No therapies or clinical trials.** See Biomarker Findings section

**No therapies or clinical trials.** See Biomarker Findings section

#### THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

none

#### THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

**Sunitinib**

**NOTE** Genomic alterations detected may be associated with activity of certain FDA-approved drugs; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type.

Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

**Sample Preparation:** 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
**Sample Analysis:** 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
**Post-Sequencing Analysis:** 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**BIOMARKER FINDINGS**
**BIOMARKER**

## Microsatellite status

**RESULT**  
MS-Stable

**POTENTIAL TREATMENT STRATEGIES**
**— Targeted Therapies —**

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors<sup>1-3</sup>, including approved therapies nivolumab and pembrolizumab<sup>4</sup>. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%,  $p=0.001$ )<sup>5</sup>.

**FREQUENCY & PROGNOSIS**

In a computational analysis of paired tumor and normal sarcomas in the TCGA dataset, of which 25% were liposarcomas, only 0.8% (2/255) of samples were MSI-high (MSI-H)<sup>6</sup>. However, reports of MSI in sarcomas in the literature are conflicting and varied due to substantial heterogeneity, lack of consensus on the markers and methods used for MSI assessment, and small sample size in most studies<sup>7</sup>. In these smaller studies of soft tissue sarcoma, reports of MSI at any level have been rare, with the highest incidences between 11% (2/18) to 25% (10/40) of cases<sup>8-13</sup>. In one study, MSI was reported to occur more frequently in high-grade soft tissue sarcomas compared with lower grade<sup>14</sup>. However, published data investigating the prognostic implications of MSI in sarcoma are limited (PubMed, Jan 2023).

**FINDING SUMMARY**

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor<sup>15</sup>. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2<sup>15-17</sup>. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers<sup>18-20</sup>. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins<sup>15,17,19-20</sup>.

**BIOMARKER**

## Tumor Mutational Burden

**RESULT**  
0 Muts/Mb

**POTENTIAL TREATMENT STRATEGIES**
**— Targeted Therapies —**

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1<sup>21-23</sup>, anti-PD-1 therapies<sup>21-24</sup>, and combination nivolumab and ipilimumab<sup>25-30</sup>. In multiple pan-tumor studies, increased tissue tumor mutational burden (TMB) was associated with sensitivity to immune checkpoint inhibitors<sup>21-24,31-35</sup>. In the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors, significant improvement in ORR was observed for patients with TMB  $\geq 10$  Muts/Mb (as measured by this assay) compared with those with TMB  $< 10$  Muts/Mb in a large cohort that included multiple tumor types<sup>31</sup>; similar findings were observed in the KEYNOTE 028 and 012 trials<sup>24</sup>. At the same TMB cutpoint, retrospective analysis of patients with solid tumors treated with any checkpoint inhibitor identified that tissue TMB scores  $\geq 10$  Muts/Mb were associated with

prolonged time to treatment failure compared with scores  $< 10$  Muts/Mb (HR=0.68)<sup>35</sup>. For patients with solid tumors treated with nivolumab plus ipilimumab in the CheckMate 848 trial, improved responses were observed in patients with a tissue TMB  $\geq 10$  Muts/Mb independent of blood TMB at any cutpoint in matched samples<sup>36</sup>. However, support for higher TMB thresholds and efficacy was observed in the prospective Phase 2 MyPathway trial of atezolizumab for patients with pan-solid tumors, where improved ORR and DCR was seen in patients with TMB  $\geq 16$  Muts/Mb than those with TMB  $\geq 10$  and  $< 16$  Muts/Mb<sup>34</sup>. Similarly, analyses across several solid tumor types reported that patients with higher TMB (defined as  $\geq 16-20$  Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy compared with patients with higher TMB treated with chemotherapy<sup>37</sup> or those with lower TMB treated with PD-1 or PD-L1-targeting agents<sup>22</sup>.

**FREQUENCY & PROGNOSIS**

Perivascular epithelioid cell tumors harbor a median TMB of 1.6 mutations per megabase (mut/Mb) in a study of 31 cases<sup>38</sup>. Soft tissue sarcomas harbor a median tumor mutational burden (TMB) of 2.4-2.5 mutations per megabase (mut/Mb), with angiosarcoma (up to 15%) and malignant peripheral nerve sheath tumor (up to 11%) having the highest percentage of cases with high TMB<sup>39-40</sup>. In a study, 3.9% of soft tissue sarcoma samples analyzed harbor TMB  $\geq 10$  mut/Mb<sup>39</sup>; in addition, increased

mutational burden has been reported in undifferentiated pleomorphic sarcomas as compared with Ewing sarcomas or rhabdomyosarcomas<sup>41-43</sup>. Published data investigating the prognostic implications of tissue TMB in sarcoma are conflicting (PubMed, Feb 2023). High tissue TMB was associated with improved PFS and metastasis-free survival in a study of undifferentiated sarcomas<sup>44</sup>, but with reduced survival in a study of patients with rhabdomyosarcoma<sup>45</sup>.

**FINDING SUMMARY**

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma<sup>46-47</sup> and cigarette smoke in lung cancer<sup>48-49</sup>, treatment with temozolomide-based chemotherapy in glioma<sup>50-51</sup>, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes<sup>52-56</sup>, and microsatellite instability (MSI)<sup>52,55-56</sup>. This sample harbors a TMB level associated with lower rates of clinical benefit from treatment with PD-1- or PD-L1-targeting immune checkpoint inhibitors compared with patients with tumors harboring higher TMB levels, based on several studies in multiple solid tumor types<sup>22-23,31</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**GENOMIC FINDINGS**
**GENE**
**TFE3**
**ALTERATION**

NONO-TFE3 fusion

**POTENTIAL TREATMENT STRATEGIES**
**— Targeted Therapies —**

Clinical benefit in certain TFE3-rearranged solid tumors has been reported in response to anti-angiogenic agents such as sunitinib<sup>57-64</sup>, sorafenib<sup>57-58</sup>, and cediranib<sup>65-66</sup>; to mTOR inhibitors<sup>58,67-72</sup>; and to MET inhibitors<sup>73-74</sup>. Clinical studies have shown high response rates to treatment with sunitinib for patients with TFE3 fusion-associated cancers, including alveolar soft part sarcoma (ASPS)<sup>62-64</sup> and translocation renal cell carcinoma (tRCC or Xp11.2 RCC)<sup>57-61</sup>. In addition, TFE3 fusions result in increased MET expression<sup>73,75-76</sup> and clinical and preclinical sensitivity to MET inhibitors<sup>73-75</sup>; a Phase 2 study of crizotinib to treat advanced or metastatic ASPS reported 1 PR and 35 SDs out of 40 MET-expressing cases<sup>73</sup>. Two patients with Xp11.2 RCC experienced clinical benefit from immune

checkpoint inhibitors, with ongoing time on treatments of 37 and 15.7 months in a retrospective analysis (n=8)<sup>67</sup>.

**FREQUENCY & PROGNOSIS**

TFE3 rearrangements have been observed in a variety of types of perivascular epithelioid cell tumors (PEComa), including those of cervical, renal, and uterine origin<sup>77-79</sup>. Due to the low frequency of PEComa cases, the incidence of TFE3 rearrangement or mutation in PEComa is unclear. However, one study tested 29 PEComa patients for TFE3 alteration and identified 4 (14%) patients with TFE3 rearrangements and concomitant increased TFE3 expression as well as an additional 4 (14%) patients with increased TFE3 expression<sup>78</sup>. Another study reported TFE3 rearrangement in 23% (9/38) of PEComa patients and identified TFE3 rearrangements as being mutually exclusive from TSC2 mutation, a frequent mutation found in PEComa<sup>80</sup>. TFE3 rearrangements are generally rare in human cancer but characterize Xp11 translocation cancers, which include renal cell carcinoma, perivascular epithelioid cell tumor (PEComa), epithelioid hemangioendothelioma, alveolar soft part sarcoma, and melanotic Xp11 neoplasm, with a variety of TFE3 fusion partners

reported in these diseases<sup>81</sup>. An analysis of 3 PEComa patients with TFE3 rearrangement and increased expression reported that 2 of the 3 patients exhibited histological malignancy and aggressive growth, while the other patient exhibited a benign course<sup>82</sup>.

**FINDING SUMMARY**

TFE3 encodes the transcription factor for immunoglobulin heavy chain enhancer 3 protein. Multiple fusions with TFE3 as a 3' partner have been reported in cancer and characterized as activating<sup>76,83-87</sup>. NONO-TFE3 fusions have been reported in renal cell carcinomas<sup>85,88-90</sup> and neoplasms with melanocytic differentiation including melanocytic perivascular epithelioid cell neoplasms (PEComas)<sup>90-92</sup> and are likely activating<sup>88,93-94</sup>, although one in vitro study found NONO-TFE3 was unable to activate transcription<sup>87</sup>.

**POTENTIAL DIAGNOSTIC IMPLICATIONS**

TFE3 rearrangements are characteristic of a group of neoplasms known as Xp11 translocation cancers, which include renal cell carcinoma, PEComa, epithelioid hemangioendothelioma, alveolar soft part sarcoma, and melanotic Xp11 neoplasm<sup>81,95-101</sup>.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**THERAPIES WITH CLINICAL BENEFIT**
**IN OTHER TUMOR TYPE**

## Sunitinib

*Assay findings association*

### TFE3

NONO-TFE3 fusion

### AREAS OF THERAPEUTIC USE

Sunitinib is a small-molecule tyrosine kinase inhibitor that targets PDGFRs, VEGFRs, KIT, FLT3, CSF-1R, and RET. It is FDA approved for the treatment of advanced or metastatic pancreatic neuroendocrine tumors, gastrointestinal stromal tumors (GISTs) in patients who have progressed on or are intolerant to imatinib, and advanced renal cell carcinoma (RCC) as well as for the adjuvant treatment of patients at high risk of recurrent RCC after nephrectomy. Please see the drug label for full prescribing information.

### GENE ASSOCIATION

TFE3 fusions may confer sensitivity to sunitinib, based on

both clinical<sup>57-64</sup> and preclinical<sup>76,102-104</sup> evidence.

### SUPPORTING DATA

Clinical data on the efficacy of sunitinib for the treatment of PEComa and angiomyolipoma are limited (PubMed, Apr 2023). A patient with malignant PEComa of the kidney treated with sunitinib for 4 weeks followed by resection exhibited a durable response before progressing<sup>105</sup>. A patient with epithelioid angiomyolipoma (EAML) of kidney treated with sunitinib had a short response for 2 months<sup>106</sup>. Another patient with kidney EAML who was treated with sunitinib for a short time and then switched to everolimus experienced favorable response for 30 months<sup>107</sup>.

**NOTE** Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

**ORDERED TEST #** ORD-1611555-01

**CLINICAL TRIALS**

**NOTE** Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://clinicaltrials.gov). Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

**GENE**  
**TFE3**
**ALTERATION**  
NONO-TFE3 fusion

**RATIONALE**  
TFE3 fusions may activate angiogenic pathways and increase expression of the MET oncogene and may therefore confer sensitivity to MET inhibitors

or to tyrosine kinase inhibitors such as sunitinib, sorafenib, or cediranib. TFE3 fusions may also predict sensitivity to mTOR inhibitors.

**NCT03175224**
**PHASE 1/2**

CBT-101 Study for Advanced Solid Tumors and c-Met Dysregulation

**TARGETS**  
MET

**LOCATIONS:** Taipei City (Taiwan), Taipei (Taiwan), New Taipei City (Taiwan), Taoyuan City (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Singapore (Singapore), Nedlands (Australia), North Adelaide (Australia), Bedford Park (Australia)

**NCT05439993**
**PHASE 1/2**

Tepotinib Plus Paclitaxel in MET Amplified or MET Exon 14 Altered Gastric and GEJ Carcinoma

**TARGETS**  
MET

**LOCATIONS:** Gyeonggi-do (Korea, Republic of)

**NCT03784014**
**PHASE 3**

MOLECULAR PROFILING OF ADVANCED SOFT-TISSUE SARCOMAS

**TARGETS**  
ABL, KIT, ROS1, ALK, MET, ERBB2, EGFR, BRAF, MEK, PARP, PD-L1, CDK4, CDK6

**LOCATIONS:** Strasbourg (France), Dijon (France), Paris (France), Villejuif (France), Lyon (France), Clermont-Ferrand (France), Marseille (France), Saint-Herblain (France), Bordeaux (France)

**NCT04817956**
**PHASE 2**

Improving Public Cancer Care by Implementing Precision Medicine in Norway

**TARGETS**  
PD-L1, VEGFA, ERBB2, ALK, RET, PARP, SMO, TRKB, TRKC, ROS1, TRKA, MEK, BRAF, PI3K-alpha, FGFR1, FGFR2, FGFR3, MET, KIT, ABL

**LOCATIONS:** Tromsø (Norway), Bodø (Norway), Hamar (Norway), Oslo (Norway), Fredrikstad (Norway), Drammen (Norway), Trondheim (Norway), Skien (Norway), Førde (Norway), Bergen (Norway)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**CLINICAL TRIALS**
**NCT04220229**
**PHASE 1/2**

Cabozantinib With Radiation Therapy for the Treatment of Sarcomas of the Extremities

**TARGETS**  
MET, ROS1, RET, VEGFRs

**LOCATIONS:** Washington

**NCT04200443**
**PHASE 2**

Cabozantinib and Temozolomide for the Treatment of Unresectable or Metastatic Leiomyosarcoma or Other Soft Tissue Sarcoma

**TARGETS**  
MET, ROS1, RET, VEGFRs

**LOCATIONS:** California, Iowa, Wisconsin, Illinois, Missouri

**NCT05135975**
**PHASE 2**

A Study of Cabozantinib as a Maintenance Agent to Prevent Progression or Recurrence in High-Risk Pediatric Solid Tumors

**TARGETS**  
MET, ROS1, RET, VEGFRs

**LOCATIONS:** Ohio

**NCT05468359**
**PHASE 1/2**

Safety and Efficacy of Cyclophosphamide, Sorafenib, Bevacizumab, and Atezolizumab in Pediatric Solid Tumor Patients

**TARGETS**  
RET, KIT, RAFs, FLT3, VEGFRs, VEGFA, PD-L1

**LOCATIONS:** Tennessee

**NCT02484404**
**PHASE 1/2**

Phase I/II Study of the Anti-Programmed Death Ligand-1 Antibody MEDI4736 in Combination With Olaparib and/or Cediranib for Advanced Solid Tumors and Advanced or Recurrent Ovarian, Triple Negative Breast, Lung, Prostate and Colorectal Cancers

**TARGETS**  
PARP, VEGFRs, PDGFRA, PDGFRB, KIT, PD-L1

**LOCATIONS:** Maryland

**NCT05038839**
**PHASE 1**

Cabozantinib and Pamiparib for the Treatment of Advanced of Refractory Solid Tumors

**TARGETS**  
MET, ROS1, RET, VEGFRs, PARP

**LOCATIONS:** Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**APPENDIX**
**Variants of Unknown Significance**

**NOTE** One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future. Please note that some VUS rearrangements between targeted genes and unknown fusion partners or intergenic regions detected by RNA sequencing may not be reported.

**APC**

NM\_000038.4: c.3374T>C  
(p.V1125A)  
chr5:112174665

**ASXL1**

NM\_015338.5: c.342T>G  
(p.N114K)  
chr20:31016020

**FLT1**

NM\_002019.4: c.2434C>T  
(p.R812W)  
chr13:28913359

**MAP3K14**

NM\_003954.4: c.295C>T  
(p.R99C)  
chr17:43366633

**NUP98**

NM\_016320.4: c.4891A>C  
(p.K1631Q)  
chr11:3704457

**PRKDC**

NM\_006904.6: c.9844C>A  
(p.R3282S)  
chr8:48715945

**SOCS1**

NM\_003745.1: c.44C>T  
(p.T15I)  
chr16:11349292

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531



ORDERED TEST # ORD-1611555-01

**APPENDIX**
**Genes Assayed in FoundationOne®Heme**

FoundationOne Heme is designed to include genes known to be somatically altered in human hematologic malignancies and sarcomas that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay utilizes DNA sequencing to interrogate 406 genes as well as selected introns of 31 genes involved in rearrangements, in addition to RNA sequencing of 265 genes. The assay will be updated periodically to reflect new knowledge about cancer biology.

**HEMATOLOGICAL MALIGNANCY DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS**

ABL1	ACTB	ADGRA2 (GPR124)	AKT1	AKT2	AKT3	ALK	AMER1 (FAM123B or WTX)
APC	APH1A	AR	ARAF	ARFRP1	ARHGAP26 (GRAF)	ARID1A	ARID2
ASMTL	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1
B2M	BAP1	BARD1	BCL10	BCL11B	BCL2	BCL2L2	BCL6
BCOR	BCORL1	BIRC3	BLM	BRAF	BRCA1	BRCA2	BRD4
BRSK1	BTG2	BTK	BTLA	CAD	CALR*	CARD11	CBFB
CCN6 (WISP3)	CCND1	CCND2	CCND3	CCNE1	CCT6B	CD22	CD274 (PD-L1)
CD58	CD70	CD79A	CD79B	CDC73	CDH1	CDK12	CDK4
CDK8	CDKN1B	CDKN2A	CDKN2B	CDKN2C	CEBPA	CHD2	CHEK1
CIC	CIITA	CKS1B	CPS1	CREBBP	CRKL	CRLF2	CSF1R
CTCF	CTNNA1	CTNNB1	CUX1	CXCR4	DAXX	DDR2	DDX3X
DNMT3A	DOT1L	DTX1	DUSP2	DUSP9	EBF1	ECT2L	EED
ELP2	EMSY (C11orf30)	EP300	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2
ERBB4	ERG	ESR1	ETS1	ETV6	EXOSC6	EZH2	FAF1
FANCC	FANCD2	FANCE	FANCF	FANCG	FANCL	FAS (TNFRSF6)	FBXO11
FBXW7	FGF10	FGF14	FGF19	FGF23	FGF3	FGF4	FGF6
FGFR2	FGFR3	FGFR4	FHIT	FLCN	FLT1	FLT3	FLT4
FOXL2	FOXO1	FOXO3	FOXP1	FRS2	GADD45B	GATA1	GATA2
GID4 (C17orf39)	GNA11	GNA12	GNA13	GNAQ	GNAS	GRIN2A	GSK3B
HDAC1	HDAC4	HDAC7	HGF	H1-2 (HIST1H1C)		H1-3 (HIST1H1D)	
H1-4 (HIST1H1E)		H2AC6 (HIST1H2AC)		H2AC11 (HIST1H2AG)		H2AC16 (HIST1H2AL)	
H2AC17 (HIST1H2AM)		H2BC4 (HIST1H2BC)		H2BC11 (HIST1H2BJ)		H2BC12 (HIST1H2BK)	
H2BC17 (HIST1H2BO)		H3C2 (HIST1H3B)		HNF1A	HRAS	HSP90AA1	ICK
IDH1	IDH2	IGF1R	IKBKE	IKZF1	IKZF2	IKZF3	IL7R
INPP4B	INPP5D (SHIP)	IRF1	IRF4	IRF8	IRS2	JAK1	JAK2
JARID2	JUN	KAT6A (MYST3)	KDM2B	KDM4C	KDM5A	KDM5C	KDM6A
KEAP1	KIT	KLHL6	KMT2A (MLL)	KMT2C (MLL3)	KMT2D (MLL2)	KRAS	LEF1
LRRK2	MAF	MAFB	MAGED1	MALT1	MAP2K1	MAP2K2	MAP2K4
MAP3K14	MAP3K6	MAP3K7	MAPK1	MCL1	MDM2	MDM4	MED12
MEF2C	MEN1	MET	MIB1	MITF	MKI67	MLH1	MPL
MSH2	MSH3	MSH6	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN
MYO18A	NCOR2	NCSTN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1
NOTCH1	NOTCH2	NPM1	NRAS	NSD2 (WHSC1 or MMSET)		NT5C2	NTRK1
NTRK3	NUP93	NUP98	P2RY8	PAG1	PAK3	PALB2	PASK
PBRM1	PC	PCBP1	PCLO	PDCD1	PDCD11	PDCD1LG2 (PD-L2)	
PDGFRB	PDK1	PHF6	PIK3CA	PIK3CG	PIK3R1	PIK3R2	PIM1
POT1	PPP2R1A	PRDM1	PRKAR1A	PRKDC	PRSS8	PTCH1	PTEN
PTPN2	PTPN6 (SHP-1)	PTPRO	RAD21	RAD50	RAD51	RAF1	RARA
RB1	RELN	RET	RHOA	RICTOR	RNF43	ROS1	RPTOR
S1PR2	SDHA	SDHB	SDHC	SDHD	SERP2	SETBP1	SETD2
SGK1	SMAD2	SMAD4	SMARCA1	SMARCA4	SMARCB1	SMC1A	SMC3
SOC3	SOC3	SOC3	SOX10	SOX2	SPEN	SPOP	SRC
STAG2	STAT3	STAT4	STAT5A	STAT5B	STAT6	STK11	SUFU
TAF1	TBL1XR1	TCF3 (E2A)	TCL1A (TCL1)	TENT5C (FAM46C)	TET2	TGFBR2	TLL2
TMSB4XP8 (TMSL3)		TNFAIP3	TNFRSF11A	TNFRSF14	TNFRSF17	TOP1	TP53
TRAF2	TRAF3	TRAF5	TSC1	TSC2	TSHR	TUSC3	TYK2
							U2AF1

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.



ORDERED TEST # ORD-1611555-01

**APPENDIX**
**Genes Assayed in FoundationOne®Heme**

U2AF2	VHL	WDR90	WT1	XBP1	XPO1	YY1AP1	ZMYM3	ZNF217
ZNF24 (ZSCAN3)	ZNF703	ZRSR2						

\*Note: the assay was updated on 11/8/2016 to include the detection of alterations in CALR

**HEMATOLOGICAL MALIGNANCY DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS**

ALK	BCL2	BCL6	BCR	BRAF	CCND1	CRLF2	EGFR	EPOR
ETV1	ETV4	ETV5	ETV6	EWSR1	FGFR2	IGH	IGK	IGL
JAK1	JAK2	KMT2A (MLL)	MYC	NTRK1	PDGFRA	PDGFRB	RAF1	RARA
RET	ROS1	TMPRSS2	TRG					

**HEMATOLOGICAL MALIGNANCY RNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS\***

ABI1	ABL1	ABL2	ACSL6	AFDN (MLLT4 or AF6)	AFF1	AFF4	ALK
ARHGAP26 (GRAF)		ARHGEF12	ARID1A	ARNT	ASXL1	ATF1	ATIC
BCL10	BCL11A	BCL11B	BCL2	BCL3	BCL6	BCL7A	BCOR
BCR	BIRC3	BRAF	BTG1	CAMTA1	CARS1 (CARS)	CBFA2T3	CBL
CCND1	CCND2	CCND3	CD274 (PD-L1)	CDK6	CDX2	CEP43 (FGFR10P)	CHN1
CIC	CIITA	CLP1	CLTC	CLTCL1	CNTRL (CEP110)	COL1A1	CREB3L1
CREBBP	CRLF2	CSF1	CTNNB1	DDIT3	DDX10	DDX6	DUSP22
EGFR	EIF4A2	ELF4	ELL	ELN	EML4	EP300	EPOR
ERBB2	ERG	ETS1	ETV1	ETV4	ETV5	ETV6	EWSR1
FCRL4	FEV	FGFR1	FGFR2	FGFR3	FLI1	FNBP1	FOXO3
FOXO4	FOXP1	FSTL3	FUS	GAS7	GLI1	GMPS	GPHN
HERPUD1	HEY1	HIP1	HLF	HMGAI	HMGAI2	HOXA11	HOXA13
HOXA9	HOXC11	HOXC13	HOXD11	HOXD13	HSP90AA1	HSP90AB1	IGH
IGL	IKZF1	IL21R	IL3	IRF4	ITK	JAK1	JAK2
JAZF1	KAT6A (MYST3)	KDSR	KIF5B	KMT2A (MLL)	LASP1	LCP1	LMO1
LPP	LYL1	MAF	MAFB	MALT1	MDS2	MECOM	MLF1
MLLT10 (AF10)	MLLT3	MLLT6	MN1	MXN1	MRTFA (MKL1)	MSI2	MSN
MYB	MYC	MYH11	MYH9	NACA	NBEAP1 (BCL8)	NCOA2	NDRG1
NF2	NFKB2	NIN	NOTCH1	NPM1	NR4A3	NSD1	NSD2 (WHSC1 or MMSET)
NSD3 (WHSC1L1)	NTRK1	NTRK2	NTRK3	NUMA1	NUP214	NUP98	NUTM2A
P2RY8	PAFAH1B2	PAX3	PAX5	PAX7	PBX1	PCM1	PCSK7
PDE4DIP	PDGFB	PDGFRA	PDGFRB	PER1	PHF1	PICALM	PIM1
PML	POU2AF1	PPP1CB	PRDM1	PRDM16	PRRX1	PSIP1	PTCH1
RABEP1	RAF1	RALGDS	RAP1GDS1	RARA	RBM15	RET	RHOH
RNF217-AS1 (STL)		ROS1	RPL22	RPN1	RUNX1	RUNX1T1 (ETO)	RUNX2
SEPTIN5 (SEPT5)	SEPTIN6 (SEPT6)	SEPTIN9 (SEPT9)	SET	SH3GL1	SLC1A2	SNX29 (RUNDC2A)	SRSF3
SS18	SSX1	SSX2	SSX4	STAT6	SYK	TAF15	TAL1
TBL1XR1	TCF3 (E2A)	TCL1A (TCL1)	TEC	TET1	TFE3	TFG	TFPT
TLX1	TLX3	TMPRSS2	TNFRSF11A	TOP1	TP63	TPM3	TPM4
TRIP11	TTL	TYK2	USP6	YPEL5	ZBTB16	ZMYM2	ZNF384
							ZNF521

\*Note: some VUS rearrangements between targeted genes and unknown fusion partners or intergenic regions detected by RNA sequencing may not be reported.

**ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS**

Microsatellite (MS) status  
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**APPENDIX**
**Performance Specifications**

The median exon coverage for this sample is 668x

**ACCURACY**

Sensitivity: Base Substitutions	At $\geq 5\%$ Minor Allele Frequency	>99.0%
Sensitivity: Insertions/Deletions (1-40bp)	At $\geq 10\%$ Minor Allele Frequency	98.0%
Sensitivity: Focal Copy Number Alterations (Homozygous Deletions or Amplifications)	At $\geq 8$ copies	>95.0%
Sensitivity: Microsatellite Instability-High (MSI-H) status	Positive Predictive Agreement (PPA)	100.0% (87.54%-100.00%)*
Sensitivity: Microsatellite Stable (MSS) status	Positive Predictive Agreement (PPA)	89.66% (81.50%, 94.46%)*
Sensitivity: Known Gene Fusions	>95.0%	
Specificity: Base Substitutions, Insertions/Deletions, and Focal Copy Number Alterations	Positive Predictive Value (PPV)	>99.0%
Specificity: Known Gene Fusions	Positive Predictive Value (PPV)	>95.0%
Specificity: Microsatellite Instability-High (MSI-H) status	Negative Predictive Agreement (NPA)	97.44% (91.12%-99.29%)*
Specificity: Microsatellite Stable (MSS) status	Negative Predictive Agreement (NPA)	94.44% (86.57%, 97.82%)*
Accuracy: Tumor Mutation Burden	At $\geq 20\%$ tumor nuclei	>90.0%
Reproducibility (average concordance between replicates)	97.0% inter-batch precision 97.0% intra-batch precision 95.0% microsatellite status precision 96.0% tumor mutation burden precision	

\* 95% Confidence Interval

Assay specifications were determined for typical median exon coverage of approximately 500X. For additional information regarding the validation of FoundationOne®Heme, please refer to the article He, J. et al. Integrated genomic DNA/RNA profiling of hematologic malignancies in the clinical setting, Blood (2016 Jun. 16).

In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the FoundationOne Heme assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by

the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.

Tumor Mutational Burden (TMB) is determined by measuring the number of somatic mutations in sequenced genes on the FoundationOne Heme test and extrapolating to the genome as a whole. TMB is assayed for all FoundationOne Heme samples and is reported as the number of mutations per megabase (Muts/Mb). Tumor Mutational Burden is reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine Tumor Mutational Burden.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**APPENDIX**
**About FoundationOne®Heme**
**ABOUT FOUNDATIONONE HEME**

FoundationOne®Heme is a comprehensive genomic profiling test for hematologic malignancies and sarcomas. The test is designed to provide physicians with clinically actionable information to help with diagnostic sub-classification, prognosis assessment, and targeted therapeutic selection. Test results provide information about clinically significant alterations, potential targeted therapies, available clinical trials, and quantitative markers that may support immunotherapy clinical trial enrollment. FoundationOne Heme is analytically validated to detect all classes of genomic alterations in more than 400 cancer-related genes. In addition to DNA sequencing, FoundationOne Heme employs RNA sequencing across 265 genes to capture a broad range of gene fusions, common drivers of hematologic malignancies and sarcomas.

FoundationOne Heme was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Heme has not been cleared or approved by the United States Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. FoundationOne Heme may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing.

**THE REPORT**

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

**Diagnostic Significance**

FoundationOne Heme identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

**Qualified Alteration Calls (Equivocal and Subclonal)**

An alteration denoted as "amplification – equivocal" implies that FoundationOne Heme data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne Heme for identifying a copy number amplification is five (5) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that FoundationOne Heme data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that FoundationOne Heme analytical methodology has identified as being present in <10% of the assayed tumor DNA.

**Ranking of Therapies and Clinical Trials**
*Ranking of Therapies in Summary Table*

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

*Ranking of Clinical Trials*

Pediatric trial qualification → Geographical proximity → Later trial phase.

**NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION**

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) ([www.nccn.org](http://www.nccn.org)). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to [NCCN.org](http://NCCN.org). NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

**LEVEL OF EVIDENCE NOT PROVIDED**

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

**NO GUARANTEE OF CLINICAL BENEFIT**

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

**NO GUARANTEE OF REIMBURSEMENT**

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Heme.

**TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN**

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. These include: subclonal alterations in heterogeneous samples, low sample quality or with homozygous losses of <3 exons; and deletions and insertions >40bp, or in repetitive/high homology sequences. FoundationOne Heme is performed using DNA and RNA derived from tumor, and as such germline events may not be reported.

The following targets typically have low coverage resulting in a reduction in sensitivity: SDHD exon 4, TNFRSF11A exon1, and TP53 exon 1.

FoundationOne Heme fulfills the requirements of the European Directive 98/79 EC for in vitro

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1611555-01

**APPENDIX**
**About FoundationOne®Heme**

diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a., Cipalstraat 3, 2440 Geel, Belgium.



## REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

## VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

## VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only.

Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

## SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

## REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.8.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Lena Stuart, M.D. | 26 April 2023  
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531  
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309  
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309  
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531  
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # **ORD-1611555-01**
**APPENDIX**
**References**

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Bonneville R, et al. JCO Precis Oncol (2017) PMID: 29850653
7. Monument MJ, et al. ISRN Oncol (2012) PMID: 23401795
8. Wooster R, et al. Nat. Genet. (1994) PMID: 8162069
9. Kawaguchi K, et al. Oncol. Rep. (2005) PMID: 15643505
10. Saito T, et al. Hum. Pathol. (2003) PMID: 14562278
11. Suwa K, et al. J Orthop Sci (1999) PMID: 10370164
12. Garcia JJ, et al. Mod. Pathol. (2006) PMID: 16619000
13. Aue G, et al. Cancer Genet. Cytogenet. (1998) PMID: 9689926
14. Rucińska M, et al. Med. Sci. Monit. (2005) PMID: 15668629
15. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
16. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
17. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
18. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
19. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
20. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
21. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
22. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
23. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
24. Cristescu R, et al. Science (2018) PMID: 30309915
25. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
26. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
27. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
28. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
29. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
30. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
31. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
32. Ott PA, et al. J. Clin. Oncol. (2019) PMID: 30557521
33. Cristescu R, et al. J Immunother Cancer (2022) PMID: 35101941
34. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
35. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
36. Schenker et al., 2022; AACR Abstract 7845
37. Legrand et al., 2018; ASCO Abstract 12000
38. Akumalla S, et al. Oncology (2020) PMID: 32966992
39. Gounder MM, et al. Nat Commun (2022) PMID: 35705558
40. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
41. Lim J, et al. Clin. Cancer Res. (2015) PMID: 26330427
42. Brohl AS, et al. PLoS Genet. (2014) PMID: 25010205
43. Chen X, et al. Cancer Cell (2013) PMID: 24332040
44. Steele CD, et al. Cancer Cell (2019) PMID: 30889380
45. Casey DL, et al. Clin Cancer Res (2020) PMID: 31699828
46. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
47. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
48. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
49. Rizvi NA, et al. Science (2015) PMID: 25765070
50. Johnson BE, et al. Science (2014) PMID: 24336570
51. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
52. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
53. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
54. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
55. Nature (2012) PMID: 22810696
56. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
57. Choueiri TK, et al. Cancer (2010) PMID: 20665500
58. Malouf GG, et al. Ann. Oncol. (2010) PMID: 20154303
59. Numakura K, et al. Int. J. Clin. Oncol. (2011) PMID: 21161315
60. Liu YC, et al. Jpn. J. Clin. Oncol. (2011) PMID: 21965162
61. Chowdhury T, et al. J. Pediatr. Hematol. Oncol. (2013) PMID: 22995920
62. Stacchiotti S, et al. Clin. Cancer Res. (2009) PMID: 19188185
63. Stacchiotti S, et al. Ann. Oncol. (2011) PMID: 21242589
64. Orbach D, et al. Pediatr Blood Cancer (2013) PMID: 23857870
65. Kummar S, et al. J. Clin. Oncol. (2013) PMID: 23630200
66. Judson I, et al. Clin. Cancer Res. (2014) PMID: 24714778
67. Marcon J, et al. Clin Cancer Res (2020) PMID: 32220885
68. Qu Y, et al. Sci Rep (2016) PMID: 27026382
69. Rua Fernández OR, et al. Urology (2018) PMID: 29702156
70. Koh Y, et al. Ann Oncol (2013) PMID: 23180114
71. Kakoki K, et al. Clin Genitourin Cancer (2017) PMID: 28190703
72. Parikh J, et al. Rare Tumors (2009) PMID: 21139932
73. Schöffski P, et al. Ann. Oncol. (2018) PMID: 29216400
74. Wedekind MF, et al. Pediatr Blood Cancer (2017) PMID: 28417541
75. Tsuda M, et al. Cancer Res. (2007) PMID: 17283122
76. Kobos R, et al. J. Pathol. (2013) PMID: 23288701
77. Liu F, et al. Int J Clin Exp Pathol (2014) PMID: 25337301
78. Argani P, et al. Am. J. Surg. Pathol. (2010) PMID: 20871214
79. Schoolmeester JK, et al. Am. J. Surg. Pathol. (2015) PMID: 25517951
80. Agaram NP, et al. Am. J. Surg. Pathol. (2015) PMID: 25651471
81. Wang XT, et al. Mod. Pathol. (2018) PMID: 29713041
82. Shen Q, et al. Virchows Arch. (2014) PMID: 25239799
83. Gotoh M, et al. Genes Chromosomes Cancer (2014) PMID: 25230976
84. Argani P, et al. Am. J. Pathol. (2001) PMID: 11438465
85. Clark J, et al. Oncogene (1997) PMID: 9393982
86. Weterman MJ, et al. Oncogene (2000) PMID: 10644981
87. Skalsky YM, et al. Oncogene (2001) PMID: 11313942
88. Sato Y, et al. Nat. Genet. (2013) PMID: 23797736
89. Xia QY, et al. Mod Pathol (2017) PMID: 27934879
90. Argani P, et al. Am. J. Surg. Pathol. (2016) PMID: 26975036
91. McGregor SM, et al. Am J Surg Pathol (2017) PMID: 28009605
92. Wang XT, et al. Histopathology (2016) PMID: 26844676
93. Davis IJ, et al. Cancer Cell (2006) PMID: 16766266
94. Kauffman EC, et al. Nat Rev Urol (2014) PMID: 25048860
95. Weterman MA, et al. Proc Natl Acad Sci U S A (1996) PMID: 8986805
96. Argani P, et al. Oncogene (2003) PMID: 12917640
97. Bruder E, et al. Am. J. Surg. Pathol. (2004) PMID: 15316311
98. Trpkov K, et al. Mod Pathol (2021) PMID: 33664427
99. Moch H, et al. Eur Urol (2016) PMID: 26935559
100. Warren TE, et al. Heart Lung (1983) PMID: 6575003
101. Cajiaba MM, et al. Cancer (2018) PMID: 29905933
102. Vistica DT, et al. J. Pediatr. Hematol. Oncol. (2009) PMID: 19636271
103. Covell DG, et al. PLoS ONE (2012) PMID: 23226201
104. Ford KM, et al. Mol. Vis. (2012) PMID: 22419845
105. AlAzab RS, et al. Res Rep Urol (2019) PMID: 32010646
106. Citak EC, et al. Can Urol Assoc J () PMID: 26279736
107. Hulová S, et al. Klin Onkol (2018) PMID: 30545225

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.