

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE	Pancreas ductal adenocarcinoma	PHYSICIAN	ORDERING PHYSICIAN	Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE	Lung
	NAME	Huang, Li-Na		MEDICAL FACILITY	Taipei Veterans General Hospital		SPECIMEN ID	S111-07932C (PF22033)
	DATE OF BIRTH	15 March 1958		ADDITIONAL RECIPIENT	None		SPECIMEN TYPE	Slide Deck
	SEX	Female		MEDICAL FACILITY ID	205872		DATE OF COLLECTION	25 February 2022
	MEDICAL RECORD #	16590628		PATHOLOGIST	Not Provided		SPECIMEN RECEIVED	09 March 2022

Biomarker Findings

Microsatellite status - MS-Stable
Tumor Mutational Burden - 3 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

CDK4 amplification - equivocal[†]

KRAS Q61R

MTAP loss

CCND3 amplification - equivocal[†]

CDKN2A/B CDKN2B loss, CDKN2A loss

EPHB4 amplification - equivocal[†]

SMAD4 loss exons 8-12

TP53 Q167fs*14

2 Disease relevant genes with no reportable alterations: **BRCA1**, **BRCA2**

[†] See About the Test in appendix for details.

Report Highlights

- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 9)

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 3 Muts/Mb

GENOMIC FINDINGS

CDK4 - amplification - equivocal

10 Trials see p. 9

KRAS - Q61R

2 Trials see p. 11

MTAP - loss

1 Trial see p. 12

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. see Biomarker Findings section

No therapies or clinical trials. see Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
none	none
none	none
none	none

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CCND3 - amplification - equivocal.....	p. 5	SMAD4 - loss exons 8-12.....	p. 7
CDKN2A/B - CDKN2B loss, CDKN2A loss.....	p. 6	TP53 - Q167fs*14.....	p. 8
EPHB4 - amplification - equivocal.....	p. 7		

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

BIOMARKER FINDINGS
BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective

analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

MSI is rare in pancreatic carcinoma, reported in less than 1% of samples ($n>1,000$)⁶⁻¹⁰. The prognostic significance of MSI in pancreatic cancer is unknown (PubMed, Aug 2021).

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive

amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹¹. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹¹⁻¹³. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹⁴⁻¹⁶. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{11,13,15-16}.

BIOMARKER

Tumor Mutational Burden

RESULT

3 Muts/Mb

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁷⁻¹⁹, anti-PD-1 therapies¹⁷⁻²⁰, and combination nivolumab and ipilimumab²¹⁻²⁶. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors^{17-20,27}. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment for patients with 9 types of advanced tumors¹⁷. Analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20

Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with chemotherapy²⁸ or those with lower TMB treated with PD-1 or PD-L1-targeting agents¹⁸. However, the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors found significant improvement in ORR for patients with TMB ≥ 10 Muts/Mb (based on this assay or others) compared to those with TMB < 10 Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials^{20,27}. Together, these studies suggest that patients with TMB ≥ 10 Muts/Mb may derive clinical benefit from PD-1 or PD-L1 inhibitors.

FREQUENCY & PROGNOSIS

Pancreatic carcinomas, including ductal and acinar subtypes, have been reported to harbor a median TMB of 2-3 mutations per megabase (mut/Mb), and 0-2% of cases have high TMB (> 20 muts/Mb)²⁹; TMB has not been assessed in pancreatic mucinous neoplasms (PubMed, Oct 2021). A study of patients with pancreatic ductal adenocarcinoma harboring mismatch repair gene mutations

reported improved prognosis for patients with high TMB measured in tissue samples (defined as > 50 mutations; survival 69-314 months) compared to those with lower TMB (average of 5.7 mutations; 10-42 months)³⁰.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³¹⁻³² and cigarette smoke in lung cancer³³⁻³⁴, treatment with temozolomide-based chemotherapy in glioma³⁵⁻³⁶, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes³⁷⁻⁴¹, and microsatellite instability (MSI)^{37,40-41}. This sample harbors a TMB level associated with lower rates of clinical benefit from treatment with PD-1- or PD-L1-targeting immune checkpoint inhibitors compared with patients with tumors harboring higher TMB levels, based on several studies in multiple solid tumor types^{18-19,27}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

GENOMIC FINDINGS
GENE

CDK4

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

CDK4 amplification or activation may predict sensitivity to CDK4/6 inhibitors such as abemaciclib, palbociclib, and ribociclib⁴²⁻⁴⁵. Clinical benefit has been reported for limited

tumor types including patients with CDK4-amplified liposarcoma and sarcoma in response to treatment with abemaciclib⁴⁶, palbociclib^{42,47}, and ribociclib⁴⁸.

FREQUENCY & PROGNOSIS

CDK4 amplification has been observed in 2% of pancreatic adenocarcinomas⁴⁹. High CDK4 expression has been associated with an increase in overall survival in patients with pancreatic cancer⁵⁰. CDK4/6 are frequently deregulated by CDKN2A/B inactivation in pancreatic carcinoma and may cooperate with activated KRAS to promote pancreatic carcinoma progression⁵¹⁻⁵³.

FINDING SUMMARY

CDK4 encodes the cyclin-dependent kinase 4, which regulates the cell cycle, senescence, and apoptosis⁵⁴. CDK4 and its functional homolog CDK6 are activated by D-type cyclins and promote cell cycle progression by inactivating the tumor suppressor Rb⁵⁵⁻⁵⁶. Amplification of the chromosomal region that includes CDK4 has been reported in multiple cancer types, including lung cancer, glioblastoma, and liposarcoma, and has been associated with overexpression of CDK4 protein^{42,57-63}.

GENE

KRAS

ALTERATION

Q61R

TRANSCRIPT ID

NM_004985

CODING SEQUENCE EFFECT

182A>G

VARIANT ALLELE FREQUENCY (% VAF)

16.3%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Preclinical evidence suggests that KRAS activation may predict sensitivity to MEK inhibitors, such as trametinib, binimetinib, cobimetinib, and selumetinib⁶⁴⁻⁶⁹. Initial Phase 1 monotherapy trials of MEK inhibitors in patients with pancreatic cancer showed promise, with DCR (PR and/or SD) up to 37%⁷⁰, response rates up to 25%⁷⁰⁻⁷⁴, and prolonged PRs in certain patients^{71,73,75}. However, subsequent clinical trials combining various MEK inhibitors with gemcitabine reported no additional benefit compared to gemcitabine alone irrespective of KRAS mutation status⁷⁶⁻⁷⁹, with refametinib and gemcitabine even showing a trend towards worse response and survival in patients with KRAS-mutant pancreatic tumors than in those with KRAS wild-type tumors (OS 6.6 months vs 18.2 months)⁷⁶. Trials combining MEK inhibitors with other targeted therapies, such as EGFR

inhibitors⁸⁰ or PI3K-AKT pathway inhibitors⁸¹⁻⁸², reported no PRs and frequent adverse events in patients with KRAS-mutant pancreatic cancer. Emerging preclinical studies suggest MEK inhibition downstream of KRAS-mutant pancreatic tumors leads to increased autophagy⁸³⁻⁸⁴. Combination MEK/autophagy inhibitors may therefore be more beneficial. A heavily pretreated patient with pancreatic cancer treated with trametinib plus hydroxychloroquine exhibited a PR⁸³. A Phase 2 trial of paclitaxel/carboplatin with or without Reolysin in patients with metastatic pancreatic adenocarcinoma reported no improvement in PFS with addition of Reolysin, regardless of KRAS mutational status⁸⁵; however a Phase 2 study of Reolysin and gemcitabine in patients with pancreatic cancer reported 1 PR, 23 SDs, and 5 PDs in 34 patients with a favorable median OS of 10.2 months⁸⁶. In a Phase 1 study evaluating the MEK-pan-RAF dual inhibitor CH5126766, 6 patients harboring KRAS mutations experienced PRs, including 3 with non-small cell lung cancer (NSCLC), 1 with low-grade serous ovarian carcinoma (LGSOC), 1 with endometrial adenocarcinoma, and 1 with multiple myeloma⁸⁷. Combination of CH5126766 with the FAK inhibitor defactinib elicited PR rates of 50% (4/8) for patients with KRAS-mutated low-grade serous ovarian cancer and 12% (2/17) for patients with KRAS-mutated non-small cell lung cancer (NSCLC) in a Phase 1 study⁸⁸⁻⁸⁹. Preclinical and clinical data suggest that KRAS mutations may predict clinical benefit from SHP2 inhibitors⁹⁰⁻⁹¹. A Phase 1 study of RMC-4630 for relapsed/refractory solid tumors reported a DCR of 58% (23/40) for patients with NSCLC and KRAS

mutations and a DCR of 75% (12/16) for patients with NSCLC and KRAS G12C mutations⁹². Interim results from a Phase 1/2 study of RMC-4630 plus cobimetinib reported tumor reduction in 3 of 8 patients with KRAS-mutated colorectal cancer⁹³. Preclinical data suggest that KRAS mutation may confer sensitivity to SOS1 inhibitors⁹⁴⁻⁹⁵. Phase 1 studies of the SOS1 inhibitor BI 1701963 alone or in combination with MEK inhibitors, KRAS G12C inhibitors, or irinotecan are recruiting for patients with solid tumors harboring KRAS mutations⁹⁶⁻⁹⁷.

FREQUENCY & PROGNOSIS

KRAS mutations have been observed in 91-95% of pancreatic ductal adenocarcinoma cases⁹⁸⁻⁹⁹, with the majority of mutations found at codon 12¹⁰⁰⁻¹⁰³. KRAS mutations, particularly G12D, have been associated with decreased median survival time in patients with pancreatic ductal adenocarcinoma¹⁰¹.

FINDING SUMMARY

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor formation^{65,104}. KRAS alterations affecting amino acids G12, G13, Q22, P34, A59, Q61, and A146, as well as mutations G10_A11insG, G10_A11insAG (also reported as G10_A11dup and G12_G13insAG), A18D, L19F, D33E, G60_A66dup/E62_A66dup, E62K, E63K, R68S, and K117N have been characterized as activating and oncogenic^{65,105-127}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

GENOMIC FINDINGS

GENE

MTAP

ALTERATION

loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical and limited clinical evidence indicate that MTAP inactivation produces specific metabolic vulnerabilities. MTAP inactivation may confer sensitivity to MAT2A inhibitors¹²⁸. A Phase 1 trial of MAT2A inhibitor AG-270 reported 1 PR and 2 SDs lasting longer than 6 months for patients with advanced solid tumors displaying MTAP loss¹²⁹. Although preclinical data have suggested that MTAP loss sensitizes cells to PRMT5 inhibition^{128,130-131}, MTAP loss may not be a biomarker of response to previously developed small-molecule SAM-uncompetitive PRMT5 inhibitors¹³²; dual PRMT1 and PRMT5 inhibition may be more effective¹³³⁻¹³⁵. In preclinical cancer models, MTAP inactivation showed increased

sensitivity to inhibitors of purine synthesis or purine analogs, especially upon addition of exogenous MTA, which is converted to adenine in normal cells, thereby providing competition to purine poisons lacking in MTAP-deficient cells¹³⁶⁻¹⁴⁶. A Phase 2 study of L-alanosine, an inhibitor of adenine synthesis, as a monotherapy for 65 patients with MTAP-deficient cancers reported no responses and stable disease in 23.6% (13/55) of patients¹⁴⁷.

FREQUENCY & PROGNOSIS

MTAP loss/homozygous deletion as well as loss of expression has been reported in a wide variety of solid tumors and hematologic cancers¹⁴⁸⁻¹⁴⁹; such events have been correlated with poor prognosis in a variety of cancer types, including hepatocellular carcinoma¹⁵⁰, gastrointestinal stromal tumors¹⁵¹, mantle cell lymphoma (MCL)¹⁵², melanoma¹⁵³⁻¹⁵⁴, gastric cancer¹⁵⁵, myxofibrosarcoma¹⁵⁶, nasopharyngeal carcinoma¹⁵⁷, ovarian carcinoma¹⁴⁸ and non-small cell lung cancer¹⁵⁸. MTAP loss was not prognostic in pediatric B-cell acute lymphocytic leukemia¹⁵⁹ or in astrocytoma¹⁶⁰. However, MTAP has also

been reported to be overexpressed in colorectal cancer (CRC) samples¹⁶¹, and MTAP retention is thought to be important for prostate cancer growth due to continuous supply of SAM¹⁶². Germline SNPs in MTAP have been correlated with the development of cutaneous melanoma¹⁶³⁻¹⁶⁴, esophageal cancer¹⁶⁵⁻¹⁶⁶, osteosarcoma¹⁶⁷, and CRC¹⁶⁸.

FINDING SUMMARY

MTAP encodes S-methyl-5'-thioadenosine (MTA) phosphorylase, a tumor suppressor involved in polyamine metabolism and methionine synthesis, although its enzymatic function is dispensable for its tumor suppressor activity¹⁶⁹⁻¹⁷⁰. Decreased expression of MTAP leads to MTA accumulation within tumor cells and their microenvironment^{150,171-172}, thereby reducing intracellular arginine methylation^{128,130,173} and altering cell signaling^{172,174}. MTAP is located at 9p21, adjacent to CDKN2A and CDKN2B, with which it is frequently co-deleted in various cancers. Other alterations in MTAP are rare and have not been extensively characterized.

GENE

CCND3

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Amplification or activation of CCND3 may predict sensitivity to CDK4/6 inhibitors¹⁷⁵⁻¹⁷⁷, such as abemaciclib, palbociclib, and ribociclib^{42-45,178-179}.

As supported by strong preclinical studies, tumors with CCND3 activation depend on CDK4/6¹⁷⁵⁻¹⁷⁷.

FREQUENCY & PROGNOSIS

Amplification of CCND3 has been reported in 2-7% of pancreatic adenocarcinoma cases (cBioPortal, Apr 2021)^{99,180-181}. Overexpression of cyclin D3 has been reported in one study in the majority of pancreatic intraepithelial neoplasia and ductal tumors analyzed, and the downregulation of cyclin D3 by siRNA has been shown to inhibit the growth of pancreatic cancer cell lines¹⁸²⁻¹⁸³. Published data investigating the prognostic

implications of CCND3 alterations in pancreatic carcinoma are limited (PubMed, Sep 2021).

FINDING SUMMARY

CCND3 encodes cyclin D3, a G1/S-specific cell cycle regulator. Cyclin D3 interacts with and regulates the cyclin-dependent kinases CDK4 and CDK6, resulting in the phosphorylation and inactivation of Rb and the progression of the cell cycle¹⁸⁴. CCND3 amplification has been associated with increased cyclin D3 expression¹⁸⁵.

ORDERED TEST # ORD-1318266-01

GENOMIC FINDINGS

GENE

CDKN2A/B

ALTERATION

CDKN2B loss, CDKN2A loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical data suggest that tumors with loss of p16INK4a function may be sensitive to CDK4/6 inhibitors, such as abemaciclib, ribociclib, and palbociclib¹⁸⁶⁻¹⁸⁹. Although case studies have reported that patients with breast cancer or uterine leiomyosarcoma harboring CDKN2A loss responded to palbociclib treatment¹⁹⁰⁻¹⁹¹, multiple other clinical studies have shown no significant correlation between p16INK4a loss or inactivation and therapeutic benefit of these agents^{45,48,178-179,192-194}; it is not known whether CDK4/6 inhibitors would be beneficial in this case. Although preclinical studies have suggested that loss of p14ARF function may be associated with reduced sensitivity to MDM2 inhibitors¹⁹⁵⁻¹⁹⁶, the clinical relevance of p14ARF as a predictive biomarker is not clear. There are no drugs that directly target the mutation or loss of CDKN2B in cancer. Because the p15INK4b protein encoded by CDKN2B is known to inhibit CDK4, tumors with CDKN2B mutation or loss may predict sensitivity to CDK4/6 inhibitors, such as ribociclib, abemaciclib, and palbociclib^{42-43,48,178-179,197}.

FREQUENCY & PROGNOSIS

CDKN2A/B loss has been reported in 27-36% of pancreatic carcinomas (cBioPortal, Sep 2021)^{99,180-181}. CDKN2A loss has been reported in 25-100% of pancreatic ductal adenocarcinomas analyzed, with a portion of those due to gene deletion^{101,198-200}. A study of multiple pancreatic cancer subtypes reported no CDKN2A mutations (0/6 samples) and loss of heterozygosity in only one of four samples of pancreatic acinar carcinomas, compared to mutation or loss of heterozygosity in 38% and 67% of pancreatic ductal carcinomas, respectively²⁰¹. Promoter methylation affecting p16INK4a and p14ARF has been reported in 43% (16/37) and 20.6% (7/34), respectively, of pancreatic fluid specimens of patients with pancreatic carcinoma²⁰². The loss or decrease of p16INK4a expression levels in pancreatic ductal adenocarcinoma has been reported in 32-80% of samples analyzed and one study reported concurrent loss of p16INK4a and p14ARF protein expression in 68% (19/28) of cases^{198,203-204}. p16INK4a expression has been associated with improved overall survival in pancreatic adenocarcinoma patients in univariate and multivariate analysis²⁰⁵. Furthermore, CDKN2A alterations (deletion or mutation) in the presence of concomitant KRAS mutation may correlate with shorter survival in patients with pancreatic ductal adenocarcinoma²⁰³.

FINDING SUMMARY

CDKN2A encodes two different, unrelated tumor suppressor proteins, p16INK4a and p14ARF, whereas CDKN2B encodes the tumor suppressor

p15INK4b²⁰⁶⁻²⁰⁷. Both p15INK4b and p16INK4a bind to and inhibit CDK4 and CDK6, thereby maintaining the growth-suppressive activity of the Rb tumor suppressor; loss or inactivation of either p15INK4b or p16INK4a contributes to dysregulation of the CDK4/6-cyclin-Rb pathway and loss of cell cycle control²⁰⁸⁻²⁰⁹. The tumor suppressive functions of p14ARF involve stabilization and activation of p53, via a mechanism of MDM2 inhibition²¹⁰⁻²¹¹. One or more alterations observed here are predicted to result in p16INK4a loss of function²¹²⁻²³³. One or more alterations seen here are predicted to result in p14ARF loss of function^{216,233-236}. CDKN2B alterations such as seen here are predicted to inactivate p15INK4b²³⁷.

POTENTIAL GERMLINE IMPLICATIONS

Germline CDKN2A mutation is associated with melanoma-pancreatic cancer syndrome, a condition marked by increased risk of developing malignant melanoma and/or pancreatic cancer²³⁸. Mutation carriers within families may develop either or both types of cancer, and melanoma cases may be referred to as familial or hereditary melanoma²³⁹⁻²⁴⁰. CDKN2A is the most implicated gene in familial melanoma, with germline mutations present in 16% to 20% of familial melanoma cases²⁴¹⁻²⁴³. CDKN2A alteration has also been implicated in familial melanoma-astrocytoma syndrome, an extremely rare tumor association characterized by dual predisposition to melanoma and nervous system tumors²⁴⁴⁻²⁴⁶. In the appropriate clinical context, germline testing of CDKN2A is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

GENOMIC FINDINGS

GENE

EPHB4

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies available to target EPHB4 alterations in cancer. sEPHB4 is a soluble monomeric extracellular domain of EPHB4 that functions as an antagonist of EphrinB2-EPHB4 interaction²⁴⁷, and fusion of sEPHB4 with human serum albumin (HSA) increases its stability²⁴⁸. Recombinant sEPHB4-HSA is under investigation in clinical trials. Preclinical studies have demonstrated that sEPHB4-HSA inhibits cell proliferation and xenograft tumor growth, including for cells expressing cancer-associated EPHB4 mutants or overexpressing wild-type EPHB4^{247,249-253}. In addition, small-molecule

inhibitors targeting multiple tyrosine kinases including EPHB4, such as JI-101 and XL647, have been under preclinical and clinical investigation²⁵⁴⁻²⁵⁶.

FREQUENCY & PROGNOSIS

Increased EPHB4 mRNA and/or protein expression has been reported in a variety of cancer types, including head and neck squamous cell carcinoma (HNSCC)²⁵⁷⁻²⁶⁰, gastric and esophageal²⁶¹⁻²⁶⁵, colorectal carcinoma (CRC)²⁶⁶⁻²⁷², breast²⁷³⁻²⁷⁷, ovarian²⁷⁷⁻²⁷⁹, endometrial²⁸⁰⁻²⁸², thyroid²⁸³⁻²⁸⁵, lung²⁸⁶⁻²⁸⁷, glioma²⁸⁸⁻²⁸⁹, and other solid tumors^{249,290-297}. In several of these studies, increased EPHB4 expression has been associated with clinicopathologic features, including disease stage^{249,257,273,278-279,287,290,292}, histological grade^{263,273,280,289}, and hormone receptor status^{276,281}. High EPHB4 expression has been associated with inferior survival in multivariate analyses for patients with CRC treated with bevacizumab [hazard ratio (HR) = 5.95]²⁷¹, HNSCC (HR = 2.95)²⁶⁰, epithelial ovarian cancer (HR =

4.53)²⁷⁷, or glioma (HR = 3.21)²⁸⁹.

FINDING SUMMARY

EPHB4 encodes a member of the EPH family of receptor tyrosine kinases²⁹⁸. Ephrin signaling has been implicated in multiple processes, including cell adhesion, cytoskeletal organization, and cell migration²⁹⁹, and signaling between EPHB4 and its ligand EphrinB2 is particularly important for angiogenesis³⁰⁰⁻³⁰¹. EPHB receptors, including EPHB4, have been shown to undergo dysregulation (amplification, mutation, under- or overexpression) in a number of different cancer types³⁰². EPHB4 amplification has been reported in several solid tumor types^{257-258,263,303-304} and was associated with advanced disease stage in head and neck squamous cell carcinoma (HNSCC)²⁵⁷. Activating missense mutations in or near the tyrosine kinase domain, including G723S, A742V, and P881S, have also been identified in lung cancer²⁵³.

GENE

SMAD4

ALTERATION

loss exons 8-12

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies to address SMAD4 alterations in cancer. Preclinical studies³⁰⁵⁻³⁰⁶ and a clinical study of pancreatic cancer suggest that low SMAD4 expression exhibit increased responsiveness to chemotherapeutic agents such as cisplatin and irinotecan³⁰⁷.

FREQUENCY & PROGNOSIS

SMAD4 mutation or homozygous deletion is most frequently observed in pancreatic adenocarcinoma (43%)⁹⁹, pancreatic acinar cell carcinoma³⁰⁸, cholangiocarcinoma (25%)³⁰⁹, appendiceal

adenocarcinoma (14-20% mutation; 57% deletion)³¹⁰⁻³¹¹, colorectal adenocarcinoma (CRC; 14%)⁴⁰, esophageal adenocarcinoma (14%)³¹², and stomach adenocarcinoma (13%)³¹³. In preclinical studies, SMAD4 loss of function has been implicated in the development of mucinous neoplasms of the pancreas, including mucinous cystic neoplasms (MCN)³¹⁴ and intraductal papillary mucinous neoplasms (IPMN)³¹⁵; in clinical samples, SMAD4 homozygous deletion has been observed in 10% of IPMNs and 8% of MCNs, and mutation was also observed in 5% of IPMNs³¹⁶. SMAD4 gene alterations have been associated with reduced overall survival for patients with pancreatic adenocarcinoma³¹⁷. Reduced SMAD4 expression has been associated with worse prognosis in various cancer types, including CRC³¹⁸⁻³²⁰, appendiceal mucinous neoplasm³²¹, gastric adenocarcinoma³²²⁻³²³, esophageal adenocarcinoma³²⁴, esophageal squamous cell carcinoma³²⁵, breast cancer³²⁶, and prostate cancer³²⁷.

FINDING SUMMARY

SMAD4, also known as DPC4, encodes a tumor suppressor that regulates transcriptional activity downstream of TGF-beta receptor signaling³²⁸⁻³²⁹. SMAD4 alterations that result in loss or disruption of the MH1 domain (aa 18-142), MH2 domain (aa 323-552), or SAD domain (aa 275-320) are predicted to be inactivating³³⁰⁻³⁴³.

POTENTIAL GERMLINE IMPLICATIONS

Germline SMAD4 mutations, including those at the R361 hotspot, have been observed in patients with juvenile polyposis syndrome³⁴⁴⁻³⁴⁶, which is associated with an increased risk of gastrointestinal cancers³⁴⁷. The penetrance of deleterious SMAD4 mutations in patients with colon cancer is estimated at 20% by age 35 and 70% by age 65³⁴⁸. In the appropriate clinical context, germline testing of SMAD4 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

Q167fs*14

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

498_499insA

VARIANT ALLELE FREQUENCY (% VAF)

51.0%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib³⁴⁹⁻³⁵², or p53 gene therapy and immunotherapeutics such as SGT-53³⁵³⁻³⁵⁷ and ALT-801³⁵⁸. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype³⁵⁹. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer³⁶⁰. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer³⁶¹. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone³⁶². In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric

cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel³⁶³. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations³⁶⁴. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring³⁶⁵. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage³⁵⁷. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies³⁶⁶⁻³⁶⁷; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies³⁶⁸⁻³⁶⁹. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

TP53 mutations have been reported in 33-75% of pancreatic carcinomas, with the majority occurring as missense mutations, while deletion of TP53 has been found in 66% of pancreatic ductal adenocarcinoma cases^{98,370-372}. TP53 mutations are common in pancreatic ductal adenocarcinomas and are known to occur in the process of pancreatic carcinogenesis³⁷³⁻³⁷⁴. Additionally, aberrant expression of p53 has been found in 54-81% of pancreatic ductal adenocarcinoma cases^{203,371,375-376}. Studies have found inconsistent results regarding the prognostic significance of p53 expression in pancreatic ductal adenocarcinoma, although one study correlated low levels of TP53 mRNA with poor patient

prognosis^{203,377-378}.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers³⁷⁹. Alterations such as seen here may disrupt TP53 function or expression³⁸⁰⁻³⁸⁴.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers³⁸⁵⁻³⁸⁷, including sarcomas³⁸⁸⁻³⁸⁹. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000³⁹⁰ to 1:20,000³⁸⁹. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30³⁹¹. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion³⁹²⁻³⁹⁷. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy³⁹²⁻³⁹³. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease³⁹⁸. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{396,399-400}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
CDK4
RATIONALE

CDK4 amplification may predict sensitivity to

CDK4/6 inhibitors.

ALTERATION

amplification - equivocal

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04282031
PHASE 1/2

A Study of BPI-1178 in Patients With Advanced Solid Tumor and HR+/HER2- Breast Cancer

TARGETS

CDK6, CDK4, ER, Aromatase

LOCATIONS: Shanghai (China)

NCT04594005
PHASE 1/2

CDK4/6 Tumor, Abemaciclib, Paclitaxel

TARGETS

CDK4, CDK6

LOCATIONS: Seoul (Korea, Republic of)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

CLINICAL TRIALS

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT03994796
PHASE 2

Genetic Testing in Guiding Treatment for Patients With Brain Metastases

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, CDK4, CDK6, PI3K, mTOR

LOCATIONS: Washington, Oregon, Idaho, Montana

NCT03310879
PHASE 2

Study of the CDK4/6 Inhibitor Abemaciclib in Solid Tumors Harboring Genetic Alterations in Genes Encoding D-type Cyclins or Amplification of CDK4 or CDK6

TARGETS

CDK4, CDK6

LOCATIONS: Massachusetts

NCT02693535
PHASE 2

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, CDK4, CDK6, FLT3, CSF1R, KIT, RET, mTOR, EGFR, ERBB2, MEK, BRAF, SMO, DDR2, PARP, PD-1, CTLA-4, ERBB4

LOCATIONS: Hawaii, Washington, Oregon, California

NCT04557449
PHASE 1

Study to Test the Safety and Tolerability of PF-07220060 in Participants With Advance Solid Tumors

TARGETS

CDK4, Aromatase, ER

LOCATIONS: Michigan, Massachusetts, Connecticut, Tennessee, Texas

NCT02896335
PHASE 2

Palbociclib In Progressive Brain Metastases

TARGETS

CDK4, CDK6

LOCATIONS: Massachusetts

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 17 March 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

CLINICAL TRIALS
GENE
KRAS
ALTERATION
Q61R
RATIONALE

KRAS activating mutations or amplification may predict sensitivity to inhibitors of MAPK pathway components, including MEK inhibitors. Multiple clinical studies have reported lack of efficacy of MEK inhibitors as monotherapy for treatment of KRAS-mutant pancreatic cancer. Emerging data

suggest patients with KRAS-mutant pancreatic cancer may be sensitive to combination MEK/autophagy inhibitors. Limited clinical and preclinical studies indicate KRAS mutations may predict sensitivity to MEK-pan-RAF dual inhibitors.

NCT03825289
PHASE 1

Trametinib and Hydroxychloroquine in Treating Patients With Pancreatic Cancer

TARGETS
MEK
LOCATIONS: Utah

NCT04132505
PHASE 1

Binimetinib and Hydroxychloroquine in Treating Patients With KRAS Mutant Metastatic Pancreatic Cancer

TARGETS
MEK
LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

CLINICAL TRIALS

GENE

MTAP

RATIONALE

MTAP loss may predict sensitivity to MAT2A inhibitors.

ALTERATION

loss

NCT03435250
PHASE 1

Study of AG-270 in Participants With Advanced Solid Tumors or Lymphoma With MTAP Loss

TARGETS
MAT2A
LOCATIONS: Villejuif Cedex (France), Barcelona (Spain), Massachusetts, Connecticut, New York, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

AR
R101H

BRCA2
E1806K

EPHB1
S766F

FBXW7
N432I

KMT2A (MLL)
A53V

MTOR
rearrangement

NOTCH3
R1175W

SNCAIP
rearrangement

ZNF217
S3L

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B)	APC
AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2
BTK	C11orf30 (EMSY)	C17orf39 (GID4)	CALR	CARD11	CASP8	CBFB	CBL	CCND1
CCND2	CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2	DIS3	DNMT3A	DOT1L	EED	EGFR	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FAM46C	FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12
FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3
FGFR4	FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3
GATA4	GATA6	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3F3A
HDAC1	HGF	HNF1A	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDMSA	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKKN1	MLH1	MPL	MRE11A	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD3 (WHSC1L1)	NTSC2	NTRK1	NTRK2	NTRK3	P2RY8	PALB2
PARK2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA
PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1	PMS2
POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	REL	RET
RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC	SDHD	SETD2
SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO	SNCAIP	SOC3
SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3	STK11	SUFU
SYK	TBX3	TEK	TET2	TGFBR2	TIPARP	TNFAIP3	TNFRSF14	TP53
TSC1	TSC2	TYRO3	U2AF1	VEGFA	VHL	WHSC1	WT1	XPO1
XRCC2	ZNF217	ZNF703						

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TMPRSS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium.



ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLES

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for ERBB2 and six (6) for all other genes. Conversely, an alteration denoted as "loss - equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX

About FoundationOne®CDx

- Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.
 - Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
 - Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine

whether the patient is a candidate for biopsy.

- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of HER2 overexpression as 20% in breast cancer. Based on the F1CDx HER2 CDx concordance study, approximately 10% of HER2 amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX

About FoundationOne®CDx

cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 6.0.0

The median exon coverage for this sample is 1,000x

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # **ORD-1318266-01**
APPENDIX
References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Hu ZI, et al. Clin. Cancer Res. (2018) PMID: 29367431
7. Campbell BB, et al. Cell (2017) PMID: 29056344
8. Pihlak R, et al. Cancers (Basel) (2018) PMID: 29329208
9. Salem ME, et al. Mol. Cancer Res. (2018) PMID: 29523759
10. Laghi L, et al. PLoS ONE (2012) PMID: 23029359
11. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
12. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
13. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
14. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
15. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
16. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
17. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
18. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
19. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
20. Cristescu R, et al. Science (2018) PMID: 30309915
21. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
22. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
23. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
24. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
25. Rozeman EA, et al. Nat. Med. (2021) PMID: 33558721
26. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
27. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
28. Legrand et al., 2018; ASCO Abstract 12000
29. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
30. Hu et al., 2017; ASCO Abstract e15791
31. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
32. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
33. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
34. Rizvi NA, et al. Science (2015) PMID: 25765070
35. Johnson BE, et al. Science (2014) PMID: 24336570
36. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
37. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
38. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
39. Heitzner E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
40. Nature (2012) PMID: 22810696
41. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
42. Dickson MA, et al. J. Clin. Oncol. (2013) PMID: 23569312
43. Flaherty KT, et al. Clin. Cancer Res. (2012) PMID: 22090362
44. Patnaik A, et al. Cancer Discov (2016) PMID: 27217383
45. Infante JR, et al. Clin. Cancer Res. (2016) PMID: 27542767
46. Dickson et al., 2019; ASCO Abstract 11004
47. Dickson MA, et al. JAMA Oncol (2016) PMID: 27124835
48. Peguero et al., 2016; ASCO Abstract 2528
49. Cancer Genome Atlas Research Network. Electronic address: andrew_aguirre@dfci.harvard.edu, et al. Cancer Cell (2017) PMID: 28810144
50. Huang X, et al. Front Cell Dev Biol (2020) PMID: 33282875
51. Aguirre AJ, et al. Genes Dev. (2003) PMID: 14681207
52. Bardeesy N, et al. Nat. Rev. Cancer (2002) PMID: 12459728
53. Ryan DP, et al. N. Engl. J. Med. (2014) PMID: 25207767
54. Choi YJ, et al. Oncogene (2014) PMID: 23644662
55. Cell (1995) PMID: 7736585
56. Musgrove EA, et al. Nat. Rev. Cancer (2011) PMID: 21734724
57. Wikman H, et al. Genes Chromosomes Cancer (2005) PMID: 15543620
58. Rao SK, et al. J. Neurooncol. (2010) PMID: 19609742
59. Chung L, et al. Am. J. Surg. Pathol. (2009) PMID: 19574885
60. Ragazzini P, et al. Histol. Histopathol. (2004) PMID: 15024701
61. Dujardin F, et al. Mod. Pathol. (2011) PMID: 21336260
62. Zhang K, et al. Cancer Res. (2013) PMID: 23393200
63. Horvai AE, et al. Mod. Pathol. (2009) PMID: 19734852
64. Nakano H, et al. Proc. Natl. Acad. Sci. U.S.A. (1984) PMID: 6320174
65. Pylayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) PMID: 21993244
66. Yamaguchi T, et al. Int. J. Oncol. (2011) PMID: 21523318
67. Watanabe M, et al. Cancer Sci. (2013) PMID: 23438367
68. Gilmartin AG, et al. Clin. Cancer Res. (2011) PMID: 21245089
69. Yeh JJ, et al. Mol. Cancer Ther. (2009) PMID: 19372556
70. Bodoky G, et al. Invest New Drugs (2012) PMID: 21594619
71. Rinehart J, et al. J. Clin. Oncol. (2004) PMID: 15483017
72. Lorusso PM, et al. J. Clin. Oncol. (2005) PMID: 16009947
73. Infante JR, et al. Lancet Oncol. (2012) PMID: 22805291
74. Weekes CD, et al. Clin. Cancer Res. (2013) PMID: 23434733
75. Garrido-Laguna I, et al. Oncoscience (2015) PMID: 25897431
76. Van Laethem JL, et al. Target Oncol (2017) PMID: 27975152
77. Infante JR, et al. Eur. J. Cancer (2013) PMID: 23583440
78. Infante JR, et al. Eur. J. Cancer (2014) PMID: 24915778
79. Van Cutsem E, et al. Int. J. Cancer (2018) PMID: 29756206
80. Ko AH, et al. Clin. Cancer Res. (2016) PMID: 26251290
81. Chung V, et al. JAMA Oncol (2017) PMID: 27978579
82. Bedard PL, et al. Clin. Cancer Res. (2015) PMID: 25500057
83. Kinsey CG, et al. Nat. Med. (2019) PMID: 30833748
84. Bryant KL, et al. Nat. Med. (2019) PMID: 30833752
85. Noonan AM, et al. Mol. Ther. (2016) PMID: 27039845
86. Mahalingam D, et al. Cancers (Basel) (2018) PMID: 29799479
87. Guo C, et al. Lancet Oncol (2020) PMID: 33128873
88. Krebs et al., 2021; AACR Abstract CT019
89. Shinde et al., 2020; AACR Abstract CT143
90. Lu H, et al. Mol Cancer Ther (2019) PMID: 31068384
91. Mainardi S, et al. Nat Med (2018) PMID: 29808006
92. Koczywas et al., 2021; AACR Abstract LB001
93. Bendell et al., 2020; EORTC-NCI-AACR Abstract 5
94. Hillig RC, et al. Proc Natl Acad Sci U S A (2019) PMID: 30683722
95. Hofmann MH, et al. Cancer Discov (2021) PMID: 32816843
96. Hofmann et al., 2021; AACR Abstract CT210
97. Gort et al., 2020; ASCO Abstract TPS3651
98. Biankin AV, et al. Nature (2012) PMID: 23103869
99. Witkiewicz AK, et al. Nat Commun (2015) PMID: 25855536
100. Feldmann G, et al. J Hepatobiliary Pancreat Surg (2007) PMID: 17520196
101. Rachakonda PS, et al. PLoS ONE (2013) PMID: 23565280
102. Hruban RH, et al. Am. J. Pathol. (1993) PMID: 8342602
103. Maitra A, et al. Best Pract Res Clin Gastroenterol (2006) PMID: 16549325
104. Kahn S, et al. Anticancer Res. () PMID: 3310850
105. Akagi K, et al. Biochem. Biophys. Res. Commun. (2007) PMID: 17150185
106. Bollag G, et al. J. Biol. Chem. (1996) PMID: 8955068
107. Buhrman G, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20194776
108. Sci. STKE (2004) PMID: 15367757
109. Edkins S, et al. Cancer Biol. Ther. (2006) PMID: 16969076
110. Feig LA, et al. Mol. Cell. Biol. (1988) PMID: 3043178
111. Gremer L, et al. Hum. Mutat. (2011) PMID: 20949621
112. Janakiraman M, et al. Cancer Res. (2010) PMID: 20570890
113. Kim E, et al. Cancer Discov (2016) PMID: 27147599
114. Lukman S, et al. PLoS Comput. Biol. (2010) PMID: 20838576
115. Naguib A, et al. J Mol Signal (2011) PMID: 21371307
116. Prior IA, et al. Cancer Res. (2012) PMID: 22589270
117. Privé GG, et al. Proc. Natl. Acad. Sci. U.S.A. (1992) PMID: 1565661
118. Scheffzek K, et al. Science (1997) PMID: 9219684
119. Scholl C, et al. Cell (2009) PMID: 19490892
120. Smith G, et al. Br. J. Cancer (2010) PMID: 20147967
121. Tyner JW, et al. Blood (2009) PMID: 19075190
122. Valencia A, et al. Biochemistry (1991) PMID: 2029511
123. White Y, et al. Nat Commun (2016) PMID: 26854029
124. Wiest JS, et al. Oncogene (1994) PMID: 8058307
125. Angeles AKJ, et al. Oncol Lett (2019) PMID: 31289513
126. Tong JH, et al. Cancer Biol. Ther. (2014) PMID: 24642870
127. Loree JM, et al. Clin Cancer Res (2021) PMID: 34117033
128. Marjon K, et al. Cell Rep (2016) PMID: 27068473
129. Heist et al., 2019; AACR-NCI-EORTC Abstract B116
130. Mavrakis KJ, et al. Science (2016) PMID: 26912361
131. Endoscopy (1989) PMID: 2691236
132. Guccione E, et al. Nat. Rev. Mol. Cell Biol. (2019) PMID: 31350521
133. Fedorow A, et al. Cancer Cell (2019) PMID: 31257072
134. Srour N, et al. Cancer Cell (2019) PMID: 31287990
135. Gao G, et al. Nucleic Acids Res. (2019) PMID: 30916320
136. Hansen LJ, et al. Cancer Res. (2019) PMID: 31040154
137. Tang B, et al. Cancer Res. (2018) PMID: 29844120
138. Munshi PN, et al. Oncologist (2014) PMID: 24928612
139. de Oliveira SF, et al. PLoS ONE (2016) PMID: 26751376
140. Lubin M, et al. PLoS ONE (2009) PMID: 19478948
141. Tang B, et al. Cancer Biol. Ther. (2012) PMID: 22825330
142. Collins CC, et al. Mol. Cancer Ther. (2012) PMID: 22252602
143. Bertino JR, et al. Cancer Biol. Ther. (2011) PMID: 21301207
144. Coulthard SA, et al. Mol. Cancer Ther. (2011) PMID: 21282358
145. Miyazaki S, et al. Int. J. Oncol. (2007) PMID: 17912432
146. Efferth T, et al. Blood Cells Mol. Dis. () PMID: 11987241
147. Kindler HL, et al. Invest New Drugs (2009) PMID: 18618081
148. Wei R, et al. Sci Rep (2016) PMID: 27929028
149. Zhao M, et al. BMC Genomics (2016) PMID: 27556634

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX
References

150. Kirovski G, et al. Am. J. Pathol. (2011) PMID: 21356366
151. Huang HY, et al. Clin. Cancer Res. (2009) PMID: 19887491
152. Marcé S, et al. Clin. Cancer Res. (2006) PMID: 16778103
153. Meyer S, et al. Exp. Dermatol. (2010) PMID: 20500769
154. Wild PJ, et al. Arch Dermatol (2006) PMID: 16618867
155. Kim J, et al. Genes Chromosomes Cancer (2011) PMID: 21412930
156. Li CF, et al. Oncotarget (2014) PMID: 25426549
157. He HL, et al. Medicine (Baltimore) (2015) PMID: 26656376
158. Su CY, et al. Eur J Surg Oncol (2014) PMID: 24969958
159. Mirebeau D, et al. Haematologica (2006) PMID: 16818274
160. Becker AP, et al. Pathobiology (2015) PMID: 26088413
161. Snezhkina AV, et al. Oxid Med Cell Longev (2016) PMID: 27433286
162. Bistulfi G, et al. Oncotarget (2016) PMID: 26910893
163. Antonopoulou K, et al. J. Invest. Dermatol. (2015) PMID: 25407435
164. Maccioni L, et al. BMC Cancer (2013) PMID: 23816148
165. Hyland PL, et al. Int J Epidemiol (2016) PMID: 26635288
166. Lin X, et al. Cancer Sci. (2017) PMID: 27960044
167. Zhi L, et al. J Cancer (2016) PMID: 27994653
168. Gu F, et al. Br. J. Cancer (2013) PMID: 23361049
169. Limm K, et al. PLoS ONE (2016) PMID: 27479139
170. Tang B, et al. G3 (Bethesda) (2014) PMID: 25387827
171. Limm K, et al. Eur. J. Cancer (2013) PMID: 23265702
172. Stevens AP, et al. J. Cell. Biochem. (2009) PMID: 19097084
173. Kryukov GV, et al. Science (2016) PMID: 26912360
174. Limm K, et al. Eur. J. Cancer (2014) PMID: 25087184
175. Schmitz R, et al. Nature (2012) PMID: 22885699
176. Choi YJ, et al. Cancer Cell (2012) PMID: 23079655
177. Sawai CM, et al. Cancer Cell (2012) PMID: 23079656
178. Finn RS, et al. Lancet Oncol. (2015) PMID: 25524798
179. DeMichele A, et al. Clin. Cancer Res. (2015) PMID: 25501126
180. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
181. Gao J, et al. Sci Signal (2013) PMID: 23550210
182. Al-Aynati MM, et al. Clin. Cancer Res. (2004) PMID: 15475449
183. Radulovich N, et al. Mol. Cancer (2010) PMID: 20113529
184. Sridhar J, et al. AAPS J (2006) PMID: 16584130
185. Kasugai Y, et al. Clin. Cancer Res. (2005) PMID: 16322284
186. Konecny GE, et al. Clin. Cancer Res. (2011) PMID: 21278246
187. Katsumi Y, et al. Biochem. Biophys. Res. Commun. (2011) PMID: 21871868
188. Cen L, et al. Neuro-oncology (2012) PMID: 22711607
189. Logan JE, et al. Anticancer Res. (2013) PMID: 23898052
190. Elvin JA, et al. Oncologist (2017) PMID: 28283584
191. Gao J, et al. Curr Oncol (2015) PMID: 26715889
192. Gopalan et al., 2014; ASCO Abstract 8077
193. Konecny et al., 2016; ASCO Abstract 5557
194. Johnson DB, et al. Oncologist (2014) PMID: 24797823
195. Van Maerken T, et al. Mol. Cancer Ther. (2011) PMID: 21460101
196. Gamble LD, et al. Oncogene (2012) PMID: 21725357
197. Shapiro et al., 2013; ASCO Abstract 2500
198. Tsiambas E, et al. J BUON () PMID: 17600882
199. Bergmann F, et al. J. Clin. Pathol. (2006) PMID: 16497872
200. Attri J, et al. BMC Gastroenterol (2005) PMID: 15985168
201. Moore PS, et al. Br. J. Cancer (2001) PMID: 11161385
202. Klump B, et al. Br. J. Cancer (2003) PMID: 12610506
203. Oshima M, et al. Ann. Surg. (2013) PMID: 23470568
204. Geradts J, et al. Mod. Pathol. (2001) PMID: 11706079
205. Chang DT, et al. Cancer (2010) PMID: 20665497
206. Quelle DE, et al. Cell (1995) PMID: 8521522
207. Mutat. Res. (2005) PMID: 15878778
208. Gazzeri S, et al. Oncogene (1998) PMID: 9484839
209. Oncogene (1999) PMID: 10498883
210. Sherr CJ, et al. Cold Spring Harb. Symp. Quant. Biol. (2005) PMID: 16869746
211. Ozenne P, et al. Int. J. Cancer (2010) PMID: 20549699
212. Ruas M, et al. Oncogene (1999) PMID: 10498896
213. Jones R, et al. Cancer Res. (2007) PMID: 17909018
214. Haferkamp S, et al. Aging Cell (2008) PMID: 18843795
215. Huot TJ, et al. Mol. Cell. Biol. (2002) PMID: 12417717
216. Rizo H, et al. J. Biol. Chem. (2001) PMID: 11518711
217. Gombart AF, et al. Leukemia (1997) PMID: 9324288
218. Yang R, et al. Cancer Res. (1995) PMID: 7780957
219. Parry D, et al. Mol. Cell. Biol. (1996) PMID: 8668202
220. Greenblatt MS, et al. Oncogene (2003) PMID: 12606942
221. Yarbrough WG, et al. J. Natl. Cancer Inst. (1999) PMID: 10491434
222. Poi MJ, et al. Mol. Carcinog. (2001) PMID: 11255261
223. Byeon IJ, et al. Mol. Cell (1998) PMID: 9660926
224. Kannengiesser C, et al. Hum. Mutat. (2009) PMID: 19260062
225. Lal G, et al. Genes Chromosomes Cancer (2000) PMID: 10719365
226. Koh J, et al. Nature (1995) PMID: 7777061
227. McKenzie HA, et al. Hum. Mutat. (2010) PMID: 20340136
228. Miller PJ, et al. Hum. Mutat. (2011) PMID: 21462282
229. Kutscher CL, et al. Physiol. Behav. (1977) PMID: 905385
230. Scaini MC, et al. Hum. Mutat. (2014) PMID: 24659262
231. Jenkins NC, et al. J. Invest. Dermatol. (2013) PMID: 23190892
232. Walker GJ, et al. Int. J. Cancer (1999) PMID: 10389768
233. Rutter JL, et al. Oncogene (2003) PMID: 12853981
234. Itahana K, et al. Cancer Cell (2008) PMID: 18538737
235. Zhang Y, et al. Mol. Cell (1999) PMID: 10360174
236. Zhang Y, et al. Cell (1998) PMID: 9529249
237. Jafri M, et al. Cancer Discov (2015) PMID: 25873077
238. Whelan AJ, et al. N Engl J Med (1995) PMID: 7666917
239. Adv Exp Med Biol (2010) PMID: 20687502
240. Hogg D, et al. J Cutan Med Surg (1998) PMID: 9479083
241. De Unamuno B, et al. Melanoma Res (2018) PMID: 29543703
242. Soura E, et al. J Am Acad Dermatol (2016) PMID: 26892650
243. Huerta C, et al. Acta Derm Venereol (2018) PMID: 29405243
244. Kaufman DK, et al. Neurology (1993) PMID: 8414022
245. Bahau M, et al. Cancer Res (1998) PMID: 9622062
246. Chan AK, et al. Clin Neuropathol () PMID: 28699883
247. Kertesz N, et al. Blood (2006) PMID: 16322467
248. Shi S, et al. J Pharm Sci (2012) PMID: 22411527
249. Li X, et al. PLoS ONE (2014) PMID: 25148033
250. Liu R, et al. BMC Cancer (2013) PMID: 23721559
251. Bhatia S, et al. Sci Rep (2016) PMID: 27941840
252. Sehnnet JS, et al. Blood (2009) PMID: 18836096
253. Ferguson BD, et al. Sci Rep (2015) PMID: 26073592
254. Werner TL, et al. Invest New Drugs (2015) PMID: 26365907
255. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22011666
256. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22722787
257. Sinha UK, et al. Arch. Otolaryngol. Head Neck Surg. (2006) PMID: 17043250
258. Masood R, et al. Int. J. Cancer (2006) PMID: 16615113
259. Ferguson BD, et al. Growth Factors (2014) PMID: 25391996
260. Yavrouian EJ, et al. Arch. Otolaryngol. Head Neck Surg. (2008) PMID: 18794445
261. Liersch-Löhn B, et al. Int. J. Cancer (2016) PMID: 26414866
262. Hu F, et al. Tumour Biol. (2014) PMID: 24771266
263. Hasina R, et al. Cancer Res. (2013) PMID: 23100466
264. Li M, et al. Dig. Dis. Sci. (2011) PMID: 20686847
265. Yin J, et al. Anticancer Res. (2017) PMID: 28739744
266. Stephenson SA, et al. BMC Mol. Biol. (2001) PMID: 11801186
267. Liu W, et al. Cancer (2002) PMID: 11920461
268. McCall JL, et al. Mol. Cell. Biol. (2016) PMID: 27273865
269. Stremtitz S, et al. Mol. Cancer Ther. (2016) PMID: 27535973
270. Lv J, et al. Exp. Mol. Pathol. (2016) PMID: 27072105
271. Guijarro-Muñoz I, et al. Med. Oncol. (2013) PMID: 23579861
272. Kumar SR, et al. Cancer Res. (2009) PMID: 19366806
273. Wu Q, et al. Pathol. Oncol. Res. (2004) PMID: 15029258
274. Berclaz G, et al. Oncol. Rep. () PMID: 12168060
275. Brantley-Sieders DM, et al. PLoS ONE (2011) PMID: 21935409
276. Huang G, et al. Int J Clin Exp Pathol (2015) PMID: 26191333
277. Pradeep S, et al. Cancer Cell (2015) PMID: 26481148
278. Alam SM, et al. Br. J. Cancer (2008) PMID: 18231102
279. Kumar SR, et al. Br. J. Cancer (2007) PMID: 17353927
280. Takai N, et al. Oncol. Rep. () PMID: 11295082
281. Dong LD, et al. Oncol Lett (2017) PMID: 28454369
282. Berclaz G, et al. Ann. Oncol. (2003) PMID: 12562648
283. Sharma GK, et al. Head Neck (2015) PMID: 24634162
284. Giaginis C, et al. Pathol. Oncol. Res. (2016) PMID: 26220827
285. Xuqing W, et al. Tumour Biol. (2012) PMID: 22528941
286. Ferguson BD, et al. PLoS ONE (2013) PMID: 23844053
287. Zheng MF, et al. Mol Med Rep (2012) PMID: 22684742
288. Chen T, et al. Tumour Biol. (2013) PMID: 23138393
289. Tu Y, et al. Clin Transl Oncol (2012) PMID: 22374425
290. Li M, et al. Mol. Biol. Rep. (2013) PMID: 23079712
291. Xia G, et al. Cancer Res. (2005) PMID: 15930280
292. Alam SM, et al. Gynecol. Oncol. (2009) PMID: 19356789
293. Özgür E, et al. Urol. Oncol. () PMID: 19272799
294. Pierscianek D, et al. Neuropathology (2017) PMID: 27388534
295. Pierscianek D, et al. Brain Tumor Pathol (2016) PMID: 26951238
296. Becerikli M, et al. Int. J. Cancer (2015) PMID: 25274141
297. Xia G, et al. Clin. Cancer Res. (2005) PMID: 15958611
298. Noren NK, et al. Cancer Res. (2007) PMID: 17483308
299. Cell (2008) PMID: 18394988
300. Salvucci O, et al. Adv. Cancer Res. (2012) PMID: 22588055
301. Pitulescu ME, et al. Genes Dev. (2010) PMID: 21078817
302. Nat. Rev. Cancer (2010) PMID: 20179713
303. Boberg DR, et al. Chem. Biol. Interact. (2013) PMID: 23063927
304. Cromer A, et al. Oncogene (2004) PMID: 14676830
305. Cui Y, et al. Clin. Cancer Res. (2012) PMID: 22753594
306. Haeger SM, et al. Oncogene (2016) PMID: 25893305

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1318266-01

APPENDIX **References**

307. Bachtet JB, et al. Ann. Oncol. (2012) pmid: 22377565
308. Jiao Y, et al. J. Pathol. (2014) pmid: 24293293
309. Churi CR, et al. PLoS ONE (2014) pmid: 25536104
310. Liu X, et al. Clin. Chem. (2014) pmid: 24821835
311. Maru D, et al. Oncogene (2004) pmid: 14647445
312. Wang K, et al. Oncologist (2015) pmid: 26336083
313. Nature (2014) pmid: 25079317
314. Izeradjene K, et al. Cancer Cell (2007) pmid: 17349581
315. Bardeesy N, et al. Genes Dev. (2006) pmid: 17114584
316. Springer S, et al. Gastroenterology (2015) pmid: 26253305
317. Blackford A, et al. Clin. Cancer Res. (2009) pmid: 19584151
318. Yan P, et al. Clin. Cancer Res. (2016) pmid: 26861460
319. Kozak MM, et al. J. Clin. Pathol. (2015) pmid: 25681512
320. Roth AD, et al. J. Natl. Cancer Inst. (2012) pmid: 23104212
321. Davison JM, et al. Am. J. Surg. Pathol. (2014) pmid: 24618609
322. Kim YH, et al. Ann. Oncol. (2004) pmid: 15033661
323. Xiangming C, et al. Clin. Cancer Res. (2001) pmid: 11234879
324. Singhi AD, et al. Am. J. Surg. Pathol. (2015) pmid: 25634752
325. Natsugoe S, et al. Clin. Cancer Res. (2002) pmid: 12060625
326. de Kruijff EM, et al. Ann. Oncol. (2013) pmid: 23022998
327. Shipitsin M, et al. Br. J. Cancer (2014) pmid: 25032733
328. Nat. Rev. Mol. Cell Biol. (2012) pmid: 22992590
329. Cell (2008) pmid: 18662538
330. Massagué J, et al. Genes Dev. (2005) pmid: 16322555
331. Morén A, et al. Oncogene (2000) pmid: 10980615
332. Xu J, et al. Proc. Natl. Acad. Sci. U.S.A. (2000) pmid: 10781087
333. Luo K, et al. Genes Dev. (1999) pmid: 10485843
334. Jones JB, et al. Nucleic Acids Res. (2000) pmid: 10871368
335. Fink SP, et al. Cancer Res. (2001) pmid: 11196171
336. De Bosscher K, et al. Biochem. J. (2004) pmid: 14715079
337. Shi Y, et al. Nature (1997) pmid: 9214508
338. Miyaki M, et al. Oncogene (1999) pmid: 10340381
339. Prokova V, et al. Biochemistry (2007) pmid: 17994767
340. Wu JW, et al. J. Biol. Chem. (2001) pmid: 11274206
341. Ding L, et al. J. Clin. Invest. (2009) pmid: 19139564
342. Kuang C, et al. Oncogene (2004) pmid: 14647410
343. Watanabe M, et al. EMBO Rep. (2000) pmid: 11265759
344. Houlston R, et al. Hum. Mol. Genet. (1998) pmid: 9811934
345. Woodford-Richens K, et al. Gut (2000) pmid: 10764709
346. Howe JR, et al. J. Med. Genet. (2004) pmid: 15235019
347. Brosens LA, et al. World J. Gastroenterol. (2011) pmid: 22171123
348. Kalia SS, et al. Genet. Med. (2017) pmid: 27854360
349. Hirai H, et al. Cancer Biol. Ther. (2010) pmid: 20107315
350. Bridges KA, et al. Clin. Cancer Res. (2011) pmid: 21799033
351. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) pmid: 21389100
352. Osman AA, et al. Mol. Cancer Ther. (2015) pmid: 25504633
353. Xu L, et al. Mol. Cancer Ther. (2002) pmid: 12489850
354. Xu L, et al. Mol. Med. (2001) pmid: 11713371
355. Camp ER, et al. Cancer Gene Ther. (2013) pmid: 23470564
356. Kim SS, et al. Nanomedicine (2015) pmid: 25240597
357. Pirolo KF, et al. Mol. Ther. (2016) pmid: 27357628
358. Hajdenberg et al., 2012; ASCO Abstract e15010
359. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27601554
360. Moore et al., 2019; ASCO Abstract 5513
361. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27998224
362. Oza et al., 2015; ASCO Abstract 5506
363. Lee J, et al. Cancer Discov (2019) pmid: 31315834
364. Méndez E, et al. Clin. Cancer Res. (2018) pmid: 29535125
365. Seligmann JF, et al. J Clin Oncol (2021) pmid: 34538072
366. Kwok M, et al. Blood (2016) pmid: 26563132
367. Boudny M, et al. Haematologica (2019) pmid: 30975914
368. Dillon MT, et al. Mol. Cancer Ther. (2017) pmid: 28062704
369. Middleton FK, et al. Cancers (Basel) (2018) pmid: 30127241
370. Morton JP, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) pmid: 20018721
371. Scarpa A, et al. Am. J. Pathol. (1993) pmid: 8494051
372. Luo Y, et al. Pathol. Oncol. Res. (2013) pmid: 22782330
373. Iacobuzio-Donahue CA, et al. Clin. Cancer Res. (2012) pmid: 22896692
374. Macgregor-Das AM, et al. J Surg Oncol (2013) pmid: 22806689
375. Ottenhof NA, et al. Cell Oncol (Dordr) (2012) pmid: 22351431
376. Tsiambas E, et al. J BUON () pmid: 20414934
377. Ansari D, et al. Br J Surg (2011) pmid: 21644238
378. Grochola LF, et al. Pancreas (2011) pmid: 21404460
379. Brown CJ, et al. Nat. Rev. Cancer (2009) pmid: 19935675
380. Joerger AC, et al. Annu. Rev. Biochem. (2008) pmid: 18410249
381. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12826609
382. Kamada R, et al. J. Biol. Chem. (2011) pmid: 20978130
383. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) pmid: 28472496
384. Yamada H, et al. Carcinogenesis (2007) pmid: 17690113
385. Bougeard G, et al. J. Clin. Oncol. (2015) pmid: 26014290
386. Sorrell AD, et al. Mol Diagn Ther (2013) pmid: 23355100
387. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) pmid: 11219776
388. Kleihues P, et al. Am. J. Pathol. (1997) pmid: 9006316
389. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
390. Lalloo F, et al. Lancet (2003) pmid: 12672316
391. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
392. Jaiswal S, et al. N. Engl. J. Med. (2014) pmid: 25426837
393. Genovese G, et al. N. Engl. J. Med. (2014) pmid: 25426838
394. Xie M, et al. Nat. Med. (2014) pmid: 25326804
395. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) pmid: 28669404
396. Severson EA, et al. Blood (2018) pmid: 29678827
397. Fuster JJ, et al. Circ. Res. (2018) pmid: 29420212
398. Hematology Am Soc Hematol Educ Program (2018) pmid: 30504320
399. Chabon JJ, et al. Nature (2020) pmid: 32269342
400. Razavi P, et al. Nat. Med. (2019) pmid: 31768066

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 17 March 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531