

ABOUT THE TEST FoundationOne®Liquid CDx is a next generation sequencing (NGS) assay that identifies clinically relevant genomic alterations in circulating cell-free DNA.

PATIENT	DISEASE Liver cholangiocarcinoma	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN ID CCL 6/20/1961
	NAME Lin, Chin-Chun		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN TYPE Blood
	DATE OF BIRTH 20 June 1961		ADDITIONAL RECIPIENT None		DATE OF COLLECTION 15 September 2022
	SEX Male		MEDICAL FACILITY ID 205872		SPECIMEN RECEIVED 21 September 2022
	MEDICAL RECORD # 21384474		PATHOLOGIST Not Provided		

Biomarker Findings

Blood Tumor Mutational Burden - 18 Muts/Mb
Microsatellite status - MSI-High Not Detected
Tumor Fraction - Elevated Tumor Fraction

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

CCNE1 amplification	KDM6A R1351*
FGFR1 amplification - equivocal†	NSD3 (WHSC1L1) amplification - equivocal†
MDM2 amplification	RAD21 R450fs*6
PTEN splice site 165-1G>C	RB1 R698S
STK11 F157fs*5	TP53 R280T
ASXL1 E635fs*15	ZNF703 amplification - equivocal†
ESR1 S341L	

† See About the Test in appendix for details.

Report Highlights

- Targeted therapies with **NCCN categories of evidence** in this tumor type: **Nivolumab** (p. 15)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 17)
- Variants that may represent **clonal hematopoiesis** and may originate from non-tumor sources: **ASXL1** E635fs*15 (p. 9)

BIOMARKER FINDINGS

Blood Tumor Mutational Burden - 18 Muts/Mb

10 Trials see p. 17

Microsatellite status - MSI-High Not Detected

Tumor Fraction - Elevated Tumor Fraction

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

Dostarlimab
 Pembrolizumab

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Nivolumab 2B
 Cemiplimab
 Nivolumab + Ipilimumab

MSI-High not detected. No evidence of microsatellite instability in this sample (see Appendix section).

Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected. There is higher sensitivity for identifying genomic alterations and a lower risk of false negative results in specimens with elevated tumor fraction; the positive percent agreement observed between liquid and tissue for defined short variants is $\geq 90\%$ (Li et al., 2021; AACR Abstract 2231) (see Biomarker Findings section).

GENOMIC FINDINGS

CCNE1 - amplification

VAF%

-

4 Trials see p. 19

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

None

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

None

☐ NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

GENOMIC FINDINGS	VAF%	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
FGFR1 - amplification - equivocal 10 Trials see p. 20	-	None	None
MDM2 - amplification 2 Trials see p. 22	-	None	None
PTEN - splice site 165-1G>C 10 Trials see p. 23	17.8%	None	None
STK11 - F157fs*5 1 Trial see p. 25	21.4%	None	None

☐ NCCN category

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

ASXL1 - E635fs*15 p. 9

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

ASXL1 - E635fs*15 p. 9	RAD21 - R450fs*6 p. 11
ESR1 - S341L p. 10	RB1 - R698S p. 11
KDM6A - R1351* p. 10	TP53 - R280T p. 12
NSD3 (WHSC1L1) - amplification - equivocal p. 10	ZNF703 - amplification - equivocal p. 13

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the therapies listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and/or exhaustive. Neither the therapies nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies. Therapies contained in this report may have been approved by the US FDA or other national authorities; however, they might not have been approved in your respective country. In the appropriate clinical context, germline testing of APC, ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MEN1, MLH1, MSH2, MSH6, MUTYH, NF1, NF2, PALB2, PMS2, POLE, PTEN, RAD51C, RAD51D, RB1, RET, SDHA, SDHB, SDHC, SDHD, SMAD4, STK11, TGFBR2, TP53, TSC1, TSC2, VHL, and WT1 is recommended.

Variant Allele Frequency is not applicable for copy number alterations.


Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01



FoundationOne®Liquid CDx
04 Oct 2022

HISTORIC PATIENT FINDINGS

ORD-1461672-01
VAF%

Blood Tumor Mutational Burden

18 Muts/Mb

Microsatellite status

MSI-High Not Detected

Tumor Fraction

24%

CCNE1	amplification	Detected
FGFR1	amplification	Detected
MDM2	amplification	Detected
PTEN	● splice site 165-1G>C	17.8%
STK11	● F157fs*5	21.4%
ASXL1	● E635fs*15	3.2%
ESR1	● S341L	16.7%
KDM6A	● R1351*	30.7%
NSD3 (WHSC1L1)	amplification	Detected
RAD21	● R450fs*6	0.31%

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

HISTORIC PATIENT FINDINGS		ORD-1461672-01 VAF%
RB1	● R698S	15.9%
TP53	● R280T	19.6%
ZNF703	amplification	Detected

NOTE This comparison table refers only to genes and biomarkers assayed by prior FoundationOne®Liquid CDx or FoundationOne®CDx tests. Up to five previous tests may be shown.

For some genes in FoundationOne Liquid CDx, only select exons are assayed. Therefore, an alteration found by a previous test may not have been confirmed despite overlapping gene lists. Please refer to the Appendix for the complete list of genes and exons assayed. The gene and biomarker list will be updated periodically to reflect new knowledge about cancer biology.

As new scientific information becomes available, alterations that had previously been listed as Variants of Unknown Significance (VUS) may become reportable.

Tissue Tumor Mutational Burden (TMB) and blood TMB (bTMB) are estimated from the number of synonymous and non-synonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.

Not Tested = not baited, not reported on test, or test preceded addition of biomarker or gene

Not Detected = baited but not detected on test

Detected = present (VAF% is not applicable)

VAF% = variant allele frequency percentage

Cannot Be Determined = Sample is not of sufficient data quality to confidently determine biomarker status

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

BIOMARKER FINDINGS

BIOMARKER

Blood Tumor Mutational Burden

RESULT

18 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased blood tumor mutational burden (bTMB) may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1³⁻⁴, and anti-PD-1/CTLA4 therapies⁵⁻⁶. A Phase 2 multi-solid-tumor trial showed that bTMB ≥ 16 Muts/Mb (as measured by this assay) was associated with improved survival from treatment with a PD-1 inhibitor alone or in combination with a CTLA-4 inhibitor⁵. In non-small cell lung cancer (NSCLC), multiple clinical trials have shown

patients with higher bTMB derive clinical benefit from immune checkpoint inhibitors following single-agent or combination treatments with either CTLA4 inhibitors or chemotherapy, with reported high bTMB cutpoints ranging from 6 Muts/Mb-16 Muts/Mb¹. In head and neck squamous cell carcinoma (HNSCC), a Phase 3 trial showed that bTMB ≥ 16 Muts/Mb (approximate equivalency ≥ 8 Muts/Mb as measured by this assay) was associated with improved survival from treatment with a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor⁷.

FREQUENCY & PROGNOSIS

Average bTMB levels in solid tumors other than NSCLC have not been evaluated (PubMed, Mar 2022). Published data investigating the prognostic implications of bTMB levels in biliary tract cancer are limited (PubMed, Jul 2022). Although cases with hypermutated biliary tract cancer were enriched in a subgroup with poor prognosis in 1 study⁸, TMB-high (≥ 10 mut/Mb) status in biliary adenocarcinoma not treated with immunotherapy

was not significantly associated with OS in another study, in which patients with TMB-high tumors experienced numerically longer OS compared with patients with TMB-low tumors (11.5 vs. 8.4 months, adjusted HR=0.65)⁹.

FINDING SUMMARY

Blood tumor mutational burden (bTMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations from circulating tumor DNA in blood. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma¹⁰⁻¹¹ and cigarette smoke in lung cancer¹²⁻¹³, treatment with temozolomide-based chemotherapy in glioma¹⁴⁻¹⁵, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes¹⁶⁻²⁰, and microsatellite instability (MSI)^{16,19-20}. This sample harbors a bTMB level that may be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents¹⁻⁴.

BIOMARKER

Tumor Fraction

RESULT

Elevated Tumor Fraction

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Specimens with elevated tumor fraction have high circulating-tumor DNA (ctDNA) content, and thus high sensitivity for identifying genomic alterations. Such specimens are at low risk of false negative results. Tumor fraction levels currently have limited implications for diagnosis, surveillance, or therapy and should not be overinterpreted or compared from one blood draw to another. There are currently no targeted approaches to address

specific tumor fraction levels. In the research setting, changes in tumor fraction estimates have been associated with treatment duration and clinical response and may be a useful indicator for future cancer management²¹⁻²⁶.

FREQUENCY & PROGNOSIS

Detectable ctDNA levels have been reported in a variety of tumor types, with higher tumor fraction levels reported for patients with metastatic (Stage 4) tumors compared with patients with localized disease (Stages 1 to 3)²⁷. Elevated tumor fraction levels have been reported to be associated with worse prognosis in a variety of cancer types, including pancreatic cancer²⁸, Ewing sarcoma and osteosarcoma²⁹, prostate cancer²⁴, breast cancer³⁰, leiomyosarcoma³¹, esophageal cancer³², colorectal cancer³³, and gastrointestinal cancer³⁴.

FINDING SUMMARY

Tumor fraction provides an estimate of the percentage of ctDNA present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate for this sample is based on the observed level of aneuploid instability. The tumor fraction algorithm utilized for FoundationOne Liquid CDx uses the allele frequencies of approximately 1,000 single-nucleotide polymorphism (SNP) sites across the genome. Unlike the maximum somatic allele frequency (MSAF) method of estimating ctDNA content³⁵, the tumor fraction metric does not take into account the allele frequency of individual variants but rather produces a more holistic estimate of ctDNA content using data from across the genome. The amount of ctDNA detected may correlate with disease burden and response to therapy³⁶⁻³⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE
CCNE1

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies that directly target CCNE1 alterations. Because amplification or overexpression of CCNE1 leads to increased genomic instability through the ATR-CHK1-WEE1 pathway³⁸⁻³⁹ and cyclin E1 promotes cell cycle progression in a complex with CDK2⁴⁰, clinical and preclinical studies have investigated inhibitors of CHK1, ATR, CDK2, and WEE1 as potential therapeutic approaches for tumors with CCNE1 activation. Clinical benefit has been reported for patients with recurrent high-grade serous ovarian carcinoma (HGSOC) with CCNE1 amplification or expression in response to treatment with the

CHK1 inhibitor prexasertib⁴¹. Studies of the WEE1 inhibitor adavosertib observed PRs in patients with CCNE1-amplified HGSOC and ovarian cancer⁴²⁻⁴³. Similarly, in a Phase 2 study of patients with CCNE1-amplified solid tumors, adavosertib elicited an ORR of 26% with PRs reported for patients with ovarian cancer, urothelial carcinoma, or melanoma⁴⁴. Preclinical studies have demonstrated that cell lines with CCNE1 amplification or overexpression were sensitive to inhibitors of ATR⁴⁵⁻⁴⁶, CDK2⁴⁷, or WEE1^{39,48}. However, other studies have shown that sensitivity of various cell lines to CDK2 inhibitors, including SNS-032, dinaciclib, and seliciclib, at clinically achievable doses, is largely independent of CCNE1 copy number or expression⁴⁹⁻⁵². One study has reported a reduction in tumor CCNE1 levels in 4/6 lung and esophageal cancer cases following treatment with the HDAC inhibitor vorinostat⁵³.

FREQUENCY & PROGNOSIS

CCNE1 amplification has been reported in 1-2% of samples in published cholangiocarcinoma

datasets⁵⁴⁻⁵⁵. Published data investigating the prognostic implications of CCNE1 alterations in biliary tract carcinoma are limited (PubMed, Mar 2022).

FINDING SUMMARY

CCNE1 encodes the protein cyclin E1, which plays a role in the regulated transition from the G1 to S phase by binding to and activating cyclin-dependent protein kinase 2 (CDK2). It also has a direct role in initiation of replication and the maintenance of genomic stability⁴⁰. Amplification of chromosomal region 19q12-q13 has been demonstrated in many types of cancer, and CCNE1 is a well-studied gene within this amplicon⁵⁶⁻⁵⁷. Increased copy number of CCNE1 is highly associated with overexpression of the cyclin E1 protein⁵⁸⁻⁵⁹. Cyclin E1 overexpression can lead to cell transformation as a result of an increase in cyclin E1 activity^{40,60}.

GENE
FGFR1

ALTERATION
amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Alterations that activate FGFR1 may predict sensitivity to selective FGFR inhibitors including erdafitinib⁶¹⁻⁶³, pemigatinib⁶⁴, infigratinib⁶⁵⁻⁶⁶, futibatinib⁶⁷⁻⁶⁹, rogaratinib⁷⁰, Debio 1347⁷¹⁻⁷², and derazantinib⁷³, or multikinase inhibitors such as pazopanib⁷⁴ and ponatinib⁷⁵⁻⁷⁷. The activity and

efficacy of selective FGFR inhibitors for FGFR1-amplified tumors has been modest with limited responses reported in FGFR1-amplified lung squamous cell carcinoma (SCC) treated with infigratinib⁷⁸ or AZD4579⁷⁹ and no responses reported among patients with FGFR1-amplified breast cancer treated with infigratinib⁷⁸. Two case studies reported PRs in patients with FGFR1-amplified breast cancer treated with pazopanib⁷⁴.

FREQUENCY & PROGNOSIS

Although FGFR1 amplification and mutation have been reported infrequently in biliary tract tumors and cholangiocarcinoma^{8,80-84}, recurrent activating FGFR2 fusions have been reported in 8-50% of

intrahepatic cholangiocarcinomas^{8,82-83,85-88}. Published data investigating the prognostic implications of FGFR1 alterations in biliary tract carcinoma or cholangiocarcinoma are limited (PubMed, Aug 2021).

FINDING SUMMARY

FGFR1 encodes the protein fibroblast growth factor receptor 1, which plays key roles in regulation of the cell cycle and angiogenesis and is an upstream regulator of the RAS, MAPK, and AKT signaling pathways⁸⁹. Amplification of FGFR1 has been correlated with protein expression⁹⁰⁻⁹¹ and may predict pathway activation and sensitivity to therapies targeting this pathway⁹²⁻⁹³.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE

MDM2

ALTERATION
amplification

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

MDM2 antagonists disrupt the MDM2-p53 interaction, thereby stabilizing p53⁹⁴. Preclinical studies have suggested that the amplification of MDM2, in the absence of concurrent TP53 mutations, may increase sensitivity to these agents⁹⁵⁻⁹⁶. Preliminary Phase 1 studies of the MDM2-p53 antagonist alrizomadlin (APG-115) reported a PR in a patient with liposarcoma harboring an MDM2 amplification and wildtype for TP53 and SD in 21%-38% (6/28 and 5/13, respectively) of patients in genomically unselected solid tumors⁹⁷⁻⁹⁸. A Phase 2 trial of alrizomadlin in combination with pembrolizumab reported a PR in 1 of 3 patients with malignant peripheral nerve sheath tumor that had failed standard therapy, as well as PRs in patients with multiple types of solid tumors that had failed immunotherapy, including 1

out of 14 patients with non-small cell lung cancer; 1 out of 5 patients with urothelial carcinoma; and 2 out of 5, 1 out of 5, and 1 out of 11 patients with mucosal, uveal, and cutaneous melanoma, respectively⁹⁹. Phase 1b studies of the MDM2 inhibitor idasanutlin for refractory AML in combination with cytarabine or venetoclax reported anti-leukemic response rates of 33% (25/75) and 37% (11/30), respectively¹⁰⁰⁻¹⁰¹; clinical benefit (58% ORR, 7/12) with idasanutlin monotherapy has been reported for patients with polycythemia vera¹⁰². The dual MDM2/MDM4 inhibitor ALRN-6924 led to an ORR of 27% (4/15) for patients with TP53 wildtype peripheral T-cell lymphoma in a Phase 2 study¹⁰³; responses have also been observed in TP53 wildtype AML, MDS, Merkel cell carcinoma, colorectal cancer, and liposarcoma¹⁰⁴⁻¹⁰⁵.

FREQUENCY & PROGNOSIS

MDM2 amplification has been observed in 10.8% of gallbladder carcinoma and 3.2% of cholangiocarcinoma samples^{54,106-107}. MDM2 overexpression has been reported to be a common event in biliary tract cancers, occurring in 75% of gallbladder adenocarcinoma and in 38-68% of ICC cases, respectively¹⁰⁸⁻¹¹¹. MDM2 overexpression in

pancreatic carcinomas and biliary tract tumors has been associated with advanced tumor stage, metastasis, and poor patient prognosis^{109-110,112}.

FINDING SUMMARY

MDM2 encodes an E3 ubiquitin protein ligase, which mediates the ubiquitination and subsequent degradation of p53, Rb1, and other proteins¹¹³⁻¹¹⁵. MDM2 acts to prevent the activity of the tumor suppressor p53; therefore, overexpression or amplification of MDM2 may be oncogenic¹¹⁶⁻¹¹⁷. Overexpression or amplification of MDM2 is frequent in cancer¹¹⁸. Although two retrospective clinical studies suggest that MDM2 amplification may predict a short time-to-treatment failure on anti-PD-1/PD-L1 immune checkpoint inhibitors, with 4/5 patients with MDM2 amplification¹¹⁹ and 2/3 patients with MDM2 or MDM4 amplification¹²⁰ experiencing tumor hyperprogression, amplification of MDM2 or MDM4 was not associated with shorter progression-free survival (PFS) in a retrospective analysis of non-small cell lung cancer (NSCLC) outcomes with immune checkpoint inhibitors (hazard ratio of 1.4, p=0.44)¹²¹. The latter study reported PFS of >2 months for 5/8 patients with MDM2/MDM4 amplification¹²¹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE

PTEN

ALTERATION

splice site 165-1G>C

TRANSCRIPT ID

NM_000314

CODING SEQUENCE EFFECT

165-1G>C

breast cancer including triple negative breast cancer¹³², ovarian cancer¹³³, uterine leiomyosarcoma¹³⁴, and endometrial cancer¹³¹ treated with PARP inhibitors. However, some studies have reported a lack of association between PTEN mutation and PARP inhibitor sensitivity¹³⁵⁻¹³⁶.

PTEN can lead to uncontrolled cell growth and suppression of apoptosis¹²³. Alterations such as seen here may disrupt PTEN function or expression¹⁴²⁻¹⁸³.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the PTEN variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with hamartoma tumor syndrome (ClinVar, Sep 2022)¹⁸⁴. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. PTEN mutations underlie several inherited disorders, collectively termed PTEN hamartoma tumor syndrome (PHTS), which include Cowden syndrome (CS) and its variant Lhermitte-Duclos disease (LD), Bannayan-Riley-Ruvalcaba syndrome (BRRS), PTEN-related Proteus syndrome (PS), and Proteus-like syndrome¹⁸⁵⁻¹⁸⁶. The mutation rate for PTEN in these disorders ranges from 20 to 85% of patients^{185,187}. The estimated incidence of Cowden syndrome is 1/200,000, which may be an underestimate due to the high variability of this disorder¹⁸⁵. Given the association between PTEN and these inherited syndromes, in the appropriate clinical context, germline testing for mutations affecting PTEN is recommended.

FREQUENCY & PROGNOSIS

PTEN mutations have been reported in 1-11% of cholangiocarcinoma samples^{81,83,137}. PTEN homozygous deletion was reported in fewer than 1% of gallbladder cancer cases¹⁰⁶. PTEN was not observed as a significantly altered gene in an analysis of biliary tract cancers⁸. In a study of 60 cholangiocarcinoma samples, 60% (36/60) showed a loss of PTEN expression¹³⁸. Loss of PTEN expression has been reported in 51.8% of gallbladder adenocarcinomas¹³⁹. Loss of PTEN has been associated with increased invasion, advanced tumor stage, and shorter survival in patients with cholangiocarcinoma^{138,140-141}, and with poor prognosis in patients with gallbladder adenocarcinoma¹³⁹.

FINDING SUMMARY

PTEN encodes an inositol phosphatase that functions as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway; loss of

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

PTEN loss or mutation leads to activation of the PI3K-AKT-mTOR pathway and may predict sensitivity to inhibitors of this pathway¹²²⁻¹²⁵. A Phase 2 trial examining treatment with the PI3K inhibitor copanlisib in combination with gemcitabine and cisplatin in biliary tract cancer reported longer response rates for patients with low PTEN expression (PFS of 8.5 months, OS of 18 months) compared with patients with high PTEN expression (PFS of 4.6 months, OS of 7.0 months); however, differences were not statistically significant ($p=0.19$)¹²⁶. Preclinical data indicate that PTEN loss or inactivation may predict sensitivity to PARP inhibitors¹²⁷⁻¹³¹, and clinical benefit has been observed for patients with PTEN-altered

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE
STK11

ALTERATION
F157fs*5

TRANSCRIPT ID
NM_000455

CODING SEQUENCE EFFECT
471_472delCT

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Increased mTOR signaling is present in LKB1-deficient tumors, suggesting therapies targeting mTOR may be relevant for tumors with STK11 alterations¹⁸⁸⁻¹⁹². Case studies have reported PRs for 2 patients with STK11-mutated pancreatic cancer following treatment with the mTOR inhibitor everolimus¹⁹³, with 1 PR observed in a PJS patient for 9 months until progression¹⁹³. However, retrospective analysis of a Phase 2 trial for patients with endometrial carcinoma found LKB1 (STK11) protein levels were not significantly correlated with response to everolimus treatment¹⁹⁴. In one

preclinical study, STK11 loss was associated with sensitivity to combination treatment including an SRC inhibitor¹⁹⁵; however, the clinical relevance of these findings has not been established.

— Potential Resistance —

STK11 alteration is associated with poorer response to immune checkpoint inhibitors for patients with NSCLC, including those with tumors harboring co-occurring KRAS mutation¹⁹⁶⁻²⁰⁸.

FREQUENCY & PROGNOSIS

STK11 mutations have been reported in 3% of biliary tract carcinomas analyzed in the COSMIC database (Mar 2022)²⁰⁹. STK11 mutation or loss have also been reported in biliary tract adenocarcinomas in the literature²¹⁰⁻²¹². Loss of STK11 protein expression in intrahepatic cholangiocarcinoma has been reported as an independent predictor of shorter OS and time to recurrence²¹³.

FINDING SUMMARY

The serine/threonine kinase STK11 (also called LKB1) activates AMPK and negatively regulates the

mTOR pathway in response to changes in cellular energy levels¹⁸⁸. LKB1 acts as a tumor suppressor in cancer, as loss of function promotes proliferation and tumorigenesis^{195,214}. Alterations such as seen here may disrupt STK11 function or expression²¹⁵⁻²²⁷.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in STK11 underlie Peutz-Jeghers syndrome (PJS), a rare autosomal dominant disorder associated with a predisposition for tumor formation²²⁸. This disorder has an estimated frequency between 1:29,000 and 1:120,000, although reported rates in the literature vary greatly²²⁸⁻²³⁰. Although gastrointestinal tumors are the most common malignancies associated with PJS, patients also exhibit an 18-fold increased risk of developing other epithelial cancers²²⁸⁻²³⁰, and individuals with this syndrome have a 30-50% risk of developing breast cancer^{228,230}. Given the association with PJS, in the appropriate clinical context testing for the presence of germline mutations in STK11 is recommended.

GENE
ASXL1

ALTERATION
E635fs*15

TRANSCRIPT ID
NM_015338

CODING SEQUENCE EFFECT
1900_1922del23

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in ASXL1.

FREQUENCY & PROGNOSIS

ASXL1 alterations occur infrequently across various solid tumor types²³¹ and are not known to act as drivers in any specific solid cancer type²³². Published data investigating the prognostic implications of ASXL1 alterations in solid tumors are limited (PubMed, May 2022). In the context of clonal hematopoiesis, ASXL1 mutations are significantly enriched in current or former smokers²³³.

FINDING SUMMARY

ASXL1 regulates epigenetic marks and transcription through interaction with polycomb complex proteins and various transcription activators and repressors²³⁴⁻²³⁶. Alterations such as seen here may disrupt ASXL1 function or expression²³⁷⁻²³⁹.

POTENTIAL CLONAL HEMATOPOIESIS

IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁴⁰⁻²⁴⁵. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²⁴⁰⁻²⁴¹. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease²⁴⁶. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{244,247-248}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE

ESR1

ALTERATION

S341L

TRANSCRIPT ID

NM_000125

CODING SEQUENCE EFFECT

1022C>T

aromatase inhibitors (AIs) that inhibit estrogen production, are approved to treat ER-positive (ER+) and/or hormone receptor-positive (HR+) breast cancer (NCCN Guidelines v4.2022). AI treatment has also been reported to provide clinical benefit in a subset of HR+ gynecologic malignancies²⁴⁹⁻²⁵³. Combinations of fulvestrant and CDK4/6 inhibitors such as abemaciclib, palbociclib, and ribociclib, have also demonstrated efficacy for patients with ESR1-mutated breast cancer²⁵⁴. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

investigating the prognostic implications of ESR1 alterations in biliary tract carcinoma are limited (PubMed, Dec 2021).

FINDING SUMMARY

ESR1 encodes estrogen receptor alpha (ER-alpha), one of the major estrogen receptor isoforms in humans. Along with co-activator proteins, the ER complex promotes transcription of genes involved in cell cycle progression and survival²⁵⁵. Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapies that directly target ER-alpha, such as selective ER modulators (SERMs) and the selective ER degrader (SERD) fulvestrant, as well as

FREQUENCY & PROGNOSIS

ESR1 mutation has not been reported in multiple cholangiocarcinoma datasets⁸⁰⁻⁸². Published data

GENE

KDM6A

ALTERATION

R1351*

TRANSCRIPT ID

NM_021140

CODING SEQUENCE EFFECT

4051C>T

alterations in cancer.

FREQUENCY & PROGNOSIS

KDM6A mutations have been reported in 3.9% of samples analyzed, with the highest incidence in tumors of the urinary tract (31%), liver (7.3%), endometrium (6.7%), salivary gland (6.0%), and pancreas (5.1%) (COSMIC, Jan 2022)²⁰⁹. KDM6A mutations or copy number alterations have also been identified in medulloblastoma (8.9%)²⁵⁶, adenoid cystic carcinoma (6.7%)²⁵⁷, and metastatic prostate cancer (10%)²⁵⁸. KDM6A inactivation has been found as a recurrent tumorigenic event in male T-cell acute lymphoblastic leukemia (T-ALL), and loss of KDM6A increased the sensitivity of T-

ALL cells to therapies targeting histone H3 lysine 27 methylation in preclinical assays²⁵⁹. However, KDM6A overexpression has been noted in breast cancer and renal cell carcinoma, and correlated with inferior prognosis in patients with breast cancer²⁶⁰⁻²⁶².

FINDING SUMMARY

KDM6A encodes a histone H3 lysine 27 demethylase UTX, which functions as a transcriptional regulator²⁶³. A significant number of inactivating KDM6A mutations have been found across multiple tumor types, suggesting a role as a tumor suppressor²⁶³.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies available to address KDM6A

GENE

NSD3 (WHSC1L1)

ALTERATION

amplification - equivocal

FREQUENCY & PROGNOSIS

In TCGA datasets, NSD3 amplification has been most frequently observed in lung squamous cell carcinoma (17%)²⁶⁴, breast invasive carcinoma (13%)²⁶⁵, bladder urothelial carcinoma (9%)²⁶⁶, and head and neck squamous cell carcinoma (9%)²⁶⁷ samples²⁶⁸⁻²⁶⁹. Amplification of at least one member of the NSD3-CHD8-BRD4 pathway has been associated with worse overall survival in ovarian high-grade serous carcinoma and endometrial cancer²⁷⁰. In endometrial cancers, amplification of this pathway was more frequent in endometrial serous and endometrioid serious-like

carcinomas compared to low-grade endometrioid endometrial adenocarcinomas²⁷⁰.

FINDING SUMMARY

NSD3, also known as WHSC1L1, encodes an enzyme that mediates histone methylation²⁷¹. NSD3 has been shown to be amplified in various cancers²⁷²⁻²⁷⁴.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in NSD3.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE

RAD21

ALTERATION

R450fs*6

TRANSCRIPT ID

NM_006265

CODING SEQUENCE EFFECT

1348delC

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies to target alterations in this gene.

FREQUENCY & PROGNOSIS

RAD21 amplifications have been reported in solid tumors, including breast cancers (7%), melanoma (5.4%), and prostate (2.4%) cancers²³¹. RAD21 overexpression has been correlated with poor prognosis in endometrial cancer²⁷⁵, breast cancer²⁷⁶⁻²⁷⁷, Ewing sarcoma²⁷⁸, and colorectal cancer (CRC), especially in KRAS-mutant CRC²⁷⁹.

FINDING SUMMARY

RAD21 encodes a protein involved in DNA double-strand break repair and sister chromatid cohesion as a part of the cohesin complex²⁸⁰⁻²⁸³. In preclinical studies, downregulation of RAD21 or other cohesin components leads to loss of expression from amplified genes, as well as amplifications themselves upon cell passaging²⁸⁴,

but also leads to an increase in deletions, insertions, and other rearrangements²⁸⁵. High RAD21 expression has also been associated with increased genomic instability²⁸⁶. Cohesin complex also organizes chromatin domains and regulates gene expression²⁸⁷⁻²⁸⁸. Both overexpression and reduction of expression of RAD21 has been reported to alter gene expression²⁸⁹. RAD21 amplification has been correlated with increased expression in breast^{276,286,290} and endometrial²⁷⁵ cancers. Other RAD21 alterations, including truncating and point mutations, have been reported in the context of cancer, but the majority have not been characterized.

GENE

RB1

ALTERATION

R698S

TRANSCRIPT ID

NM_000321

CODING SEQUENCE EFFECT

2094G>T

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of limited clinical data²⁹¹ and strong preclinical data²⁹²⁻²⁹⁵, RB1 inactivation may be associated with sensitivity to inhibitors of Aurora kinase A, particularly in small cell lung cancer (SCLC). A clinical study evaluating the Aurora kinase A inhibitor alisertib for patients with prostate cancer did not find an association between RB1 deletion and clinical benefit²⁹⁶. Other approaches to target RB1 inactivation under investigation in preclinical studies include inhibitors of BCL-2 family members²⁹⁷ and

activation of the NOTCH pathway²⁹⁸. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

FREQUENCY & PROGNOSIS

A study of 18 intrahepatic cholangiocarcinoma patients did not detect loss of heterozygosity at the chromosomal region 13q32 (where RB1 is located) in any cases²⁹⁹. However, studies have demonstrated that 12-41% of analyzed cholangiocarcinomas lack expression of Rb protein³⁰⁰⁻³⁰¹. RB1 mutations have been reported across solid tumors including small cell lung cancer (60%), bladder (13%), uterine sarcoma (10%), non-melanoma skin cancer (9.5%), and gastrointestinal neuroendocrine tumors (9.3%)³⁰². One analysis of 50 cholangiocarcinoma cases did not find a correlation between RB1 gene expression and disease metastasis³⁰³.

FINDING SUMMARY

RB1 encodes the retinoblastoma protein (Rb), a tumor suppressor and negative regulator of the cell cycle³⁰⁴⁻³⁰⁵. Although alterations such as seen here

have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the RB1 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with retinoblastoma (ClinVar, Sep 2022)¹⁸⁴. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Mutations in RB1 underlie the development of retinoblastoma (RB), a rare tumor that arises at a rate of approximately 1:20,000 live births, with nearly 5,000 new cases worldwide per year³⁰⁶. Germline mutations in RB1 account for approximately 40% of RB tumors³⁰⁷ and are associated with an increased risk of developing secondary malignancies that include soft tissue and bone sarcoma and malignant melanoma³⁰⁸⁻³⁰⁹. In the appropriate clinical context, germline testing of RB1 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

R280T

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

839G>C

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib³¹⁰⁻³¹³ or p53 gene therapy such as SGT53³¹⁴⁻³¹⁸. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype³¹⁹. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer³²⁰. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer³²¹. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone³²². In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel³²³. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck

squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations³²⁴. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring³²⁵. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage³¹⁸. Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR³²⁶. A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)³²⁷.

FREQUENCY & PROGNOSIS

Inactivation of p53, through mutation, deletion, or loss of heterozygosity (LOH), has been observed in 25-63% of gallbladder carcinomas and 10-61% of cholangiocarcinomas^{8,80-83,85,328-331}. TP53 mutations occur more frequently in tumors caused by liver fluke (*O. viverrini*) infection (40%) than in cholangiocarcinoma cases not related to infection (9%)⁸⁰. Aberrant TP53 expression, which is indicative of TP53 dysregulation, has been observed in 20-62% of gallbladder carcinomas and 25% (5/20) of cholangiocarcinomas³³²⁻³³⁴. Data regarding the prognostic significance of TP53 mutation in cholangiocarcinoma are conflicting^{107,335-342}. Overexpression of p53 protein has been associated with reduced patient survival in poorly differentiated gallbladder adenocarcinomas and biliary tract cancers³⁴³⁻³⁴⁴; however, another study did not find such a correlation³³⁷.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers¹¹⁶. Alterations such as seen here may disrupt TP53 function or expression³⁴⁵⁻³⁴⁹.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers³⁵⁰⁻³⁵², including sarcomas³⁵³⁻³⁵⁴. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000³⁵⁵ to 1:20,000³⁵⁴. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30³⁵⁶. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁴⁰⁻²⁴⁵. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²⁴⁰⁻²⁴¹. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease²⁴⁶. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{244,247-248}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

GENOMIC FINDINGS

GENE
ZNF703

ALTERATION
amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no available targeted therapies to directly address ZNF703 alterations in cancer. One preclinical study suggested that ZNF703 expression in breast cancer cell lines is associated with reduced sensitivity to tamoxifen through AKT-

mTOR activation³⁵⁷, although these findings have not been verified in the clinical setting.

FREQUENCY & PROGNOSIS

Amplification and high expression of ZNF703 has been observed in luminal B breast tumors, a subtype associated with aggressive disease progression and poor patient outcomes³⁵⁸⁻³⁶⁰. ZNF703 expression has also been linked with aggressive tumor characteristics in patients with gastric and colorectal cancers³⁶¹⁻³⁶². Putative high-level amplification of ZNF703 has been reported with the highest frequency in breast carcinoma, bladder urothelial carcinoma, uterine carcinosarcoma, lung squamous cell carcinoma

(SCC), esophageal carcinoma and head and neck SCC (5-13% of samples)(cBioPortal, 2022)²⁶⁸⁻²⁶⁹.

FINDING SUMMARY

ZNF703 encodes a transcriptional repressor that plays roles in stem cell proliferation, cell cycle progression, and other key cellular functions^{359,363}. Amplification of ZNF703 has been correlated with protein expression³⁵⁸⁻³⁵⁹. ZNF703 was established as a breast cancer oncoprotein by studies showing that ZNF703 expression resulted in transformation and increased proliferation of cultured cells^{358-359,364}, as well as increased lung metastases in a breast cancer xenograft model³⁶⁴.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Dostarlimab

Assay findings association

Blood Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Dostarlimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor response. It is FDA approved to treat patients with mismatch repair deficient recurrent or advanced endometrial cancer or solid tumors. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors⁵, blood tumor mutational burden (bTMB) ≥ 16 Muts/Mb (based on this assay) may predict sensitivity to immune checkpoint

inhibitors targeting PD-1.

SUPPORTING DATA

In the Phase 1 GARNET trial of dostarlimab as a single agent for patients with mismatch repair-deficient tumors, 1 patient with a gallbladder tumor and 1 patient with a biliary neoplasm each exhibited a CR³⁶⁵. Dostarlimab has been studied primarily in recurrent and advanced mismatch repair-deficient (dMMR) endometrial and non-endometrial cancers³⁶⁵⁻³⁶⁷. In the Phase 1 GARNET trial, single-agent dostarlimab elicited an ORR of 39% (41/106) and an immune-related ORR of 46% (50/110) for patients with non-endometrial dMMR solid tumors^{365,368}.

Pembrolizumab

Assay findings association

Blood Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Pembrolizumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved for patients with tumor mutational burden (TMB)-high (≥ 10 Muts/Mb), microsatellite instability-high (MSI-H), or mismatch repair-deficient (dMMR) solid tumors; as monotherapy for PD-L1-positive non-small cell lung cancer (NSCLC), head and neck squamous cell cancer (HNSCC), cervical cancer, or esophageal cancer; and in combination with chemotherapy for PD-L1-positive triple-negative breast cancer (TNBC) or cervical cancer. It is also approved in various treatment settings as monotherapy for patients with melanoma, HNSCC, urothelial carcinoma, hepatocellular carcinoma, Merkel cell carcinoma, cutaneous squamous cell carcinoma, endometrial carcinoma that is MSI-H or dMMR, classical Hodgkin lymphoma, or primary mediastinal large B-cell lymphoma; and in combination with chemotherapy or targeted therapy for NSCLC, HNSCC, esophageal or gastroesophageal junction cancer, renal cell carcinoma, TNBC, or endometrial carcinoma that is not MSI-H or dMMR. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors⁵, blood tumor mutational burden (bTMB) ≥ 16 Muts/Mb (based on this assay) may predict sensitivity to immune checkpoint

inhibitors targeting PD-1.

SUPPORTING DATA

A Phase 2 study of single-agent pembrolizumab reported a 5.8% (6/104) ORR for patients with advanced biliary tract cancer; a Phase 1b study for patients with PD-L1-positive advanced biliary tract cancer reported an ORR of 13% (3/23)³⁶⁹. Clinical benefit has been observed from pembrolizumab in combination with other therapies. The combination of pembrolizumab or nivolumab with the antiangiogenic multikinase inhibitor lenvatinib achieved an ORR of 21.4% (3/14) and median PFS (mPFS) of 5 months for patients with intrahepatic cholangiocarcinoma³⁷⁰, as well as an ORR of 10% (3/31) and mPFS of 6.1 months for patients with advanced biliary cancers³⁷¹. Combination of pembrolizumab with granulocyte-macrophage colony-stimulating factor (GM-CSF) achieved an ORR of 19% (5/27) for patients with advanced biliary cancer; a longer mPFS was observed for patients with intermediate or high tumor mutational burden (TMB) compared with low TMB (2.1 vs. 12.8 months, $p=0.012$)³⁷². Pembrolizumab combined with the VEGFR2-targeting antibody ramucirumab elicited an ORR of 4.2% (1/24) for patients with advanced biliary tract carcinoma; an improved median OS (11.3 vs. 6.1 months), but not mPFS (1.5 vs. 1.6 months), was reported for patients with PD-L1-positive tumors compared with those with PD-L1-negative tumors (combined positive score [CPS] $\geq 1\%$)³⁷³.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Cemiplimab

Assay findings association

Blood Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Cemiplimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved to treat patients with NSCLC with high PD-L1 expression (TPS \geq 50%), cutaneous squamous cell carcinoma (CSCC), or basal cell carcinoma (BCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors⁵, blood tumor mutational burden (bTMB) \geq 16 Muts/Mb (based on this assay) may predict sensitivity to immune checkpoint inhibitors targeting PD-1.

SUPPORTING DATA

Clinical data on the efficacy of cemiplimab for the treatment of biliary tract carcinoma are limited (PubMed, Jun 2022). Cemiplimab has been studied primarily in advanced cutaneous squamous cell carcinoma (CSCC), where it elicited a combined ORR of 48% (41/85) in Phase 1 and 2 studies³⁷⁴. A Phase 2 trial of cemiplimab in patients with basal cell carcinoma (BCC) reported ORRs of 31% (5 CRs and 21 PRs) in patients with locally advanced BCC and 21% (6 PRs) in patients with metastatic BCC³⁷⁵⁻³⁷⁶. The Phase 3 EMPOWER-Lung 1 trial for advanced non-small cell lung cancer (NSCLC) with PD-L1 expression \geq 50% reported that cemiplimab is associated with improved PFS (8.2 vs. 5.7 months), OS (not reached vs. 14.2 months), and ORR (37% vs. 21%) compared with chemotherapy³⁷⁷.

Nivolumab

Assay findings association

Blood Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response. It is FDA approved as a monotherapy in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), urothelial carcinoma, colorectal cancer (CRC), classical Hodgkin lymphoma (cHL), gastric cancer, gastroesophageal junction cancer, or esophageal adenocarcinoma or squamous cell carcinoma (ESCC). It is also approved in combination with chemotherapy to treat ESCC, in combination with cabozantinib to treat RCC, and in combination with relatlimab to treat melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors⁵, blood tumor mutational burden (bTMB) \geq 16 Muts/Mb (based on

this assay) may predict sensitivity to immune checkpoint inhibitors targeting PD-1.

SUPPORTING DATA

In the Phase 2 CheckMate 848 multi-solid-tumor trial, treatment with the PD-1 inhibitor nivolumab led to improved ORR for patients with a blood tumor mutational burden (bTMB) of 16 Muts/Mb or higher (based on this assay) compared with those with bTMB of 10 Muts/Mb or higher but $<$ 16 Muts/MB (22% [5/23] vs. 9.1% [2/22])⁵. A Phase 2 study of nivolumab for patients with advanced biliary tract cancers reported an ORR of 10.9% (5/46) and a DCR of 58.7% (27/46) by independent central review; median PFS and OS were 3.7 and 14.2 months, respectively³⁷⁸. This study reported a significant association between PD-L1 tumor cell expression $>$ 1% and improved median PFS (10.4 vs. 2.3 months, HR=0.23)³⁷⁸. A case report described an ongoing, 1+ year response to nivolumab in a patient with intrahepatic cholangiocarcinoma and a high TMB³⁷⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Nivolumab + Ipilimumab

Assay findings association

Blood Tumor Mutational Burden

18 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response, and ipilimumab is a cytotoxic T-lymphocyte antigen 4 (CTLA-4)-blocking antibody. The combination is FDA approved in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), hepatocellular carcinoma (HCC), pleural mesothelioma, and esophageal squamous cell carcinoma (ESCC). Furthermore, nivolumab is approved in combination with ipilimumab to treat patients with mismatch repair-deficient (dMMR) or microsatellite instability-high (MSI-H) colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors⁵, blood

tumor mutational burden (bTMB) of ≥ 16 Muts/Mb (based on this assay) may predict sensitivity to combination nivolumab and ipilimumab treatment.

SUPPORTING DATA

In the Phase 2 CheckMate 848 multi-solid tumor trial, combination treatment with nivolumab and ipilimumab led to improved ORR for patients with blood tumor mutational burden (bTMB) ≥ 16 Muts/Mb (based on this assay) compared with those with bTMB ≥ 10 Muts/Mb but < 16 Muts/Mb (34% [13/38] vs 12% [5/42])⁵. A Phase 2 study evaluating nivolumab combined with ipilimumab for patients with biliary cancers, including intrahepatic cholangiocarcinoma and gallbladder carcinoma, reported ORR of 23% (9/39), median PFS of 2.9 months, and median OS of 5.7 months³⁸⁰.

NOTE Genomic alterations detected may be associated with activity of certain US FDA or other specific country approved therapies; however, the therapies listed in this report may have varied evidence in the patient's tumor type. The listed therapies are not ranked in order of potential or predicted efficacy for this patient or in order of level of evidence for this patient's tumor type. The therapies listed in this report may not be complete and/or exhaustive. Furthermore, the listed therapies are limited to US FDA approved pharmaceutical drug products that are linked to a specific genomic alteration. There may also be US FDA approved pharmaceutical drug products that are not linked to a genomic alteration. Further there may also exist pharmaceutical drug products that are not approved by the US FDA or other national authorities. There may also be other treatment modalities available than pharmaceutical drug products.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS

IMPORTANT Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and should be investigated by the physician or

research staff. This is not a comprehensive list of all available clinical trials. There may also be compassionate use or early access programs available, which are not listed in this report. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials are not ranked in order of potential or predicted efficacy for this patient or

in order of level of evidence for this patient's tumor type. Clinical trials listed here may have additional enrollment criteria that may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. However, clinicaltrials.gov does not list all clinical trials that might be available.

BIOMARKER

Blood Tumor Mutational Burden

RESULT

18 Muts/Mb

RATIONALE

Increased tumor mutational burden may predict response to anti-PD-1 (alone or in combination

with anti-CTLA-4) or anti-PD-L1 immune checkpoint inhibitors.

NCT04237649
PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS

ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Sunto Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT05007106
PHASE 2

MK-7684A With or Without Other Anticancer Therapies in Participants With Selected Solid Tumors (MK-7684A-005)

TARGETS

PD-1, KIT, VEGFRs, FGFRs, PDGFRA, RET, TIGIT

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Tokyo (Japan), Kashiwa (Japan), Alaska, Adana (Turkey), Ankara (Turkey), Ramat Gan (Israel)

NCT04152018
PHASE 1

Study of PF-06940434 in Patients With Advanced or Metastatic Solid Tumors.

TARGETS

PD-1

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Seoul (Korea, Republic of), Wollongong (Australia), Washington, Arizona, Missouri, Texas

NCT03396445
PHASE 1

Safety and Pharmacokinetics Study of MK-5890 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Adults With Advanced Solid Tumors (MK-5890-001)

TARGETS

PD-1, CD27

LOCATIONS: Taipei (Taiwan), Seoul (Korea, Republic of), Be'er Sheva (Israel), Amsterdam (Netherlands), Rotterdam (Netherlands), Barcelona (Spain), Madrid (Spain), Pozuelo de Alarcon (Spain), Santiago (Chile)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
 PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Melbourne (Australia), Amsterdam (Netherlands), Ravenna (Italy)

NCT02628067
PHASE 2

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Makati (Philippines), Seoul (Korea, Republic of), Beijing (China), North Ryde (Australia), Moscow (Russian Federation), Hod Hasharon (Israel), Drammen (Norway), Glostrup (Denmark), Haar (Germany)

NCT03861793
PHASE 1/2

A Dose Escalation and Cohort Expansion Study of Subcutaneously-Administered Cytokine (ALKS 4230) as a Single Agent and in Combination With Anti-PD-1 Antibody (Pembrolizumab) in Subjects With Select Advanced or Metastatic Solid Tumors (ARTISTRY-2)

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Suwon (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Edmonton (Canada), Badalona (Spain), Rotterdam (Netherlands), Valencia (Spain), Madrid (Spain)

NCT04047862
PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS
 PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Guangdong (China), Changsha (China), Wuhan (China)

NCT04261439
PHASE 1

A Phase I/Ib Study of NIZ985 Alone and in Combination With Spartalizumab

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Chuo ku (Japan), Essen (Germany), Napoli (Italy), Leuven (Belgium), Barcelona (Spain), California, Texas

NCT04892498
PHASE 2

Hypofractionated Radiotherapy Combined With PD-1 Inhibitor Sequential GM-CSF and IL-2 for the Treatment of Advanced Refractory Solid Tumors (PRaG2.0)

TARGETS
 PD-1

LOCATIONS: Hangzhou (China), Suzhou (China), Wuxi (China), Hefei (China), Xuzhou (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
GENE
CCNE1
RATIONALE

Strong preclinical and clinical data suggest that CCNE1 amplification may predict sensitivity to

WEE1 inhibitors.

ALTERATION

amplification

NCT04768868
PHASE 1

The Safety and Pharmacokinetics Preliminary Efficacy of IMP7068 in Patients With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Shanghai (China), Wuhan (China), Beijing (China), Chengdu (China), Kansas, Texas

NCT05128825
PHASE 2

A Study of ZN-c3 in Subjects With Malignant Tumors

TARGETS
WEE1
LOCATIONS: Nevada, Colorado, Texas, Ohio, Pennsylvania, Maryland, Virginia

NCT03968653
PHASE 1

Study of Oral Debio 0123 in Combination With Carboplatin in Participants With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Groningen (Netherlands), Nijmegen (Netherlands), Leiden (Netherlands), Barcelona (Spain)

NCT05109975
PHASE 1

A Study to Evaluate Safety and Preliminary Anti-tumor Activity of Debio 0123 as Monotherapy in Adult Participants With Advanced Solid Tumors

TARGETS
WEE1
LOCATIONS: Bellinzona (Switzerland), Zürich (Switzerland), Michigan, Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
GENE
FGFR1
RATIONALE

FGFR inhibitors may be relevant in tumors with alterations that activate FGFR1.

ALTERATION

amplification - equivocal

NCT05024214
PHASE 1/2

Phase Ib/II Trial of Envafolelimab Plus Lenvatinib for Subjects With Solid Tumors

TARGETS

PD-L1, FGFRs, RET, PDGFRA, VEGFRs, KIT, FLT3, CSF1R

LOCATIONS: Hangzhou (China), Shanghai (China), Dongguan (China), Guangzhou (China), Zhuhai (China), Benbu (China), Zhengzhou (China), Jinan (China), Dalian (China), Tianjin (China)

NCT05254847
PHASE 2

Capecitabine Combined With Lenvatinib and Tislelizumab as Adjuvant Treatment After Resection in Patients With BTC

TARGETS

FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Shanghai (China)

NCT05156788
PHASE 2

Tislelizumab (Anti PD-1), Lenvatinib and GEMOX Transformation in the Treatment of Potentially Resectable, Locally Advanced Biliary Tract Cancer

TARGETS

FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Shanghai (China)

NCT05010668
PHASE 2

Cryoablation Combined With Sintilimab Plus Lenvatinib in Patients With Advanced Intrahepatic Cholangiocarcinoma

TARGETS

FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Shanghai (China)

NCT04169672
PHASE 2

Study of Surufatinib Combined With Toripalimab in Patients With Advanced Solid Tumors

TARGETS

FGFR1, CSF1R, VEGFRs, PD-1

LOCATIONS: Shanghai (China), Beijing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
NCT05010681
PHASE 2

Lenvatinib Plus Sintilimab in Patients With Immune Checkpoint Inhibitor Previously Treated Advanced Liver Cancer

TARGETS
 PD-1, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Shanghai (China)

NCT05098847
PHASE 2

Cryoablation Combined With Sintilimab Plus Lenvatinib In Previously Treated Unresectable Liver Metastasis From Solid Tumors

TARGETS
 FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Shanghai (China)

NCT04550624
PHASE 2

Pembrolizumab in Combination With Lenvatinib in Patients With Advanced Cholangiocarcinoma

TARGETS
 PD-1, KIT, VEGFRs, FGFRs, PDGFRA, RET

LOCATIONS: Shanghai (China)

NCT05215665
PHASE NULL

GEMOX Combined With Targeted Therapy and Immunotherapy for Patients With Advanced Cholangiocarcinoma

TARGETS
 FGFRs, RET, PDGFRA, VEGFRs, KIT, PD-1

LOCATIONS: Tianjin (China)

NCT04976634
PHASE 2

Pembrolizumab Plus Lenvatinib in Combination With Belzutifan in Solid Tumors (MK-6482-016)

TARGETS
 HIF2a, PD-1, KIT, VEGFRs, FGFRs, PDGFRA, RET

LOCATIONS: Seoul (Korea, Republic of), Gosford (Australia), Westmead (Australia), Epping (Australia), Malvern (Australia), Haifa (Israel), Jerusalem (Israel), Ramat Gan (Israel), Tel Aviv (Israel), Utrecht (Netherlands)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
GENE
MDM2
ALTERATION

amplification

RATIONALE

Inhibitors of the MDM2-p53 interaction are being tested in clinical trials. Overexpression or

amplification of MDM2 may increase sensitivity to these agents, but more data are required.

NCT04785196
PHASE 1/2

APG-115 in Combination With PD-1 Inhibitor in Patients With Advanced Liposarcoma or Advanced Solid Tumors

TARGETS
 PD-1, MDM2

LOCATIONS: Shanghai (China), Guangzhou (China)

NCT03611868
PHASE 1/2

A Study of APG-115 in Combination With Pembrolizumab in Patients With Metastatic Melanomas or Advanced Solid Tumors

TARGETS
 MDM2, PD-1

LOCATIONS: Brisbane (Australia), South Brisbane (Australia), Bedford Park (Australia), Heidelberg (Australia), California, Arizona, Missouri, Arkansas, Ohio, Pennsylvania

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
GENE
PTEN
ALTERATION

splice site 165-1G>C

RATIONALE

PTEN loss or inactivating mutations may lead to increased activation of the PI3K-AKT-mTOR pathway and may indicate sensitivity to inhibitors

of this pathway. PTEN loss or inactivation may also predict sensitivity to PARP inhibitors.

NCT04644068
PHASE 1/2

Study of AZD5305 as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Malignancies

TARGETS

ERBB2, TROP2, PARP

LOCATIONS: Shanghai (China), Seoul (Korea, Republic of), Chongqing (China), Chuo-ku (Japan), Koto-ku (Japan), Melbourne (Australia), Warszawa (Poland), Gdynia (Poland), Grzebnica (Poland), Budapest (Hungary)

NCT04341259
PHASE 1

A Study Of The Pharmacokinetics And Safety Of Ipatasertib In Chinese Participants With Locally Advanced Or Metastatic Solid Tumors.

TARGETS

AKTs

LOCATIONS: Shanghai City (China)

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS

ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT04001569
PHASE 1/2

AZD8186 and Paclitaxel in Advanced Gastric Cancer

TARGETS

PI3K-beta

LOCATIONS: Seongnam-si (Korea, Republic of)

NCT04298021
PHASE 2

DDR-Umbrella Study of DDR Targeting Agents in Advanced Biliary Tract Cancer

TARGETS

PD-L1, ATR, PARP

LOCATIONS: Seoul (Korea, Republic of)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
NCT05035745
PHASE 1/2

Selinexor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
 XPO1, PARP

LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS
 CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT04497116
PHASE 1/2

Study of RP-3500 in Advanced Solid Tumors

TARGETS
 ATR, PARP

LOCATIONS: Copenhagen (Denmark), Newcastle Upon Tyne (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Illinois, Toronto (Canada), Massachusetts, Rhode Island, New York, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

CLINICAL TRIALS
GENE
STK11
RATIONALE

 Increased mTOR signaling is present in
 LKB1-deficient tumors, suggesting therapies

 targeting mTOR may be relevant for tumors with
 STK11 alterations.

ALTERATION

F157fs*5

NCT04337463
PHASE NULL

ATG-008 Combined With Toripalimab in Advanced Solid Tumors

TARGETS

mTORC1, mTORC2, PD-1

LOCATIONS: Chongqing (China), Chengdu (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ALOX12B
 C341R

ATR
 I1145T and L196F

CUL4A
 E277K

ERBB4
 P98L

ERRFI1
 V359D

FGFR4
 L546P

GRM3
 E55K

MLL2
 G872R

MST1R
 D172N

PTPRO
 E722K

RARA
 Y2F

SMAD4
 H92Y and P320L

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

ABL1 Exons 4-9	ACVR1B	AKT1 Exon 3	AKT2	AKT3	ALK Exons 20-29, Introns 18, 19	ALOX12B	AMER1 (FAM123B or WTX)	APC
AR	ARAF Exons 4, 5, 7, 11, 13, 15, 16	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BCR* Introns 8, 13, 14	BRAF Exons 11-18, Introns 7-10	BRCA1 Introns 2, 7, 8, 12, 16, 19, 20	BRCA2 Intron 2	BRD4	BRIP1	BTG1
BTG2	BTK Exons 2, 15	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD70	CD74* Introns 6-8	CD79A	CD79B	CD274 (PD-L1)	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1 Exon 3	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2 Exons 5, 17, 18	DIS3	DNMT3A	DOT1L	EED	EGFR Introns 7, 15, 24-27	EMSY (C11orf30)	EP300	EPHA3
EPHB1	EPHB4	ERBB2	ERBB3 Exons 3, 6, 7, 8, 10, 12, 20, 21, 23, 24, 25	ERBB4	ERCC4	ERG	ERRF1	ESR1 Exons 4-8
ETV4* Intron 8	ETV5* Introns 6, 7	ETV6* Introns 5, 6	EWSR1* Introns 7-13	EZH2 Exons 4, 16, 17, 18	EZR* Introns 9-11	FANCA	FANCC	FANCG
FANCL	FAS	FBXW7	FGF10	FGF12	FGF14	FGF19	FGF23	FGF3
FGF4	FGF6	FGFR1 Introns 1, 5, Intron 17	FGFR2 Intron 1, Intron 17	FGFR3 Exons 7, 9 (alternative designation exon 10), 14, 18, Intron 17	FGFR4	FH	FLCN	FLT1
FLT3 Exons 14, 15, 20	FOXL2	FUBP1	GABRA6	GATA3	GATA4	GATA6	GID4 (C17orf39)	GNA11 Exons 4, 5
GNA13	GNAQ Exons 4, 5	GNAS Exons 1, 8	GRM3	GSK3B	H3-3A (H3F3A)	HDAC1	HGF	HNFI1A
HRAS Exons 2, 3	HSD3B1	ID3	IDH1 Exon 4	IDH2 Exon 4	IGF1R	IKBKE	IKZF1	INPP4B
IRF2	IRF4	IRS2	JAK1	JAK2 Exon 14	JAK3 Exons 5, 11, 12, 13, 15, 16	JUN	KDM5A	KDM5C
KDM6A	KDR	KEAP1	KEL	KIT Exons 8, 9, 11, 12, 13, 17, Intron 16	KLHL6	KMT2A (MLL) Introns 6, 8-11, Intron 7	KMT2D (MLL2)	KRAS

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1461672-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

LTK	LYN	MAF	MAP2K1 (MEK1) Exons 2, 3	MAP2K2 (MEK2) Exons 2-4, 6, 7	MAP2K4	MAP3K1	MAP3K13	MAPK1
MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MERTK	MET	MITF
MKNK1	MLH1	MPL Exon 10	MRE11 (MRE11A)	MSH2 Intron 5	MSH3	MSH6	MST1R	MTAP
MTOR Exons 19, 30, 39, 40, 43-45, 47, 48, 53, 56	MUTYH	MYB* Intron 14	MYC Intron 1	MYCL (MYCL1)	MYCN	MYD88 Exon 4	NBN	NF1
NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2 Intron 26	NOTCH3	NPM1 Exons 4-6, 8, 10	NRAS Exons 2, 3
NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1 Exons 14, 15, Introns 8-11	NTRK2 Intron 12	NTRK3 Exons 16, 17	NUTM1* Intron 1	P2RY8	PALB2
PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA Exons 12, 18, Introns 7, 9, 11	PDGFRB Exons 12-21, 23 9, 11
PDK1	PIK3C2B	PIK3C2G	PIK3CA Exons 2, 3, 5-8, 10, 14, 19, 21 (Coding Exons 1, 2, 4-7, 9, 13, 18, 20)	PIK3CB	PIK3R1	PIM1	PMS2	POLD1
POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PRKN (PARK2)	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1 Exons 3, 4, 6, 7, 10, 14, 15, 17, Introns 4-8	RARA Intron 2	RB1	RBM10	REL	RET Introns 7, 8, Exons 11, 13-16, Introns 9-11
RICTOR	RNF43	ROS1 Exons 31, 36-38, 40, Introns 31-35	RPTOR	RSP02* Intron 1	SDC4* Intron 2	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SLC34A2* Intron 4	SMAD2	SMAD4	SMARCA4	SMARCB1
SMO	SNCAIP	SOCS1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2
STAT3	STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TERC* ncRNA	TERT* Promoter
TET2	TGFBR2	TIPARP	TMPRSS2* Introns 1-3	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2
TYRO3	U2AF1	VEGFA	VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite (MS) status

Blood Tumor Mutational Burden (bTMB)

Tumor Fraction

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX

About FoundationOne® Liquid CDx

FoundationOne Liquid CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplastraat 3, 2440 Geel, Belgium. The CE-IVD regulatory status of FoundationOne Liquid CDx is applicable in countries that accept and/or recognize the CE mark.



ABOUT FOUNDATIONONE LIQUID CDx

FoundationOne Liquid CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Liquid CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details.

INTENDED USE

FoundationOne Liquid CDx is a next generation sequencing based *in vitro* diagnostic device that analyzes 324 genes. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The test also detects the genomic signatures blood tumor mutational burden (bTMB), microsatellite instability (MSI), and tumor fraction. FoundationOne Liquid CDx utilizes circulating cell-free DNA (cfDNA) isolated from plasma derived from the anti-coagulated peripheral whole blood of cancer patients. The test is intended to be used as a companion diagnostic to identify patients who may benefit from treatment with targeted therapies in accordance with the approved therapeutic product labeling. Additionally, FoundationOne Liquid CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with malignant neoplasms.

TEST PRINCIPLES

The FoundationOne Liquid CDx assay is performed exclusively as a laboratory service using circulating cell-free DNA (cfDNA) isolated from plasma derived from anti-coagulated peripheral whole blood from patients with solid malignant neoplasms. The assay employs a single DNA extraction method to obtain cfDNA from plasma from whole blood. Extracted

cfDNA undergoes whole-genome shotgun library construction and hybridization-based capture of 324 cancer-related genes including coding exons and select introns of 309 genes, as well as only select intronic regions or non-coding regions of 15 genes. Hybrid-capture selected libraries are sequenced with deep coverage using the NovaSeq® 6000 platform. Sequence data are processed using a customized analysis pipeline designed to accurately detect genomic alterations, including base substitutions, indels, select copy number variants, and select genomic rearrangements. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The assay also reports tumor fraction, and genomic signatures including MSI and bTMB. A subset of targeted regions in 75 genes is baited for increased sensitivity.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. *Note:* A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

QUALIFIED ALTERATION CALLS (EQUIVOCAL)

All equivocal calls, regardless of alteration type, imply that there is adequate evidence to call the alteration with confidence. However, the repeatability of equivocal calls may be lower than non-equivocal calls.

RANKING OF THERAPIES AND CLINICAL TRIALS

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

LIMITATIONS

1. For *in vitro* diagnostic use.
2. For prescription use only. This test must be ordered by a qualified medical professional in accordance with clinical laboratory regulations.
3. A negative result does not rule out the presence of a mutation below the limits of detection of the assay. Patients for whom no companion diagnostic alterations are detected should be considered for confirmation with an appropriately validated tumor tissue test, if available.
4. The FoundationOne Liquid CDx assay does not detect heterozygous deletions.
5. The test is not intended to provide information on cancer predisposition.
6. Performance has not been validated for cfDNA input below the specified minimum input.
7. Tissue TMB and blood TMB (bTMB) are estimated from the number of synonymous and nonsynonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.
8. Tumor fraction is the percentage of circulating tumor DNA (ctDNA) present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate is computationally derived from the observed level of aneuploidy in the sample. Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected and is significantly distinct from that typically found in non-tumor samples.
9. Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the tumor genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor. The MSI algorithm is based on genome wide analysis of 1765 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines for solid tissue testing.
10. Genomic findings from circulating cell-free DNA (cfDNA) may originate from circulating tumor DNA fragments, germline alterations, or non-tumor somatic alterations, such as clonal hematopoiesis of indeterminate potential (CHIP). Genes with alterations that may be derived from CHIP include, but are not limited to: *ASXL1*, *ATM*, *CBL*, *CHEK2*, *DNMT3A*, *JAK2*,

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX

About FoundationOne® Liquid CDx

KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, TP53, and U2AF1.

11. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. If a reported alteration is suspected to be germline, confirmatory testing should be considered in the appropriate clinical context.
12. The test is not intended to replace germline testing or to provide information about cancer predisposition.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >30%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MLH1, MSH2, MSH6, MUTYH, PALB2, PMS2, POLE, RAD51C, RAD51D, RET, SDHA, SDHB, SDHC, SDHD, TSC2, and VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1, ATM, CBL, CHEK2, DNMT3A, IDH2, JAK2, KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, and U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This report makes no promises or guarantees that a particular drug will be effective in the treatment of

disease in any patient. This report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Liquid CDx.

TREATMENT DECISIONS ARE THE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test or the information contained in this report.

Certain sample of variant characteristics may result in reduced sensitivity. These include: low sample quality, deletions and insertions >40bp, or repetitive/high homology sequences. FoundationOne Liquid CDx is performed using cell-free DNA, and as such germline events may not be reported.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX

About FoundationOne®Liquid CDx

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
Muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.1.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1461672-01

APPENDIX
References

1. Gandara DR, et al. Nat. Med. (2018) PMID: 30082870
2. Wang Z, et al. JAMA Oncol (2019) PMID: 30816954
3. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
4. Aggarwal C, et al. Clin. Cancer Res. (2020) PMID: 32102950
5. Schenker et al., 2022; AACR Abstract CT022
6. Saori et al., 2021; ESMO Abstract 80P
7. Li et al., 2020; ASCO Abstract 6511
8. Nakamura H, et al. Nat. Genet. (2015) PMID: 26258846
9. Shao C, et al. JAMA Netw Open (2020) PMID: 33119110
10. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
11. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
12. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
13. Rizvi NA, et al. Science (2015) PMID: 25765070
14. Johnson BE, et al. Science (2014) PMID: 24336570
15. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
16. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
17. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
18. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
19. Nature (2012) PMID: 22810696
20. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
21. Bronkhorst AJ, et al. Biomol Detect Quantif (2019) PMID: 30923679
22. Raja R, et al. Clin. Cancer Res. (2018) PMID: 30093454
23. Hrebien S, et al. Ann. Oncol. (2019) PMID: 30860573
24. Choudhury AD, et al. JCI Insight (2018) PMID: 30385733
25. Goodall J, et al. Cancer Discov (2017) PMID: 28450425
26. Goldberg SB, et al. Clin. Cancer Res. (2018) PMID: 29330207
27. Bettgowda C, et al. Sci Transl Med (2014) PMID: 24553385
28. Lapin M, et al. J Transl Med (2018) PMID: 30400802
29. Shulman DS, et al. Br. J. Cancer (2018) PMID: 30131550
30. Stover DG, et al. J. Clin. Oncol. (2018) PMID: 29298117
31. Hemming ML, et al. JCO Precis Oncol (2019) PMID: 30793095
32. Egyud M, et al. Ann. Thorac. Surg. (2019) PMID: 31059681
33. Fan G, et al. PLoS ONE (2017) PMID: 28187169
34. Vu et al., 2020; DOI: 10.1200/PO.19.00204
35. Li G, et al. J Gastrointest Oncol (2019) PMID: 31602320
36. Zhang EW, et al. Cancer (2020) PMID: 32757294
37. Butler TM, et al. Cold Spring Harb Mol Case Stud (2019) PMID: 30833418
38. Lin AB, et al. Clin. Cancer Res. (2017) PMID: 28331049
39. Chen X, et al. Clin Cancer Res (2018) PMID: 30181387
40. Mörry T, et al. Int. J. Biochem. Cell Biol. (2004) PMID: 15147722
41. Lee JM, et al. Lancet Oncol. (2018) PMID: 29361470
42. Lheureux S, et al. Lancet (2021) PMID: 33485453
43. Oza AM, et al. Clin Cancer Res (2020) PMID: 32611648
44. Fu et al., 2021; AACR abstract 974
45. Toledo LI, et al. Nat. Struct. Mol. Biol. (2011) PMID: 21552262
46. Buisson R, et al. Mol. Cell (2015) PMID: 26365377
47. Yang L, et al. Oncotarget (2015) PMID: 26204491
48. Kok YP, et al. Oncogenesis (2020) PMID: 33028815
49. Taylor-Harding B, et al. Oncotarget (2015) PMID: 25557169
50. Etemadmoghadam D, et al. Clin. Cancer Res. (2013) PMID: 24004674
51. Scaltriti M, et al. Proc. Natl. Acad. Sci. U.S.A. (2011) PMID: 21321214
52. Nanos-Webb A, et al. Breast Cancer Res. Treat. (2012) PMID: 21695458
53. Ma T, et al. Mol. Cancer Ther. (2013) PMID: 23686769
54. Lowery MA, et al. Clin. Cancer Res. (2018) PMID: 29848569
55. Sanchez-Vega F, et al. Cell (2018) PMID: 29625050
56. Leung SY, et al. Mod. Pathol. (2006) PMID: 16575401
57. Lin L, et al. Cancer Res. (2000) PMID: 11156406
58. Mayr D, et al. Am. J. Clin. Pathol. (2006) PMID: 16753589
59. Nakayama N, et al. Cancer (2010) PMID: 20336784
60. Stamatakis M, et al. World J Surg Oncol (2010) PMID: 21176227
61. Lorient Y, et al. N. Engl. J. Med. (2019) PMID: 31340094
62. Tabernero J, et al. J. Clin. Oncol. (2015) PMID: 26324363
63. Karkera JD, et al. Mol. Cancer Ther. (2017) PMID: 28416604
64. Necchi et al., 2018; ESMO Abstract 900P
65. Pal SK, et al. Cancer Discov (2018) PMID: 29848605
66. Pal SK, et al. Cancer (2020) PMID: 32208524
67. Bahleda R, et al. Ann Oncol (2020) PMID: 32622884
68. Meric-Bernstam F, et al. Cancer Discov (2022) PMID: 34551969
69. Kasbekar M, et al. Blood Adv (2020) PMID: 32649766
70. Schuler M, et al. Lancet Oncol. (2019) PMID: 31405822
71. Farouk Sait S, et al. JCO Precis Oncol (2021) PMID: 34250399
72. Voss MH, et al. Clin. Cancer Res. (2019) PMID: 30745300
73. Papadopoulos KP, et al. Br. J. Cancer (2017) PMID: 28972963
74. Cheng FT, et al. J Natl Compr Canc Netw (2017) PMID: 29223982
75. Khodadoust MS, et al. Leukemia (2016) PMID: 26055304
76. Tanasi I, et al. Blood (2019) PMID: 31434701
77. Strati P, et al. Leuk. Lymphoma (2018) PMID: 29119847
78. Nogova L, et al. J. Clin. Oncol. (2017) PMID: 27870574
79. Aggarwal C, et al. J Thorac Oncol (2019) PMID: 31195180
80. Chan-On W, et al. Nat. Genet. (2013) PMID: 24185513
81. Ong CK, et al. Nat. Genet. (2012) PMID: 22561520
82. Jiao Y, et al. Nat. Genet. (2013) PMID: 24185509
83. Ross JS, et al. Oncologist (2014) PMID: 24563076
84. Lee H, et al. J. Clin. Pathol. (2016) PMID: 26500333
85. Javle M, et al. Cancer (2016) PMID: 27622582
86. Borad MJ, et al. PLoS Genet. (2014) PMID: 24550739
87. Arai Y, et al. Hepatology (2014) PMID: 24122810
88. Sia D, et al. Nat Commun (2015) PMID: 25608663
89. Turner N, et al. Nat. Rev. Cancer (2010) PMID: 20094046
90. Kohler LH, et al. Virchows Arch. (2012) PMID: 22648708
91. Kim HR, et al. J. Clin. Oncol. (2013) PMID: 23182986
92. André F, et al. Lancet Oncol. (2014) PMID: 24508104
93. Dienstmann R, et al. Ann. Oncol. (2014) PMID: 24265351
94. Cheok CF, et al. Nat Rev Clin Oncol (2011) PMID: 20975744
95. Ohnstad HO, et al. Cancer (2013) PMID: 23165797
96. Gamble LD, et al. Oncogene (2012) PMID: 21725357
97. Zhang et al., 2019; ASCO Abstract 3124
98. Rasco et al., 2019; ASCO Abstract 3126
99. Tolcher et al., 2021; ASCO Abstract 2506
100. Martinelli et al., 2016; EHA21 Abstract S504
101. Daver et al., 2018; ASH Abstract 767
102. Mascarenhas et al., 2019; ASH Abstract 134
103. Shustov et al., 2018; ASH Abstract 1623
104. Sallman et al., 2018; ASH Abstract 4066
105. Meric-Bernstam et al., 2017; ASCO Abstract 2505
106. Narayan RR, et al. Cancer (2019) PMID: 30427539
107. Boerner T, et al. Hepatology (2021) PMID: 33765338
108. Momoi H, et al. J. Hepatol. (2001) PMID: 11580146
109. Horie S, et al. Virchows Arch. (2000) PMID: 10963376
110. Jarnagin WR, et al. J. Clin. Oncol. (2006) PMID: 16505435
111. Della Torre G, et al. Diagn. Mol. Pathol. (2000) PMID: 10718212
112. Grochola LF, et al. Pancreas (2011) PMID: 21404460
113. Sdek P, et al. Mol. Cell (2005) PMID: 16337594
114. Brady M, et al. Mol. Cell. Biol. (2005) PMID: 15632057
115. Li M, et al. Mol. Cell (2004) PMID: 15053880
116. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
117. Cordon-Cardo C, et al. Cancer Res. (1994) PMID: 8306343
118. Beroukhi R, et al. Nature (2010) PMID: 20164920
119. Kato S, et al. Clin. Cancer Res. (2017) PMID: 28351930
120. Singavi et al., 2017; ESMO Abstract 1140PD
121. Rizvi H, et al. J. Clin. Oncol. (2018) PMID: 29337640
122. Courtney KD, et al. J. Clin. Oncol. (2010) PMID: 20085938
123. Simpson L, et al. Exp. Cell Res. (2001) PMID: 11237521
124. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
125. Milella M, et al. Sci Rep (2017) PMID: 28220839
126. Tan ES, et al. Cancer (2021) PMID: 33289918
127. Mendes-Pereira AM, et al. EMBO Mol Med (2009) PMID: 20049735
128. Shen Y, et al. Clin. Cancer Res. (2013) PMID: 23881923
129. Chatterjee P, et al. PLoS ONE (2013) PMID: 23565244
130. McCormick A, et al. Int. J. Gynecol. Cancer (2016) PMID: 26905328
131. Forster MD, et al. Nat Rev Clin Oncol (2011) PMID: 21468130
132. Eikesdal HP, et al. Ann Oncol (2021) PMID: 33242536
133. Dougherty et al., 2014; ASCO Abstract 5536
134. Pan M, et al. Perm J (2021) PMID: 33970096
135. Sandhu SK, et al. Lancet Oncol. (2013) PMID: 23810788
136. Romero I, et al. Gynecol Oncol (2020) PMID: 32988624
137. Jang S, et al. Mod. Pathol. (2014) PMID: 24186137
138. Kim JH, et al. Gut Liver (2012) PMID: 22844568
139. Liu DC, et al. Pathol. Res. Pract. (2011) PMID: 21719208
140. Lee D, et al. Mod. Pathol. (2012) PMID: 21874010
141. Chung JY, et al. Clin. Cancer Res. (2009) PMID: 19147772
142. Campbell RB, et al. J. Biol. Chem. (2003) PMID: 12857747
143. Rodríguez-Escudero I, et al. Hum. Mol. Genet. (2011) PMID: 21828076
144. He X, et al. Cancer Res. (2013) PMID: 23475934
145. Han SY, et al. Cancer Res. (2000) PMID: 10866302
146. Myers MP, et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PMID: 9811831
147. Pradella LM, et al. BMC Cancer (2014) PMID: 24498881
148. Kim JS, et al. Mol. Cell. Biol. (2011) PMID: 21536651
149. Denning G, et al. Oncogene (2007) PMID: 17213812
150. Hlobilkova A, et al. Anticancer Res. (2010) PMID: 16619501
151. Redfern RE, et al. Protein Sci. (2010) PMID: 20718038
152. Shenoy S, et al. PLoS ONE (2012) PMID: 22505997
153. Wang Y, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19329485
154. Okumura K, et al. J. Biol. Chem. (2006) PMID: 16829519
155. Lee JO, et al. Cell (1999) PMID: 10555148
156. Maxwell GL, et al. Cancer Res. (1998) PMID: 9635567
157. Risinger JI, et al. Clin. Cancer Res. (1998) PMID: 9865913

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1461672-01**
APPENDIX
References

158. Kato H, et al. Clin. Cancer Res. (2000) PMID: 11051241
159. Fenton TR, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22891331
160. Ngeow J, et al. J. Clin. Endocrinol. Metab. (2012) PMID: 23066114
161. Lobo GP, et al. Hum. Mol. Genet. (2009) PMID: 19457929
162. Liu J, et al. Oncogene (2014) PMID: 23995781
163. Maehama T, et al. Annu. Rev. Biochem. (2001) PMID: 11395408
164. De Vivo I, et al. J. Med. Genet. (2000) PMID: 10807691
165. Ramaswamy S, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10051603
166. Liu JL, et al. Mol. Cell. Biol. (2005) PMID: 15988030
167. Karoui M, et al. Br. J. Cancer (2004) PMID: 2526806
168. Gil A, et al. PLoS ONE (2015) PMID: 25875300
169. Furnari FB, et al. Cancer Res. (1998) PMID: 9823298
170. Spinelli L, et al. J. Med. Genet. (2015) PMID: 25527629
171. Mingo J, et al. Eur. J. Hum. Genet. (2018) PMID: 29706633
172. Wang Q, et al. J. Mol. Graph. Model. (2010) PMID: 20538496
173. Andrés-Pons A, et al. Cancer Res. (2007) PMID: 17942903
174. Butler MG, et al. J. Med. Genet. (2005) PMID: 15805158
175. Georgescu MM, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10468583
176. Staal FJ, et al. Br. J. Cancer (2002) PMID: 12085208
177. Nguyen HN, et al. Oncogene (2014) PMID: 24292679
178. Rahdar M, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19114656
179. Das S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12808147
180. Wang X, et al. Biochem. J. (2008) PMID: 18498243
181. Valiente M, et al. J. Biol. Chem. (2005) PMID: 15951562
182. Nguyen HN, et al. Oncogene (2015) PMID: 25263454
183. Shan L, et al. Cell Discov (2020) PMID: 32704382
184. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
185. Blumenthal GM, et al. Eur. J. Hum. Genet. (2008) PMID: 18781191
186. Orloff MS, et al. Oncogene (2008) PMID: 18794875
187. Zbuk KM, et al. Nat. Rev. Cancer (2007) PMID: 17167516
188. Shaw RJ, et al. Cancer Cell (2004) PMID: 15261145
189. Ji H, et al. Nature (2007) PMID: 17676035
190. Contreras CM, et al. Cancer Res. (2008) PMID: 18245476
191. Gurumurthy S, et al. Cancer Res. (2008) PMID: 18172296
192. Shackelford DB, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19541609
193. Klumpen HJ, et al. J. Clin. Oncol. (2011) PMID: 21189378
194. Trédan O, et al. Target Oncol (2013) PMID: 23238879
195. Carretero J, et al. Cancer Cell (2010) PMID: 20541700
196. Skoulidis F, et al. Cancer Discov (2018) PMID: 29773717
197. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
198. Scheel AH, et al. Oncoimmunology (2016) PMID: 27467949
199. Skoulidis F, et al. Cancer Discov (2015) PMID: 26069186
200. Koyama S, et al. Cancer Res. (2016) PMID: 26833127
201. Schabath MB, et al. Oncogene (2016) PMID: 26477306
202. Dong ZY, et al. Clin. Cancer Res. (2017) PMID: 28039262
203. Kadara H, et al. Ann. Oncol. (2017) PMID: 27687306
204. Herter-Sprie GS, et al. JCI Insight (2016) PMID: 27699275
205. Skoulidis et al., 2017; IASLC 17th World Congress on Lung Cancer Abstract MA04.07
206. Jure-Kunkel et al., 2018; ASCO Abstract 3028
207. Stephens, 2017; AACR Abstract SY40-02
208. Skoulidis et al., 2015; ASCO Abstract 11002
209. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
210. Olschwang S, et al. J. Med. Genet. (2001) PMID: 11389158
211. Su GH, et al. Am. J. Pathol. (1999) PMID: 10362809
212. Sahin F, et al. Mod. Pathol. (2003) PMID: 12861065
213. Wang J, et al. Oncotarget (2015) PMID: 26056085
214. Ollila S, et al. J Mol Cell Biol (2011) PMID: 21926085
215. Qiu W, et al. Oncogene (2006) PMID: 16407837
216. Mehenni H, et al. Am. J. Hum. Genet. (1998) PMID: 9837816
217. Karuman P, et al. Mol. Cell (2001) PMID: 11430832
218. Baas AF, et al. EMBO J. (2003) PMID: 12805220
219. Zeng PY, et al. Cancer Res. (2006) PMID: 17108107
220. Boudeau J, et al. J. Cell. Sci. (2004) PMID: 15561763
221. Scott KD, et al. Cancer Res. (2007) PMID: 17575127
222. Xie Z, et al. Mol. Cell. Biol. (2009) PMID: 19414597
223. Boudeau J, et al. Hum. Mutat. (2003) PMID: 12552571
224. Forcet C, et al. Hum. Mol. Genet. (2005) PMID: 15800014
225. Zhang L, et al. Sci Rep (2015) PMID: 25960268
226. Berger AH, et al. Cancer Cell (2016) PMID: 27478040
227. Donnelly LL, et al. Carcinogenesis (2021) PMID: 34849607
228. Amos CI, et al. J. Med. Genet. (2004) PMID: 15121768
229. Hearle N, et al. Clin. Cancer Res. (2006) PMID: 16707622
230. van der Groep P, et al. Cell Oncol (Dordr) (2011) PMID: 21336636
231. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
232. Bailey MH, et al. Cell (2018) PMID: 29625053
233. Bolton KL, et al. Nat Genet (2020) PMID: 33106634
234. Scheuermann JC, et al. Nature (2010) PMID: 20436459
235. Cho YS, et al. J. Biol. Chem. (2006) PMID: 16606617
236. Park UH, et al. J. Biol. Chem. (2011) PMID: 21047783
237. Inoue D, et al. J. Clin. Invest. (2013) PMID: 24216483
238. Abdel-Wahab O, et al. Cancer Cell (2012) PMID: 22897849
239. Br. J. Cancer (2013) PMID: 23736028
240. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
241. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
242. Xie M, et al. Nat. Med. (2014) PMID: 25326804
243. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
244. Severson EA, et al. Blood (2018) PMID: 29678827
245. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
246. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
247. Chabon JJ, et al. Nature (2020) PMID: 32269342
248. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
249. Straubhar A, et al. Gynecol Oncol Rep (2017) PMID: 28560298
250. Thangavelu A, et al. Gynecol. Oncol. (2013) PMID: 24076063
251. Gershenson DM, et al. J. Clin. Oncol. (2017) PMID: 28221866
252. Eshani K, et al. BMJ Case Rep (2014) PMID: 24925537
253. Ramirez PT, et al. Gynecol. Oncol. (2008) PMID: 18457865
254. Turner NC, et al. N. Engl. J. Med. (2018) PMID: 30345905
255. Pearce ST, et al. Crit. Rev. Oncol. Hematol. (2004) PMID: 15094156
256. Robinson G, et al. Nature (2012) PMID: 22722829
257. Ho AS, et al. Nat. Genet. (2013) PMID: 23685749
258. Grasso CS, et al. Nature (2012) PMID: 22722839
259. Van der Meulen J, et al. Blood (2015) PMID: 25320243
260. Wang L, et al. Nat Commun (2013) PMID: 23792809
261. Kim JH, et al. Cancer Res. (2014) PMID: 24491801
262. Shen Y, et al. BMC Cancer (2012) PMID: 23057811
263. van Haaften G, et al. Nat. Genet. (2009) PMID: 19330029
264. Nature (2012) PMID: 22960745
265. Ciriello G, et al. Cell (2015) PMID: 26451490
266. Nature (2014) PMID: 24476821
267. Nature (2015) PMID: 25631445
268. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
269. Gao J, et al. Sci Signal (2013) PMID: 23550210
270. Jones DH, et al. Mol Clin Oncol (2017) PMID: 28781807
271. Kim SM, et al. Biochem. Biophys. Res. Commun. (2006) PMID: 16682010
272. Kang D, et al. Genes Chromosomes Cancer (2013) PMID: 23011637
273. Chen Y, et al. PLoS ONE (2014) PMID: 24874471
274. Morishita M, et al. Biochim. Biophys. Acta (2011) PMID: 21664949
275. Supernat A, et al. Oncol Lett (2012) PMID: 23205091
276. Xu H, et al. Breast Cancer Res. (2011) PMID: 21255398
277. Sharaf R, et al. Genome Med (2022) PMID: 35227290
278. Su XA, et al. Genes Dev (2021) PMID: 33766983
279. Deb S, et al. Br. J. Cancer (2014) PMID: 24548858
280. Xu H, et al. Nat. Rev. Cancer (2011) PMID: 21326324
281. Hill VK, et al. Biochim. Biophys. Acta (2016) PMID: 27207471
282. Solomon DA, et al. BMB Rep (2014) PMID: 24856830
283. Bauerschmidt C, et al. Nucleic Acids Res. (2010) PMID: 19906707
284. Yun J, et al. Nucleic Acids Res. (2016) PMID: 26420833
285. Gelot C, et al. Nucleus (2016) PMID: 27326661
286. Yan M, et al. Breast Cancer Res. (2012) PMID: 22537934
287. Sofueva S, et al. EMBO J. (2013) PMID: 24185899
288. Deng Z, et al. EMBO J. (2012) PMID: 23010778
289. Yun J, et al. EMBO Rep. (2016) PMID: 27466323
290. Mahmood SF, et al. Carcinogenesis (2014) PMID: 24148822
291. Owonikoko et al., 2016; ESMO Abstract 14230
292. Hook KE, et al. Mol. Cancer Ther. (2012) PMID: 22222631
293. Gong X, et al. Cancer Discov (2019) PMID: 30373917
294. Oser MG, et al. Cancer Discov (2019) PMID: 30373918
295. Yang W, et al. Kaohsiung J Med Sci (2022) PMID: 34741392
296. Beltran H, et al. Clin. Cancer Res. (2019) PMID: 30232224
297. Allaman-Pillet N, et al. Ophthalmic Genet. () PMID: 21955141
298. Viatour P, et al. J. Exp. Med. (2011) PMID: 21875955
299. Kawaki J, et al. Int. J. Cancer (2000) PMID: 11004673
300. Kang YK, et al. Hum. Pathol. (2002) PMID: 12378511
301. Tannapfel A, et al. J. Pathol. (2002) PMID: 12210082
302. Nguyen B, et al. Cell (2022) PMID: 35120664
303. Boonjarasapinyo S, et al. Pathol. Oncol. Res. (2012) PMID: 21750953
304. Burkhart DL, et al. Nat. Rev. Cancer (2008) PMID: 18650841
305. Knudsen ES, et al. Nat. Rev. Cancer (2008) PMID: 19143056
306. Chen Z, et al. Hum. Mutat. (2014) PMID: 24282159
307. Yun J, et al. Int J Ophthalmol (2011) PMID: 22553621
308. Houston SK, et al. Int Ophthalmol Clin (2011) PMID: 21139478
309. Ng AK, et al. Semin Radiat Oncol (2010) PMID:

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1461672-01

APPENDIX
References

- 19959033
310. Hirai H, et al. Cancer Biol. Ther. (2010) pmid: 20107315
311. Bridges KA, et al. Clin. Cancer Res. (2011) pmid: 21799033
312. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) pmid: 21389100
313. Osman AA, et al. Mol. Cancer Ther. (2015) pmid: 25504633
314. Xu L, et al. Mol. Cancer Ther. (2002) pmid: 12489850
315. Xu L, et al. Mol. Med. (2001) pmid: 11713371
316. Camp ER, et al. Cancer Gene Ther. (2013) pmid: 23470564
317. Kim SS, et al. Nanomedicine (2015) pmid: 25240597
318. Pirolo KF, et al. Mol. Ther. (2016) pmid: 27357628
319. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27601554
320. Moore et al., 2019; ASCO Abstract 5513
321. Leijen S, et al. J. Clin. Oncol. (2016) pmid: 27998224
322. Oza et al., 2015; ASCO Abstract 5506
323. Lee J, et al. Cancer Discov (2019) pmid: 31315834
324. Méndez E, et al. Clin. Cancer Res. (2018) pmid: 29535125
325. Seligmann JF, et al. J Clin Oncol (2021) pmid: 34538072
326. Gourley et al., 2016; ASCO Abstract 5571
327. Park H, et al. ESMO Open (2022) pmid: 36084396
328. Li M, et al. Nat. Genet. (2014) pmid: 24997986
329. Borger DR, et al. Oncologist (2012) pmid: 22180306
330. Suto T, et al. J Surg Oncol (2000) pmid: 10738270
331. Nault JC, et al. Semin. Liver Dis. (2011) pmid: 21538283
332. Won HS, et al. BMC Cancer (2010) pmid: 20955617
333. Chaube A, et al. BMC Cancer (2006) pmid: 16686942
334. Chuang SC, et al. World J Surg (2004) pmid: 15573254
335. Ruzzenente A, et al. Ann. Surg. Oncol. (2016) pmid: 26717940
336. Xiaofang L, et al. World J Surg Oncol (2012) pmid: 22230750
337. Ajiki T, et al. Hepatogastroenterology () pmid: 8799388
338. J Surg Oncol (2006) pmid: 16724348
339. Guo R, et al. Hum. Pathol. (2014) pmid: 24746206
340. Conci S, et al. Updates Surg (2020) pmid: 32020551
341. Simbolo M, et al. Sci Rep (2018) pmid: 29740198
342. Churi CR, et al. PLoS ONE (2014) pmid: 25536104
343. Lee CS, et al. Pathology (1995) pmid: 7567135
344. Ahrendt SA, et al. J Hepatobiliary Pancreat Surg (2000) pmid: 11180865
345. Joerger AC, et al. Annu. Rev. Biochem. (2008) pmid: 18410249
346. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) pmid: 12826609
347. Kamada R, et al. J. Biol. Chem. (2011) pmid: 20978130
348. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) pmid: 28472496
349. Yamada H, et al. Carcinogenesis (2007) pmid: 17690113
350. Bougeard G, et al. J. Clin. Oncol. (2015) pmid: 26014290
351. Sorrell AD, et al. Mol Diagn Ther (2013) pmid: 23355100
352. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) pmid: 11219776
353. Kleihues P, et al. Am. J. Pathol. (1997) pmid: 9006316
354. Gonzalez KD, et al. J. Clin. Oncol. (2009) pmid: 19204208
355. Laloo F, et al. Lancet (2003) pmid: 12672316
356. Mandelker D, et al. Ann. Oncol. (2019) pmid: 31050713
357. Zhang X, et al. PLoS ONE (2013) pmid: 23991038
358. Holland DG, et al. EMBO Mol Med (2011) pmid: 21337521
359. Sircoulomb F, et al. EMBO Mol Med (2011) pmid: 21328542
360. Reynisdottir I, et al. Cancer Med (2013) pmid: 24156016
361. Yang G, et al. Oncol. Rep. (2014) pmid: 24481460
362. Ma F, et al. Oncol. Rep. (2014) pmid: 25017610
363. Bazarov AV, et al. Breast Cancer Res. (2011) pmid: 21635707
364. Slorach EM, et al. Genes Dev. (2011) pmid: 21317240
365. Andre et al., 2021; ASCO GI Abstract 9
366. Oaknin A, et al. JAMA Oncol (2020) pmid: 33001143
367. Berton et al., 2021; ASCO Abstract 2564
368. Andre et al., 2021; ESMO GI Abstract SO-9
369. Bang et al., 2019; ASCO Abstract 4079
370. Lin et al., 2018; ASCO Abstract 500
371. Villanueva et al., 2021; ASCO Abstract 321
372. Javle et al., 2019; ASCO Abstract 4087
373. Arkenau HT, et al. Oncologist (2018) pmid: 29853658
374. Migden MR, et al. N. Engl. J. Med. (2018) pmid: 29863979
375. Stratigos et al., 2020; EMSO Abstract LBA47
376. Lewis et al. 2020; doi: 10.1136/jitc-2020-SITC2020.0428
377. Sezer et al., 2020; ESMO Abstract LBA52
378. Kim RD, et al. JAMA Oncol (2020) pmid: 32352498
379. Gbolahan O, et al. J Natl Compr Canc Netw (2019) pmid: 31200359
380. Klein O, et al. JAMA Oncol (2020) pmid: 32729929

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Douglas A. Mata, MD, MPH | 04 October 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531