

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Lung adenocarcinoma	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Liver
	NAME Wu, Hung Ying		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S110-14026C (PF22104)
	DATE OF BIRTH 02 October 1955		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Female		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 22 April 2021
	MEDICAL RECORD # 38161329		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 08 September 2022

Biomarker Findings

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

ARID1A S11fs*91

KRAS G12D

CDK8 amplification - equivocal[†]

FLT3 amplification - equivocal[†]

JAK1V310I

SMAD2 loss exons 3-11

TP53 V157D

7 Disease relevant genes with no reportable alterations: **ALK, BRAF, EGFR, ERBB2, MET, RET, ROS1**

[†] See About the Test in appendix for details.

Report Highlights

- Evidence-matched clinical trial options based on this patient's genomic findings: (p. [9](#))
- Variants with prognostic implications for this tumor type that may impact treatment decisions: **KRAS G12D** (p. [5](#))

BIOMARKER FINDINGS

Microsatellite status - MS-Stable

Tumor Mutational Burden - 1 Muts/Mb

GENOMIC FINDINGS

ARID1A - S11fs*91

9 Trials see p. [9](#)

KRAS - G12D

10 Trials see p. [11](#)

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

No therapies or clinical trials. See Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

none

none

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

none

none

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

CDK8 - amplification - equivocal	p. 6	SMAD2 - loss exons 3-11	p. 7
FLT3 - amplification - equivocal	p. 6	TP53 - V157D	p. 8
JAK1 - V310I	p. 7		

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT
MS-Stable

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors¹⁻³, including approved therapies nivolumab and pembrolizumab⁴. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and

experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵.

FREQUENCY & PROGNOSIS

MSI-H is generally infrequent in NSCLC, reported in fewer than 1% of samples across several large studies⁶⁻¹¹, whereas data on the reported incidence of MSI-H in SCLC has been limited and conflicting¹²⁻¹⁵. One study reported MSI-H in lung adenocarcinoma patients with smoking history, and 3 of 4 MSI-H patients examined also had metachronous carcinomas in other organs, although this has not been investigated in large scale studies⁶. Published data investigating the prognostic implications of MSI in NSCLC are limited (PubMed, Oct 2021).

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor¹⁶. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2¹⁶⁻¹⁸. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers¹⁹⁻²¹. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{16,18,20-21}.

BIOMARKER

Tumor Mutational Burden

RESULT
1 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1²²⁻²⁴, anti-PD-1 therapies²²⁻²⁵, and combination nivolumab and ipilimumab²⁶⁻³¹. Multiple clinical trials of PD-1- or PD-L1-targeting immune checkpoint inhibitors or combination of PD-1 and CTLA-4 inhibitors in NSCLC have reported that patients with tumors harboring TMB ≥ 10 Muts/Mb derive greater clinical benefit from these therapies than those with TMB < 10 Muts/Mb (based on this assay or others); similarly, higher efficacy of anti-PD-1 or anti-PD-L1 immunotherapy for treatment of patients with NSCLC, compared with the use of chemotherapy, has been observed more significantly in cases of TMB ≥ 10 Muts/Mb (based on this assay or others)^{22-23,26-28,32-39}. Improved OS of patients with NSCLC treated with pembrolizumab plus chemotherapy relative to chemotherapy only⁴⁰, or those treated with nivolumab plus ipilimumab also relative to

chemotherapy⁴¹, has been observed across all TMB levels.

FREQUENCY & PROGNOSIS

A large-scale genomic analysis found that unspecified lung non-small cell lung carcinoma (NSCLC), lung adenocarcinoma, and lung squamous cell carcinoma (SCC) samples harbored median TMBs between 6.3 and 9 Muts/Mb, and 12% to 17% of cases had an elevated TMB of greater than 20 Muts/Mb⁴². Lower TMB is observed more commonly in NSCLCs harboring known driver mutations (EGFR, ALK, ROS1, or MET) with the exception of BRAF or KRAS mutations, which are commonly observed in elevated TMB cases⁴³. Although some studies have reported a lack of association between smoking and mutational burden in NSCLC⁴⁴⁻⁴⁵, several other large studies did find a strong association with increased TMB⁴⁶⁻⁴⁹. TMB > 10 muts/Mb was found to be more frequent in NSCLC metastases compared with primary tumors for both adenocarcinoma (38% vs. 25%) and SCC (41% vs. 35%) subtypes⁵⁰. A meta-analysis of 19 studies of immune checkpoint inhibitor-treated NSCLC ($n = 2,315$ patients) demonstrated that high TMB predicted a significantly longer OS than low TMB (HR = 0.70), and within the high TMB group, immunotherapy was associated with an improved PFS (HR = 0.62, $P < 0.001$), OS (HR = 0.67, $P < 0.001$) and a higher response rate (OR = 2.35, $P < 0.001$) compared to chemotherapy⁵¹. In contrast, a large study of Chinese patients with untreated lung

adenocarcinoma reported a shorter median OS for tumors with a higher number of mutations in a limited gene set compared with a lower mutation number (48.4 vs. 61.0 months)⁴⁴. Another study of patients with NSCLC treated with EGFR inhibitors or platinum doublet chemotherapy found elevated TMB to be correlated with poorer prognosis, as well as finding lower TMB in combination with PD-L1 negative status to be significantly associated with longer median survival in patients with lung adenocarcinoma⁵². However, no significant prognostic association of TMB and/or PD-L1 status with survival has been reported in patients with lung SCC⁵²⁻⁵³.

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁵⁴⁻⁵⁵ and cigarette smoke in lung cancer^{32,56}, treatment with temozolomide-based chemotherapy in glioma⁵⁷⁻⁵⁸, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁵⁹⁻⁶³, and microsatellite instability (MSI)^{59,62-63}. This sample harbors a TMB below levels that would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents^{22-23,26-28,32-39,64}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

GENOMIC FINDINGS

GENE

ARID1A

ALTERATION

S11fs*91

TRANSCRIPT ID

NM_006015

CODING SEQUENCE EFFECT

31_56del26

VARIANT ALLELE FREQUENCY (% VAF)

11.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies approved to address the mutation or loss of ARID1A in cancer. However, on the basis of limited clinical and preclinical evidence, ARID1A inactivating mutations may lead to sensitivity to ATR inhibitors such as M6620 and ceralasertib⁶⁵. In a Phase 2 study of ceralasertib in solid tumors, 2 patients with endometrial carcinoma in the cohort with loss of ARID1A expression achieved CRs on ceralasertib monotherapy; at least 1 of these 2 patients carried an inactivating ARID1A mutation. In contrast, no responses were observed for patients with normal ARID1A expression treated with ceralasertib combined with olaparib⁶⁶. One patient with small

cell lung cancer harboring an ARID1A mutation experienced a PR when treated with M6620 combined with topotecan⁶⁷. In a Phase 1 trial, a patient with metastatic colorectal cancer harboring both an ARID1A mutation and ATM loss treated with single-agent M6620 achieved a CR that was ongoing at 29 months⁶⁸. On the basis of limited preclinical evidence from studies in ovarian cancer, ARID1A inactivation may predict sensitivity to EZH2 inhibitors⁶⁹⁻⁷⁰, which are under investigation in clinical trials. Other studies have reported that the loss of ARID1A may activate the PI3K-AKT pathway and be linked with sensitivity to inhibitors of this pathway⁷¹⁻⁷³. Patients with ARID1A alterations in advanced or metastatic solid tumors may derive benefit from treatment with anti-PD-1 or anti-PD-L1 immunotherapy⁷⁴. Loss of ARID1A expression has been associated with chemoresistance to platinum-based therapy for patients with ovarian clear cell carcinoma⁷⁵⁻⁷⁶ and to 5-fluorouracil in colorectal cancer cell lines⁷⁷.

FREQUENCY & PROGNOSIS

ARID1A alterations are particularly prevalent in ovarian clear cell carcinoma (46-50%), ovarian and uterine endometrioid carcinomas (24-44%), and cholangiocarcinoma (27%); they are also reported in up to 27% of gastric carcinoma, esophageal adenocarcinoma, Waldenstrom macroglobulinemia, pediatric Burkitt lymphoma, hepatocellular

carcinoma, colorectal carcinoma, and urothelial carcinoma samples analyzed (COSMIC, cBioPortal, Jan 2022)⁷⁸⁻⁸⁶. ARID1A loss is associated with microsatellite instability in ovarian and endometrial endometrioid adenocarcinomas^{74,87-90}, CRC^{74,91-93}, and gastric cancer^{74,94-98}. ARID1A protein loss is associated with tumors of poor histological grade for many tumor types, including colorectal cancer (CRC)⁹¹⁻⁹³, cervical cancer⁹⁹⁻¹⁰⁰, gastric cancer⁹⁴⁻⁹⁸, urothelial carcinoma¹⁰¹⁻¹⁰³, ovarian and endometrial cancers^{76,87-90,104-108}, breast carcinoma¹⁰⁹⁻¹¹¹, and clear cell renal cell carcinoma¹¹²; ARID1A mutation has been associated with poor outcomes for patients with cholangiocarcinoma¹¹³⁻¹¹⁶. However, prognostic data regarding patient survival are often mixed and conflicting.

FINDING SUMMARY

ARID1A encodes the AT-rich interactive domain-containing protein 1A, also known as Baf250a, a member of the SWI/SNF chromatin remodeling complex. Mutation, loss, or inactivation of ARID1A has been reported in many cancers, and the gene is considered a tumor suppressor^{82,97,110,117-122}. ARID1A mutations, which are mostly truncating, have been identified along the entire gene and often correlate with ARID1A protein loss^{82,95,118-119,123}, whereas ARID1A missense mutations are mostly uncharacterized.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

GENOMIC FINDINGS

GENE

KRAS

ALTERATION

G12C

TRANSCRIPT ID

NM_004985

CODING SEQUENCE EFFECT

35G>A

VARIANT ALLELE FREQUENCY (% VAF)

60.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical evidence suggests that KRAS activation may predict sensitivity to MEK inhibitors, such as trametinib, binimetinib, cobimetinib, and selumetinib¹²⁴⁻¹²⁹. In a Phase 1 study evaluating the MEK-pan-RAF dual inhibitor CH5126766, 6 patients harboring KRAS mutations experienced PRs, including 3 with non-small cell lung cancer (NSCLC), 1 with low-grade serous ovarian carcinoma (LGSOC), 1 with endometrial adenocarcinoma, and 1 with multiple myeloma¹³⁰. Combination of CH5126766 with the FAK inhibitor defactinib elicited PR rates of 50% (4/8) for patients with KRAS-mutated low-grade serous ovarian cancer and 12% (2/17) for patients with KRAS-mutated non-small cell lung cancer (NSCLC) in a Phase 1 study¹³¹⁻¹³². Preclinical and clinical data suggest that KRAS mutations may predict clinical benefit from SHP2 inhibitors¹³³⁻¹³⁴. A Phase 1 study of RMC-4630 for relapsed/refractory solid tumors reported a DCR of 58% (23/40) for patients with NSCLC and KRAS mutations and a DCR of 75% (12/16) for patients with NSCLC and KRAS G12C mutations¹³⁵. Interim results from a Phase 1/2 study of RMC-4630 plus cobimetinib reported tumor reduction in 3 of 8 patients with KRAS-mutated colorectal cancer¹³⁶. Combination approaches of MEK inhibitors with chemotherapy have been investigated in KRAS-mutated NSCLC¹³⁷⁻¹³⁸; a Phase 3 study reported that addition of selumetinib to docetaxel did not improve PFS or OS for patients with KRAS-mutated NSCLC compared with docetaxel alone¹³⁷. Multiple clinical studies have reported either low response rates or response rates similar to those of chemotherapy in patients with KRAS-mutated NSCLC receiving MEK inhibitors as a monotherapy¹³⁹⁻¹⁴¹. Combinatorial approaches involving MEK inhibitors and other targeted therapies, including

PI3K or EGFR inhibitors, have generally had limited clinical efficacy for patients with NSCLC and have been associated with high toxicity¹⁴²⁻¹⁴⁵. Preclinical data suggest that KRAS mutation may confer sensitivity to SOS1 inhibitors¹⁴⁶⁻¹⁴⁷. Phase 1 studies of the SOS1 inhibitor BI 1701963 alone or in combination with MEK inhibitors, KRAS G12C inhibitors, or irinotecan are recruiting for patients with solid tumors harboring KRAS mutations¹⁴⁸⁻¹⁴⁹. Immune checkpoint inhibitors (ICIs) have been associated with benefit for patients with KRAS-mutated non-small cell lung cancer (NSCLC). Phase 3 studies of nivolumab¹⁵⁰ or atezolizumab¹⁵¹ versus docetaxel in NSCLC post-platinum doublet therapy reported benefit for their respective KRAS-mutated subcohorts in addition to the cohorts overall; KRAS mutations in NSCLC are reported to correlate with PD-(L)1 status¹⁵². A retrospective analysis of pooled data from 12 clinical trials investigating ICI with or without chemotherapy for patients with NSCLC reported that patients with KRAS-mutated NSCLC, including KRAS G12C, benefited from first-line chemotherapy plus ICI similar to those with KRAS-wildtype NSCLC (median OS [mOS] of 22.4 vs. 18.7 months, ORR of 46% vs. 51%); the ICI plus chemotherapy combination exhibited improved outcomes compared with ICI alone or chemotherapy alone (mOS of 22.4 vs. 16.2 vs. 17.1 months)¹⁵³. CDK4/6 inhibitors have been investigated in KRAS-mutated NSCLC; in a Phase 3 study of third-line platinum-refractory NSCLC, abemaciclib failed to improve mOS (7.4 months vs. 7.8 months, HR=0.97) relative to erlotinib in KRAS-mutated disease, despite increased ORR and PFS rates¹⁵⁴. A Phase 1 combination trial of the CDK4/6 inhibitor palbociclib with MEK inhibitor mirdametinib included 17 patients with KRAS-mutated NSCLC and reported 1 PR, >50% SD, and 5 patients with PFS >6 months; clinical benefit was seen among patients with tumors harboring KRAS mutation alone or together with inactivation of TP53 or CDKN2a/B, but not among patients with tumors harboring KRAS mutation and STK11 inactivation¹⁵⁵. Although some studies have suggested that KRAS mutation status may predict a lack of response to the EGFR inhibitors erlotinib and gefitinib for patients with lung cancer¹⁵⁶⁻¹⁵⁸, a retrospective study reported no statistically significant difference in response to EGFR tyrosine kinase inhibitors among KRAS-wildtype and KRAS-mutated disease independent from EGFR mutation status¹⁵⁹. Co-occurring KRAS and STK11 alterations are associated with poorer response to ICIs for patients with NSCLC. Following anti-

PD-1-based regimens, retrospective analyses have reported shorter OS for patients with KRAS- and STK11-mutated tumors than for those whose KRAS-mutated tumors were STK11-wildtype (6.4 vs. 16.1 months, HR=1.99), as well as markedly fewer objective responses for patients with KRAS- and STK11-mutated versus KRAS- and TP53-mutated tumors in the CheckMate-057 (0% [0/6] vs. 57% [4/7])¹⁶⁰ and GEMINI (0% [0/6] vs. 53% [9/17])¹⁶¹ studies. Another study observed that patients with NSCLC and KRAS-mutated tumors without STK11 alteration who were treated with second-line immunotherapy experienced similar median PFS (2.8 vs. 2.2 months, HR=1.64) and numerically longer median OS (7.7 vs. 3.5 months, HR=2.3; p=0.09) compared with patients harboring mutations in both KRAS and STK11¹⁶².

FREQUENCY & PROGNOSIS

Studies have reported KRAS mutations in 10-38% of non-small cell lung cancers (NSCLC), including 27-37% of lung adenocarcinomas^{47-48,163-172}, 10.5-33% of lung adenosquamous carcinomas¹⁷³⁻¹⁷⁵, 22% of lung large cell carcinoma without neuroendocrine features, and 6% of lung large cell neuroendocrine carcinomas¹⁷⁶. KRAS mutation in lung adenocarcinoma has been correlated with disease progression, poorly differentiated tumors, and aggressive tumor behavior (NCCN NSCLC Guidelines, v3.2022)^{166,172,177}. However, the prognostic value of KRAS mutation in lung adenocarcinoma may differ among ethnic groups and may depend upon the specific allelic variant present¹⁷⁸. KRAS mutation was associated with shorter PFS (7.0 vs. 8.6 months, p=0.026) and OS (14.2 vs. 21.6 months, p=0.019) with first-line treatment with bevacizumab plus chemotherapy in a retrospective study¹⁷⁹ and a lower major pathological response rate (0% [0/10] vs. 35.5% [11/31]) after neoadjuvant bevacizumab plus chemotherapy followed by adjuvant bevacizumab in a Phase 2 trial¹⁸⁰, relative to those patients lacking KRAS mutation. However, addition of atezolizumab to first-line bevacizumab and chemotherapy improved PFS regardless of KRAS status in the Phase 3 IMpower150 study (HR=0.50 for KRAS mutant vs. 0.47 for KRAS wild-type vs. 0.67 for KRAS unknown)¹⁸¹. In one study of 55 patients with lung adenocarcinoma, KRAS mutations, especially in combination with TP53 alterations, correlated with improved clinical outcomes to PD-1 inhibitors pembrolizumab and nivolumab, likely as a consequence of association with some immunogenic features such as tumor mutation burden¹⁸².

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

GENOMIC FINDINGS

FINDING SUMMARY

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor

formation^{125,183}. KRAS alterations affecting amino acids G12, G13, Q22, P34, A59, Q61, and A146, as well as mutations G10_A11insG, G10_A11insAG (also reported as G10_A11dup and

G12_G13insAG), A18D, L19F, D33E, G60_A66dup/E62_A66dup, E62K, E63K, R68S, and K117N have been characterized as activating and oncogenic^{125,184-206}.

GENE

CDK8

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies approved or in clinical trials that directly address genomic alterations in CDK8. Investigational inhibitors that selectively target CDK8 and the related CDK19 have shown activity against WNT-dependent tumors in preclinical assays²⁰⁷⁻²¹⁰.

FREQUENCY & PROGNOSIS

CDK8 amplification has been identified in various solid tumor types, including colorectal adenocarcinoma (3%)⁶², pancreatic adenocarcinoma (3%)²¹¹, stomach adenocarcinoma (less than 2%)²¹², prostate adenocarcinoma (0-6%)²¹³⁻²¹⁵, and breast carcinoma (1-2%)²¹⁶⁻²¹⁷. In the TCGA datasets, CDK8 mutation was observed most frequently (2%) in bladder urothelial carcinoma²¹⁸, lung adenocarcinoma¹⁶³, and lung squamous cell carcinoma²¹⁹. In colorectal cancer, CDK8 expression has been detected in 70% of cases and correlated with beta-catenin activation; patients with CDK8-positive colon cancer experienced significantly shorter survival (hazard ratio of 2.05 by multivariate analysis), which was not seen for patients with rectal cancer²²⁰⁻²²². The association of CDK8 alterations with the prognosis of other tumor types has not been firmly established²²³⁻²²⁴.

FINDING SUMMARY

CDK8 encodes a member of the cyclin-dependent kinase (CDK) family. The CDK8 protein regulates gene expression as part of the Mediator complex, and its overexpression can dysregulate various cancer pathways, including the WNT/beta-catenin and NOTCH pathways²²⁵⁻²²⁶. CDK8 acts as oncogene in colon cancer and melanoma²²⁷⁻²²⁹ but has also been described to have tumor suppressor functions²³⁰⁻²³². Whereas CDK8 mutations have not been extensively characterized, amplification or overexpression of CDK8, often as a result of copy number increases in chromosomal region 13q12.13, have been reported in the scientific literature and may be associated with shorter survival for patients with colon cancer^{220,227,233}.

GENE

FLT3

ALTERATION

amplification - equivocal

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapies targeting FLT3 are under clinical investigation, including crenolanib, gilteritinib, luseptinib, midostaurin, pacritinib, pexidartinib, ponatinib, quizartinib, sorafenib, and sunitinib. The TKIs midostaurin²³⁴⁻²³⁷ and gilteritinib²³⁸⁻²⁴⁰ have shown significant clinical activity for patients with

relapsed/refractory acute myeloid leukemia (AML) harboring FLT3-ITD or FLT3-TKD mutations. In the Phase 1 study for the FLT3/BTK inhibitor luseptinib, a patient with FLT3-ITD AML experienced a minimal residual disease (MRD)-negative CR²⁴¹. A patient with FLT3-amplified and KRAS-mutated colorectal cancer (CRC) achieved significant benefit from sorafenib treatment²⁴². Similarly, another patient with FLT3-amplified and KRAS-mutated CRC achieved a PR to regorafenib; two other patients with FLT3-amplified CRC treated with regorafenib experienced SD and PD, respectively²⁴³. However, a Phase 2 clinical trial of 10 patients with metastatic CRC and FLT3 amplification treated with sunitinib did not show significant clinical benefit²⁴⁴.

FREQUENCY & PROGNOSIS

FLT3 amplification has been reported in 1% of lung adenocarcinoma samples in the TCGA dataset¹⁶³. Published data investigating the prognostic implications of FLT3 alterations in lung NSCLC are limited (PubMed, Oct 2021).

FINDING SUMMARY

FLT3 encodes a receptor tyrosine kinase that potentiates signaling through the RAS and PI3K pathways²⁴⁵⁻²⁴⁷. FLT3 has been reported to be amplified in cancer⁸⁰ and may be biologically relevant in this context²⁴⁸⁻²⁴⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

GENOMIC FINDINGS

GENE

JAK1

ALTERATION

V310I

TRANSCRIPT ID

NM_002227

CODING SEQUENCE EFFECT

928G>A

VARIANT ALLELE FREQUENCY (% VAF)

34.4%

efficacy in reducing symptoms in Phase 1 and 2 trials in patients with myeloproliferative disorders²⁵⁰⁻²⁵². Other small molecule inhibitors of JAK1 are being investigated in preclinical studies in some types of solid tumors²⁵³⁻²⁵⁴. HSP90 inhibitors are also being investigated in preclinical studies to target components of the JAK-STAT pathway such as JAK1²⁵⁵. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

expression of p-JAK1 in NSCLC tumors has been significantly correlated with reduced patient survival²⁵⁶.

FINDING SUMMARY

The JAK1 (Janus kinase 1) gene encodes a tyrosine kinase that regulates signals triggered by cytokines and growth factors²⁵⁷. Dysregulation of JAK-STAT signaling has been implicated in a variety of epithelial tumors²⁵⁸. However, JAK-STAT signaling is required for the antiviral and antiproliferative effects of interferons²⁵⁹. Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Inhibitors of the JAK-STAT pathway are under development. The JAK1/JAK2 inhibitor ruxolitinib is approved to treat myelofibrosis, and has shown

FREQUENCY & PROGNOSIS

In the lung adenocarcinoma TCGA and COSMIC datasets, JAK1 mutations have been reported in 1.6-3.0% of cases analyzed (Sep 2022)⁷⁸⁻⁸⁰. Published data investigating the prognostic implications of JAK1 alterations in NSCLC are limited (PubMed, May 2022). In one study

GENE

SMAD2

ALTERATION

loss exons 3-11

considered for each patient. In preclinical studies, several novel small-molecule TGF-beta pathway inhibitors have been shown to reduce SMAD2 phosphorylation and tumor invasion in breast cancer cells and to increase survival in mouse xenograft models²⁶⁰⁻²⁶¹.

as well as loss of expression or phosphorylation (p-SMAD2), has been reported in several cancers²⁶⁵⁻²⁷¹ and has been correlated with poor prognosis in hepatocellular, gastric, breast, and colorectal cancer²⁶⁶⁻²⁷⁰ but good prognosis in HNSCC²⁷²⁻²⁷³.

FINDING SUMMARY

SMAD2 encodes an intracellular transducer that is activated by TGF-beta or activin to dimerize with SMAD4 and regulate transcription of TGF-beta or activin-activated genes²⁷⁴⁻²⁷⁵. SMAD2 alterations that disrupt the MH1 domain (amino acids 10-176), MH2 domain (amino acids 274-467), or critical phosphorylation sites are predicted to result in a loss of function^{263,271,276-279}.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies that target the loss of SMAD2 or loss of TGF-beta signaling in cancer. Because TGF-beta can exert both tumor suppressor as well as pro-tumor effects, the development of therapies aimed at this pathway must be carefully

FREQUENCY & PROGNOSIS

SMAD2 mutations have been reported in various tumor types, including cutaneous squamous cell carcinoma (SCC; 6-10%)²⁶², colorectal adenocarcinoma (CRC; 3-7%)^{62,263}, endometrial cancers (2-5%)⁵⁹, stomach adenocarcinoma (2-3%)²¹², lung SCC (1-2%)²¹⁹, and lung adenocarcinoma (1-2%)^{47-48,163,264}. SMAD2 deletion,

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

V157D

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

470T>A

VARIANT ALLELE FREQUENCY (% VAF)

59.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁸⁰⁻²⁸³, or p53 gene therapy and immunotherapeutics such as SGT-53²⁸⁴⁻²⁸⁸ and ALT-801²⁸⁹. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²⁹⁰. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁹¹. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²⁹². The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁹³. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel²⁹⁴. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck

squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations²⁹⁵. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²⁹⁶. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁸⁸. Missense mutations leading to TP53 inactivation may also be sensitive to therapies that reactivate mutated p53 such as APR-246²⁹⁷⁻²⁹⁹. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, APR-246 combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR³⁰⁰. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies³⁰¹⁻³⁰²; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies³⁰³⁻³⁰⁴. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

TP53 is one of the most commonly mutated genes in lung cancer; mutations have been reported in 43-80% of non-small cell lung cancers (NSCLCs)^{163,219,305-310}, including 42-52% of lung adenocarcinomas and 58-83% of lung squamous cell carcinomas (cBioPortal, COSMIC, Feb 2022)^{48-49,163,219}. TP53 homozygous deletion has been observed in 1.4% of lung adenocarcinoma and <1% of lung squamous cell carcinoma cases (cBioPortal, Feb 2022)⁷⁹⁻⁸⁰. In one study of 55 patients with lung adenocarcinoma, TP53 alterations correlated with immunogenic features including PD-L1 expression, tumor mutation burden and neoantigen presentation; likely as a consequence of this association TP53 mutations correlated with improved clinical outcomes to

PD-1 inhibitors pembrolizumab and nivolumab in this study¹⁸². Mutations in TP53 have been associated with lymph node metastasis in patients with lung adenocarcinoma³¹¹.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers³¹². Alterations such as seen here may disrupt TP53 function or expression³¹³⁻³¹⁷.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers³¹⁸⁻³²⁰, including sarcomas³²¹⁻³²². Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000³²³ to 1:20,000³²². For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30³²⁴. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion³²⁵⁻³³⁰. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy³²⁵⁻³²⁶. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease³³¹. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{329,332-333}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually

updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that

may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.clinicaltrials.gov). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

GENE
ARID1A
RATIONALE

ARID1A loss or inactivation may predict

sensitivity to ATR inhibitors.

ALTERATION

S11fs*91

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT04802174
PHASE 1/2

Lurbinectedin With Berzosertib, an ATR Kinase Inhibitor in Small Cell Cancers and High-Grade Neuroendocrine Cancers

TARGETS
ATR

LOCATIONS: Maryland

NCT04657068
PHASE 1/2

A Study of ART0380 for the Treatment of Advanced or Metastatic Solid Tumors

TARGETS
ATR

LOCATIONS: London (United Kingdom), Colorado, Oklahoma, Texas, Pennsylvania, Tennessee, Florida

NCT04491942
PHASE 1

Testing the Addition of an Anti-cancer Drug, BAY 1895344, to the Usual Chemotherapy Treatment (Cisplatin, or Cisplatin and Gemcitabine) for Advanced Solid Tumors With Emphasis on Urothelial Cancer

TARGETS
ATR

LOCATIONS: California, Wisconsin, Ohio, Pennsylvania, New York, Maryland

NCT04514497
PHASE 1

Testing the Addition of an Anti-cancer Drug, BAY 1895344, to Usual Chemotherapy for Advanced Stage Solid Tumors, With a Specific Focus on Patients With Small Cell Lung Cancer, Poorly Differentiated Neuroendocrine Cancer, and Pancreatic Cancer

TARGETS
ATR, TOP1

LOCATIONS: Arizona, Minnesota, Oklahoma, Pennsylvania, Connecticut, Tennessee, Florida

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

CLINICAL TRIALS
NCT04616534
PHASE 1

Testing the Addition of an Anti-cancer Drug, BAY 1895344 ATR Inhibitor, to the Chemotherapy Treatment (Gemcitabine) for Advanced Pancreatic and Ovarian Cancer, and Advanced Solid Tumors

TARGETS
ATR

LOCATIONS: Massachusetts, Maryland

NCT04266912
PHASE 1/2

Avelumab and M6620 for the Treatment of DDR Deficient Metastatic or Unresectable Solid Tumors

TARGETS
ATR, PD-L1

LOCATIONS: Texas

NCT03669601
PHASE 1

AZD6738 & Gemcitabine as Combination Therapy

TARGETS
ATR

LOCATIONS: Cambridge (United Kingdom)

NCT02595931
PHASE 1

ATR Kinase Inhibitor VX-970 and Irinotecan Hydrochloride in Treating Patients With Solid Tumors That Are Metastatic or Cannot Be Removed by Surgery

TARGETS
ATR

LOCATIONS: California, Missouri, Pennsylvania, Massachusetts, Connecticut, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

CLINICAL TRIALS
GENE
KRAS
ALTERATION
G12D
RATIONALE

KRAS activating mutations or amplification may predict sensitivity to inhibitors of MAPK pathway components, including MEK inhibitors. Limited clinical and preclinical studies indicate KRAS mutations may predict sensitivity to MEK-pan-RAF dual inhibitors. KRAS alterations are not

predictive biomarkers for MEK inhibitor monotherapy in NSCLC and combinatorial approaches may yield improved efficacy. Clinical evidence suggests that patients with KRAS-mutant NSCLC may be sensitive to the CDK4/6 inhibitor abemaciclib.

NCT03337698
PHASE 1/2

A Study Of Multiple Immunotherapy-Based Treatment Combinations In Participants With Metastatic Non-Small Cell Lung Cancer (Morpheus- Non-Small Cell Lung Cancer)

TARGETS

PD-L1, MEK, CEA, CXCR4, EZH2, MDM2, ADORA2A

LOCATIONS: Taipei City (Taiwan), Seoul (Korea, Republic of), Blacktown (Australia), Haifa (Israel), Petach Tikva (Israel), Ramat Gan (Israel), Newcastle upon Tyne (United Kingdom), Dijon (France), London (United Kingdom), Sutton (United Kingdom)

NCT04967079
PHASE 1

MEK Inhibitor Combined With Anlotinib in the Treatment of KRAS-mutated Advanced Non-small Cell Lung Cancer

TARGETS

MEK, FGFRs, KIT, VEGFRs

LOCATIONS: Shanghai (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT03284502
PHASE 1

Cobimetinib and HM95573 in Patients With Locally Advanced or Metastatic Solid Tumors

TARGETS

MEK, RAFs

LOCATIONS: Hwasun (Korea, Republic of), Pusan (Korea, Republic of), Seongnam (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of)

NCT04870034
PHASE NULL

Binimetinib and Palbociclib Before Surgery for the Treatment of Operable KRAS-Positive Lung, Colorectal, or Pancreatic Cancer

TARGETS

MEK, CDK4, CDK6

LOCATIONS: New York

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

CLINICAL TRIALS
NCT04801966
PHASE NULL

Safety and Oversight of the Individually Tailored Treatment Approach: A Novel Pilot Study

TARGETS

CDK4, CDK6, PI3K-alpha, PD-L1, MEK, PARP, PD-1, BRAF

LOCATIONS: Melbourne (Australia)

NCT05159245
PHASE 2

The Finnish National Study to Facilitate Patient Access to Targeted Anti-cancer Drugs

TARGETS

BRAf, VEGFRs, RET, KIT, ERBB2, TRKB, ALK, TRKC, ROS1, TRKA, SMO, PD-L1, MEK, CDK4, CDK6

LOCATIONS: Kuopio (Finland), Helsinki (Finland), Tampere (Finland), Turku (Finland)

NCT02079740
PHASE 1/2

Trametinib and Navitoclax in Treating Patients With Advanced or Metastatic Solid Tumors

TARGETS

BCL2, BCL-XL, BCL-W, MEK

LOCATIONS: Massachusetts

NCT03905148
PHASE 1/2

Study of the Safety and Pharmacokinetics of BGB-283 and PD-0325901 in Patients With Advanced or Refractory Solid Tumors

TARGETS

RAF, EGFR, MEK

LOCATIONS: Nedlands (Australia), Blacktown (Australia), Randwick (Australia), Melbourne (Australia), California, Texas

NCT04620330
PHASE 2

A Study of VS-6766 v. VS-6766 + Defactinib in Recurrent G12V or Other KRAS-Mutant Non-Small Cell Lung Cancer

TARGETS

FAK, RAF, MEK

LOCATIONS: Chemnitz (Germany), Leipzig (Germany), Verona (Italy), Orbassano (Italy), Paris (France), Villejuif (France), Oregon, Lyon (France), Barcelona (Spain), Coruña (Spain)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

APC
A2778S

BRCA2
I2615T

FLT1
amplification

NKX2-1
E96K

SF3B1
S851_R852insIVDTTVELANKV
GAAEIIIS

TET2
I1897L

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAS	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFA1	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFB2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Homologous Recombination status
Loss of Heterozygosity (LOH) score
Microsatellite (MS) status
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.


Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplstraat 3, 2440 Geel, Belgium. 

ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical

methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

APPENDIX

About FoundationOne®CDx

- analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score ≥ 16 will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and

extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
- Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal

hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1452195-01

APPENDIX

About FoundationOne®CDx

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.1.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. The median exon coverage for this sample is 713x. Foundation Medicine Inc. does not guarantee the suitability of use.

ORDERED TEST # **ORD-1452195-01**
APPENDIX References

1. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
2. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
3. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
4. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
5. Ayers et al., 2016; ASCO-SITC Abstract P60
6. Warth A, et al. Virchows Arch. (2016) PMID: 26637197
7. Ninomiya H, et al. Br. J. Cancer (2006) PMID: 16641899
8. Vanderwalde A, et al. Cancer Med (2018) PMID: 29436178
9. Zang YS, et al. Cancer Med (2019) PMID: 31270941
10. Dudley JC, et al. Clin. Cancer Res. (2016) PMID: 26880610
11. Takamochi K, et al. Lung Cancer (2017) PMID: 28676214
12. Pyllkkänen L, et al. Environ. Mol. Mutagen. (1997) PMID: 9329646
13. Gonzalez R, et al. Ann. Oncol. (2000) PMID: 11061602
14. Chen XQ, et al. Nat. Med. (1996) PMID: 8782463
15. Merlo A, et al. Cancer Res. (1994) PMID: 8174113
16. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
17. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
18. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
19. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
20. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
21. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
22. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
23. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
24. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
25. Cristescu R, et al. Science (2018) PMID: 30309915
26. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
27. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
28. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
29. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
30. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
31. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
32. Rizvi NA, et al. Science (2015) PMID: 25765070
33. Colli LM, et al. Cancer Res. (2016) PMID: 27197178
34. Wang VE, et al. J Immunother Cancer (2017) PMID: 28923100
35. Carbone DP, et al. N. Engl. J. Med. (2017) PMID: 28636851
36. Rizvi H, et al. J. Clin. Oncol. (2018) PMID: 29337640
37. Forde PM, et al. N. Engl. J. Med. (2018) PMID: 29658848
38. Miao D, et al. Nat. Genet. (2018) PMID: 30150660
39. Chae YK, et al. Clin Lung Cancer (2019) PMID: 30425022
40. Paz-Ares et al., 2019; ESMO Abstract LBA80
41. Hellmann MD, et al. N. Engl. J. Med. (2019) PMID: 31562796
42. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
43. Spigel et al., 2016; ASCO Abstract 9017
44. Xiao D, et al. Oncotarget (2016) PMID: 27009843
45. Shim HS, et al. J Thorac Oncol (2015) PMID: 26200269
46. Govindan R, et al. Cell (2012) PMID: 22980976
47. Ding L, et al. Nature (2008) PMID: 18948947
48. Imielinski M, et al. Cell (2012) PMID: 22980975
49. Kim Y, et al. J. Clin. Oncol. (2014) PMID: 24323028
50. Stein et al., 2019; DOI: 10.1200/PO.18.00376
51. Meng G, et al. PLoS One (2022) PMID: 35113949
52. Chen Y, et al. J. Exp. Clin. Cancer Res. (2019) PMID: 31088500
53. Yu H, et al. J Thorac Oncol (2019) PMID: 30253973
54. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
55. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
56. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
57. Johnson BE, et al. Science (2014) PMID: 24336570
58. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
59. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
60. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
61. Heitz E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
62. Nature (2012) PMID: 22810696
63. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
64. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
65. Williamson CT, et al. Nat Commun (2016) PMID: 27958275
66. Aggarwal et al., 2021; ESMO Abstract 5120
67. Thomas A, et al. J. Clin. Oncol. (2018) PMID: 29252124
68. Yap TA, et al. J Clin Oncol (2020) PMID: 32568634
69. Bitler BG, et al. Nat. Med. (2015) PMID: 25686104
70. Kim KH, et al. Nat. Med. (2015) PMID: 26552009
71. Wiegand KC, et al. BMC Cancer (2014) PMID: 24559118
72. Huang HN, et al. Mod. Pathol. (2014) PMID: 24336158
73. Samartzis EP, et al. Oncotarget (2014) PMID: 24979463
74. Okamura R, et al. J Immunother Cancer (2020) PMID: 32111729
75. Yokoyama Y, et al. J Gynecol Oncol (2014) PMID: 24459582
76. Katagiri A, et al. Mod. Pathol. (2012) PMID: 22101352
77. Xie C, et al. Tumour Biol. (2014) PMID: 24833095
78. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
79. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
80. Gao J, et al. Sci Signal (2013) PMID: 23550210
81. Wu RC, et al. Cancer Biol. Ther. (2014) PMID: 24618703
82. Jones S, et al. Hum. Mutat. (2012) PMID: 22009941
83. Dulak AM, et al. Nat. Genet. (2013) PMID: 23525077
84. Streppel MM, et al. Oncogene (2014) PMID: 23318448
85. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
86. Ross JS, et al. Oncologist (2014) PMID: 24563076
87. Huang HN, et al. Histopathology (2015) PMID: 25195947
88. Hussein YR, et al. Mod. Pathol. (2015) PMID: 25394778
89. Bosse T, et al. Mod. Pathol. (2013) PMID: 23702729
90. Allo G, et al. Mod. Pathol. (2014) PMID: 23887303
91. Chou A, et al. Hum. Pathol. (2014) PMID: 24925223
92. Ye J, et al. Hum. Pathol. (2014) PMID: 25311944
93. Wei XL, et al. World J. Gastroenterol. (2014) PMID: 25561809
94. Chen K, et al. Proc. Natl. Acad. Sci. U.S.A. (2015) PMID: 25583476
95. Wang K, et al. Nat. Genet. (2011) PMID: 22037554
96. Abe H, et al. Virchows Arch. (2012) PMID: 22915242
97. Wang DD, et al. PLoS ONE (2012) PMID: 22808142
98. Wiegand KC, et al. Hum. Pathol. (2014) PMID: 24767857
99. Katagiri A, et al. Int. J. Gynecol. Cancer (2012) PMID: 22274316
100. Cho H, et al. Hum. Pathol. (2013) PMID: 23427874
101. Gui Y, et al. Nat. Genet. (2011) PMID: 21822268
102. Balbás-Martínez C, et al. PLoS ONE (2013) PMID: 23650517
103. Faraj SF, et al. Hum. Pathol. (2014) PMID: 25175170
104. Rahman M, et al. Hum. Pathol. (2013) PMID: 22939958
105. Maeda D, et al. Int J Mol Sci (2010) PMID: 21614196
106. Lowery WJ, et al. Int. J. Gynecol. Cancer (2012) PMID: 22193641
107. Fadare O, et al. Mod. Pathol. (2013) PMID: 23524907
108. Mao TL, et al. Am. J. Surg. Pathol. (2013) PMID: 24076775
109. Zhang X, et al. Cancer Epidemiol (2012) PMID: 21889920
110. Mamo A, et al. Oncogene (2012) PMID: 21892209
111. Zhao J, et al. Tumour Biol. (2014) PMID: 24430365
112. Lichner Z, et al. Am. J. Pathol. (2013) PMID: 23416164
113. Feng F, et al. Int J Clin Oncol (2021) PMID: 33387086
114. Conci S, et al. Updates Surg (2020) PMID: 32020551
115. Simbolo M, et al. Sci Rep (2018) PMID: 29740198
116. Ruzzenente A, et al. Ann. Surg. Oncol. (2016) PMID: 26717940
117. Guan B, et al. Cancer Res. (2011) PMID: 21900401
118. Wiegand KC, et al. N. Engl. J. Med. (2010) PMID: 20942669
119. Jones S, et al. Science (2010) PMID: 20826764
120. Yan HB, et al. Carcinogenesis (2014) PMID: 24293408
121. Huang J, et al. Nat. Genet. (2012) PMID: 22922871
122. Chan-On W, et al. Nat. Genet. (2013) PMID: 24185513
123. Zang ZJ, et al. Nat. Genet. (2012) PMID: 22484628
124. Nakano H, et al. Proc. Natl. Acad. Sci. U.S.A. (1984) PMID: 6320174
125. Pylayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) PMID: 21993244
126. Yamaguchi T, et al. Int. J. Oncol. (2011) PMID: 21523318
127. Watanabe M, et al. Cancer Sci. (2013) PMID: 23438367
128. Gilmartin AG, et al. Clin. Cancer Res. (2011) PMID: 21245089
129. Yeh JJ, et al. Mol. Cancer Ther. (2009) PMID: 19372556
130. Guo C, et al. Lancet Oncol (2020) PMID: 33128873
131. Krebs et al., 2021; AACR Abstract CT019
132. Shinde et al., 2020; AACR Abstract CT143
133. Lu H, et al. Mol. Cancer Ther. (2019) PMID: 31068384
134. Mainardi S, et al. Nat Med (2018) PMID: 29808006
135. Koczywas et al., 2021; AACR Abstract LB001
136. Bendell et al., 2020; EORTC-NCI-AACR Abstract 5
137. Jänne PA, et al. JAMA (2017) PMID: 28492898
138. Gandara DR, et al. J Thorac Oncol (2017) PMID: 27876675
139. Blumenschein GR, et al. Ann. Oncol. (2015) PMID: 25722381
140. Leijen S, et al. Clin. Cancer Res. (2012) PMID: 22767668
141. Zimmer L, et al. Clin. Cancer Res. (2014) PMID: 24947927
142. van Geel RMJM, et al. Br. J. Cancer (2020) PMID: 32147669
143. Carter CA, et al. Ann. Oncol. (2016) PMID: 26802155
144. Bedard PL, et al. Clin. Cancer Res. (2015) PMID: 25500057
145. Tolcher AW, et al. Ann. Oncol. (2015) PMID: 25344362
146. Hillig RC, et al. Proc Natl Acad Sci U S A (2019) PMID: 30683722
147. Hofmann MH, et al. Cancer Discov (2021) PMID: 32816843
148. Hofmann et al., 2021; AACR Abstract CT210
149. Gort et al., 2020; ASCO Abstract TPS3651
150. Borghaei H, et al. N. Engl. J. Med. (2015) PMID: 26412456
151. Rittmeyer A, et al. Lancet (2017) PMID: 27979383
152. Liu C, et al. Cancer Lett (2020) PMID: 31644929
153. Nakajima et al., 2022; ASCO Abstract 9001

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1452195-01**
APPENDIX
References

154. Goldman JW, et al. *Front Oncol* (2020) PMID: 33194700
155. Shapiro et al., 2017; AACR Abstract CT046
156. Mao C, et al. *Lung Cancer* (2010) PMID: 20022659
157. Pao W, et al. *PLoS Med.* (2005) PMID: 15696205
158. Ludovini V, et al. *J Thorac Oncol* (2011) PMID: 21258250
159. Sun JM, et al. *PLoS ONE* (2013) PMID: 23724098
160. Skoulidis F, et al. *Cancer Discov* (2018) PMID: 29773717
161. Skoulidis et al., 2017; IASLC WCLC Abstract MA04.07
162. Marmarelis et al., 2018; IASLC WCLC Abstract P1.01-64
163. *Nature* (2014) PMID: 25079552
164. Aviel-Ronen S, et al. *Clin Lung Cancer* (2006) PMID: 16870043
165. Villaruz LC, et al. *Cancer* (2013) PMID: 23526491
166. Rekhman N, et al. *Mod. Pathol.* (2013) PMID: 23619604
167. Ragusa M, et al. *Am. J. Clin. Oncol.* (2014) PMID: 23357969
168. Kim ST, et al. *Med. Oncol.* (2013) PMID: 23307237
169. Russell PA, et al. *J Thorac Oncol* (2013) PMID: 23486266
170. Stella GM, et al. *J. Cancer Res. Clin. Oncol.* (2013) PMID: 23644698
171. Cai G, et al. *Cancer Cytopathol* (2013) PMID: 23495083
172. Yip PY, et al. *J Thorac Oncol* (2013) PMID: 23392229
173. Tochigi N, et al. *Am. J. Clin. Pathol.* (2011) PMID: 21502435
174. Shu C, et al. *Mod. Pathol.* (2013) PMID: 22996376
175. Wang R, et al. *J Thorac Oncol* (2014) PMID: 24481316
176. Karlsson A, et al. *Oncotarget* (2015) PMID: 26124082
177. Scoccianti C, et al. *Eur. Respir. J.* (2012) PMID: 22267755
178. *Curr Opin Oncol* (2014) PMID: 24463346
179. Ghimessy AK, et al. *Cancers (Basel)* (2019) PMID: 31600989
180. Haft JE, et al. *J Thorac Oncol* (2013) PMID: 23857398
181. Socinski MA, et al. *N. Engl. J. Med.* (2018) PMID: 29863955
182. Dong ZY, et al. *Clin. Cancer Res.* (2017) PMID: 28039262
183. Kahn S, et al. *Anticancer Res.* () PMID: 3310850
184. Akagi K, et al. *Biochem. Biophys. Res. Commun.* (2007) PMID: 17150185
185. Bollag G, et al. *J. Biol. Chem.* (1996) PMID: 8955068
186. Buhrman G, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2010) PMID: 20194776
187. *Sci. STKE* (2004) PMID: 15367757
188. Edkins S, et al. *Cancer Biol. Ther.* (2006) PMID: 16969076
189. Feig LA, et al. *Mol. Cell. Biol.* (1988) PMID: 3043178
190. Gremer L, et al. *Hum. Mutat.* (2011) PMID: 20949621
191. Janakiram M, et al. *Cancer Res.* (2010) PMID: 20570890
192. Kim E, et al. *Cancer Discov* (2016) PMID: 27147599
193. Lukman S, et al. *PLoS Comput. Biol.* (2010) PMID: 20838576
194. Naguib A, et al. *J Mol Signal* (2011) PMID: 21371307
195. Prior IA, et al. *Cancer Res.* (2012) PMID: 22589270
196. Privé GG, et al. *Proc. Natl. Acad. Sci. U.S.A.* (1992) PMID: 1565661
197. Scheffzek K, et al. *Science* (1997) PMID: 9219684
198. Scholl C, et al. *Cell* (2009) PMID: 19490892
199. Smith G, et al. *Br. J. Cancer* (2010) PMID: 20147967
200. Tyner JW, et al. *Blood* (2009) PMID: 19075190
201. Valencia A, et al. *Biochemistry* (1991) PMID: 2029511
202. White Y, et al. *Nat Commun* (2016) PMID: 26854029
203. Wiest JS, et al. *Oncogene* (1994) PMID: 8058307
204. Angeles AKJ, et al. *Oncol Lett* (2019) PMID: 31289513
205. Tong JH, et al. *Cancer Biol. Ther.* (2014) PMID: 24642870
206. Loree JM, et al. *Clin Cancer Res* (2021) PMID: 34117033
207. Dale T, et al. *Nat. Chem. Biol.* (2015) PMID: 26502155
208. Mallinger A, et al. *ACS Med Chem Lett* (2016) PMID: 27326329
209. Koehler MF, et al. *ACS Med Chem Lett* (2016) PMID: 26985305
210. Schiemann K, et al. *Bioorg. Med. Chem. Lett.* (2016) PMID: 26852363
211. Witkiewicz AK, et al. *Nat Commun* (2015) PMID: 25855536
212. *Nature* (2014) PMID: 25079317
213. Barbieri CE, et al. *Nat. Genet.* (2012) PMID: 22610119
214. *Cell* (2015) PMID: 26544944
215. Kumar A, et al. *Nat. Med.* (2016) PMID: 26928463
216. Ciriello G, et al. *Cell* (2015) PMID: 26451490
217. *Nature* (2012) PMID: 23000897
218. *Nature* (2014) PMID: 24476821
219. *Nature* (2012) PMID: 22960745
220. Firestein R, et al. *Int. J. Cancer* (2010) PMID: 19790197
221. Seo JO, et al. *Oncol. Rep.* (2010) PMID: 20514474
222. He SB, et al. *J. Exp. Clin. Cancer Res.* (2011) PMID: 22104393
223. Porter DC, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2012) PMID: 22869755
224. Kim MY, et al. *Int. J. Oncol.* (2011) PMID: 21344156
225. Galbraith MD, et al. *Transcription ()* PMID: 21327159
226. Rzymiski T, et al. *Biochim. Biophys. Acta* (2015) PMID: 26006748
227. Firestein R, et al. *Nature* (2008) PMID: 18794900
228. Morris EJ, et al. *Nature* (2008) PMID: 18794899
229. Adler AS, et al. *Cancer Res.* (2012) PMID: 22345154
230. McClelland ML, et al. *J. Pathol.* (2015) PMID: 26235356
231. Li N, et al. *Nat. Cell Biol.* (2014) PMID: 25344755
232. Gu W, et al. *Cell Cycle* (2013) PMID: 23454913
233. Xu W, et al. *J Genet Genomics* (2011) PMID: 22035865
234. Cooper BW, et al. *Clin Lymphoma Myeloma Leuk* (2015) PMID: 25776192
235. Williams CB, et al. *Pharmacotherapy* (2013) PMID: 23798029
236. Heidel F, et al. *Blood* (2006) PMID: 16150941
237. Stone RM, et al. *Blood* (2005) PMID: 15345597
238. Perl AE, et al. *N. Engl. J. Med.* (2019) PMID: 31665578
239. Perl AE, et al. *Lancet Oncol.* (2017) PMID: 28645776
240. Usuki K, et al. *Cancer Sci.* (2018) PMID: 30039554
241. et al., 2021; ASH Abstract 1272
242. Moreira RB, et al. *Case Rep Oncol ()* PMID: 25848357
243. Lim SH, et al. *Oncotarget* (2017) PMID: 27906677
244. Alvarez et al., 2019; AACR Abstract CT146/12
245. Zhang S, et al. *J. Leukoc. Biol.* (1999) PMID: 10080542
246. Mizuki M, et al. *Blood* (2000) PMID: 11090077
247. Chen W, et al. *Mol. Cancer* (2010) PMID: 21067588
248. Zack TI, et al. *Nat. Genet.* (2013) PMID: 24071852
249. Beroukhir R, et al. *Nature* (2010) PMID: 20164920
250. Verstovsek S, et al. *Br. J. Haematol.* (2013) PMID: 23480528
251. Naqvi K, et al. *Expert Opin Investig Drugs* (2011) PMID: 21635221
252. Verstovsek S, et al. *Blood* (2012) PMID: 22718840
253. Swiatek-Machado K, et al. *Cancer Biol. Ther.* (2012) PMID: 22555804
254. Yan S, et al. *Oncotarget* (2013) PMID: 23531921
255. Wang Y, et al. *Curr Opin Investig Drugs* (2010) PMID: 21154128
256. Liu D, et al. *Oncol Lett* (2017) PMID: 28989534
257. Jatiati SS, et al. *Genes Cancer* (2010) PMID: 21442038
258. *Curr. Top. Microbiol. Immunol.* (2012) PMID: 21823028
259. Pansky A, et al. *Int. J. Cancer* (2000) PMID: 10699955
260. Ehata S, et al. *Cancer Sci.* (2007) PMID: 17129361
261. Rausch MP, et al. *Anticancer Res.* (2009) PMID: 19528470
262. Li YY, et al. *Clin. Cancer Res.* (2015) PMID: 25589618
263. Fleming NI, et al. *Cancer Res.* (2013) PMID: 23139211
264. Uchida K, et al. *Cancer Res.* (1996) PMID: 8971158
265. Boulay JL, et al. *Genes Chromosomes Cancer* (2001) PMID: 11391794
266. Xie W, et al. *Cancer J ()* PMID: 12967141
267. Koumoundourou D, et al. *Anticancer Res. ()* PMID: 17695423
268. Xie W, et al. *Cancer Res.* (2002) PMID: 11809701
269. Liapis G, et al. *APMIS* (2007) PMID: 17295676
270. Wu SK, et al. *World J. Gastroenterol.* (2007) PMID: 17729405
271. Macias-Silva M, et al. *Cell* (1996) PMID: 8980228
272. Xie W, et al. *Head Neck* (2013) PMID: 22275186
273. Mangone FR, et al. *Mol. Cancer* (2010) PMID: 20462450
274. Lebrun JJ, et al. *Mol. Endocrinol.* (1999) PMID: 9892009
275. Lin X, et al. *Cell* (2006) PMID: 16751101
276. Funaba M, et al. *J. Biol. Chem.* (2002) PMID: 12193595
277. Massagué J, et al. *Genes Dev.* (2005) PMID: 16322555
278. Matsuzaki K, et al. *Cancer Res.* (2009) PMID: 19531654
279. Wu Y, et al. *PLoS ONE* (2012) PMID: 22539990
280. Hirai H, et al. *Cancer Biol. Ther.* (2010) PMID: 20107315
281. Bridges KA, et al. *Clin. Cancer Res.* (2011) PMID: 21799033
282. Rajeshkumar NV, et al. *Clin. Cancer Res.* (2011) PMID: 21389100
283. Osman AA, et al. *Mol. Cancer Ther.* (2015) PMID: 25504633
284. Xu L, et al. *Mol. Cancer Ther.* (2002) PMID: 12489850
285. Xu L, et al. *Mol. Med.* (2001) PMID: 11713371
286. Camp ER, et al. *Cancer Gene Ther.* (2013) PMID: 23470564
287. Kim SS, et al. *Nanomedicine* (2015) PMID: 25240597
288. Pirollo KF, et al. *Mol. Ther.* (2016) PMID: 27357628
289. Hajdenberg et al., 2012; ASCO Abstract e15010
290. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27601554
291. Moore et al., 2019; ASCO Abstract 5513
292. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27998224
293. Oza et al., 2015; ASCO Abstract 5506
294. Lee J, et al. *Cancer Discov* (2019) PMID: 31315834
295. Méndez E, et al. *Clin. Cancer Res.* (2018) PMID: 29535125
296. Seligmann JF, et al. *J Clin Oncol* (2021) PMID: 34538072
297. Lehmann S, et al. *J. Clin. Oncol.* (2012) PMID: 22965953
298. Mohell N, et al. *Cell Death Dis* (2015) PMID: 26086967
299. Franssen Å, et al. *J Ovarian Res* (2016) PMID: 27179933
300. Gourley et al., 2016; ASCO Abstract 5571
301. Kwok M, et al. *Blood* (2016) PMID: 26563132
302. Boudny M, et al. *Haematologica* (2019) PMID: 30975914
303. Dillon MT, et al. *Mol. Cancer Ther.* (2017) PMID: 28062704
304. Middleton FK, et al. *Cancers (Basel)* (2018) PMID: 30127241
305. Mogi A, et al. *J. Biomed. Biotechnol.* (2011) PMID: 21331359
306. Tekpli X, et al. *Int. J. Cancer* (2013) PMID: 23011884
307. Vignot S, et al. *J. Clin. Oncol.* (2013) PMID: 23630207
308. Maeng CH, et al. *Anticancer Res.* (2013) PMID: 24222160
309. Cortot AB, et al. *Clin Lung Cancer* (2014) PMID: 24169260
310. Itakura M, et al. *Br. J. Cancer* (2013) PMID: 23922113

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1452195-01**
APPENDIX
References

- | | | |
|---|--|--|
| <p>311. Seo JS, et al. Genome Res. (2012) PMID: 22975805</p> <p>312. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675</p> <p>313. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249</p> <p>314. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609</p> <p>315. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130</p> <p>316. Zerdoumi Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496</p> <p>317. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113</p> <p>318. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290</p> | <p>319. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23355100</p> <p>320. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776</p> <p>321. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316</p> <p>322. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208</p> <p>323. Laloo F, et al. Lancet (2003) PMID: 12672316</p> <p>324. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713</p> <p>325. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837</p> <p>326. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838</p> | <p>327. Xie M, et al. Nat. Med. (2014) PMID: 25326804</p> <p>328. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404</p> <p>329. Severson EA, et al. Blood (2018) PMID: 29678827</p> <p>330. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212</p> <p>331. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320</p> <p>332. Chabon JJ, et al. Nature (2020) PMID: 32269342</p> <p>333. Razavi P, et al. Nat. Med. (2019) PMID: 31768066</p> |
|---|--|--|

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 20 September 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531