

ACT Onco[®] + Report

PATIENT		
Name: 許雅琪		Patient ID: 48261070
Date of Birth: Mar 14, 1982		Gender: Female
Diagnosis: Lung adenocarcinoma		
ORDERING PHYSICIAN		
Name: 江起陸醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: 1114591	Collection site: Brain	Type: FFPE tissue
Date received: Jul 13, 2022	Lab ID: AA-22-04031	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 31 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
CHEK2 R145W	-	-	Olaparib
EGFR E746_A750del (Exon 19 deletion)	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	-	-

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
CHEK2 R145W	Rucaparib	-
STAG2 R614*	Olaparib, Rucaparib	Vemurafenib
CDKN2A Homozygous deletion	Abemaciclib, Palbociclib, Ribociclib	-

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACT Onco[®] + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
APC	R1273*	51.1%
CHEK2	R145W	30.2%
EGFR	E746_A750del (Exon 19 deletion)	64.5%
STAG2	R614*	20.8%

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr9	CDKN2A	Homozygous deletion	0
Chr10	PTEN	Heterozygous deletion	1
Chr13	BRCA2	Heterozygous deletion	1
Chr15	RAD51	Heterozygous deletion	1
Chr17	FLCN	Heterozygous deletion	1
Chr3	MLH1	Heterozygous deletion	1
Chr9	PTCH1, TSC1	Heterozygous deletion	1

- Fusions

Fusion Gene & Exon	Transcript ID
Not tested	

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	2.6 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 68% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco[®] + to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 1		
EGFR E746_A750del (Exon 19 deletion)	Afatinib, Dacomitinib, Erlotinib, Gefitinib, Osimertinib	sensitive
Level 3A		
CHEK2 R145W	Olaparib	sensitive
Level 3B		
CDKN2A Homozygous deletion	Abemaciclib, Palbociclib, Ribociclib	sensitive
CHEK2 R145W	Rucaparib	sensitive
Level 4		
STAG2 R614*	Olaparib, Rucaparib	sensitive
STAG2 R614*	Vemurafenib	resistant

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
EGFR aberration	Likely associated with WORSE response to ICIs

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

APC R1273*

Biological Impact

APC (adenomatous polyposis coli) gene encodes a negative regulator of the WNT/ β -catenin signaling pathway. It binds to β -catenin, leading to its degradation and subsequently inhibits transcriptional activation^[1]. APC is also associated with cell migration and adhesion, apoptosis, and DNA repair^{[2][3]}. APC mutations are commonly observed in colorectal cancer and are also reported in lung, breast, prostate, uterine, skin, bladder, stomach and head and neck cancers (cBioPortal, MSKCC, April 2015). R1273* mutation results in a premature truncation of the APC protein at amino acid 1273 (UniProtKB). This mutation is predicted to lead to a loss of APC function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

A study of colorectal cancer patients (n= 468) indicated that MSS tumors without any APC mutation carry a worse prognosis than single APC mutation tumors. However, tumors with two APC, KRAS, and TP53 mutations confer the poorest survival among all the subgroups examined^[4].

CHEK2 R145W

Biological Impact

The checkpoint kinase 2 (CHEK2 or CHK2) gene encodes a serine/threonine protein kinase involved in transducing DNA damage signals that are required for both the intra-S phase and G2/M checkpoints^[5]. CHEK2 heterozygosity has been shown to cause haploinsufficient phenotypes that can contribute to tumorigenesis through inappropriate S phase entry, accumulation of DNA damage during replication, and failure to restrain mitotic entry^{[6][7]}. CHEK2 aberrations are associated with glioblastoma, breast, ovarian, prostate, colorectal, gastric, thyroid, and lung cancers^{[8][9][10][11][12]}.

CHEK2 R145W lies within the FHA domain of the CHEK2 protein (UniProtKB). R145W confers a loss of function to the CHEK2 protein as demonstrated by reduced protein stability and kinase activity, failure to bind several proteins, and cannot be activated by gamma irradiation in vitro^{[13][14]}.

Therapeutic and prognostic relevance

In May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[15].

In addition, CHEK2 has been determined as an inclusion criterion for the trials evaluating rucaparib efficacy in ovarian cancer or prostate cancer(NCT03533946)^{[16][17]}, niraparib efficacy in melanoma (NCT03925350), pancreatic cancer (NCT03553004), prostate cancer (NCT02854436), and any malignancy, except prostate (NCT03207347), and talazoparib efficacy in HER2-negative breast cancer (NCT02401347), prostate cancer (NCT03148795), and lung cancer (NCT03377556), respectively.

In a phase 2 trial, two prostate cancer patients harboring CHEK2 homozygous deletion was enrolled. One of the two patients had a response to olaparib^[18].

EGFR E746_A750del (Exon 19 deletion)

Biological Impact

The EGFR gene encodes for the Epidermal Growth Factor Receptor, a receptor tyrosine kinase which binds to its ligands, including Epidermal Growth Factor (EGF) and Transforming Growth Factor-alpha (TGF-alpha), activates

ACT Onco[®] + Report

downstream signaling pathways, including the canonical oncogenic MAPK and PI3K/AKT/mTOR signaling cascades^[19]. Increased EGFR activity by mutations and/or amplification of the EGFR gene has been described in a wide range of cancers, such as lung, brain, colorectal and head and neck cancer^[20]. Mutations in the kinase domain of EGFR are commonly observed in non-small cell lung cancer (NSCLC), resulting in a constitutively activated form of the receptor^[21]. On the other hand, in the brain and colorectal cancers, the most prevalent EGFR alteration is copy number amplification that results in receptor overexpression^[22].

EGFR E746_A750del is located within the protein kinase domain of the EGFR protein, resulting in the deletion of five amino acids from amino acids 746 to 750 (UniProtKB)^[23]. E746_A750del confers a gain of function to the EGFR protein, as demonstrated by increased EGFR kinase activity, activation of p44/42 MAPK and AKT, oncogenic transformation of the cells in vitro and promoting tumor growth in xenograft models^{[24][25][26]}.

EGFR exon 19 deletions are in-frame deletions of 9–24 nucleotides in exon 19 centred around codons 746–750 of the kinase domain of EGFR. The two most common EGFR alterations, L858R mutation and exon 19 deletions can result in constitutive activation of signal transduction pathways, leading to cell proliferation or anti-apoptosis without ligand binding^[27].

Therapeutic and prognostic relevance

There is accumulated clinical evidence suggested that patients with MDM2/MDM4 amplification or EGFR aberrations exhibited poor clinical outcome and demonstrated a significantly increased rate of tumor growth (hyper-progression) after receiving immune checkpoint (PD-1/PD-L1) inhibitors therapies^[28] (Annals of Oncology (2017) 28 (suppl_5): v403-v427. 10.1093/annonc/mdx376).

In a retrospective study, treatment with gefitinib resulted in partial response in 7 patients and stable disease in 1 patient who are with lung adenocarcinoma harboring EGFR E746_A750del^[29]. Preclinically, EGFR E746_A750del is sensitive to gefitinib, erlotinib, afatinib and osimertinib, demonstrated by the inhibition of EGFR signaling and cell proliferation in vitro^{[30][31][32]}.

The first- and second-generation EGFR tyrosine kinase inhibitors (EGFR-TKIs), dacomitinib, erlotinib, gefitinib and afatinib have been approved by the U.S. Food and Drug Administration (FDA) as the first-line treatment in non-small cell lung cancer (NSCLC) patients whose tumor carries EGFR exon 19 deletion or L858R mutation^{[33][34][35]}, as detected by a U.S. FDA-approved test. A Phase III clinical trial (NCT01774721) show that dacomitinib significantly improved progression-free survival over gefitinib in first-line treatment of patients with EGFR-mutation-positive NSCLC^[33]. Another Phase III clinical trial (NCT00949650) demonstrated that median progression-free survival (PFS) among lung cancer patients with exon 19 deletion or L858R EGFR mutation (n=308) was 13.6 months for afatinib and 6.9 months for chemotherapy. The EGFR T790M mutation has been demonstrated to confer resistance to TKIs (dacomitinib, gefitinib, erlotinib, and afatinib) in preclinical and clinical studies^{[36][37][38][39]}.

Osimertinib, a third-generation irreversible EGFR-TKI that selectively inhibits both EGFR-TKI-sensitizing and EGFR T790M resistance mutations, has been approved by the U.S. FDA for NSCLC patient harboring T790M mutation-positive tumor^{[40][41][42]}. Results from a double-blind, Phase 3 trial further showed that osimertinib significantly demonstrated longer PFS than standard EGFR-TKIs (18.9 months vs. 10.2 months) in previously untreated EGFR mutation-positive (exon 19 deletion or L858R) advanced NSCLC^[43].

STAG2 R614*

Biological Impact

The Stromal antigen 2 (STAG2) gene encodes a subunit of the cohesin complex, which regulates the separation of sister chromatids during cell division^{[44][45]}. Inactivation of STAG2 has been shown to promote tumorigenesis by delaying telomere shortening and postponed senescence onset^[46].

ACT Onco[®] + Report

STAG2 mutations are the most common in bladder cancer, but also found in Ewing sarcoma, melanoma, glioblastoma and other cancers^{[47][44]}.

R614* mutation results in a premature truncation of the STAG2 protein at amino acid 614 (UniProtKB). This mutation is predicted to lead to a loss of STAG2 function, despite not having characterized in the literature.

Therapeutic and prognostic relevance

Preclinical data showed that glioblastomas harboring STAG2 mutations are more sensitive to PARP inhibition^[48]. Both clinical and preclinical studies demonstrated that inactivation of STAG2 confers resistance to BRAF inhibitor vemurafenib in melanoma via the reactivation of MAPK signaling cascade^[49].

BRCA2 Heterozygous deletion

Biological Impact

The BRCA2 gene encodes a tumor suppressor involved in the homologous recombination pathway for double-strand DNA repair^[50]. BRCA2 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[51]. BRCA2 germline mutations confer an increased lifetime risk of developing breast, ovarian, prostate and pancreatic cancer, limited reports of related gastric cancer, and Fanconi anemia subtype D1-associated risk of brain cancer, medulloblastoma, pharyngeal cancer, chronic lymphocytic leukemia and acute myeloid leukemia^[52]. Somatic mutations in BRCA2 are highest in colorectal, non-small cell lung cancer (NSCLC), and ovarian cancers^[53].

Therapeutic and prognostic relevance

The U.S. FDA has approved olaparib in advanced ovarian cancer under several settings including (1) first-line maintenance treatment for patients with deleterious or suspected deleterious germline or somatic BRCA mutation who are in complete or partial response to first-line platinum-based chemotherapy^[54]; (2) in combination with bevacizumab as first-line maintenance treatment for patients with homologous recombination deficiency (HRD)-positive status^[55]; (3) maintenance treatment for patients with germline BRCA-mutated recurrent ovarian cancer who are in complete or partial response to platinum-based chemotherapy^{[56][57]}; (4) treatment for patients with germline BRCA-mutated advanced ovarian cancer who have been treated with three or more prior lines of chemotherapy^[58]. In addition, olaparib has also been approved in patients with deleterious or suspected deleterious germline BRCA-mutated, HER2-negative metastatic breast cancer who have been treated with chemotherapy in either neoadjuvant, adjuvant, or metastatic setting^[59] and germline BRCA-mutated metastatic pancreatic cancer^[60]. Of note, in May 2020, the U.S. FDA approved olaparib for the treatment of adult patients with metastatic castration-resistant prostate cancer (mCRPC) who carry mutations in homologous recombination repair (HRR) genes, including BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, RAD54L, and progressed following prior treatment with enzalutamide or abiraterone acetate^[15].

Rucaparib has been approved for the maintenance treatment of adult patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in a complete or partial response to platinum-based chemotherapy and patients with BRCA-mutated epithelial ovarian, fallopian tube, or primary peritoneal cancer, who have been treated with two or more chemotherapies^{[16][61]}. In May 2020, the U.S. FDA also approved rucaparib to treat adult patients with a deleterious BRCA mutation-associated metastatic castration-resistant prostate cancer (mCRPC) who have been treated with androgen receptor-directed therapy and a taxane-based chemotherapy (TRITON2, NCT02952534).

The U.S. FDA also approved niraparib for the maintenance treatment of patients with recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer who are in response to platinum-based chemotherapy and patients who have been treated with three or more prior lines of chemotherapy and associated with HRD positive status^{[62][63][64]}. In addition, talazoparib for patients with deleterious or suspected deleterious germline BRCA-mutated, HER2 negative locally advanced or metastatic breast cancer^[65].

ACT Onco[®] + Report

CDKN2A Homozygous deletion

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^{[66][67][68]}. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[69]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[70][71]}.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[72][73]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[74][75][76]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[77][78][79]}. However, CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[73][80][81]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[75]. Administration of combined palbociclib and MEK inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[82].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[83].

FLCN Heterozygous deletion

Biological Impact

The FLCN gene encodes the tumor suppressor, Folliculin, a GTPase activating protein (GAP) for RagC/D GTPase proteins involved in amino acid sensing and signaling to mTORC1^[84]. FLCN has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[85][86]}. Inactivation of the FLCN gene by mutation or deletion results in the activation of the mTOR pathway and AKT signaling^{[87][88]}. Germline mutation of the FLCN gene causes the Birt-Hogg-Dubé syndrome, a rare disorder that is characterized by benign hamartomatous skin lesions and an increased risk of pneumothorax and renal tumors^[89].

Therapeutic and prognostic relevance

In a prospective Phase 2 study, four anaplastic thyroid cancer (ATC)/ poorly differentiated thyroid cancer (PDTC) patients who had PI3K/mTOR/AKT alterations, including TSC2, FLCN or NF1, showed impressive progression-free survival (PFS) of 15.2 months after receiving everolimus^[90]. mTOR inhibition via rapamycin also demonstrated potential in inhibiting the growth of renal cells deficient in FLCN in the preclinical setting^[91].

ACT Onco[®] + Report

MLH1 Heterozygous deletion

Biological Impact

The MutL protein homolog 1 (MLH1) gene encodes a tumor suppressor that dimerizes with PMS2 protein to form a component of the DNA mismatch repair (MMR) system^[92]. Deletion of one copy of the MLH1 gene resulted in haploinsufficiency in the correction of small insertions/deletions (indels), and could be a driving force in pancreatic and renal carcinogenesis^[93]. Genetic alterations such as mutation, loss of heterozygosity or epigenetic silencing could lead to inactivation of MLH1 and are associated with a broad spectrum of cancers, including a subset of sporadic colon, gastric and endometrial cancers, as well as the hereditary non-polyposis colon cancer (HNPCC, also known as Lynch syndrome)^{[94][95][96]}.

Therapeutic and prognostic relevance

Currently, there are no FDA-approved medications specifically targeting MLH1. A screening test for microsatellite instability (MSI) is commonly used to identify an MMR-deficient tumor in the clinic^{[97][98]}. Pembrolizumab (KEYTRUDA), an inhibitor targeting programmed cell death 1 (PD-1), has been approved by the U.S. FDA for adult and pediatric patients with unresectable or metastatic, microsatellite instability-high (MSI-H) or mismatch repair-deficient cancer. Down-regulation of genes involved in the MMR pathway such as MLH1, MSH2 and, MSH6 in high-grade serous epithelial ovarian cancer cell lines rendered cells sensitive to PARP inhibitors^[99].

PTCH1 Heterozygous deletion

Biological Impact

The PTCH1 (protein patched homolog 1) gene encodes a multi-pass transmembrane receptor for sonic hedgehog (shh), a tumor suppressor that acts to repress shh signaling in the absence of ligand^[100]. Inactivation of PTCH1 results in hedgehog ligand-independent activation of SMO, causing a downstream activation of the pathway and lead to the neoplastic growth^{[101][102]}. Recurrent PTCH1 mutations have been reported in sporadic basal cell carcinoma (BCCs) and medulloblastoma^{[103][104][105][106]}. Germline PTCH1 mutations are associated with the nevoid basal cell carcinoma syndrome (NBCCS, Gorlin syndrome), predisposing patients to basal cell carcinoma and medulloblastoma^[104]. PTCH1 is a haploinsufficient tumor suppressor gene with one copy loss may be sufficient to promote tumor development in mice^{[101][107]}.

Therapeutic and prognostic relevance

Vismodegib and sonidegib are small molecule inhibitors of SMO approved by the U.S. FDA for the treatment of patients with basal cell carcinoma^{[108][109][110][111]}. A heavily pretreated patient with metastatic medulloblastoma harboring loss-of-heterozygosity and somatic mutation of PTCH1 showed rapid regression of the tumor after treated with vismodegib^[112]. Furthermore, a phase II study demonstrated that vismodegib treatment results in extended progression-free survival (PFS) in patients with loss-of-heterozygosity, SHH-driven medulloblastoma^[113]. In the phase II MyPathway trial, three advanced solid tumors patients harboring PTCH1 loss-of-function mutations had partial responses to vismodegib treatment^[114].

PTEN Heterozygous deletion

Biological Impact

The phosphatase and tensin homolog deleted on chromosome ten (PTEN) gene encodes a lipid/protein phosphatase that is important for the regulation of cell proliferation, survival, homologous recombination and maintenance of genomic integrity^{[115][116]}. PTEN acts as an essential tumor suppressor by antagonizing the PI3K/AKT/mTOR signaling pathway^[117]. PTEN is a haploinsufficient tumor suppressor gene, in which having only one copy of the wild-type allele does not produce enough protein product to execute wild-type functions^{[118][119][120]}. Germline loss-of-function PTEN mutations are found in approximately 80% of patients with Cowden syndrome, a disorder that is associated with high-penetrance breast and thyroid cancer^{[121][122][123]}. Somatic mutations or monoallelic loss of PTEN is regularly observed

ACT Onco[®] + Report

in a significant fraction of human cancers, including sporadic breast cancer, colon cancer, endometrial cancer, prostate cancer, and glioblastoma^{[124][125][126][127][128]}.

Therapeutic and prognostic relevance

Somatic loss of PTEN results in aberrant activation of PI3K/AKT/mTOR signaling pathway and provides a mechanistic rationale for PI3K pathway inhibitors treatment^{[129][130]}. Preclinical studies demonstrated that PTEN deficiency was associated with increased sensitivity to PI3K pathway inhibitors in selected cancer subtypes^{[131][132][133][134][135][136]}. Moreover, early clinical data also indicated that PTEN loss was associated with improved response and longer PFS in patients with advanced breast cancer^[137], advanced pancreatic neuroendocrine tumors^[138], and metastatic castration-resistant prostate cancer treated with mTORC1 inhibitor, everolimus^[139].

Several groups found that PTEN loss was generally associated with poor response to trastuzumab therapy, whether this agent was administered in the neoadjuvant, adjuvant, or metastatic settings^{[140][141][142][143][144]}.

Loss of PTEN expression in advanced colorectal cancer (CRC) has been linked with resistance to anti-EGFR mAbs like cetuximab and panitumumab^{[145][146][147][148][149][150]}. However, encouraging anti-tumor activity of the combination of an EGFR antibody and a mTORC1 inhibitor (everolimus or temsirolimus) have been reported in early-phase clinical studies (J Clin Oncol. 2011;29 (suppl): abstr 3587; J Clin Oncol. 2013;31 (suppl): abstr 608). Ongoing phase I/II studies testing combinations of EGFR antibodies and PI3K/AKT/mTOR pathway inhibitors (e.g., NCT01816984, NCT01252628, NCT01719380) will provide larger numbers of patients to assess the role of PTEN status in therapeutic response.

Preclinical studies showed that loss of PTEN expression in EGFR mutant cells was associated with decreased sensitivity to EGFR TKIs, erlotinib and gefitinib^{[151][152]}. Inhibition of the PI3K/AKT/mTOR signal pathway has been shown to be an effective strategy to radiosensitize NSCLC cells harboring the EGFR activating mutation that acquires resistance to both TKIs due to PTEN loss or inactivation mutations^[153].

Loss or biallelic inactivation of PTEN is associated with resistance to anti-PD-1 checkpoint blockade therapies, including pembrolizumab and nivolumab in melanoma and leiomyosarcoma patients^{[154][155][156]}.

PTEN loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in metastatic biliary tract cancer (NCT04042831); talazoparib efficacy in HER2-negative breast cancer (NCT02401347), and niraparib efficacy in breast cancer (NCT04508803) or any malignancy (except prostate) cancer (NCT03207347). Clinical data also suggested that PTEN deficient cancers may be sensitive to olaparib^[157].

RAD51 Heterozygous deletion

Biological Impact

The RAD51 gene encodes a recombinase that is crucial for homologous recombination (HR)-mediated repair of double-strand DNA breaks (DSBs) by forming complexes with known tumor suppressors including BRCA1, BRCA2, and PALB2^{[158][159][160]}. RAD51 has been characterized as a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[161]. Overexpression of RAD51 has been observed in many cancer cells, including pancreatic cancer and breast cancer and its hyperexpression is implicated in drug resistance^{[162][163][164][165][166][167][168]}. Germline mutations in RAD51 are associated with increased susceptibility to breast cancer^{[169][170][171][172]}.

Therapeutic and prognostic relevance

RAD51 loss of function mutation has been determined as an inclusion criterion for the trial evaluating olaparib efficacy in ovarian cancer^[173]; rucaparib efficacy in solid tumor (NCT04171700); talazoparib efficacy in lung cancer (NCT03377556); niraparib efficacy in pancreatic cancer (NCT03553004) or any malignancy (except prostate cancer) (NCT03207347).

ACT Onco[®] + Report

Preclinical studies showed that decreased RAD51 expression could sensitize cells to olaparib-induced tumor cell cytotoxicity^{[174][175]}.

TSC1 Heterozygous deletion

Biological Impact

The tuberous sclerosis complex 1 (TSC1) gene encodes a tumor suppressor, hamartin, a key negative regulator of the mammalian target of rapamycin (mTOR) pathway^{[176][177]}. Mutations in TSC1/TSC2 tumor suppressor genes that result in inactivation of the complex are commonly found in patients with tuberous sclerosis^{[178][179][180]}, while LOH in TSC1/TSC2 has been identified in head and neck squamous cell carcinoma (HNSCC)^[181] and endometrial cancer^[182]. Loss of single TSC1 allele (haploinsufficiency) may provide a growth advantage to bladder epithelial cells, contributing to bladder cancer development^[183]. Both TSC1 and TSC2 mutations cause the autosomal dominant genetic disorder tuberous sclerosis complex (TSC), in which individuals develop a variety of benign but often progressive neoplasms^[184].

Therapeutic and prognostic relevance

Genomic alterations with activating effects of the mTOR signaling pathway (including deletion/inactivation of TSC1/TSC2) have been shown to confer sensitivity to everolimus across multiple neoplasms, such as bladder tumors^[185], gastric, sarcoma, thyroid cancer, and HNSCC^[186]. There were case reports demonstrated the efficacy of sirolimus in malignant uterine perivascular epithelioid cell tumors (PEComa) patients harboring mutations/deletions in TSC1 and TSC2 genes, and temsirolimus in PEComa patients with hyperactivated mTOR pathway. Genomic profiling analysis of GOG248, a Phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer showed that mutations in AKT1, TSC1 and TSC2 may predict clinical benefit from temsirolimus^[187]. Recent studies indicate that there are mTORC1-independent signaling pathways downstream of hamartin-tuberin, which may represent new therapeutic targets^[188].

Everolimus has been approved by the U.S. FDA for Tuberous Sclerosis Complex (TSC)-associated renal angiomyolipoma and Tuberous Sclerosis Complex (TSC)-associated subependymal giant cell astrocytoma (SEGA). This approval is based on the results from EXIST-1, EXIST-2, and Study 2485 trials (NCT00789828, NCT00790400, and NCT00411619).

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Abemaciclib (VERZENIO)

Abemaciclib is a cyclin-dependent kinase 4/6 (CDK4/6) inhibitor. Abemaciclib is developed and marketed by Eli Lilly under the trade name VERZENIO.

- FDA Approval Summary of Abemaciclib (VERZENIO)

monarchE NCT03155997	Breast cancer (Approved on 2021/10/12)
	HR-positive, HER2-negative Abemaciclib + tamoxifen/aromatase inhibitor vs. Tamoxifen/aromatase inhibitor [IDFS at 36 months(%): 86.1 vs. 79.0]
MONARCH 3 ^[189] NCT02246621	Breast cancer (Approved on 2018/02/26)
	HR-positive, HER2-negative Abemaciclib + anastrozole/letrozole vs. Placebo + anastrozole/letrozole [PFS(M): 28.2 vs. 14.8]
MONARCH 2 ^[81] NCT02107703	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative Abemaciclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 16.4 vs. 9.3]
MONARCH 1 ^[190] NCT02102490	Breast cancer (Approved on 2017/09/28)
	HR-positive, HER2-negative Abemaciclib [ORR(%): 19.7 vs. 17.4]

Afatinib (GILOTRIF)

Afatinib acts as an irreversible covalent inhibitor of the ErbB family of receptor tyrosine kinases, including epidermal growth factor receptor (EGFR) and erbB-2 (HER2). Afatinib is developed and marketed by Boehringer Ingelheim under the trade name GILOTRIF (United States) and GIOTRIF (Europe).

- FDA Approval Summary of Afatinib (GILOTRIF)

LUX-Lung 8 ^[191] NCT01523587	Non-small cell lung carcinoma (Approved on 2016/04/15)
	EGFR Del19/L858R Afatinib vs. Erlotinib [PFS(M): 2.4 vs. 1.9]
LUX-Lung 3 ^[192] NCT00949650	Non-small cell lung carcinoma (Approved on 2013/07/13)
	EGFR Del19/L858R Afatinib vs. Pemetrexed + cisplatin [PFS(M): 11.1 vs. 6.9]

Dacomitinib (VIZIMPRO)

Dacomitinib is an oral kinase inhibitor that targets EGFR. Dacomitinib is developed and marketed by Pfizer under the trade name VIZIMPRO.

- FDA Approval Summary of Dacomitinib (VIZIMPRO)

ARCHER 1050 ^[33] NCT01774721	Non-small cell lung carcinoma (Approved on 2018/09/27)
	EGFR Del 19/ L858R Dacomitinib vs. Gefitinib [PFS(M): 14.7 vs. 9.2]

ACT Onco[®] + Report

Erlotinib (TARCEVA)

Erlotinib is a small molecule, reversible inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Erlotinib is developed by OSI Pharmaceuticals, Genentech and Roche, and marketed by Astellas Pharm Global Development under the trade name TARCEVA.

- FDA Approval Summary of Erlotinib (TARCEVA)

RELAY NCT02411448	Non-small cell lung carcinoma (Approved on 2020/05/29)
	EGFR exon 19 deletion or exon 21 (L858R) Erlotinib + ramucirumab vs. Erlotinib + placebo [PFS(M): 19.4 vs. 12.4]
EURTAC^[193] NCT00446225	Non-small cell lung carcinoma (Approved on 2013/05/14)
	Exon 19 Del/Exon 21 substitution (L858R) Erlotinib vs. Cisplatin + gemcitabine or cisplatin + docetaxel or carboplatin + gemcitabine or carboplatin + docetaxel [PFS(M): 10.4 vs. 5.2]
PA.3^[194] NCT00026338	Pancreatic cancer (Approved on 2005/11/02)
	- Gemcitabine vs. Placebo [OS(M): 6.4 vs. 6]

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4^[195] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
BOLERO-2^[196] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2- Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
EXIST-2 NCT00790400	Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26)
	- Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
RADIANT-3^[138] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	- Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1^[197] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	- Everolimus vs. Placebo [ORR(%): 35.0]
RECORD-1^[198] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	- Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

ACT Onco[®] + Report

Gefitinib (IRESSA)

Gefitinib is a small molecule inhibitor of epidermal growth factor receptor (EGFR), a receptor tyrosine kinase. Gefitinib is developed and marketed by AstraZeneca under the trade name IRESSA.

- FDA Approval Summary of Gefitinib (IRESSA)

IFUM ^[199] NCT01203917	Non-small cell lung carcinoma (Approved on 2015/07/13)
	Exon 19 Del/Exon 21 substitution (L858R)
	Gefitinib [ORR(%): 50.0]

Niraparib (Zejula)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (Zejula)

PRIMA NCT02655016	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29)
	-
	Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
QUADRA ^[64] NCT02354586	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2019/10/23)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)
	Niraparib [ORR(%): 24.0, DOR(M): 8.3]
NOVA ^[63] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	-
	Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

Olaparib (Lynparza)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (Lynparza)

OlympiA NCT02032823	Her2-negative high-risk early breast cancer (Approved on 2022/03/11)
	gBRCA
	Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):]
PROfound ^[15] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	ATMm, BRCA1m, BRCA2m, BARD1m, BRIP1m, CDK12m, CHEK1m, CHEK2m, FANCLm, PALB2m, RAD51Bm, RAD51Cm, RAD51Dm, RAD54Lm
	Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1 ^[55] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD-positive (defined by either a deleterious or suspected deleterious BRCA mutation, and/or genomic instability)
	Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]

ACT Onco[®] + Report

POLO ^[60] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	Germline BRCA mutation (deleterious/suspected deleterious)
	Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1 ^[54] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	Germline or somatic BRCA-mutated (gBRCAm or sBRCAm)
	Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
OlympiAD ^[59] NCT02000622	Breast cancer (Approved on 2018/02/06)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative
	Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21 ^[200] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA+
	Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
Study19 ^[201] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	-
	Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]
Study 42 ^[202] NCT01078662	Ovarian cancer (Approved on 2014/12/19)
	Germline BRCA mutation (deleterious/suspected deleterious)
	Olaparib [ORR(%): 34.0, DOR(M): 7.9]

Osimertinib (TAGRISSO)

Osimertinib is a third-generation tyrosine kinase inhibitor (TKI) for patients with tumors harboring EGFR T790M mutation. Osimertinib is developed and marketed by AstraZeneca under the trade name TAGRISSO.

- FDA Approval Summary of Osimertinib (TAGRISSO)

ADAURA NCT02511106	Non-small cell lung carcinoma (Approved on 2020/12/18)
	EGFR exon 19 deletions or exon 21 L858R mutations
	Osimertinib vs. Placebo + adjuvant chemotherapy [DFS(M): NR vs. 19.6]
FLAURA ^[43] NCT02296125	Non-small cell lung carcinoma (Approved on 2018/04/18)
	EGFR Del19/L858R
	Osimertinib vs. Gefitinib or erlotinib [PFS(M): 18.9 vs. 10.2]
AURA3 ^[203] NCT02151981	Non-small cell lung carcinoma (Approved on 2017/03/30)
	EGFR T790M+
	Osimertinib vs. Chemotherapy [PFS(M): 10.1 vs. 4.4]
AURA ^[42] NCT01802632	Non-small cell lung carcinoma (Approved on 2015/11/13)
	EGFR T790M+
	Osimertinib [ORR(%): 59.0]

ACT Onco[®] + Report

Palbociclib (IBRANCE)

Palbociclib is an oral, cyclin-dependent kinase (CDK) inhibitor specifically targeting CDK4 and CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Palbociclib is developed and marketed by Pfizer under the trade name IBRANCE.

- FDA Approval Summary of Palbociclib (IBRANCE)

PALOMA-2 ^[204] NCT01740427	Breast cancer (Approved on 2017/03/31)
	ER+, HER2-
	Palbociclib + letrozole vs. Placebo + letrozole [PFS(M): 24.8 vs. 14.5]
PALOMA-3 ^[205] NCT01942135	Breast cancer (Approved on 2016/02/19)
	ER+, HER2-
	Palbociclib + fulvestrant vs. Placebo + fulvestrant [PFS(M): 9.5 vs. 4.6]

Ribociclib (KISQALI)

Ribociclib is a cyclin-dependent kinase (CDK) inhibitor specifically targeting cyclin D1/CDK4 and cyclin D3/CDK6, thereby inhibiting retinoblastoma (Rb) protein phosphorylation. Ribociclib is developed by Novartis and Astex Pharmaceuticals and marketed by Novartis under the trade name KISQALI.

- FDA Approval Summary of Ribociclib (KISQALI)

MONALEESA-2 ^[80] NCT01958021	Breast cancer (Approved on 2017/03/13)
	HR+, HER2-
	Ribociclib vs. Letrozole [PFS(M): NR vs. 14.7]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA+, sBRCA
	Rucaparib [ORR(%): 44.0, DOR(M): NE]
ARIEL3 ^[16] NCT01968213	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	All HRD tBRCA
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]
ARIEL2 ^[206] NCT01482715, NCT01891344	Ovarian cancer (Approved on 2016/12/19)
	Germline and/or somatic BRCA mutation
	Rucaparib [ORR(%): 54.0]

ACT Onco[®] + Report

Sonidegib (ODOMZO)

Sonidegib is a Hedgehog signaling pathway inhibitor by blocking its key component, smoothened (smo). Sonidegib is developed and marketed by Novartis under the trade name ODOMZO.

- FDA Approval Summary of Sonidegib (ODOMZO)

BOLT ^[110] NCT01327053	Basal cell carcinoma (Approved on 2015/07/24)
	-
	Sonidegib [ORR(%): 58.0]

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

EMBRACA ^[65] NCT01945775	Breast cancer (Approved on 2018/10/16)
	Germline BRCA mutation (deleterious/suspected deleterious) HER2-negative
	Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

[207] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	-
	Temsirolimus vs. IFN- α [OS(M): 10.9 vs. 7.3]

Vismodegib (ERIVEDGE)

Vismodegib is a cyclopamine-competitive antagonist and acts as a first-in-class Hedgehog signaling pathway inhibitor by blocking its key component smoothened (smo). Vismodegib is developed by Genentech and marketed by Roche under the trade name ERIVEDGE.

- FDA Approval Summary of Vismodegib (ERIVEDGE)

ERIVANCE BCC ^[108] NCT00833417	Basal cell carcinoma (Approved on 2012/01/30)
	-
	Vismodegib [ORR (mBCC)(%): 30.3, ORR (laBCC)(%): 42.9]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
APC	R1273*	16	c.3817A>T	NM_000038	-	51.1%	131
CHEK2	R145W	3	c.433C>T	NM_007194	COSM2935997	30.2%	377
EGFR	E746_A750del (Exon 19 deletion)	19	c.2236_2250del	NM_005228	COSM6225	64.5%	2244
STAG2	R614*	20	c.1840C>T	NM_001042751	COSM166815	20.8%	144

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ATM	N1650S	33	c.4949A>G	NM_000051	-	48.7%	832
CDK12	V605A	2	c.1814T>C	NM_016507	COSM9248152	22.0%	2872
CDKN1A	Splice region	2	c.-3G>A	NM_000389	COSM3076223	98.9%	2351
E2F3	T68I	1	c.203C>T	NM_001949	-	45.1%	752
ESR1	Splice region	-	c.452+6C>G	NM_000125	-	45.5%	145
IL7R	S105N	3	c.314G>A	NM_002185	-	36.3%	347
KEAP1	G9R	2	c.25G>A	NM_203500	COSM2812662	49.9%	2095
PTCH1	D1305Y	23	c.3913G>T	NM_000264	COSM22562	29.1%	770
PTGS2	F343L	8	c.1029C>G	NM_000963	-	54.8%	168
RAD54L	Splice region	-	c.477+6T>C	NM_003579	-	63.7%	1231
ROS1	H1999N	37	c.5995C>A	NM_002944	-	46.7%	471
UBR5	Splice region	-	c.2152-5T>C	NM_015902	-	42.2%	102
USH2A	V2333A	37	c.6998T>C	NM_206933	-	40.9%	1714


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: May 2022
- Facility retrieved: 高雄榮總
- H&E-stained section No.: 1114591
- Collection site: Brain
- Examined by: Dr. Chien-Ta Chiang
 1. The percentage of viable tumor cells in total cells in the whole slide (%): 25%
 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 60%
 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 20%
 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 10%
 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 822x
- Target Base Coverage at 100x: 90%

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 31 genes sequenced. Therefore, fusion in genes not covered by this test would

ACT Onco[®] + Report

not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at 100x $\geq 85\%$ with a mean coverage $\geq 500x$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

ACT Onco[®] + Report

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師黃靖婷 博士
Ching-Ting Huang Ph.D.
檢字第 016511 號

CT Huang

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMP1R1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTX	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRAS*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBKB	IKBKE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC1B1*
SLC1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOCS1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Gene	Therapies	Possible effect
<i>PTEN</i>	Everolimus, Niraparib, Olaparib, Rucaparib, Talazoparib, Temsirolimus	sensitive
<i>FLCN</i>	Everolimus, Temsirolimus	sensitive
<i>TSC1</i>	Everolimus, Temsirolimus	sensitive
<i>BRCA2</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>MLH1</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>RAD51</i>	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
<i>PTCH1</i>	Sonidegib, Vismodegib	sensitive
<i>PTEN</i>	Cetuximab, Erlotinib, Gefitinib, Panitumumab, Trastuzumab	resistant


SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Olaparib, Niraparib, Rucaparib, Talazoparib

ACT Onco[®] + Report

Hedgehog Signalling




1: Sonidegib, Vismodegib




1: Abemaciclib, Palbociclib, Ribociclib

ACT Onco[®] + Report



1: Everolimus, Temsirolimus

Receptor Tyrosine Kinase/Growth Factor Signalling



1: Gefitinib, Afatinib, Erlotinib, Osimertinib, Dacomitinib; 2: Afatinib

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 24200292; 2014, Curr Drug Targets;15(1):90-102
Exploiting APC function as a novel cancer therapy.
2. PMID: 18848448; 2008, Trends Cell Biol;18(12):587-96
APC shuttling to the membrane, nucleus and beyond.
3. PMID: 18662849; 2008, Cancer Lett;271(2):272-80
A novel function of adenomatous polyposis coli (APC) in regulating DNA repair.
4. PMID: 27302369; 2016, Nat Commun;7():11743
A multigene mutation classification of 468 colorectal cancers reveals a prognostic role for APC.
5. PMID: 21088254; 2011, Clin Cancer Res;17(3):401-5
Tumor suppressor CHK2: regulator of DNA damage response and mediator of chromosomal stability.
6. PMID: 15261141; 2004, Cancer Cell;6(1):45-59
Chk1 is haploinsufficient for multiple functions critical to tumor suppression.
7. PMID: 15539958; 2005, Cell Cycle;4(1):131-9
Chk1 is essential for tumor cell viability following activation of the replication checkpoint.
8. PMID: 23296741; 2013, Fam Cancer;12(3):473-8
The risk of gastric cancer in carriers of CHEK2 mutations.
9. PMID: 24713400; 2014, Hered Cancer Clin Pract;12(1):10
A risk of breast cancer in women - carriers of constitutional CHEK2 gene mutations, originating from the North - Central Poland.
10. PMID: 25583358; 2015, Int J Cancer;137(3):548-52
CHEK2 mutations and the risk of papillary thyroid cancer.
11. PMID: 12052256; 2002, Breast Cancer Res;4(3):R4
Mutation analysis of the CHK2 gene in breast carcinoma and other cancers.
12. PMID: 15125777; 2004, Mol Cancer;3():14
CHK2 kinase expression is down-regulated due to promoter methylation in non-small cell lung cancer.
13. PMID: 11053450; 2001, J Biol Chem;276(4):2971-4
Characterization of tumor-associated Chk2 mutations.
14. PMID: 12049740; 2002, Mol Cell;9(5):1045-54
Structural and functional versatility of the FHA domain in DNA-damage signaling by the tumor suppressor kinase Chk2.
15. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
16. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
17. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
18. PMID: 26510020; 2015, N Engl J Med;373(18):1697-708
DNA-Repair Defects and Olaparib in Metastatic Prostate Cancer.
19. PMID: 18045542; 2007, Cell;131(5):1018

ACT Onco[®] + Report

SnapShot: EGFR signaling pathway.

20. PMID: 10880430; 2000, EMBO J;19(13):3159-67
The ErbB signaling network: receptor heterodimerization in development and cancer.
21. PMID: 15329413; 2004, Proc Natl Acad Sci U S A;101(36):13306-11
EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib.
22. PMID: 11426640; 2000, Oncogene;19(56):6550-65
The EGF receptor family as targets for cancer therapy.
23. PMID: 17318210; 2007, Nat Rev Cancer;7(3):169-81
Epidermal growth factor receptor mutations in lung cancer.
24. PMID: 16373402; 2006, FASEB J;20(2):311-3
Dimerization and the signal transduction pathway of a small in-frame deletion in the epidermal growth factor receptor.
25. PMID: 16912195; 2006, Cancer Res;66(16):8163-71
Kinetic analysis of epidermal growth factor receptor somatic mutant proteins shows increased sensitivity to the epidermal growth factor receptor tyrosine kinase inhibitor, erlotinib.
26. PMID: 29533785; 2018, Cancer Cell;33(3):450-462.e10
Systematic Functional Annotation of Somatic Mutations in Cancer.
27. PMID: 22263017; 2010, J Thorac Dis;2(1):48-51
Epidermal growth factor receptor (EGFR) in lung cancer: an overview and update.
28. PMID: 28351930; 2017, Clin Cancer Res;23(15):4242-4250
Hyperprogressors after Immunotherapy: Analysis of Genomic Alterations Associated with Accelerated Growth Rate.
29. PMID: 18981003; 2008, Clin Cancer Res;14(21):7060-7
Effects of erlotinib in EGFR mutated non-small cell lung cancers with resistance to gefitinib.
30. PMID: 29467275; 2018, Mol Cancer Ther;17(4):740-750
Pharmacological and Structural Characterizations of Naquotinib, a Novel Third-Generation EGFR Tyrosine Kinase Inhibitor, in EGFR-Mutated Non-Small Cell Lung Cancer.
31. PMID: 31314158; 2019, FEBS Open Bio;9(10):1689-1704
Characterization of epidermal growth factor receptor (EGFR) P848L, an unusual EGFR variant present in lung cancer patients, in a murine Ba/F3 model.
32. PMID: 33395611; 2021, Lung Cancer;152():135-142
Extensive functional evaluation of exon 20 insertion mutations of EGFR.
33. PMID: 28958502; 2017, Lancet Oncol;18(11):1454-1466
Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial.
34. PMID: 24868098; 2014, Oncologist;19(7):774-9
U.S. Food and Drug Administration approval summary: Erlotinib for the first-line treatment of metastatic non-small cell lung cancer with epidermal growth factor receptor exon 19 deletions or exon 21 (L858R) substitution mutations.
35. PMID: 23982599; 2013, Drugs;73(13):1503-15
Afatinib: first global approval.
36. PMID: 27912760; 2016, J Biomed Sci;23(1):86
Update on recent preclinical and clinical studies of T790M mutant-specific irreversible epidermal growth factor receptor tyrosine kinase inhibitors.

ACT Onco[®] + Report

37. PMID: 15728811; 2005, N Engl J Med;352(8):786-92
EGFR mutation and resistance of non-small-cell lung cancer to gefitinib.
38. PMID: 26862733; 2016, Oncotarget;7(11):12404-13
The mechanism of acquired resistance to irreversible EGFR tyrosine kinase inhibitor-afatinib in lung adenocarcinoma patients.
39. PMID: 29410323; 2018, J Thorac Oncol;13(5):727-731
EGFR T790M and C797S Mutations as Mechanisms of Acquired Resistance to Dacomitinib.
40. PMID: 24893891; 2014, Cancer Discov;4(9):1046-61
AZD9291, an irreversible EGFR TKI, overcomes T790M-mediated resistance to EGFR inhibitors in lung cancer.
41. PMID: 27071706; 2016, J Hematol Oncol;9():34
Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer.
42. PMID: 25923549; 2015, N Engl J Med;372(18):1689-99
AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer.
43. PMID: 29151359; 2018, N Engl J Med;378(2):113-125
Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer.
44. PMID: 24856830; 2014, BMB Rep;47(6):299-310
Cohesin gene mutations in tumorigenesis: from discovery to clinical significance.
45. PMID: 24854081; 2014, Nat Rev Cancer;14(6):389-93
Cohesin in cancer: chromosome segregation and beyond.
46. PMID: 28819029; 2017, Cancer Res;77(20):5530-5542
Loss of Tumor Suppressor STAG2 Promotes Telomere Recombination and Extends the Replicative Lifespan of Normal Human Cells.
47. PMID: 24930350; 2014, MMW Fortschr Med;156 Spec No 1(1):64-7; quiz 68
[Patient with fever].
48. PMID: 24356817; 2014, Mol Cancer Ther;13(3):724-32
Glioblastoma cells containing mutations in the cohesin component STAG2 are sensitive to PARP inhibition.
49. PMID: 27500726; 2016, Nat Med;22(9):1056-61
Loss of cohesin complex components STAG2 or STAG3 confers resistance to BRAF inhibition in melanoma.
50. PMID: 11239455; 2001, Mol Cell;7(2):263-72
BRCA2 is required for homology-directed repair of chromosomal breaks.
51. PMID: 17597348; 2007, Ann Surg Oncol;14(9):2510-8
Heterogenic loss of the wild-type BRCA allele in human breast tumorigenesis.
52. PMID: 22193408; 2011, Nat Rev Cancer;12(1):68-78
BRCA1 and BRCA2: different roles in a common pathway of genome protection.
53. PMID: 27283171; 2016, J Natl Compr Canc Netw;14(6):795-806
The Relevance of Hereditary Cancer Risks to Precision Oncology: What Should Providers Consider When Conducting Tumor Genomic Profiling?
54. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
55. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
56. PMID: 28884698; 2017, Lancet Oncol;18(9):e510
Correction to Lancet Oncol 2017; 18: 1274-84.

ACT Onco[®] + Report

57. PMID: 22452356; 2012, N Engl J Med;366(15):1382-92
Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer.
58. PMID: 26187614; 2015, Clin Cancer Res;21(19):4257-61
FDA Approval Summary: Olaparib Monotherapy in Patients with Deleterious Germline BRCA-Mutated Advanced Ovarian Cancer Treated with Three or More Lines of Chemotherapy.
59. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
60. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
61. PMID: 28882436; 2017, Gynecol Oncol;147(2):267-275
Antitumor activity and safety of the PARP inhibitor rucaparib in patients with high-grade ovarian carcinoma and a germline or somatic BRCA1 or BRCA2 mutation: Integrated analysis of data from Study 10 and ARIEL2.
62. PMID: 31562799; 2019, N Engl J Med;381(25):2391-2402
Niraparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
63. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
64. PMID: 30948273; 2019, Lancet Oncol;20(5):636-648
Niraparib monotherapy for late-line treatment of ovarian cancer (QUADRA): a multicentre, open-label, single-arm, phase 2 trial.
65. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talazoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
66. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
67. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
68. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
69. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.
70. PMID: 7550353; 1995, Nat Genet;11(2):210-2
Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.
71. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
72. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
73. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
74. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
75. PMID: 27217383; 2016, Cancer Discov;6(7):740-53

ACT Onco[®] + Report

Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.

76. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
77. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
78. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
79. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.
80. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.
81. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
82. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
83. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
84. PMID: 24095279; 2013, Mol Cell;52(4):495-505
The folliculin tumor suppressor is a GAP for the RagC/D GTPases that signal amino acid levels to mTORC1.
85. PMID: 26342594; 2016, Fam Cancer;15(1):127-32
Birt-Hogg-Dubé syndrome: novel FLCN frameshift deletion in daughter and father with renal cell carcinomas.
86. PMID: 23223565; 2013, J Clin Pathol;66(3):178-86
Birt-Hogg-Dubé syndrome: clinicopathological features of the lung.
87. PMID: 19850877; 2009, Proc Natl Acad Sci U S A;106(44):18722-7
Homozygous loss of BHD causes early embryonic lethality and kidney tumor development with activation of mTORC1 and mTORC2.
88. PMID: 24908670; 2014, Hum Mol Genet;23(21):5706-19
Folliculin (Flcn) inactivation leads to murine cardiac hypertrophy through mTORC1 deregulation.
89. PMID: 15956655; 2005, J Natl Cancer Inst;97(12):931-5
High frequency of somatic frameshift BHD gene mutations in Birt-Hogg-Dubé-associated renal tumors.
90. PMID: 29301825; 2018, Clin Cancer Res;24(7):1546-1553
Genomic Correlates of Response to Everolimus in Aggressive Radioiodine-refractory Thyroid Cancer: A Phase II Study.
91. PMID: 26418749; 2015, Oncotarget;6(32):32761-73
Flcn-deficient renal cells are tumorigenic and sensitive to mTOR suppression.
92. PMID: 16873062; 2006, Cell;126(2):297-308
Endonucleolytic function of MutLalpha in human mismatch repair.
93. PMID: 22156295; 2012, Genome Res;22(2):208-19
Whole-exome sequencing of human pancreatic cancers and characterization of genomic instability caused by MLH1 haploinsufficiency and complete deficiency.

ACT Onco[®] + Report

94. PMID: 8484122; 1993, Science;260(5109):816-9
Microsatellite instability in cancer of the proximal colon.
95. PMID: 8040889; 1994, J Natl Cancer Inst;86(16):1216-21
Microsatellite instability in sporadic endometrial carcinoma.
96. PMID: 8261393; 1993, Cancer Res;53(24):5853-5
Microsatellite instability is associated with tumors that characterize the hereditary non-polyposis colorectal carcinoma syndrome.
97. PMID: 9823339; 1998, Cancer Res;58(22):5248-57
A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer.
98. PMID: 12454837; 2002, Gastroenterology;123(6):1804-11
Evaluation of tumor microsatellite instability using five quasimonomorphic mononucleotide repeats and pentaplex PCR.
99. PMID: 27374179; 2017, Oncotarget;8(25):40152-40168
Cumulative defects in DNA repair pathways drive the PARP inhibitor response in high-grade serous epithelial ovarian cancer cell lines.
100. PMID: 8906794; 1996, Nature;384(6605):176-9
Biochemical evidence that patched is the Hedgehog receptor.
101. PMID: 12016144; 2002, Carcinogenesis;23(5):727-33
Unbalanced overexpression of the mutant allele in murine Patched mutants.
102. PMID: 11130178; 2000, Cell Mol Life Sci;57(12):1720-31
Hedgehog signalling in cancer.
103. PMID: 8782823; 1996, Nat Genet;14(1):78-81
The role of the human homologue of Drosophila patched in sporadic basal cell carcinomas.
104. PMID: 8658145; 1996, Science;272(5268):1668-71
Human homolog of patched, a candidate gene for the basal cell nevus syndrome.
105. PMID: 9422511; 1998, Nature;391(6662):90-2
Activating Smoothened mutations in sporadic basal-cell carcinoma.
106. PMID: 22832583; 2012, Nature;488(7409):100-5
Dissecting the genomic complexity underlying medulloblastoma.
107. PMID: 10738305; 2000, Genes Chromosomes Cancer;28(1):77-81
Evidence that haploinsufficiency of Ptch leads to medulloblastoma in mice.
108. PMID: 22670903; 2012, N Engl J Med;366(23):2171-9
Efficacy and safety of vismodegib in advanced basal-cell carcinoma.
109. PMID: 28511673; 2017, BMC Cancer;17(1):332
Long-term safety and efficacy of vismodegib in patients with advanced basal cell carcinoma: final update of the pivotal ERIVANCE BCC study.
110. PMID: 25981810; 2015, Lancet Oncol;16(6):716-28
Treatment with two different doses of sonidegib in patients with locally advanced or metastatic basal cell carcinoma (BOLT): a multicentre, randomised, double-blind phase 2 trial.
111. PMID: 31545507; 2020, Br J Dermatol;182(6):1369-1378
Long-term efficacy and safety of sonidegib in patients with advanced basal cell carcinoma: 42-month analysis of the phase II randomized, double-blind BOLT study.
112. PMID: 19726761; 2009, N Engl J Med;361(12):1173-8
Treatment of medulloblastoma with hedgehog pathway inhibitor GDC-0449.

ACT Onco[®] + Report

113. PMID: 26169613; 2015, J Clin Oncol;33(24):2646-54
Vismodegib Exerts Targeted Efficacy Against Recurrent Sonic Hedgehog-Subgroup Medulloblastoma: Results From Phase II Pediatric Brain Tumor Consortium Studies PBTC-025B and PBTC-032.
114. PMID: 29320312; 2018, J Clin Oncol;36(6):536-542
Targeted Therapy for Advanced Solid Tumors on the Basis of Molecular Profiles: Results From MyPathway, an Open-Label, Phase IIa Multiple Basket Study.
115. PMID: 17218262; 2007, Cell;128(1):157-70
Essential role for nuclear PTEN in maintaining chromosomal integrity.
116. PMID: 18794879; 2008, Oncogene;27(41):5443-53
PTEN: a new guardian of the genome.
117. PMID: 18767981; 2009, Annu Rev Pathol;4():127-50
PTEN and the PI3-kinase pathway in cancer.
118. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
119. PMID: 11553783; 2001, Proc Natl Acad Sci U S A;98(20):11563-8
Haploinsufficiency of the Pten tumor suppressor gene promotes prostate cancer progression.
120. PMID: 20400965; 2010, Nat Genet;42(5):454-8
Subtle variations in Pten dose determine cancer susceptibility.
121. PMID: 9467011; 1998, Hum Mol Genet;7(3):507-15
Mutation spectrum and genotype-phenotype analyses in Cowden disease and Bannayan-Zonana syndrome, two hamartoma syndromes with germline PTEN mutation.
122. PMID: 24136893; 2013, J Natl Cancer Inst;105(21):1607-16
Cowden syndrome and the PTEN hamartoma tumor syndrome: systematic review and revised diagnostic criteria.
123. PMID: 21430697; 2011, Nat Rev Cancer;11(4):289-301
PTEN loss in the continuum of common cancers, rare syndromes and mouse models.
124. PMID: 18455982; 2008, Cell;133(3):403-14
Tenets of PTEN tumor suppression.
125. PMID: 9393738; 1997, Cancer Res;57(23):5221-5
MMAC1/PTEN mutations in primary tumor specimens and tumor cell lines.
126. PMID: 9829719; 1998, Clin Cancer Res;4(11):2577-83
Loss of heterozygosity and mutational analysis of the PTEN/MMAC1 gene in synchronous endometrial and ovarian carcinomas.
127. PMID: 9582022; 1998, Oncogene;16(13):1743-8
Analysis of PTEN and the 10q23 region in primary prostate carcinomas.
128. PMID: 9671321; 1998, Oncogene;17(1):123-7
Allelic loss of chromosome 10q23 is associated with tumor progression in breast carcinomas.
129. PMID: 11504908; 2001, Proc Natl Acad Sci U S A;98(18):10314-9
Enhanced sensitivity of PTEN-deficient tumors to inhibition of FRAP/mTOR.
130. PMID: 23714559; 2013, Am Soc Clin Oncol Educ Book;():
Targeting the PI3K/AKT/mTOR pathway: biomarkers of success and tribulation.
131. PMID: 20231295; 2010, J Biol Chem;285(20):14980-9
Phosphoinositide 3-kinase pathway activation in phosphate and tensin homolog (PTEN)-deficient prostate cancer cells is independent of

ACT Onco[®] + Report

receptor tyrosine kinases and mediated by the p110beta and p110delta catalytic subunits.

132. PMID: 23287563; 2013, Clin Cancer Res;19(7):1760-72
Targeting activated Akt with GDC-0068, a novel selective Akt inhibitor that is efficacious in multiple tumor models.
133. PMID: 17047067; 2006, Cancer Res;66(20):10040-7
Inhibition of mammalian target of rapamycin or apoptotic pathway induces autophagy and radiosensitizes PTEN null prostate cancer cells.
134. PMID: 22422409; 2012, Clin Cancer Res;18(6):1777-89
PIK3CA/PTEN mutations and Akt activation as markers of sensitivity to allosteric mTOR inhibitors.
135. PMID: 22662154; 2012, PLoS One;7(5):e37431
Genotype-dependent efficacy of a dual PI3K/mTOR inhibitor, NVP-BEZ235, and an mTOR inhibitor, RAD001, in endometrial carcinomas.
136. PMID: 23136191; 2012, Clin Cancer Res;18(24):6771-83
Phosphoinositide 3-kinase (PI3K) pathway alterations are associated with histologic subtypes and are predictive of sensitivity to PI3K inhibitors in lung cancer preclinical models.
137. PMID: 27091708; 2016, J Clin Oncol;34(18):2115-24
Molecular Alterations and Everolimus Efficacy in Human Epidermal Growth Factor Receptor 2-Overexpressing Metastatic Breast Cancers: Combined Exploratory Biomarker Analysis From BOLERO-1 and BOLERO-3.
138. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
139. PMID: 23582881; 2013, Eur Urol;64(1):150-8
Phase 2 trial of single-agent everolimus in chemotherapy-naïve patients with castration-resistant prostate cancer (SAKK 08/08).
140. PMID: 15324695; 2004, Cancer Cell;6(2):117-27
PTEN activation contributes to tumor inhibition by trastuzumab, and loss of PTEN predicts trastuzumab resistance in patients.
141. PMID: 20813970; 2010, Am J Pathol;177(4):1647-56
PTEN, PIK3CA, p-AKT, and p-p70S6K status: association with trastuzumab response and survival in patients with HER2-positive metastatic breast cancer.
142. PMID: 21135276; 2011, J Clin Oncol;29(2):166-73
Loss of phosphatase and tensin homolog or phosphoinositide-3 kinase activation and response to trastuzumab or lapatinib in human epidermal growth factor receptor 2-overexpressing locally advanced breast cancers.
143. PMID: 21594665; 2011, Breast Cancer Res Treat;128(2):447-56
Evaluation of the association of PIK3CA mutations and PTEN loss with efficacy of trastuzumab therapy in metastatic breast cancer.
144. PMID: 17936563; 2007, Cancer Cell;12(4):395-402
A functional genetic approach identifies the PI3K pathway as a major determinant of trastuzumab resistance in breast cancer.
145. PMID: 18700047; 2008, BMC Cancer;8():234
Potential value of PTEN in predicting cetuximab response in colorectal cancer: an exploratory study.
146. PMID: 17940504; 2007, Br J Cancer;97(8):1139-45
PTEN loss of expression predicts cetuximab efficacy in metastatic colorectal cancer patients.
147. PMID: 19398573; 2009, J Clin Oncol;27(16):2622-9
PTEN expression and KRAS mutations on primary tumors and metastases in the prediction of benefit from cetuximab plus irinotecan for patients with metastatic colorectal cancer.
148. PMID: 19953097; 2010, Br J Cancer;102(1):162-4
PTEN status in advanced colorectal cancer treated with cetuximab.
149. PMID: 27605871; 2016, World J Gastroenterol;22(28):6345-61
Mechanisms of resistance to anti-epidermal growth factor receptor inhibitors in metastatic colorectal cancer.

ACT Onco[®] + Report

150. PMID: 24666267; 2014, Acta Oncol;53(7):852-64
The predictive value of KRAS, NRAS, BRAF, PIK3CA and PTEN for anti-EGFR treatment in metastatic colorectal cancer: A systematic review and meta-analysis.
151. PMID: 19351834; 2009, Cancer Res;69(8):3256-61
PTEN loss contributes to erlotinib resistance in EGFR-mutant lung cancer by activation of Akt and EGFR.
152. PMID: 23133538; 2012, PLoS One;7(10):e48004
Modeling of tumor progression in NSCLC and intrinsic resistance to TKI in loss of PTEN expression.
153. PMID: 23592446; 2013, J Cell Biochem;114(6):1248-56
mTOR inhibitors radiosensitize PTEN-deficient non-small-cell lung cancer cells harboring an EGFR activating mutation by inducing autophagy.
154. PMID: 26645196; 2016, Cancer Discov;6(2):202-16
Loss of PTEN Promotes Resistance to T Cell-Mediated Immunotherapy.
155. PMID: 28228279; 2017, Immunity;46(2):197-204
Loss of PTEN Is Associated with Resistance to Anti-PD-1 Checkpoint Blockade Therapy in Metastatic Uterine Leiomyosarcoma.
156. PMID: 30150660; 2018, Nat Genet;50(9):1271-1281
Genomic correlates of response to immune checkpoint blockade in microsatellite-stable solid tumors.
157. PMID: 21468130; 2011, Nat Rev Clin Oncol;8(5):302-6
Treatment with olaparib in a patient with PTEN-deficient endometrioid endometrial cancer.
158. PMID: 20930833; 2010, Nature;467(7316):667-8
DNA repair: A protein giant in its entirety.
159. PMID: 20729858; 2010, Nat Struct Mol Biol;17(10):1263-5
The breast cancer tumor suppressor BRCA2 promotes the specific targeting of RAD51 to single-stranded DNA.
160. PMID: 20729832; 2010, Nature;467(7316):678-83
Purified human BRCA2 stimulates RAD51-mediated recombination.
161. PMID: 22305526; 2012, Am J Hum Genet;90(2):301-7
RAD51 haploinsufficiency causes congenital mirror movements in humans.
162. PMID: 18243065; 2008, DNA Repair (Amst);7(5):686-93
The consequences of Rad51 overexpression for normal and tumor cells.
163. PMID: 24811120; 2014, Oncotarget;5(10):3261-72
Rad51 supports triple negative breast cancer metastasis.
164. PMID: 26317153; 2015, Cell Cycle;14(19):3190-202
High levels of RAD51 perturb DNA replication elongation and cause unscheduled origin firing due to impaired CHK1 activation.
165. PMID: 21807066; 2011, Biochim Biophys Acta;1816(2):209-18
RAD51 as a potential biomarker and therapeutic target for pancreatic cancer.
166. PMID: 10851081; 2000, Oncogene;19(23):2791-5
DNA repair and recombination factor Rad51 is over-expressed in human pancreatic adenocarcinoma.
167. PMID: 24741789; 2014, Rev Med Chir Soc Med Nat Iasi;118(1):133-40
Rad51 overexpression and resistance to genotoxic agents. A study in the fission yeast Schizosaccharomyces pombe.
168. PMID: 18618591; 2009, Mol Carcinog;48(2):105-9
Rad51 overexpression rescues radiation resistance in BRCA2-defective cancer cells.
169. PMID: 10807537; 2000, J Hum Genet;45(3):133-7

ACT Onco[®] + Report

Identification of Rad51 alteration in patients with bilateral breast cancer.

170. PMID: 26108708; 2015, Sci Rep;5():11588
RAD51 135G>C substitution increases breast cancer risk in an ethnic-specific manner: a meta-analysis on 21,236 cases and 19,407 controls.
171. PMID: 11248061; 2001, Proc Natl Acad Sci U S A;98(6):3232-6
A single nucleotide polymorphism in the RAD51 gene modifies cancer risk in BRCA2 but not BRCA1 carriers.
172. PMID: 17999359; 2007, Am J Hum Genet;81(6):1186-200
RAD51 135G-->C modifies breast cancer risk among BRCA2 mutation carriers: results from a combined analysis of 19 studies.
173. PMID: 30353044; 2018, Br J Cancer;119(11):1401-1409
Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes.
174. PMID: 24577941; 2014, Mol Cancer Ther;13(5):1170-80
The use of Olaparib (AZD2281) potentiates SN-38 cytotoxicity in colon cancer cells by indirect inhibition of Rad51-mediated repair of DNA double-strand breaks.
175. PMID: 28759753; 2017, Biomed Pharmacother;94():165-168
Inhibition of Rad51 sensitizes breast cancer cells with wild-type PTEN to olaparib.
176. PMID: 21157483; 2011, Nat Rev Mol Cell Biol;12(1):21-35
mTOR: from growth signal integration to cancer, diabetes and ageing.
177. PMID: 12271141; 2002, Proc Natl Acad Sci U S A;99(21):13571-6
Tuberous sclerosis complex-1 and -2 gene products function together to inhibit mammalian target of rapamycin (mTOR)-mediated downstream signaling.
178. PMID: 9242607; 1997, Science;277(5327):805-8
Identification of the tuberous sclerosis gene TSC1 on chromosome 9q34.
179. PMID: 8269512; 1993, Cell;75(7):1305-15
Identification and characterization of the tuberous sclerosis gene on chromosome 16.
180. PMID: 1303246; 1992, Nat Genet;2(1):37-41
Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease.
181. PMID: 18538015; 2008, BMC Cancer;8():163
Involvement of TSC genes and differential expression of other members of the mTOR signaling pathway in oral squamous cell carcinoma.
182. PMID: 28339086; 2017, Int J Oncol;50(5):1778-1784
Identification of novel mutations in endometrial cancer patients by whole-exome sequencing.
183. PMID: 20610279; 2010, Urol Oncol;28(4):409-28
Bladder cancer or bladder cancers? Genetically distinct malignant conditions of the urothelium.
184. PMID: 17005952; 2006, N Engl J Med;355(13):1345-56
The tuberous sclerosis complex.
185. PMID: 22923433; 2012, Science;338(6104):221
Genome sequencing identifies a basis for everolimus sensitivity.
186. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
187. PMID: 27016228; 2016, Gynecol Oncol;141(1):43-8
Tumor mutational analysis of GOG248, a phase II study of temsirolimus or temsirolimus and alternating megestrol acetate and tamoxifen for advanced endometrial cancer (EC): An NRG Oncology/Gynecologic Oncology Group study.
188. PMID: 26412398; 2015, Sci Rep;5():14534

ACT Onco[®] + Report

PAK2 is an effector of TSC1/2 signaling independent of mTOR and a potential therapeutic target for Tuberous Sclerosis Complex.

189. PMID: 28968163; 2017, J Clin Oncol;35(32):3638-3646
MONARCH 3: Abemaciclib As Initial Therapy for Advanced Breast Cancer.
190. PMID: 28533223; 2017, Clin Cancer Res;23(17):5218-5224
MONARCH 1, A Phase II Study of Abemaciclib, a CDK4 and CDK6 Inhibitor, as a Single Agent, in Patients with Refractory HR+/HER2-Metastatic Breast Cancer.
191. PMID: 26156651; 2015, Lancet Oncol;16(8):897-907
Afatinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label randomised controlled phase 3 trial.
192. PMID: 23816960; 2013, J Clin Oncol;31(27):3327-34
Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations.
193. PMID: 22285168; 2012, Lancet Oncol;13(3):239-46
Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTAC): a multicentre, open-label, randomised phase 3 trial.
194. PMID: 17452677; 2007, J Clin Oncol;25(15):1960-6
Erlotinib plus gemcitabine compared with gemcitabine alone in patients with advanced pancreatic cancer: a phase III trial of the National Cancer Institute of Canada Clinical Trials Group.
195. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
196. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
197. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
198. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
199. PMID: 24263064; 2014, Br J Cancer;110(1):55-62
First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study.
200. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.
201. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
202. PMID: 25366685; 2015, J Clin Oncol;33(3):244-50
Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation.
203. PMID: 27959700; 2017, N Engl J Med;376(7):629-640
Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer.
204. PMID: 27959613; 2016, N Engl J Med;375(20):1925-1936
Palbociclib and Letrozole in Advanced Breast Cancer.
205. PMID: 26030518; 2015, N Engl J Med;373(3):209-19
Palbociclib in Hormone-Receptor-Positive Advanced Breast Cancer.

ACT Onco[®] + Report

206. PMID: 27908594; 2017, Lancet Oncol;18(1):75-87
Rucaparib in relapsed, platinum-sensitive high-grade ovarian carcinoma (ARIEL2 Part 1): an international, multicentre, open-label, phase 2 trial.
207. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsirrolimus, interferon alfa, or both for advanced renal-cell carcinoma.