



Sample Information

Patient Name: 曾守智
Gender: Male
ID No.: U120212568
History No.: 25630904
Age: 48

Ordering Doctor: DOC3069L 孫瑞璘
Ordering REQ.: 0BCMZZX
Signing in Date: 2021/02/24

Path No.: S110-98261
MP No.: F21016
Assay: Oncomine Focus Assay
Sample Type: FFPE
Block No.: S110-75021C
Percentage of tumor cells: 50%
Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	0 Therapies Available
Relevant Therapy Summary	2	6 Clinical Trials
Clinical Trials Summary	3	

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	Not detected	NTRK1	Not detected
BRAF	Not detected	NTRK2	Not detected
EGFR	Not detected	NTRK3	Not detected
ERBB2	Not detected	RET	Not detected
KRAS	Not detected	ROS1	Not detected
MET	Not detected		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	CCND1 amplification cyclin D1	None	None	6

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

Copy Number Variations

Gene	Locus	Copy Number
CCND1	chr11:69456942	7.92

Biomarker Descriptions

CCND1 (cyclin D1)

Background: The CCND1 gene encodes the cyclin D1 protein, a member of the highly conserved D-cyclin family that also includes CCND2 and CCND3^{1,2,3}. D-type cyclins are known to regulate cell cycle progression by binding to and activating cyclin dependent kinases (CDKs), specifically CDK4 and CDK6, which leads to the phosphorylation and inactivation of the retinoblastoma (RB1) protein^{1,2}. Consequently, RB1 inactivation results in E2F transcription factor activation and cellular G1/S phase transition thereby resulting in cell cycle progression, a common event observed in tumorigenesis^{1,2,4}. Aberrations in the D-type cyclins have been observed to promote tumor progression suggesting an oncogenic role for CCND1^{3,5}.

Alterations and prevalence: Recurrent somatic alterations to CCND1, including mutations, amplifications, and chromosomal translocations, are observed in many cancer types. A common mechanism of these alterations is to increase the expression and nuclear localization of the cyclin D1 protein. Recurrent somatic mutations include missense mutations at codons T286 and P287 and c-terminal truncating mutations that are enriched in about 33% of uterine cancer, and missense mutations at Y44 that are enriched in about 50% of Mantle cell lymphoma (MCL)^{6,7,8,9}. These mutations block phosphorylation-dependent nuclear export and proteolysis^{10,11,12,13}. CCND1 is recurrently amplified in many cancer types, including up to 35% of esophageal cancer, 20-30% of head and neck cancer, and 10-20% of breast, squamous lung, and bladder cancers^{6,8,14}. MCL is genetically characterized by the t(11;14) (q13;q13) translocation, a rearrangement that juxtaposes CCND1 to the immunoglobulin heavy (IgH) chain gene. This rearrangement leads to constitutive expression of cyclin D1 and plays an important role in MCL pathogenesis^{15,16}.

Potential relevance: Currently, no therapies are approved for CCND1 aberrations.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

CCND1 amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
abemaciclib	×	×	×	×	● (II)
palbociclib	×	×	×	×	● (II)
siremadlin, ribociclib	×	×	×	×	● (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Clinical Trials Summary

CCND1 amplification

NCT ID	Title	Phase
NCT02664935	National Lung Matrix Trial: Multi-drug, Genetic Marker-directed, Non-comparative, Multi-centre, Multi-arm Phase II Trial in Non-small Cell Lung Cancer	II
NCT03310879	A Phase II Study of the CDK4/6 Inhibitor Abemaciclib in Patients With Solid Tumors Harboring Genetic Alterations in Genes Encoding D-type Cyclins or Amplification of CDK4 or CDK6	II
NCT04116541	MegaMOST - A Multicenter, Open-label, Biology Driven, Phase II Study Evaluating the Activity of Anti-cancer Treatments Targeting Tumor Molecular Alterations /Characteristics in Advanced / Metastatic Tumors.	II
NCT03297606	Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR): A Phase II Basket Trial	II
NCT03155620	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice) Screening Protocol	II
NCT03526250	NCI-COG Pediatric MATCH (Molecular Analysis for Therapy Choice) - Phase 2 Subprotocol of Palbociclib in Patients With Tumors Harboring Activating Alterations in Cell Cycle Genes	II

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Malumbres et al. Cell cycle, CDKs and cancer: a changing paradigm. *Nat. Rev. Cancer*. 2009 Mar;9(3):153-66. PMID: 19238148
2. Koyama-Nasu et al. The critical role of cyclin D2 in cell cycle progression and tumorigenicity of glioblastoma stem cells. *Oncogene*. 2013 Aug 15;32(33):3840-5. PMID: 22964630
3. Ding et al. Prognostic role of cyclin D2/D3 in multiple human malignant neoplasms: A systematic review and meta-analysis. *Cancer Med*. 2019 Jun;8(6):2717-2729. PMID: 30950241
4. Bartek et al. Pathways governing G1/S transition and their response to DNA damage. *FEBS Lett*. 2001 Feb 16;490(3):117-22. PMID: 11223026
5. Shan et al. Cyclin D1 overexpression correlates with poor tumor differentiation and prognosis in gastric cancer. *Oncol Lett*. 2017 Oct;14(4):4517-4526. PMID: 28943959
6. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
7. Cancer et al. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013 May 2;497(7447):67-73. PMID: 23636398
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
9. Beà et al. Landscape of somatic mutations and clonal evolution in mantle cell lymphoma. *Proc. Natl. Acad. Sci. U.S.A.* 2013 Nov 5;110(45):18250-5. PMID: 24145436
10. Diehl et al. Glycogen synthase kinase-3beta regulates cyclin D1 proteolysis and subcellular localization. *Genes Dev*. 1998 Nov 15;12(22):3499-511. PMID: 9832503
11. Alt et al. Phosphorylation-dependent regulation of cyclin D1 nuclear export and cyclin D1-dependent cellular transformation. *Genes Dev*. 2000 Dec 15;14(24):3102-14. PMID: 11124803
12. Moreno-Bueno et al. Cyclin D1 gene (CCND1) mutations in endometrial cancer. *Oncogene*. 2003 Sep 4;22(38):6115-8. PMID: 12955092
13. Benzeno et al. Identification of mutations that disrupt phosphorylation-dependent nuclear export of cyclin D1. *Oncogene*. 2006 Oct 12;25(47):6291-303. PMID: 16732330
14. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
15. Kim et al. Nuclear cyclin D1: an oncogenic driver in human cancer. *J. Cell. Physiol*. 2009 Aug;220(2):292-6. PMID: 19415697
16. Jares et al. Genetic and molecular pathogenesis of mantle cell lymphoma: perspectives for new targeted therapeutics. *Nat. Rev. Cancer*. 2007 Oct;7(10):750-62. PMID: 17891190