

ABOUT THE TEST FoundationOne® Heme is a comprehensive genomic profiling test designed to identify genomic alterations within hundreds of cancer-related genes in hematologic malignancies and sarcomas.

PATIENT	DISEASE Unknown primary malignant neoplasm	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Mouth
	NAME Lin, Mei-Chen		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S111-07951 A (PF22046)
	DATE OF BIRTH 30 January 1981		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Female		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 23 February 2022
	MEDICAL RECORD # 43982770		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 04 April 2022

Biomarker Findings

Tumor Mutational Burden - 63 Muts/Mb
Microsatellite status - MS-Stable

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

MET amplification

ERBB3 V104L

ETV6 inversion intron 5, rearrangement intron 5

FLT4 V160L

HIST1H1D Q182*

KEAP1 E41*

TET2 V1417F

TP53 splice site 375G>T

TRAF3 Q492*

Report Highlights

- Targeted therapies with potential clinical benefit **approved in this patient's tumor type**: Dostarlimab (p. 11), Pembrolizumab (p. 11)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 18)
- Variants that may represent **clonal hematopoiesis** and may originate from non-tumor sources: **TET2 V1417F** (p. 8)

BIOMARKER FINDINGS

Tumor Mutational Burden - 63 Muts/Mb

10 Trials *see p. 18*

Microsatellite status - MS-Stable

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

Dostarlimab

Pembrolizumab

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Atezolizumab

Avelumab

Cemiplimab

Durvalumab

Nivolumab

Nivolumab +
Ipilimumab

No therapies or clinical trials. see Biomarker Findings section

GENOMIC FINDINGS	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
MET - amplification	none	Cabozantinib
		Capmatinib
		Crizotinib
		Tepotinib
6 Trials see p. 21		
ERBB3 - V104L	none	none
1 Trial see p. 20		

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

TET2 - V1417F p. 8

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

ETV6 - inversion intron 5, rearrangement intron 5.....	p. 5	KEAP1 - E41*.....	p. 7
FLT4 - V160L.....	p. 6	TET2 - V1417F.....	p. 8
HIST1H1D - Q182*.....	p. 6	TP53 - splice site 375G>T.....	p. 9
		TRAF3 - Q492*.....	p. 10

NOTE Genomic alterations detected may be associated with activity of certain FDA-approved drugs; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

BIOMARKER FINDINGS
BIOMARKER

Tumor Mutational Burden

RESULT

63 Muts/Mb

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1 therapies¹⁻⁴, and combination nivolumab and ipilimumab⁵⁻¹⁰. In multiple pan-tumor studies, higher TMB has been reported to be associated with increased ORR and OS from treatment with immune checkpoint inhibitors^{1-4,11}. Higher TMB was found to be significantly associated with improved OS upon immune checkpoint inhibitor treatment for patients with 9 types of advanced tumors¹. Analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 –20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy, compared with patients with higher TMB treated with

chemotherapy¹² or those with lower TMB treated with PD-1 or PD-L1-targeting agents². However, the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors found significant improvement in ORR for patients with TMB ≥ 10 Muts/Mb (based on this assay or others) compared to those with TMB < 10 Muts/Mb, in a large cohort that included multiple tumor types; similar findings were observed in the KEYNOTE 028 and 012 trials^{4,11}. Together, these studies suggest that patients with TMB ≥ 10 Muts/Mb may derive clinical benefit from PD-1 or PD-L1 inhibitors.

FREQUENCY & PROGNOSIS

High or elevated TMB has been most frequently reported in melanoma (17–42%)¹³⁻¹⁵, colorectal cancer (CRC; 8–25%)¹⁶⁻¹⁹, endometrial carcinoma (7–24%)²⁰⁻²², intestinal-type stomach adenocarcinoma (20%)²⁰, and non-small cell lung carcinoma (NSCLC; 8–13%)²³⁻²⁴. In patients with NSCLC, increased TMB is associated with higher tumor grade and poor prognosis²⁵, as well as with a decreased frequency of known driver mutations in EGFR, ALK, ROS1, or MET (1% of high-TMB samples each), but not BRAF (10.3%) or KRAS (9.4%)²³. Although some studies have reported a lack of association between smoking and increased TMB in NSCLC²⁵⁻²⁶, several other large

studies did find a strong link²⁷⁻³⁰. In CRC, elevated TMB is associated with a higher frequency of BRAF V600E driver mutations¹⁸⁻¹⁹ and with microsatellite instability (MSI)¹⁹, which in turn has been reported to correlate with better prognosis³¹⁻³⁸. Although increased TMB is associated with increased tumor grade in endometrioid endometrial carcinoma^{22,39-41} and bladder cancer⁴², it is also linked with improved prognosis in patients with these tumor types²².

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁴³⁻⁴⁴ and cigarette smoke in lung cancer⁴⁵⁻⁴⁶, treatment with temozolomide-based chemotherapy in glioma⁴⁷⁻⁴⁸, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{18,22,49-51}, and microsatellite instability (MSI)^{18,22,51}. This sample harbors a TMB level that may be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors in multiple solid tumor types^{2-4,11}.

BIOMARKER

Microsatellite status

RESULT

MS-Stable

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical evidence, MSS tumors are significantly less likely than MSI-H tumors to respond to anti-PD-1 immune checkpoint inhibitors⁵²⁻⁵⁴, including approved therapies nivolumab and pembrolizumab⁵⁵. In a retrospective analysis of 361 patients with solid tumors treated with pembrolizumab, 3% were MSI-H and experienced a significantly higher ORR compared with non-MSI-H cases (70% vs. 12%, $p=0.001$)⁵⁶.

FREQUENCY & PROGNOSIS

MSI-high (MSI-H) has been observed at high frequency in endometrial cancers (14–33%)⁵⁷⁻⁶⁴, colorectal cancers (CRCs; 10–15%)^{18,37,54,65-66}, and gastric cancers (12–35%)⁶⁷⁻⁷⁰ and at lower frequencies in many other tumor types, including esophageal⁷¹, small bowel⁷²⁻⁷⁶, hepatobiliary⁷⁷⁻⁸³, prostate⁸⁴⁻⁸⁶, and urinary tract carcinomas⁸⁷⁻⁸⁹. In one study, MSI-H status was associated with a positive prognostic effect in patients with gastric cancer treated with surgery alone and a negative predictive effect in patients treated with chemotherapy⁹⁰. Data regarding the role of MSI-H on prognosis and survival in endometrial cancer are conflicting^{57,60-61,63,91-93}. However, studies specifically analyzing early stage endometrial cancer have reported a correlation between MSI-H and decreased survival^{59,62,64,92}, thereby suggesting that MSI-H predicts for poor prognosis in this subset of endometrial tumors.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor⁶⁶. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2^{66,94-95}. This sample is microsatellite-stable (MSS), equivalent to the clinical definition of an MSS tumor: one with mutations in none of the tested microsatellite markers^{65,96-97}. MSS status indicates MMR proficiency and typically correlates with intact expression of all MMR family proteins^{65-66,95,97}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE
MET
ALTERATION
 amplification

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of extensive clinical evidence, MET amplification or activating mutations may predict sensitivity to MET-targeting therapies such as kinase inhibitors crizotinib, capmatinib, tepotinib, and cabozantinib. Crizotinib has benefited patients with MET-amplified non-small cell lung cancer (NSCLC) of varied histologies⁹⁸⁻¹⁰¹, gastroesophageal cancer¹⁰², glioblastoma¹⁰³, and carcinoma of unknown primary¹⁰⁴. Capmatinib has demonstrated clinical efficacy for patients with MET-amplified NSCLC both as a monotherapy¹⁰⁵⁻¹⁰⁶ and in combination with an EGFR-TKI for patients with concurrent activating EGFR mutations¹⁰⁷⁻¹⁰⁹. Tepotinib has demonstrated efficacy for patients with MET-amplified hepatocellular carcinoma¹¹⁰ and NSCLC¹¹¹ as a monotherapy, as well as in combination with gefitinib for patients with MET-amplified and EGFR-mutated NSCLC¹¹²⁻¹¹⁴. Savolitinib elicited responses in patients with MET-amplified papillary renal cell carcinoma¹¹⁵ and gastric cancer either alone or in combination with docetaxel¹¹⁶⁻¹¹⁷. AMG 337 elicited an ORR of

50% (5/10), including 1 CR, for patients with MET-amplified gastric, esophageal, or gastroesophageal junction cancer¹¹⁸. Patients with MET-amplified NSCLC¹¹⁹ or MET-amplified gastric cancer¹²⁰ treated with the MET-targeting antibody onartuzumab (MetMab) achieved clinical responses. In addition, high MET expression has been suggested to predict patient response to therapies such as the monoclonal HGF-targeting antibody rilotumumab, as well as the combination of ramucirumab and the monoclonal MET-targeting antibody emibetuzumab¹²¹. A first-in-human Phase 1 trial of telisotuzumab vedotin (teliso-V), a MET antibody-drug conjugate, reported activity in a subset of patients with MET-positive NSCLC, with an ORR of 19% (3/16) and a DCR of 56% (9/16); no responses were observed in any other patients¹²². A subsequent Phase 2 trial of teliso-V in patients with MET-positive NSCLC reported a 35% (13/37) ORR in patients with non-squamous, EGFR-wildtype tumors, which met the prespecified criteria for transition to the next stage; lower ORRs were observed in patients with squamous (14%; 3/21) or non-squamous EGFR-mutated (13%; 4/30) tumors¹²³.

FREQUENCY & PROGNOSIS

In the TCGA datasets, amplification of MET has been found in several tumor types, with the highest incidences in ovarian serous cystadenocarcinoma (5.5%), esophageal carcinoma (3.4%), stomach adenocarcinoma (3.7%), and lung

adenocarcinoma (3.0%) datasets; lower incidences were observed in various other tumor types (cBioPortal, Sep 2021)¹²⁴⁻¹²⁵. Overexpression of MET mRNA and protein has been observed in a number of cancers¹²⁶⁻¹²⁸. MET amplification has been associated with poor prognosis in gastroesophageal adenocarcinoma, gastric and esophageal cancer^{102,129-134}. Increased MET expression has been associated with poor prognosis in cutaneous malignant melanoma¹³⁵, gallbladder adenocarcinoma¹³⁶, lung large cell neuroendocrine carcinoma¹³⁷, and breast cancer¹³⁸⁻¹⁴⁰. The prognostic value of MET amplification or expression in non-small cell lung cancer (NSCLC)¹⁴¹⁻¹⁴⁸, endometrial carcinoma¹⁴⁹⁻¹⁵² or colon cancer¹⁵³⁻¹⁵⁵ have yielded conflicting results, although concurrent MET amplification and EGFR mutation have been correlated with reduced disease-free survival in NSCLC¹⁵⁶.

FINDING SUMMARY

MET encodes a receptor tyrosine kinase, also known as c-MET or hepatocyte growth factor receptor (HGFR), that is activated by the ligand HGF; MET activation results in signaling mediated partly by the RAS-RAF-MAPK and PI3K pathways to promote proliferation¹⁵⁷⁻¹⁵⁸. MET has been reported to be amplified in cancer¹²⁵, with amplification positively correlating with protein expression in some cancer types^{126-128,142,159} and associating with therapeutic response to MET inhibitors in a variety of cancer types^{98-100,102-104,160-161}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE
ERBB3
ALTERATION

V104L

TRANSCRIPT ID

NM_001982

CODING SEQUENCE EFFECT

310G>T

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

ERBB3 cooperates with other ERBB family members, in particular ERBB2, for efficient signaling¹⁶²⁻¹⁶⁵. Therefore, ERBB3 amplification or activating mutation may predict sensitivity to therapies targeting ERBB2, including antibodies such as trastuzumab, pertuzumab, and ado-trastuzumab emtansine (T-DM1), and dual EGFR/HER2 TKIs such as lapatinib and afatinib. Preclinical studies support the sensitivity of cells with ERBB3 activating mutations to various anti-

ERBB2 agents^{164,166-167}. In a Phase 2 study of afatinib in platinum-refractory urothelial cancer, 2 patients with activating ERBB3 mutations but no EGFR or HER2 activating alterations experienced clinical benefit and PFS > 6 months¹⁶⁸. Other studies in solid tumors have reported mixed efficacy for afatinib for patients with uncharacterized ERBB3 mutations¹⁶⁹⁻¹⁷⁰. Case studies report clinical benefit from lapatinib combined with either capecitabine¹⁶⁹ or trastuzumab^{169,171} for patients with breast cancer harboring activating ERBB3 mutations. However, Phase 2 trials have suggested limited efficacy of other ERBB2-targeting TKIs against ERBB3 mutations, with no objective response to neratinib in any of 16 patients with ERBB3-mutated solid tumors¹⁷² or dacomitinib in either of 2 patients with ERBB3-mutated cutaneous squamous cell carcinoma¹⁷³.

FREQUENCY & PROGNOSIS

ERBB3 mutations have been reported with highest incidence in cancers of the meninges (31%), urinary tract (11%), stomach (7.5%), skin

(7.2%), endometrium (6.8%), prostate (5.3%), large intestine (5.2%), biliary tract (4.7%), and cervix (4%) and at lower frequencies in other tumor types (COSMIC, Apr 2022)¹⁷⁴. High expression of ERBB3 mRNA or ERBB3 protein has been found in a number of tumor types, including those of the breast, lung, ovary, stomach, colon, and bladder¹⁷⁵⁻¹⁸⁵. Elevated ERBB3 expression has been associated with increased disease severity or negative prognostic indicators in glioblastoma¹⁸⁶, gastric cancer¹⁸⁷⁻¹⁸⁸, non-small cell lung cancer¹⁷⁸, and colorectal cancer¹⁸⁹. ERBB3 amplification in breast cancer has been correlated with reduced disease-free survival, although some studies have found no association with prognosis¹⁹⁰⁻¹⁹⁷. The data associating ERBB3 expression and invasion and metastasis in urothelial carcinoma were similarly conflicting¹⁹⁸⁻²⁰³.

FINDING SUMMARY

ERBB3 (also known as HER3) encodes a member of the epidermal growth factor receptor (EGFR) family²⁰⁴. ERBB3 mutations such as observed here have been shown to be activating^{164,167,205-206}.

GENE
ETV6
ALTERATION

inversion intron 5, rearrangement intron 5

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no approved therapies that address ETV6 mutation or inactivation.

FREQUENCY & PROGNOSIS

ETV6 mutation has been observed in up to 7% of cutaneous squamous cell carcinomas²⁰⁷, 6% of colorectal adenocarcinomas²⁰⁸, 5.5% of bladder cancers²⁰⁹, 4% each of melanomas and endometrial carcinomas, and in lower frequencies in other tumor types cBioPortal, Apr 2022¹²⁴⁻¹²⁵. High ETV6 expression associated with poor prognosis in one study²¹⁰, and low ETV6 expression associated with improved OS and disease-free survival in patients with NSCLC²¹¹.

FINDING SUMMARY

ETV6 encodes an ETS family transcription factor

required for hematopoiesis. The portion of chromosome 12 that encodes ETV6 is frequently involved in translocation events, and at least 30 fusion partners have been identified to date²¹²⁻²¹³. ETV6 rearrangements have been implicated in various hematopoietic malignancies²¹⁴ and have been reported to be pathogenic by a number of different mechanisms including activation of a partner oncogene, ETV6 inactivation, or aberrant transcription factor activity²¹³. Alterations such as seen here may disrupt ETV6 function or expression²¹⁵⁻²¹⁹.

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE

FLT4

ALTERATION

V160L

TRANSCRIPT ID

NM_002020

CODING SEQUENCE EFFECT

478G>T

sorafenib; responses to axitinib have been observed in patients with FLT4-amplified differentiated thyroid carcinoma²²⁰, and responses to sorafenib have been reported in patients with FLT4-amplified angiosarcoma²²¹. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

FREQUENCY & PROGNOSIS

In the TCGA datasets, FLT4 mutation has been reported in 11.6% of cutaneous melanomas, 7.5% of endometrial carcinomas, 5.6% of colorectal adenocarcinomas, 5.5% of stomach adenocarcinomas, and at lower incidence across a range of other tumor types (cBioPortal, Apr 2022)¹²⁴⁻¹²⁵. Published data investigating the

prognostic implications of FLT4 alterations in solid tumors are generally limited (PubMed, Apr 2022).

FINDING SUMMARY

FLT4 encodes the protein fms-related tyrosine kinase 4, also known as VEGFR-3 (vascular endothelial growth factor receptor 3). Targeting of VEGF receptors has been a major therapeutic strategy in cancer as growth of new blood and lymph vessels is a critical determinant of tumor growth and metastasis²²²⁻²²³. Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Based on clinical studies, FLT4 amplification may be associated with sensitivity to kinase inhibitors with activity against VEGFR3 such as axitinib and

GENE

HIST1H1D

ALTERATION

Q182*

TRANSCRIPT ID

NM_005320

CODING SEQUENCE EFFECT

544C>T

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are currently no therapies available to directly address genomic alterations in HIST1D.

FREQUENCY & PROGNOSIS

Recurrent missense mutations in the genes that encode H1 isoforms B-E, have been reported at high frequencies in B cell lymphomas, including 30-40% of diffuse large B cell lymphomas (DLBCLs), 30% of follicular lymphomas and 50% of Hodgkin lymphomas, and lower frequencies across other tumor types²²⁴⁻²²⁸. Published data investigating the prognostic implications of

HIST1H1D alterations in cancer are limited (PubMed, Apr 2022).

FINDING SUMMARY

HIST1H1D encodes histone H1.3 (H1D), a linker histone that interacts with DNA between nucleosomes and functions in the regulation of gene expression and compaction of chromatin into higher order structures²²⁹⁻²³⁰. Missense mutations in linker histones affecting the globular and C-terminal domains have been identified in cancer and often correlate with loss of function²³⁰, whereas other alterations are mostly uncharacterized.

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE

KEAP1

ALTERATION

E41*

TRANSCRIPT ID

NM_012289

CODING SEQUENCE EFFECT

121G>T

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

A study of patients with localized non-small cell lung cancer (NSCLC) identified pathogenic KEAP1 and NFE2L2 mutations as predictors of local recurrence following radiotherapy but not surgery; limited preclinical data also showed that treatment with a glutaminase inhibitor sensitized KEAP1-mutated NSCLC cells to radiation²³¹. In other preclinical studies, treatment with AKT inhibitors sensitized lung cancer cells harboring KEAP1 or NFE2L2 mutations to both chemotherapy and radiation therapy²³²⁻²³³. Mixed clinical data have been reported for the association between KEAP1 mutations and the response to immunotherapy. A pan-cancer study of immunotherapy showed that patients with KEAP1 mutations had shorter OS (10 vs. 20 months) than those without²³⁴. However, another study across solid tumors showed that KEAP1 mutations were associated with higher tumor mutational burden (TMB) and PD-L1 expression, as well as improved

survival outcomes with immunotherapy compared with other treatments (20.0 vs. 11.5 months)²³⁵. For patients with non-small cell lung cancer (NSCLC), a study of PD-L1 inhibitors showed that patients with concurrent mutations of STK11 and KEAP1 (n=39) experienced significantly shorter PFS (1.6 vs. 2.5 months, HR=1.5) and OS (4 vs. 11 months, HR=1.9) compared with patients with STK11- and KEAP1-wildtype tumors (n=210) despite significantly higher TMB in the group harboring STK11 and KEAP1 mutations (median 9.4 vs. 6.1 Muts/Mb)²³⁶. Retrospective analyses of patients with NSCLC who received immunotherapy reported reduced OS (p=0.040) for patients harboring KEAP1- or NFE2L2-mutated tumors²³⁷ or STK11- or KEAP1-mutated tumors (p < 0.001)²³⁸ compared with those without. Studies of immune checkpoint inhibitors for patients with lung adenocarcinoma showed that coexisting mutations between KEAP1, PBRM1, SMARCA4, STK11, and KRAS were associated with worse OS²³⁹. An exploratory analysis of a subset of patients with PD-L1-positive NSCLC treated in the first-line setting with pembrolizumab showed similar ORR, PFS, and OS when comparing patients with STK11 or KEAP1 mutations and those without²⁴⁰. In addition, preclinical data suggest that KEAP1 inactivation increases tumor demand for glutamine and increases tumor sensitivity to glutaminase inhibitors like telaglenastat²⁴¹⁻²⁴³. Limited clinical data suggest that KEAP1 mutations may predict improved clinical benefit from combinations of glutaminase inhibitors and

anti-PD-1 inhibitors²⁴⁴; a Phase 1/2 study of the glutaminase inhibitor telaglenastat (CB-839) plus nivolumab to treat advanced NSCLC reported better clinical benefit rates and median PFS for patients with KEAP1 mutations (75% [3/4] vs. 15% [2/13], 6.4 vs. 3.7 months), KRAS mutations (38% [3/8] vs. 20% [2/10], 4.5 vs. 3.7 months), or KEAP1 and KRAS concurrent mutations (100% [2/2] vs. 13% [1/8], 7.2 vs. 3.7 months) compared with patients without these mutations²⁴⁴. The KEAP1 mutation has also been identified as a potential biomarker for sensitivity to combined AKT and TXNRD1 inhibition in lung cancer²⁴⁵.

FREQUENCY & PROGNOSIS

Somatic mutation of KEAP1 occurs in a range of solid tumors, including gastric, hepatocellular, colorectal, and lung cancers²⁴⁶. KEAP1 mutations are rare in hematological malignancies, occurring in fewer than 1% of samples analyzed (COSMIC, 2022)¹⁷⁴. In a retrospective analysis of the pan-solid MSKCC dataset, KEAP1 mutation correlated with reduced OS (13.28 vs. 26.53 months)²³⁵.

FINDING SUMMARY

KEAP1 encodes a substrate adaptor protein that regulates the cellular response to oxidative stress by providing substrate specificity for a CUL3-dependent ubiquitin ligase²⁴⁷. KEAP1 exerts anti-tumor effects through negative regulation of NRF2, a transcription factor encoded by NFE2L2²⁴⁸⁻²⁵⁰; KEAP1 inactivation promotes cancer progression through NRF2-mediated chemoresistance and cell growth²⁴⁹⁻²⁵⁰.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE
TET2
ALTERATION

V1417F

TRANSCRIPT ID

NM_001127208

CODING SEQUENCE EFFECT

4249G>T

low frequencies in solid tumors and are more prevalent in hematological malignancies (cBioPortal, Jan 2022)¹²⁴⁻¹²⁵. Published data investigating the prognostic implications of TET2 alterations in solid tumors are limited (PubMed, Jan 2022).

FINDING SUMMARY

TET2 encodes a tumor suppressor involved in reversing DNA methylation marks, a process critical for proper gene regulation²⁵¹⁻²⁵². Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁵³⁻²⁵⁸. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²⁵³⁻²⁵⁴. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease²⁵⁹. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{257,260-261}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in TET2 in solid tumors.

FREQUENCY & PROGNOSIS

TET2 alterations have been reported at relatively

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE
TP53
ALTERATION

splice site 375G>T

TRANSCRIPT ID

NM_000546

CODING SEQUENCE EFFECT

375G>T

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib²⁶²⁻²⁶⁵, or p53 gene therapy and immunotherapeutics such as SGT-53²⁶⁶⁻²⁷⁰ and ALT-801²⁷¹. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype²⁷². A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer²⁷³. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer²⁷⁴. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone²⁷⁵. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel¹⁷⁶. A Phase 1 trial of neoadjuvant adavosertib in combination

with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations²⁷⁶. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87 months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring²⁷⁷. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage²⁷⁰. ATR inhibitor treatment of chronic lymphocytic leukemia (CLL) cells with biallelic inactivation of TP53 suppressed cell viability, promoted DNA damage, and attenuated xenograft growth in preclinical studies²⁷⁸⁻²⁷⁹; however, ATR inhibitors as monotherapy had little effect on these parameters in solid tumor models in other preclinical studies²⁸⁰⁻²⁸¹. Therefore, it is unclear whether TP53 inactivation predicts sensitivity to ATR inhibition.

FREQUENCY & PROGNOSIS

Pan-cancer analysis of the TCGA datasets across 12 cancer types identified TP53 as the most frequently mutated gene, with 42% of more than 3,000 tumors harboring a TP53 mutation; in this study TP53 mutation occurred most frequently in ovarian serous carcinoma (95%), lung squamous cell carcinoma (SCC) (79%), head and neck SCC (70%), colorectal adenocarcinoma (59%), lung adenocarcinoma (52%), and bladder urothelial carcinoma (50%)²⁸². In one study of 55 patients with lung adenocarcinoma, TP53 alterations correlated with immunogenic features including PD-L1 expression, tumor mutation burden and neoantigen presentation; likely as a consequence of this association TP53 mutations correlated with improved clinical outcomes to PD-1 inhibitors pembrolizumab and nivolumab in this study²⁸³. TP53 mutation has not been consistently demonstrated to be a significant independent

prognostic marker in the context of CRC²⁸⁴.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers²⁸⁵. Alterations such as seen here may disrupt TP53 function or expression²⁸⁶⁻²⁹⁰.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers²⁹¹⁻²⁹³, including sarcomas²⁹⁴⁻²⁹⁵. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000²⁹⁶ to 1:20,000²⁹⁵. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30²⁹⁷. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion²⁵³⁻²⁵⁸. CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy²⁵³⁻²⁵⁴. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease²⁵⁹. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{257,260-261}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

ORDERED TEST # ORD-1336448-02

GENOMIC FINDINGS
GENE

TRAF3

ALTERATION

Q492*

TRANSCRIPT ID

NM_003300

CODING SEQUENCE EFFECT

1474C>T

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no approved therapies that directly target TRAF3 inactivation or the activation of the NF-kB pathway, although these are under

development. The Bruton's tyrosine kinase inhibitor ibrutinib has been reported to inhibit NF-kB pathway activation in chronic lymphocytic leukemia samples and myeloma cells²⁹⁸⁻³⁰⁰. However, preclinical data in mantle cell lymphoma cells suggest that inactivation of TRAF3 or the related TRAF2 may correlate with insensitivity to ibrutinib and to the PKC inhibitor sotrastaurin³⁰¹. This insensitivity was attributed to activation of alternative NF-kB signaling, which could be inhibited by targeting the kinase NIK (MAP3K14) in preclinical assays³⁰¹.

FREQUENCY & PROGNOSIS

TRAF3 mutations have been reported with the highest incidences in the uterine corpus endometrial carcinoma (3.4%), skin cutaneous

melanoma (2.7%), cervical squamous cell carcinoma (2.4%), stomach adenocarcinoma (2.3%), and colorectal adenocarcinoma (2.0%) TCGA datasets (cBioPortal, Mar 2022)¹²⁴⁻¹²⁵. Published data investigating the prognostic implications of TRAF3 alterations in solid tumors are limited (PubMed, Mar 2022).

FINDING SUMMARY

TRAF3, a member of the TRAF family of adaptor proteins, has E3 ubiquitin ligase activity and is a critical determinant of B-cell survival³⁰²⁻³⁰³.

TRAF3 loss leads to the activation of the NF-kB pathway and is thought to function as a tumor suppressor in a variety of B-cell lineage neoplasms³⁰⁴⁻³⁰⁸. Alterations such as seen here may disrupt TRAF3 function or expression.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN PATIENT'S TUMOR TYPE

Dostarlimab

Assay findings association
Tumor Mutational Burden
 63 Muts/Mb

AREAS OF THERAPEUTIC USE

Dostarlimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor response. It is FDA approved to treat patients with mismatch repair deficient recurrent or advanced endometrial cancer or solid tumors. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors

targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Dostarlimab has been studied primarily in recurrent and advanced mismatch repair-deficient (dMMR) endometrial and non-endometrial cancers³¹⁰⁻³¹². In the Phase 1 GARNET trial, single-agent dostarlimab elicited an ORR of 39% (41/106) and an immune-related ORR of 46% (50/110) for patients with non-endometrial dMMR solid tumors^{310,313}.

Pembrolizumab

Assay findings association
Tumor Mutational Burden
 63 Muts/Mb

AREAS OF THERAPEUTIC USE

Pembrolizumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved for patients with tumor mutational burden (TMB)-high (≥ 10 Muts/Mb), microsatellite instability-high (MSI-H), or mismatch repair-deficient (dMMR) solid tumors; as monotherapy for PD-L1-positive non-small cell lung cancer (NSCLC), head and neck squamous cell cancer (HNSCC), cervical cancer, or esophageal cancer; and in combination with chemotherapy for PD-L1-positive triple-negative breast cancer (TNBC) or cervical cancer. It is also approved in various treatment settings as monotherapy for patients with melanoma, HNSCC, urothelial carcinoma, hepatocellular carcinoma, Merkel cell carcinoma, cutaneous squamous cell carcinoma, endometrial carcinoma that is MSI-H or dMMR, classical Hodgkin lymphoma, or primary mediastinal large B-cell lymphoma; and in combination with chemotherapy or targeted therapy for NSCLC, HNSCC, esophageal or gastroesophageal junction cancer, renal cell carcinoma, TNBC, or endometrial carcinoma that is not MSI-H or dMMR. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

In the Phase 2 KEYNOTE 158 multi-solid tumor trial, treatment with the PD-1 inhibitor pembrolizumab led to

improved ORR for patients with TMB of 10 Muts/Mb or higher compared those with TMB < 10 Muts/Mb (28.3% [34/120] vs. 6.5% [41/635])¹¹. In the KEYNOTE 028/012 pan-solid tumor trials, a similar improvement in ORR was reported for patients with > 103 non-synonymous mutations/exome (\sim equivalency > 8 Muts/Mb as measured by this assay) compared to those with < 103 non-synonymous mutations/exome (30.6% [11/36] vs. 6.5% [5/77])⁴. Pembrolizumab has achieved significant clinical benefit for patients with PD-L1-expressing solid tumors including gastric carcinoma (ORR=22.0%)³¹⁴, esophageal carcinoma (ORR=22.0%)³¹⁵, endometrial carcinoma (ORR=13.0%)^{39,316}, lung carcinoma (median OS=14.9 months)³¹⁷, urothelial tract carcinoma (ORR=24.0%)³¹⁸, head and neck carcinoma (ORR=12.0-26.0%)³¹⁹⁻³²², breast carcinoma (ORR=16.0-25.0%)³²³⁻³²⁴, cervical squamous cell carcinoma (ORR=12.5%)³²⁵, thyroid carcinoma (ORR=9.0%)³²⁶, and pleural mesothelioma (ORR=24.0%)³²⁷. In the KEYNOTE-051 Phase 1/2 study of pembrolizumab for the treatment of pediatric patients with advanced cancer, PRs were reported in 8 patients including 2 adrenocortical carcinomas, 2 mesotheliomas, 1 malignant ganglioglioma, 1 epithelioid sarcoma, 1 lymphoepithelial carcinoma, and 1 malignant rhabdoid³²⁸. Clinical studies have reported responses to pembrolizumab in combination with other immunotherapies such as ipilimumab in recurrent advanced non-small cell lung carcinoma (NSCLC)(ORR=24.0%)³²⁹⁻³³⁰, and nivolumab in glioma (2/12 PRs, 4/12 SDs)³³¹. Clinical benefit has also been achieved with pembrolizumab in combination with chemotherapy for advanced non-squamous NSCLC³³², IDO-1 inhibitor epacadostat for solid tumors³³³, HDAC inhibitor entinostat for breast and endometrial cancer¹²³, bevacizumab for glioblastoma³³⁴, and lenvatinib for patients with advanced endometrial carcinoma that is not MSI-H or dMMR³³⁵.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Atezolizumab

Assay findings association

Tumor Mutational Burden

63 Muts/Mb

AREAS OF THERAPEUTIC USE

Atezolizumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC) and urothelial carcinoma, depending on treatment setting. Atezolizumab is also approved in combination with other therapies to treat patients with non-squamous NSCLC lacking EGFR or ALK alterations, small cell lung cancer, hepatocellular carcinoma, and BRAF V600-positive melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

In the prospective Phase 2a MyPathway basket study evaluating atezolizumab for patients with TMB-High solid tumors, patients with TMB ≥ 16 Muts/Mb achieved improved ORR (38% [16/42] vs. 2.1% [1/48]), DCR (62% [26/42] vs. 23% [11/48]), mPFS (5.7 vs. 1.8 months, HR 0.34), and mOS (19.8 vs. 11.4, HR 0.53) as compared to those with TMB ≥ 10 and < 16 Muts/Mb³³⁶. In a retrospective analysis of patients with 17 solid tumor types (comprised of 47% NSCLC, 40% urothelial carcinoma, and 13% encompassing 15 other solid tumors),

TMB of 16 Muts/Mb or greater was reported to be associated with an improved ORR to atezolizumab compared to chemotherapy (30% vs. 14%)¹². Atezolizumab has been studied primarily for the treatment of non-small cell lung cancer (NSCLC)³³⁷⁻³⁴² and urothelial carcinoma³⁴³⁻³⁴⁶. A study of atezolizumab as monotherapy for patients with advanced solid tumors reported a median PFS of 18 weeks and an ORR of 21%, including confirmed responses in 25.6% (11/43) of melanomas, 12.5% (7/56) of renal cell carcinomas (RCC) and 16.7% (1/6) of colorectal cancers (CRCs)³⁴². As single-agent therapy in genomically unselected young patients (< 30 years old) with relapsed or refractory cancers, atezolizumab elicited an ORR of 1.5% (1/67) for patients with solid tumors, with similar safety and pharmacokinetics as seen in adults³⁴⁷. A Phase 1a study of atezolizumab reported an ORR of 14.5% (9/62), a median PFS of 5.6 months, and a median OS of 28.9 months for patients with clear cell RCC³⁴⁸. A Phase 1b study evaluated atezolizumab combined with nab-paclitaxel for patients with previously treated metastatic triple-negative breast cancer (mTNBC) and reported confirmed objective responses for 41.7% (10/24) of patients; no dose-limiting toxicities were observed³⁴⁹. A Phase 1b study that evaluated atezolizumab in combination with the MEK inhibitor cobimetinib for advanced solid tumors reported an ORR of 8.3% (7/84) in patients with CRC, 40.9% (9/22) in patients with melanoma, 17.9% (5/28) in patients with NSCLC, and 18.8% (3/16) in patients with other tumors (ovarian cancer, clear-cell sarcoma, and RCC); there was no association between BRAF or KRAS mutation status and response rate in any disease setting, and no dose-limiting toxicities were encountered³⁵⁰⁻³⁵¹.

Avelumab

Assay findings association

Tumor Mutational Burden

63 Muts/Mb

AREAS OF THERAPEUTIC USE

Avelumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 in order to enhance antitumor immune responses. It is FDA approved to treat patients 12 years and older with Merkel cell carcinoma, or for urothelial carcinoma in various treatment settings. The combination of avelumab and axitinib is FDA approved for patients with renal cell carcinoma (RCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

The JAVELIN Phase 1b study has demonstrated clinical benefit from single-agent avelumab in a variety of solid tumor types, including non-small cell lung carcinoma (NSCLC)³⁵², gastric carcinoma and gastroesophageal junction (GEJ) adenocarcinoma³⁵³, urothelial carcinoma³⁵⁴, mesothelioma³⁵⁵, ovarian carcinoma³⁵⁶, and breast cancer³⁵⁷, and from avelumab combined with axitinib in renal cell carcinoma³⁵⁸. Emerging clinical data show a positive trend toward the association of tumor cell PD-L1 expression and improved ORR, PFS, or OS in NSCLC in the first-line setting and in ovarian and breast cancer^{352,356-357}. Limited clinical data indicate activity of avelumab in adrenocortical carcinoma, metastatic castration-resistant prostate cancer, and thymic cancer³⁵⁹⁻³⁶¹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 | CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Cabozantinib

Assay findings association
MET
 amplification

AREAS OF THERAPEUTIC USE

Cabozantinib inhibits multiple tyrosine kinases, including MET, RET, VEGFRs, and ROS1. It is FDA approved as monotherapy to treat patients with renal cell carcinoma (RCC), hepatocellular carcinoma (HCC), medullary thyroid cancer (MTC), and differentiated thyroid cancer (DTC). It is also approved in combination with nivolumab to treat RCC. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Sensitivity of MET alterations to cabozantinib is suggested by clinical responses in patients with non-small cell lung cancer (NSCLC) harboring MET mutations associated with MET exon 14 skipping, with or without concurrent MET amplification³⁶²⁻³⁶³, as well as by extensive preclinical data³⁶⁴⁻³⁷⁰.

SUPPORTING DATA

A randomized Phase 2 discontinuation study of cabozantinib in 9 solid tumor types reported ORRs of 0% to 22% and response durations of 3.3 to 11.2 months across cohorts with ORRs of 10% or greater observed for patients with ovarian cancer (22% [15/69, 1 CR]), metastatic breast cancer (14% [6/44]), and non-small cell lung cancer (NSCLC) (10% [6/60])³⁷¹⁻³⁷². A Phase 1 study of cabozantinib for advanced solid tumors reported a 17%

(4/23) ORR in the dose escalation cohort and an ORR of 20% (4/20) and a DCR of 100% (20/20) in the expansion cohort for Japanese patients with NSCLC³⁷³. In the context of studies for specific solid tumors, the randomized Phase 3 EXAM study for patients with advanced medullary thyroid cancer reported an association of cabozantinib with improved PFS compared with placebo (11.2 vs. 4.0 months, HR=0.28) and a higher ORR (28% vs. 0%), with PFS improvement observed regardless of RET mutation status³⁷⁴. The randomized Phase 3 CELESTIAL study for patients with advanced hepatocellular carcinoma (HCC) previously treated with sorafenib reported significantly longer OS (10.2 vs. 8.0 months, HR=0.76) and PFS (5.2 vs. 1.9 months, HR=0.44) as well as an increased ORR (3.8% vs. 0.4%) with cabozantinib when compared to placebo³⁷⁵. The Phase 2 CABOSUN trial of first line cabozantinib versus sunitinib for patients with intermediate- or poor-risk advanced clear cell renal cell carcinoma demonstrated significantly improved median PFS (8.2 vs. 5.6 months, HR=0.66), prolonged median OS (30.3 vs. 21.8 months), and higher ORR (33% [26/79] vs. 12% [9/78]) with cabozantinib compared with sunitinib³⁷⁶. The Phase 2 CABONE study of cabozantinib reported ORRs of 26% (10/39) for patients with advanced Ewing sarcoma and 12% (5/42) for patients with advanced osteosarcoma³⁷⁷.

Capmatinib

Assay findings association
MET
 amplification

AREAS OF THERAPEUTIC USE

Capmatinib is a selective MET tyrosine kinase inhibitor that is FDA approved to treat patients with non-small cell lung cancer harboring MET exon 14 skipping-associated alterations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data in non-small cell lung cancer^{105,111-114,378}, hepatocellular carcinoma¹¹⁰, renal cell carcinoma¹¹⁵, and gastric cancer¹¹⁶, MET amplification may predict sensitivity to selective MET inhibitors.

SUPPORTING DATA

Capmatinib has been investigated primarily for the treatment of NSCLC, demonstrating efficacy as monotherapy for patients with MET amplification^{106,379-380} or MET exon 14 skipping alterations³⁸⁰⁻³⁸¹ as well as in combination with EGFR inhibitors for patients with MET amplification¹⁰⁷⁻¹⁰⁹. Multiple Phase 1 and 2 clinical studies have reported limited efficacy for capmatinib monotherapy in non-NSCLC indications, with no responses observed for patients with MET-amplified glioblastoma (n=10)³⁸², MET-overexpressing gastric cancer (n=9)³⁸³, or other advanced solid tumors with MET amplification or overexpression (n=11)³⁸³⁻³⁸⁴.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Cemiplimab

Assay findings association
Tumor Mutational Burden
 63 Muts/Mb

AREAS OF THERAPEUTIC USE

Cemiplimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved to treat patients with NSCLC with high PD-L1 expression (TPS $\geq 50\%$), cutaneous squamous cell carcinoma (CSCC), or basal cell carcinoma (BCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been

observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Cemiplimab has been studied primarily in advanced cutaneous squamous cell carcinoma (CSCC), where it elicited a combined ORR of 48% (41/85) in Phase 1 and 2 studies³⁸⁵. A Phase 2 trial of cemiplimab in patients with basal cell carcinoma (BCC) reported ORRs of 31% (5 CRs and 21 PRs) in patients with locally advanced BCC and 21% (6 PRs) in patients with metastatic BCC³⁸⁶⁻³⁸⁷. The Phase 3 EMPOWER-Lung 1 trial for advanced non-small cell lung cancer (NSCLC) with PD-L1 expression $\geq 50\%$ reported that cemiplimab is associated with improved PFS (8.2 vs. 5.7 months), OS (not reached vs. 14.2 months), and ORR (37% vs. 21%) compared with chemotherapy³⁸⁸.

Crizotinib

Assay findings association
MET
 amplification

AREAS OF THERAPEUTIC USE

Crizotinib is an inhibitor of the kinases MET, ALK, ROS1, and RON. It is FDA approved to treat patients with ALK rearrangement- or ROS1 rearrangement-positive non-small cell lung cancer (NSCLC), and to treat pediatric and young adult patients with ALK rearrangement-positive anaplastic large cell lymphoma (ALCL). Please see the drug label for full prescribing information.

GENE ASSOCIATION

Sensitivity of MET alterations to crizotinib is suggested by extensive clinical data in patients with MET-amplified cancers, including non-small cell lung cancer (NSCLC)^{98-100,389-390}, gastric cancer¹⁶⁰, gastroesophageal cancer¹⁰², glioblastoma¹⁰³, and carcinoma of unknown primary¹⁰⁴, as well as in patients with MET-mutated cancers, including NSCLC^{362,391-395}, renal cell carcinoma (RCC)³⁹⁶, and histiocytic sarcoma³⁹¹. Crizotinib has also benefited patients with NSCLC or histiocytic sarcoma tumors harboring various alterations associated with MET exon 14 skipping^{362,391,393-395,397}.

SUPPORTING DATA

Crizotinib has demonstrated efficacy in patients with

NSCLC and ALK rearrangements³⁹⁸⁻⁴⁰², ROS1 rearrangements⁴⁰³⁻⁴⁰⁷, an NTRK1 fusion⁴⁰⁸, or MET activation^{98-100,362,389-390,392-395,409-415}. Crizotinib has also benefited patients with MET-mutated renal cell carcinoma⁴¹⁶ and patients with MET-amplified gastroesophageal cancer, glioblastoma, and carcinoma of unknown primary¹⁰²⁻¹⁰⁴. While a Phase 1b study evaluating crizotinib for the treatment of patients with ALK-positive malignancies, reported ORR of 52.9% (9/17) and 66.7% (6/9) in patients with lymphoma and inflammatory myofibroblastic tumors (IMT), respectively, an ORR of 11.8% (2/17) was reported for patients with other types of tumors⁴¹⁷. Whereas median PFS and median OS were not reached for patients with lymphoma or IMT, median PFS was 1.3 months and median OS was 8.3 months for patients with other tumor types, and the median duration of treatment was ~1 month relative to 1-3 years for patients with lymphoma or IMT⁴¹⁷. A Phase 1 clinical trial of crizotinib in pediatric solid tumors reported objective responses in 14/79 patients, including nine CRs and five PRs; response was enriched in patients with activating alterations in ALK⁴¹⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Durvalumab

Assay findings association
Tumor Mutational Burden
63 Muts/Mb

AREAS OF THERAPEUTIC USE

Durvalumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC) and small cell lung cancer (SCLC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Single-agent durvalumab has demonstrated efficacy in non-small cell lung cancer⁴¹⁹⁻⁴²⁰, and head and neck squamous cell carcinoma⁴²¹⁻⁴²². In patients with advanced solid tumors, durvalumab monotherapy has elicited disease control rates (DCRs) of 36.8–46.2% (7/19 to 12/26) in Phase 1/2 studies⁴²³⁻⁴²⁴. Durvalumab is also under

investigation in combination with other agents in Phase 1/2 trials. In advanced melanoma, durvalumab in combination with trametinib and dabrafenib elicited ORRs and DCRs of 76.2% (16/21) and 100% (21/21) in patients with BRAF-mutant tumors, and durvalumab with trametinib elicited ORRs and DCRs of 21.4% (3/14) and 64.3% (9/14) in patients whose tumors were BRAF wild-type⁴²⁵. Durvalumab in combination with the PARP inhibitor olaparib has shown activity in patients with metastatic castration-resistant prostate cancer and progression on enzalutamide and/or abiraterone⁴²⁶ and in patients with BRCA-wild-type breast or gynecological cancer⁴²⁷. Durvalumab in combination with the anti-CTLA4 antibody tremelimumab, but not durvalumab as a single-agent, has shown activity in patients with previously treated advanced germ cell tumors⁴²⁸. Responses have also been reported for patients with solid tumors treated with durvalumab in combination with the anti-PD-1 antibody MEDI0680⁴²⁹, the CXCR2 antagonist AZD5069⁴³⁰, or the ATR inhibitor AZD6738⁴³¹. In patients with treatment-refractory solid tumors, concurrent durvalumab and radiotherapy achieved an ORR of 60% (6/10) for in-field evaluable lesions, including 2 CRs and 4 PRs⁴³².

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Nivolumab

Assay findings association
Tumor Mutational Burden
 63 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response. It is FDA approved as a monotherapy in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), urothelial carcinoma, colorectal cancer (CRC), classical Hodgkin lymphoma (cHL), gastric cancer, gastroesophageal junction cancer, and esophageal adenocarcinoma or squamous cell carcinoma (ESCC). It is also approved in combination with cabozantinib to treat RCC, and in combination with relatlimab to treat melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{2-4,12,309}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors²⁻³.

SUPPORTING DATA

Nivolumab monotherapy has been reported to elicit clinical benefit for patients with multiple types of solid tumors, including melanoma (27-31% ORR)⁴³³⁻⁴³⁷, non-

small cell lung cancer (NSCLC; 17-20% ORR and 9-10 months median OS [mOS] for unselected patients)⁴³⁸⁻⁴⁴⁴, urothelial carcinoma (20-26% ORR for unselected patients)⁴⁴⁵⁻⁴⁴⁶, renal cell carcinoma (RCC; 26% ORR)⁴⁴⁷⁻⁴⁵², microsatellite instability-high (MSI-High) colorectal cancer (CRC; 58% ORR)⁴⁵³⁻⁴⁵⁵, head and neck squamous cell carcinoma (11-17% ORR)⁴⁵⁶⁻⁴⁵⁸, ovarian cancer (6-15% ORR)⁴⁵⁹⁻⁴⁶¹, small cell lung cancer (SCLC; 10-12% ORR)^{355,462-463}, gastroesophageal carcinoma (12-18% ORR)⁴⁶⁴⁻⁴⁶⁶, and cancer of unknown primary (CUP; 22% ORR for unselected, previously treated patients)⁴⁶⁷, as well as with Hodgkin lymphoma (66-87% ORR)⁴⁶⁸⁻⁴⁷⁰. Combination treatment with nivolumab plus the CTLA-4 inhibitor ipilimumab has achieved further efficacy in melanoma (up to 61% ORR; mOS >60 months for the combination vs. 37 months for nivolumab monotherapy)^{433,471-473}, NSCLC (mOS of 17 months)⁴⁷⁴, MSI-High CRC (64% ORR)⁴⁷⁵, RCC (42% ORR)⁴⁷⁶⁻⁴⁷⁷, SCLC (19-25% ORR)^{355,463}, urothelial carcinoma (38% ORR for unselected patients; 58% ORR for patients with $\geq 1\%$ tumor PD-L1 expression)⁴⁴⁵, and other solid tumors. Combinations of nivolumab with various targeted therapies, such as pazopanib (1 PR for a patient with epithelioid sarcoma)⁴⁷⁸, sunitinib (9% ORR for unselected patients with sarcoma)⁴⁷⁹, cabozantinib (29% ORR for patients with immunotherapy-refractory urothelial carcinoma)⁴⁸⁰, or vemurafenib (1 durable PR for a patient with BRAF-V600E-mutated and PD-L1-positive anaplastic thyroid cancer)⁴⁸¹, have also shown evidence of efficacy and continue to undergo clinical investigation.

Nivolumab + Ipilimumab

Assay findings association
Tumor Mutational Burden
 63 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response, and ipilimumab is a cytotoxic T-lymphocyte antigen 4 (CTLA-4)-blocking antibody. The combination is FDA approved in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), hepatocellular carcinoma (HCC), and pleural mesothelioma. Furthermore, nivolumab is approved in combination with ipilimumab to treat patients with mismatch repair-deficient (dMMR) or microsatellite instability-high (MSI-H) colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{5-6,482}, a TMB score of ≥ 10 Muts/Mb (as measured by this assay) may predict sensitivity to combination nivolumab and ipilimumab treatment.

SUPPORTING DATA

Combination treatment with nivolumab plus the CTLA-4 inhibitor ipilimumab has achieved efficacy in melanoma (up to 61% ORR; mOS >60 months for the combination vs. 37 months for nivolumab monotherapy)^{433,471-473}, NSCLC (17 months mOS)⁴⁷⁴, MSI-High CRC (64% ORR)⁴⁷⁵, RCC (42% ORR)⁴⁷⁶⁻⁴⁷⁷, SCLC (19-25% ORR)^{355,463}, urothelial carcinoma (38% ORR in unselected patients; 58% ORR in patients with $\geq 1\%$ tumor PD-L1 expression)⁴⁴⁵, and other solid tumors.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

THERAPIES WITH CLINICAL BENEFIT
IN OTHER TUMOR TYPE

Tepotinib

Assay findings association
MET
 amplification

AREAS OF THERAPEUTIC USE

Tepotinib is a selective MET tyrosine kinase inhibitor that is FDA approved to treat patients with non-small cell lung cancer harboring MET exon 14 skipping alterations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data in non-small cell lung cancer^{105,111-114,378}, hepatocellular carcinoma¹¹⁰, renal cell carcinoma¹¹⁵, and gastric cancer¹¹⁶, MET amplification may predict sensitivity to selective MET inhibitors.

SUPPORTING DATA

Tepotinib has primarily been investigated in non-small cell lung cancer and has demonstrated efficacy as a single agent for patients with MET amplification¹¹¹ and MET

exon 14-skipping alterations⁴⁸³⁻⁴⁸⁴. Tepotinib has also been shown to be efficacious in combination with gefitinib for patients with concurrent EGFR mutation and MET amplification or MET overexpression in Phase 2 studies¹¹³⁻¹¹⁴. In advanced hepatocellular carcinoma, Phase 2 studies of tepotinib reported improved ORR and PFS for both treatment-naïve and previously treated patients with MET protein overexpression^{110,485-487}. In a Phase 1 study of advanced solid tumors, tepotinib monotherapy yielded an ORR of 1.3% and a DCR of 24%, with 2 confirmed PRs observed for patients with esophageal or lung cancer and 2 unconfirmed PRs for patients with colorectal or nasopharyngeal cancer⁴⁸⁸. In another Phase 1 study of solid tumors, tepotinib yielded a DCR of 17% (2/12), with 2 SD of ≥12 weeks observed in a patient with gastric cancer and another with urachal cancer⁴⁸⁹.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and

should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that may require

medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. Or visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

BIOMARKER

Tumor Mutational Burden

RESULT

63 Muts/Mb

RATIONALE

Increased tumor mutational burden may predict response to anti-PD-1 (alone or in combination

with anti-CTLA-4) or anti-PD-L1 immune checkpoint inhibitors.

NCT04237649
PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS

ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Sunto Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS

TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Xi'an (China), Tianjin (China), Beijing (China), Chengdu City (China), Changchun (China)

NCT04521621
PHASE 1/2

A Study of V937 in Combination With Pembrolizumab (MK-3475) in Participants With Advanced/Metastatic Solid Tumors (V937-013)

TARGETS

PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Tainan (Taiwan), Kaohsiung (Taiwan), Kashiwa (Japan), Afula (Israel), Jerusalem (Israel), Tel Aviv (Israel), Warszawa (Poland), Oslo (Norway)

NCT04261439
PHASE 1

A Phase I/Ib Study of NIZ985 Alone and in Combination With Spartalizumab

TARGETS

PD-1

LOCATIONS: Taipei (Taiwan), Chuo ku (Japan), Essen (Germany), Napoli (Italy), Leuven (Belgium), Barcelona (Spain), California, Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

CLINICAL TRIALS
NCT03799003
PHASE 1

A Study of ASP1951 in Subjects With Advanced Solid Tumors

TARGETS
 PD-1, TNFRSF18

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Daegu (Korea, Republic of), Chungcheongbukdo (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Washington, California, Utah

NCT03861793
PHASE 1/2

A Dose Escalation and Cohort Expansion Study of Subcutaneously-Administered Cytokine (ALKS 4230) as a Single Agent and in Combination With Anti-PD-1 Antibody (Pembrolizumab) in Subjects With Select Advanced or Metastatic Solid Tumors (ARTISTRY-2)

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Suwon (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Edmonton (Canada), Badalona (Spain), Rotterdam (Netherlands), Valencia (Spain), Madrid (Spain)

NCT04047862
PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS
 PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Guangdong (China), Changsha (China), Wuhan (China), Jinju-si (Korea, Republic of)

NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
 PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Amsterdam (Netherlands), Ravenna (Italy), Meldola (Italy)

NCT03396445
PHASE 1

Safety and Pharmacokinetics Study of MK-5890 as Monotherapy and in Combination With Pembrolizumab (MK-3475) in Adults With Advanced Solid Tumors (MK-5890-001)

TARGETS
 PD-1, CD27

LOCATIONS: Taipei (Taiwan), Seoul (Korea, Republic of), Jerusalem (Israel), Ramat Gan (Israel), Be'er Sheva (Israel), Amsterdam (Netherlands), Rotterdam (Netherlands), Barcelona (Spain), Madrid (Spain), Pozuelo de Alarcon (Spain)

NCT04892498
PHASE 2

Hypofractionated Radiotherapy Combined With PD-1 Inhibitor Sequential GM-CSF and IL-2 for the Treatment of Advanced Refractory Solid Tumors (PRaG2.O)

TARGETS
 PD-1

LOCATIONS: Hangzhou (China), Suzhou (China), Wuxi (China), Hefei (China), Xuzhou (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

CLINICAL TRIALS
GENE
ERBB3
ALTERATION

V104L

RATIONALE

Clinical and preclinical data support sensitivity of ERBB3 activating mutations to HER2-targeting TKIs, including afatinib and lapatinib. ERBB3

amplification or activating mutations may confer sensitivity to therapies targeting ERBB3.

NCT03810872
PHASE 2

An Explorative Study of Afatinib in the Treatment of Advanced Cancer Carrying an EGFR, a HER2 or a HER3 Mutation

TARGETS

EGFR, ERBB4, ERBB2

LOCATIONS: Liège (Belgium), Brussels (Belgium), Gent (Belgium)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

CLINICAL TRIALS
GENE
MET
RATIONALE

Activating MET alterations may confer sensitivity to MET inhibitors.

ALTERATION

amplification

NCT03175224
PHASE 1/2

CBT-101 Study for Advanced Solid Tumors and c-Met Dysregulation

TARGETS
 MET

LOCATIONS: Taipei City (Taiwan), Taipei (Taiwan), New Taipei City (Taiwan), Taoyuan City (Taiwan), Tainan (Taiwan), Singapore (Singapore), Nedlands (Australia), Saransk (Russian Federation), North Adelaide (Australia), Bedford Park (Australia)

NCT04647838
PHASE 2

Tepotinib in Solid Tumors Harboring MET Alterations

TARGETS
 MET

LOCATIONS: Cheonan (Korea, Republic of), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
 VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT04116541
PHASE 2

A Study Evaluating the Activity of Anti-cancer Treatments Targeting Tumor Molecular Alterations/ Characteristics in Advanced / Metastatic Tumors.

TARGETS
 CDK6, CDK4, MDM2, MET, ROS1, RET, VEGFRs

LOCATIONS: Nice (France), Lyon (France), Marseille (France), Toulouse (France), Bordeaux (France)

NCT04887194
PHASE 1

PK Study to Assess Drug-drug Interaction and QTc Between Sitravatinib and a Cocktail of Substrates

TARGETS
 AXL, KIT, DDR2, VEGFRs, PDGFRA, TRKA, MET, FLT3, RET, TRKB, PD-1

LOCATIONS: Texas, Virginia

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

 Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

CLINICAL TRIALS
NCT04693468
PHASE 1

Talazoparib and Palbociclib, Axitinib, or Crizotinib for the Treatment of Advanced or Metastatic Solid Tumors, TalaCom Trial

TARGETS
PARP, CDK4, CDK6, VEGFRs, ALK,
ROS1, AXL, TRKA, MET, TRKC

LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future. Please note that some VUS rearrangements between targeted genes and unknown fusion partners or intergenic regions detected by RNA sequencing may not be reported.

ABL1 S1007*	ALK E201Q	ARID2 H48Y	ASXL1 I980N
BCORL1 D77N and T1111M	BRCA1 G1232L	BRD4 E657Q	BRIP1 R814C
CD22 S105N	CIC A531T	CIITA A462S and A648E	EPHA5 E881D and T176A
EPHB1 I236M	ERBB4 A345V, D1131Y and P1132H	FAM123B G327C	FBXO11 S34N
FBXO31 R19P	FLT4 A634D and Y435C	FRS2 P240S	GPR124 R369Q
HGF G31*	HIST1H1C E3K	HIST1H3B A92P	IL7R K104N and L123V
JUN R221Q	LRP1B S1198Y and S913R	MAPK1 D210H	MKI67 I2578V
MYST3 A1442S and P1766L	NOTCH1 P1884Q	NSD1 R1074L	NTRK2 A187E
P2RY8 P341H	PCLO E3034D and P995A	PDGFRA A7P	PLCG2 E1208K
POT1 T425A	PPP2R1A V259F	PRKDC D3411H	SMARCA4 D1183Y
SMC3 E998*	SMO R703P	SOCS1 L150Q	SOX2 A30S
SPEN S2185F	SUZ12 S53L	TAF1 R123T	TNFAIP3 E451Q
TSHR G235R	WDR90 R690Q	ZNF217 P823L	

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX
Genes Assayed in FoundationOne®Heme

FoundationOne Heme is designed to include genes known to be somatically altered in human hematologic malignancies and sarcomas that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay utilizes DNA sequencing to interrogate 406 genes as well as selected introns of 31 genes involved in rearrangements, in addition to RNA sequencing of 265 genes. The assay will be updated periodically to reflect new knowledge about cancer biology.

HEMATOLOGICAL MALIGNANCY DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACTB	AKT1	AKT2	AKT3	ALK	AMER1 (FAM123B or WTX)	APC
APH1A	AR	ARAF	ARFRP1	ARHGAP26 (GRAF)		ARID1A	ASMTL
ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	B2M
BAP1	BARD1	BCL10	BCL11B	BCL2	BCL2L2	BCL6	BCOR
BCORL1	BIRC3	BLM	BRAF	BRCA1	BRCA2	BRD4	BRIP1
BTG2	BTK	BTLA	C11orf30 (EMSY)	CAD	CALR*	CARD11	CBFB
CCND1	CCND2	CCND3	CCNE1	CCT6B	CD22	CD274 (PD-L1)	CD36
CD70	CD79A	CD79B	CDC73	CDH1	CDK12	CDK4	CDK6
CDKN1B	CDKN2A	CDKN2B	CDKN2C	CEBPA	CHD2	CHEK1	CHEK2
CIITA	CKS1B	CPS1	CREBBP	CRKL	CRLF2	CSF1R	CSF3R
CTNNA1	CTNNB1	CUX1	CXCR4	DAXX	DDR2	DDX3X	DNM2
DOT1L	DTX1	DUSP2	DUSP9	EBF1	ECT2L	EED	EGFR
EP300	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4
ESR1	ETS1	ETV6	EXOSC6	EZH2	FAF1	FAM46C	FANCA
FANCD2	FANCE	FANCF	FANCG	FANCL	FAS (TNFRSF6)	FBXO11	FBXO31
FGF10	FGF14	FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1
FGFR3	FGFR4	FHIT	FLCN	FLT1	FLT3	FLT4	FLYWCH1
FOXO1	FOXO3	FOXP1	FRS2	GADD45B	GATA1	GATA2	GATA3
GNA11	GNA12	GNA13	GNAQ	GNAS	GPR124	GRIN2A	GSK3B
HDAC1	HDAC4	HDAC7	HGF	HIST1H1C	HIST1H1D	HIST1H1E	HIST1H2AC
HIST1H2AL	HIST1H2AM	HIST1H2BC	HIST1H2BJ	HIST1H2BK	HIST1H2BO	HIST1H3B	HNF1A
HSP90AA1	ICK	ID3	IDH1	IDH2	IGF1R	IKBKE	IKZF1
IKZF3	IL7R	INHBA	INPP4B	INPP5D (SHIP)	IRF1	IRF4	IRF8
JAK1	JAK2	JAK3	JARID2	JUN	KAT6A (MYST3)	KDM2B	KDM4C
KDM5C	KDM6A	KDR	KEAP1	KIT	KLHL6	KMT2A (MLL)	KMT2C (MLL3)
KRAS	LEF1	LRP1B	LRRK2	MAF	MAFB	MAGED1	MALT1
MAP2K2	MAP2K4	MAP3K1	MAP3K14	MAP3K6	MAP3K7	MAPK1	MCL1
MDM4	MED12	MEF2B	MEF2C	MEN1	MET	MIB1	MITF
MLH1	MPL	MRE11A	MSH2	MSH3	MSH6	MTOR	MUTYH
MYCL (MYCL1)	MYCN	MYD88	MYO18A	NCOR2	NCSTN	NF1	NF2
NFKBIA	NKX2-1	NOD1	NOTCH1	NOTCH2	NPM1	NRAS	NTSC2
NTRK2	NTRK3	NUP93	NUP98	P2RY8	PAG1	PAK3	PALB2
PAX5	PBRM1	PC	PCBP1	PCLO	PDCD1	PDCD11	PDCD1LG2 (PD-L2)
PDGFRB	PDK1	PHF6	PIK3CA	PIK3CG	PIK3R1	PIK3R2	PIM1
POT1	PPP2R1A	PRDM1	PRKAR1A	PRKDC	PRSS8	PTCH1	PTEN
PTPN2	PTPN6 (SHP-1)	PTPRO	RAD21	RAD50	RAD51	RAF1	RARA
RB1	RELN	RET	RHOA	RICTOR	RNF43	ROS1	RPTOR
SIPR2	SDHA	SDHB	SDHC	SDHD	SERP2	SETBP1	SETD2
SGK1	SMAD2	SMAD4	SMARCA1	SMARCA4	SMARCB1	SMC1A	SMC3
SOC3	SOC3	SOC3	SOX10	SOX2	SPEN	SPOP	SRC
STAG2	STAT3	STAT4	STAT5A	STAT5B	STAT6	STK11	SUFU
TAF1	TBL1XR1	TCF3 (E2A)	TCL1A (TCL1)	TET2	TGFB2	TLL2	TMEM30A
TMSB4XP8 (TMSL3)	TNFAIP3	TNFAIP3	TNFRSF11A	TNFRSF14	TNFRSF17	TOP1	TP53
TRAF2	TRAF3	TRAF5	TSC1	TSC2	TSHR	TUSC3	TYK2
U2AF2	VHL	WDR90	WHSC1 (MMSET or NSD2)		WISP3	WT1	XBP1
YY1AP1	ZMYM3	ZNF217	ZNF24 (ZSCAN3)	ZNF703	ZRSR2		XPO1

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX
Genes Assayed in FoundationOne®Heme

*Note: the assay was updated on 11/8/2016 to include the detection of alterations in CALR

HEMATOLOGICAL MALIGNANCY DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCL6	BCR	BRAF	CCND1	CRLF2	EGFR	EPOR
ETV1	ETV4	ETV5	ETV6	EWSR1	FGFR2	IGH	IGK	IGL
JAK1	JAK2	KMT2A (MLL)	MYC	NTRK1	PDGFRA	PDGFRB	RAF1	RARA
RET	ROS1	TMPRSS2	TRG					

HEMATOLOGICAL MALIGNANCY RNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS*

ABL1	ABL1	ABL2	ACSL6	AFF1	AFF4	ALK	ARHGAP26 (GRAF)	
ARHGEF12	ARID1A	ARNT	ASXL1	ATF1	ATG5	AT1C	BCL10	BCL11A
BCL11B	BCL2	BCL3	BCL6	BCL7A	BCL9	BCOR	BCR	BIRC3
BRAF	BTG1	CAMTA1	CARS	CBFA2T3	CBFB	CBL	CCND1	CCND2
CCND3	CD274 (PD-L1)	CDK6	CDX2	CHIC2	CHN1	CIC	CIITA	CLP1
CLTC	CLTCL1	CNTRL (CEP110)	COL1A1	CREB3L1	CREB3L2	CREBBP	CRLF2	CSF1
CTNNB1	DDIT3	DDX10	DDX6	DEK	DUSP22	EGFR	EIF4A2	ELF4
ELL	ELN	EML4	EP300	EPOR	EPS15	ERBB2	ERG	ETS1
ETV1	ETV4	ETV5	ETV6	EWSR1	FCGR2B	FCRL4	FEV	FGFR1
FGFR1OP	FGFR2	FGFR3	FLI1	FNBP1	FOXO1	FOXO3	FOXO4	FOXP1
FSTL3	FUS	GAS7	GLI1	GMPS	GPHN	HERPUD1	HEY1	HIP1
HIST1H4I	HLF	HMGA1	HMGA2	HOXA11	HOXA13	HOXA3	HOXA9	HOXC11
HOXC13	HOXD11	HOXD13	HSP90AA1	HSP90AB1	IGH	IGK	IGL	IKZF1
IL21R	IL3	IRF4	ITK	JAK1	JAK2	JAK3	JAZF1	KAT6A (MYST3)
KDSR	KIF5B	KMT2A (MLL)	LASP1	LCP1	LMO1	LMO2	LPP	LYL1
MAF	MAFB	MALT1	MDS2	MECOM	MKL1	MLF1	MLLT1 (ENL)	MLLT10 (AF10)
MLLT3	MLLT4 (AF6)	MLLT6	MN1	MNX1	MSI2	MSN	MUC1	MYB
MYC	MYH11	MYH9	NACA	NBEAP1 (BCL8)	NCOA2	NDRG1	NF1	NF2
NFKB2	NIN	NOTCH1	NPM1	NR4A3	NSD1	NTRK1	NTRK2	NTRK3
NUMA1	NUP214	NUP98	NUTM2A	OMD	P2RY8	PAFAH1B2	PAX3	PAX5
PAX7	PBX1	PCM1	PCSK7	PDCD1LG2 (PD-L2)	PDE4DIP	PDGFB	PDGFRA	PDGFRB
PER1	PHF1	PICALM	PIM1	PLAG1	PML	POU2AF1	PPP1CB	PRDM1
PRDM16	PRRX1	PSIP1	PTCH1	PTK7	RABEP1	RAF1	RALGDS	RAP1GDS1
RARA	RBM15	RET	RHOH	RNF213	ROS1	RPL22	RPN1	RUNX1
RUNX1T1 (ETO)	RUNX2	SEC31A	SEPT5	SEPT6	SEPT9	SET	SH3GL1	SLC1A2
SNX29 (RUNDC2A)	SRSF3	SS18	SSX1	SSX2	SSX4	STAT6	STL	SYK
TAF15	TAL1	TAL2	TBL1XR1	TCF3 (E2A)	TCL1A (TCL1)	TEC	TET1	TFE3
TFG	TFPT	TFRC	TLX1	TLX3	TMPRSS2	TNFRSF11A	TOP1	TP63
TPM3	TPM4	TRIM24	TRIP11	TTL	TYK2	USP6	WHSC1 (MMSET or NSD2)	
WHSC1L1	YPEL5	ZBTB16	ZMYM2	ZNF384	ZNF521			

*Note: some VUS rearrangements between targeted genes and unknown fusion partners or intergenic regions detected by RNA sequencing may not be reported.

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite (MS) status

Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX
Performance Specifications

The median exon coverage for this sample is 761x

ACCURACY

Sensitivity: Base Substitutions	At $\geq 5\%$ Minor Allele Frequency	>99.0%
Sensitivity: Insertions/Deletions (1-40bp)	At $\geq 10\%$ Minor Allele Frequency	98.0%
Sensitivity: Focal Copy Number Alterations (Homozygous Deletions or Amplifications)	At ≥ 8 copies	>95.0%
Sensitivity: Microsatellite Instability-High (MSI-H) status	Positive Predictive Agreement (PPA)	100.0% (87.54%-100.00%)*
Sensitivity: Microsatellite Stable (MSS) status	Positive Predictive Agreement (PPA)	89.66% (81.50%, 94.46%)*
Sensitivity: Known Gene Fusions	>95.0%	
Specificity: Base Substitutions, Insertions/Deletions, and Focal Copy Number Alterations	Positive Predictive Value (PPV)	>99.0%
Specificity: Known Gene Fusions	Positive Predictive Value (PPV)	>95.0%
Specificity: Microsatellite Instability-High (MSI-H) status	Negative Predictive Agreement (NPA)	97.44% (91.12%-99.29%)*
Specificity: Microsatellite Stable (MSS) status	Negative Predictive Agreement (NPA)	94.44% (86.57%, 97.82%)*
Accuracy: Tumor Mutation Burden	At $\geq 20\%$ tumor nuclei	>90.0%
Reproducibility (average concordance between replicates)	97.0% inter-batch precision 97.0% intra-batch precision 95.0% microsatellite status precision 96.0% tumor mutation burden precision	

*95% Confidence Interval

Assay specifications were determined for typical median exon coverage of approximately 500X. For additional information regarding the validation of FoundationOne®Heme, please refer to the article He, J. et al. Integrated genomic DNA/RNA profiling of hematologic malignancies in the clinical setting, Blood (2016 Jun. 16).

In the fractional-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the FoundationOne Heme assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by analytical concordance to a PCR comparator

assay using a pan-tumor sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.

Tumor Mutational Burden (TMB) is determined by measuring the number of somatic mutations in sequenced genes on the FoundationOne Heme test and extrapolating to the genome as a whole. TMB is assayed for all FoundationOne Heme samples and is reported as the number of mutations per megabase (Muts/Mb). Tumor Mutational Burden is reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine Tumor Mutational Burden.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX

About FoundationOne®Heme

ABOUT FOUNDATIONONE HEME

FoundationOne®Heme is a comprehensive genomic profiling test for hematologic malignancies and sarcomas. The test is designed to provide physicians with clinically actionable information to help with diagnostic sub-classification, prognosis assessment, and targeted therapeutic selection. Test results provide information about clinically significant alterations, potential targeted therapies, available clinical trials, and quantitative markers that may support immunotherapy clinical trial enrollment. FoundationOne Heme is analytically validated to detect all classes of genomic alterations in more than 400 cancer-related genes. In addition to DNA sequencing, FoundationOne Heme employs RNA sequencing across 265 genes to capture a broad range of gene fusions, common drivers of hematologic malignancies and sarcomas.

FoundationOne Heme was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Heme has not been cleared or approved by the United States Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. FoundationOne Heme may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high-complexity clinical testing.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne Heme identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

(Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that FoundationOne Heme data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne Heme for identifying a copy number amplification is five (5) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that FoundationOne Heme data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that FoundationOne Heme analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2022. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source

or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Heme.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. These include: subclonal alterations in heterogeneous samples, low sample quality or with homozygous losses of <3 exons; and deletions and insertions >40bp, or in repetitive/high homology sequences. FoundationOne Heme is performed using DNA and RNA derived from tumor, and as such germline events may not be reported.

The following targets typically have low coverage resulting in a reduction in sensitivity: SDHD exon 4, TNFRSF11A exon1, and TP53 exon 1.

FoundationOne Heme fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplastraat 3, 2440 Geel, Belgium.

© 2022 Foundation Medicine, Inc. All rights reserved.

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX

About FoundationOne®Heme

CE
REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*),

MPL, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

MR Suite Version 6.1.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX
References

1. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
2. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
3. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
4. Cristescu R, et al. Science (2018) PMID: 30309915
5. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
6. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
7. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
8. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
9. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
10. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
11. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
12. Legrand et al., 2018; ASCO Abstract 12000
13. Johnson DB, et al. Cancer Immunol Res (2016) PMID: 27671167
14. Snyder A, et al. N. Engl. J. Med. (2014) PMID: 25409260
15. Van Allen EM, et al. Science (2015) PMID: 26359337
16. George et al., 2016; ASCO Abstract 3587
17. Nagahashi et al., 2016; ASCO Abstract e15103
18. Nature (2012) PMID: 22810696
19. Stadler ZK, et al. J. Clin. Oncol. (2016) PMID: 27022117
20. Frampton et al., 2016; ASCO Abstract 11558
21. Santin et al., 2016; ASCO Abstract 414
22. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
23. Spigel et al., 2016; ASCO Abstract 9017
24. Jiang et al., 2016; ASCO Abstract e23128
25. Xiao D, et al. Oncotarget (2016) PMID: 27009843
26. Shim HS, et al. J Thorac Oncol (2015) PMID: 26200269
27. Govindan R, et al. Cell (2012) PMID: 22980976
28. Ding L, et al. Nature (2008) PMID: 18948947
29. Imielinski M, et al. Cell (2012) PMID: 22980975
30. Kim Y, et al. J. Clin. Oncol. (2014) PMID: 24323028
31. Samowitz WS, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11535541
32. Elsaleh H, et al. Clin Colorectal Cancer (2001) PMID: 12445368
33. Brueckl WM, et al. Anticancer Res. (2012) PMID: 22820457
34. Guidoboni M, et al. Am. J. Pathol. (2001) PMID: 11438476
35. Gryfe R, et al. N. Engl. J. Med. (2000) PMID: 10631274
36. Sinicrope FA, et al. Gastroenterology (2006) PMID: 16952542
37. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
38. Laghi L, et al. Dig Dis (2012) PMID: 22722556
39. Mehnert JM, et al. J. Clin. Invest. (2016) PMID: 27159395
40. Hussein YR, et al. Mod. Pathol. (2015) PMID: 25394778
41. Church DN, et al. Hum. Mol. Genet. (2013) PMID: 23528559
42. Cazier JB, et al. Nat Commun (2014) PMID: 24777035
43. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
44. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
45. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
46. Rizvi NA, et al. Science (2015) PMID: 25765070
47. Johnson BE, et al. Science (2014) PMID: 24336570
48. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
49. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
50. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
51. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
52. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
53. Kroemer G, et al. Oncoimmunology (2015) PMID: 26140250
54. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
55. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
56. Ayers et al., 2016; ASCO-SITC Abstract P60
57. Zigelboim I, et al. J. Clin. Oncol. (2007) PMID: 17513808
58. Hampel H, et al. Cancer Res. (2006) PMID: 16885385
59. Stelloo E, et al. Clin. Cancer Res. (2016) PMID: 27006490
60. Kanopienė D, et al. Medicina (Kaunas) (2014) PMID: 25458958
61. Black D, et al. J. Clin. Oncol. (2006) PMID: 16549821
62. Nout RA, et al. Gynecol. Oncol. (2012) PMID: 22609107
63. Steinbakk A, et al. Cell Oncol (Dordr) (2011) PMID: 21547578
64. Bilbao C, et al. Int. J. Radiat. Oncol. Biol. Phys. (2010) PMID: 20005452
65. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
66. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
67. Hiyama T, et al. J. Gastroenterol. Hepatol. (2004) PMID: 15209621
68. Wu MS, et al. Cancer Res. (1998) PMID: 9537253
69. dos Santos NR, et al. Gastroenterology (1996) PMID: 8536886
70. Fang WL, et al. Biomed Res Int (2013) PMID: 23555086
71. Farris AB, et al. Am. J. Surg. Pathol. (2011) PMID: 21422910
72. Agaram NP, et al. Am. J. Clin. Pathol. (2010) PMID: 20395525
73. Ruemmele P, et al. Am. J. Surg. Pathol. (2009) PMID: 19252434
74. Planck M, et al. Cancer (2003) PMID: 12627520
75. Hibi K, et al. Jpn. J. Cancer Res. (1995) PMID: 7775257
76. Muneyuki T, et al. Dig. Dis. Sci. (2000) PMID: 11117578
77. Zhang SH, et al. World J. Gastroenterol. (2005) PMID: 15918185
78. Chiappini F, et al. Carcinogenesis (2004) PMID: 14656944
79. Suto T, et al. J Surg Oncol (2001) PMID: 11223838
80. Momoi H, et al. J. Hepatol. (2001) PMID: 11580146
81. Liengswangwong U, et al. Int. J. Cancer (2003) PMID: 14506736
82. Moy AP, et al. Virchows Arch. (2015) PMID: 25680569
83. Yoshida T, et al. J. Gastroenterol. (2000) PMID: 11063221
84. Pritchard CC, et al. Nat Commun (2014) PMID: 25255306
85. Azzouzi AR, et al. BJU Int. (2007) PMID: 17233803
86. Burger M, et al. J. Mol. Med. (2006) PMID: 16924473
87. Bai S, et al. Am. J. Clin. Pathol. (2013) PMID: 23690119
88. Giedl J, et al. Am. J. Clin. Pathol. (2014) PMID: 25319978
89. Yamamoto Y, et al. Clin. Cancer Res. (2006) PMID: 16675567
90. Smyth et al., 2015; ASCO Gastrointestinal Cancers Symposium Abstract 62
91. Bilbao-Sieyro C, et al. Oncotarget (2014) PMID: 25026289
92. Mackay HJ, et al. Eur. J. Cancer (2010) PMID: 20304627
93. Arabi H, et al. Gynecol. Oncol. (2009) PMID: 19275958
94. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
95. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
96. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
97. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
98. Ou SH, et al. J Thorac Oncol (2011) PMID: 21623265
99. Schwab R, et al. Lung Cancer (2014) PMID: 24192513
100. Le X, et al. Clin Lung Cancer (2015) PMID: 25922291
101. Schrock AB, et al. J Thorac Oncol (2017) PMID: 28315738
102. Lennerz JK, et al. J. Clin. Oncol. (2011) PMID: 22042947
103. Chi AS, et al. J. Clin. Oncol. (2012) PMID: 22162573
104. Palma NA, et al. Case Rep Oncol (2014) PMID: 25232318
105. Schuler et al., 2016; ASCO Abstract 9067
106. Wu et al., 2018; WCLC Abstract P1.01-97
107. Wu YL, et al. J. Clin. Oncol. (2018) PMID: 30156984
108. Gainer JF, et al. J Thorac Oncol (2020) PMID: 31864558
109. Gautschi O, et al. J Thorac Oncol (2020) PMID: 31864554
110. Faivre et al., 2021; ASCO GI Abstract 329
111. Le et al., 2021; ASCO Abstract 9021
112. Yang et al., 2019; AACR Abstract CT193
113. Park et al., 2019; ESMO Abstract 4770
114. Wu et al., 2019; IASLC Abstract MA09.09
115. Gan HK, et al. Clin. Cancer Res. (2019) PMID: 30952639
116. Lee J, et al. Cancer Discov (2019) PMID: 31315834
117. Kim ST, et al. Transl Oncol (2019) PMID: 30695737
118. Kwak et al., 2015; ASCO GI Abstract 01
119. Spigel DR, et al. J. Clin. Oncol. (2013) PMID: 24101053
120. Catenacci DV, et al. Cancer Discov (2011) PMID: 22389872
121. Harding JJ, et al. Clin. Cancer Res. (2019) PMID: 31142504
122. Strickler JH, et al. J. Clin. Oncol. (2018) PMID: 30285518
123. Camidge et al., 2021; AACR Abstract CT179
124. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
125. Gao J, et al. Sci Signal (2013) PMID: 23550210
126. Ang CS, et al. Anticancer Res. (2013) PMID: 23898085
127. Abou-Bakr AA, et al. Gulf J Oncolog (2013) PMID: 23996864
128. Ho JC, et al. Semin Respir Crit Care Med (2013) PMID: 24258573
129. Miller CT, et al. Oncogene (2006) PMID: 16186806
130. Tuyenman JB, et al. Br. J. Cancer (2008) PMID: 18349821
131. Anderson MR, et al. Clin. Cancer Res. (2006) PMID: 17062664
132. Graziano F, et al. J. Clin. Oncol. (2011) PMID: 22042954
133. Lee J, et al. Oncol. Rep. (2011) PMID: 21424128
134. Lee HE, et al. Br. J. Cancer (2012) PMID: 22644302
135. Cruz J, et al. Oncology (2003) PMID: 12837985
136. Yang L, et al. Hepatogastroenterology (2012) PMID: 22172411
137. Rossi G, et al. J. Clin. Oncol. (2005) PMID: 16314638
138. Camp RL, et al. Cancer (1999) PMID: 10590366
139. Lengyel E, et al. Int. J. Cancer (2005) PMID: 15455388
140. Ghoussoub RA, et al. Cancer (1998) PMID: 9554529
141. Yang JJ, et al. Lung Cancer (2013) PMID: 23079155
142. Dziadziuszko R, et al. J Thorac Oncol (2012) PMID: 22237262
143. Cappuzzo F, et al. J. Clin. Oncol. (2009) PMID: 19255323
144. Park S, et al. Histol. Histopathol. (2012) PMID: 22207554
145. Chen YT, et al. J Thorac Oncol (2011) PMID: 22052229
146. Kanteti R, et al. J. Environ. Pathol. Toxicol. Oncol. (2009) PMID: 19817696
147. To C, et al. Exp. Cell Res. (2002) PMID: 11795945
148. Tsuta K, et al. J Thorac Oncol (2012) PMID: 22198430
149. Felix AS, et al. Br. J. Cancer (2012) PMID: 22617129
150. Wagatsuma S, et al. Cancer (1998) PMID: 9452270
151. Bishop EA, et al. Gynecol. Oncol. (2011) PMID: 21168200
152. Felix AS, et al. Eur. J. Gynaecol. Oncol. (2010) PMID: 20527227
153. Garouniatis A, et al. Int J Colorectal Dis (2013) PMID: 22733437

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 | CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 | CLIA: 22D2027531

ORDERED TEST # **ORD-1336448-02**
APPENDIX
References

154. Liu Y, et al. Tumori () PMID: 22495710
155. Resnick MB, et al. Clin. Cancer Res. (2004) PMID: 15131045
156. Tanaka A, et al. Lung Cancer (2012) PMID: 21733594
157. J. Clin. Oncol. (2011) PMID: 22042966
158. Jung KH, et al. Arch. Pharm. Res. (2012) PMID: 22553051
159. Madoz-Gúrpide J, et al. J Transl Med (2015) PMID: 26319934
160. Ali SM, et al. Oncologist (2015) PMID: 25882375
161. Kwak EL, et al. Cancer Discov (2015) PMID: 26432108
162. Black LE, et al. Am. J. Pathol. (2019) PMID: 31351986
163. Baselga J, et al. Nat. Rev. Cancer (2009) PMID: 19536107
164. Jaiswal BS, et al. Cancer Cell (2013) PMID: 23680147
165. Jura N, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 20007378
166. Umelo I, et al. Oncotarget (2016) PMID: 26689995
167. Mishra R, et al. Oncotarget (2018) PMID: 29963236
168. Choudhury NJ, et al. J. Clin. Oncol. (2016) PMID: 27044931
169. Verlingue L, et al. Eur. J. Cancer (2018) PMID: 29413684
170. Goss GD, et al. JAMA Oncol (2018) PMID: 29902295
171. Bidard FC, et al. Ann. Oncol. (2015) PMID: 25953157
172. Hyman DM, et al. Nature (2018) PMID: 29420467
173. Cavalieri S, et al. Eur. J. Cancer (2018) PMID: 29734047
174. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
175. Gunes S, et al. Pathol. Res. Pract. (2013) PMID: 23333248
176. Bellezza G, et al. Virchows Arch. (2013) PMID: 24013863
177. Skrzypski M, et al. Clin Lung Cancer (2013) PMID: 23870818
178. Müller-Tidow C, et al. Cancer Res. (2005) PMID: 15753374
179. Yoon HH, et al. Cancer (2014) PMID: 24151090
180. Peng D, et al. Int. J. Cancer (2008) PMID: 18000824
181. Wei Q, et al. Int. J. Oncol. (2007) PMID: 17671674
182. Paterson AL, et al. J. Pathol. (2013) PMID: 22733579
183. Leng J, et al. Chin. Med. Sci. J. (1997) PMID: 11324501
184. Tanner B, et al. J. Clin. Oncol. (2006) PMID: 16896008
185. Lédel F, et al. Eur. J. Cancer (2014) PMID: 24300455
186. Song K, et al. Am J Cancer Res (2018) PMID: 29888103
187. Begnami MD, et al. J. Clin. Oncol. (2011) PMID: 21709195
188. Hayashi M, et al. Clin. Cancer Res. (2008) PMID: 19047113
189. Baiocchi G, et al. Int J Colorectal Dis (2009) PMID: 19390858
190. Sassen A, et al. Breast Cancer Res. (2008) PMID: 18182100
191. Witton CJ, et al. J. Pathol. (2003) PMID: 12845624
192. Kouttras AK, et al. Br. J. Cancer (2008) PMID: 18985033
193. Pawlowski V, et al. Clin. Cancer Res. (2000) PMID: 11106235
194. Travis A, et al. Br. J. Cancer (1996) PMID: 8688326
195. Quinn CM, et al. Histopathology (1994) PMID: 7821892
196. Gasparini G, et al. Eur. J. Cancer (1994) PMID: 7908213
197. Lemoine NR, et al. Br. J. Cancer (1992) PMID: 1333787
198. Rajkumar T, et al. J. Pathol. (1996) PMID: 8869284
199. Junttila TT, et al. Clin. Cancer Res. (2003) PMID: 14614020
200. Røtterud R, et al. BJU Int. (2005) PMID: 15892828
201. Tsai YS, et al. Urology (2005) PMID: 16360440
202. Chow NH, et al. Virchows Arch. (1997) PMID: 9230911
203. Chow NH, et al. Clin. Cancer Res. (2001) PMID: 11448910
204. Sheng Q, et al. Br. J. Cancer (2011) PMID: 21364581
205. Li M, et al. Nat. Genet. (2014) PMID: 24997986
206. Collier TS, et al. J. Biol. Chem. (2013) PMID: 23843458
207. Li YY, et al. Clin. Cancer Res. (2015) PMID: 25589618
208. Seshagiri S, et al. Nature (2012) PMID: 22895193
209. Kim PH, et al. Eur. Urol. (2015) PMID: 25092538
210. Ke L, et al. Proc. Natl. Acad. Sci. U.S.A. (2017) PMID: 28831010
211. Liang JZ, et al. Int J Clin Exp Pathol (2015) PMID: 26045802
212. Kralik JM, et al. Diagn Pathol (2011) PMID: 21401966
213. De Braekeleer E, et al. Leuk. Res. (2012) PMID: 22578774
214. Włodarska I, et al. Leuk. Lymphoma (1996) PMID: 9031109
215. Montpetit A, et al. Oncogene (2001) PMID: 11423994
216. Donaldson LW, et al. EMBO J. (1996) PMID: 8598195
217. Green SM, et al. J. Biol. Chem. (2010) PMID: 20400516
218. Tognon CE, et al. Mol. Cell. Biol. (2004) PMID: 15143160
219. Kim CA, et al. EMBO J. (2001) PMID: 11483520
220. Schechter RB, et al. Cancer Lett. (2015) PMID: 25641339
221. Guo T, et al. Genes Chromosomes Cancer (2011) PMID: 20949568
222. Koch S, et al. Biochem. J. (2011) PMID: 21711246
223. Weryńska B, et al. Folia Histochem. Cytobiol. (2009) PMID: 20164015
224. Yusufova N, et al. Nature (2021) PMID: 33299181
225. Krysiak K, et al. Blood (2017) PMID: 28064239
226. Okosun J, et al. Nat. Genet. (2014) PMID: 24362818
227. Li H, et al. Blood (2014) PMID: 24435047
228. Lohr JG, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22343534
229. Sancho M, et al. PLoS Genet. (2008) PMID: 18927631
230. Soshnev AA, et al. Cancer Res (2021) PMID: 34580064
231. Binkley MS, et al. Cancer Discov (2020) PMID: 33071215
232. Chowdhry S, et al. Oncogene (2013) PMID: 22964642
233. Abazeed ME, et al. Cancer Res. (2013) PMID: 23980093
234. Chen X, et al. Ann Transl Med (2020) PMID: 32175433
235. Xu X, et al. Oncologist (2020) PMID: 32272498
236. Arbour et al., 2018; IASLC WCLC Abstract MA19.09
237. Zhang C, et al. J Thorac Oncol (2020) PMID: 32471565
238. Shang et al., 2020; WCLC Abstract P75.02
239. Marinelli D, et al. Ann Oncol (2020) PMID: 32866624
240. Cho et al., 2020; AACR Abstract CT084
241. Gwinn DM, et al. Cancer Cell (2018) PMID: 29316436
242. Sayin VI, et al. Elife (2017) PMID: 28967864
243. Romero R, et al. Nat. Med. (2017) PMID: 28967920
244. Skoulidis et al., 2021; ASCO Abstract TPS9627
245. Dai B, et al. Cancer Res. (2013) PMID: 23824739
246. Yoo NJ, et al. Histopathology (2012) PMID: 22348534
247. Lo SC, et al. J. Biol. Chem. (2006) PMID: 17046835
248. Wakabayashi N, et al. Nat. Genet. (2003) PMID: 14517554
249. Kansanen E, et al. Redox Biol (2013) PMID: 24024136
250. Hast BE, et al. Cancer Res. (2013) PMID: 23382044
251. Ito S, et al. Nature (2010) PMID: 20639862
252. Guo JU, et al. Cell (2011) PMID: 21496894
253. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
254. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
255. Xie M, et al. Nat. Med. (2014) PMID: 25326804
256. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
257. Severson EA, et al. Blood (2018) PMID: 29678827
258. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
259. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
260. Chabon JJ, et al. Nature (2020) PMID: 32269342
261. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
262. Hirai H, et al. Cancer Biol. Ther. (2010) PMID: 20107315
263. Bridges KA, et al. Clin. Cancer Res. (2011) PMID: 21799033
264. Rajeshkumar NV, et al. Clin. Cancer Res. (2011) PMID: 21389100
265. Osman AA, et al. Mol. Cancer Ther. (2015) PMID: 25504633
266. Xu L, et al. Mol. Cancer Ther. (2002) PMID: 12489850
267. Xu L, et al. Mol. Med. (2001) PMID: 11713371
268. Camp ER, et al. Cancer Gene Ther. (2013) PMID: 23470564
269. Kim SS, et al. Nanomedicine (2015) PMID: 25240597
270. Pirolo KF, et al. Mol. Ther. (2016) PMID: 27357628
271. Hajdenberg et al., 2012; ASCO Abstract e15010
272. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27601554
273. Moore et al., 2019; ASCO Abstract 5513
274. Leijen S, et al. J. Clin. Oncol. (2016) PMID: 27998224
275. Oza et al., 2015; ASCO Abstract 5506
276. Méndez E, et al. Clin. Cancer Res. (2018) PMID: 29535125
277. Seligmann JF, et al. J Clin Oncol (2021) PMID: 34538072
278. Kwok M, et al. Blood (2016) PMID: 26563132
279. Boudny M, et al. Haematologica (2019) PMID: 30975914
280. Dillon MT, et al. Mol. Cancer Ther. (2017) PMID: 28062704
281. Middleton FK, et al. Cancers (Basel) (2018) PMID: 30127241
282. Kandath C, et al. Nature (2013) PMID: 24132290
283. Dong ZY, et al. Clin. Cancer Res. (2017) PMID: 28039262
284. Russo A, et al. J. Clin. Oncol. (2005) PMID: 16172461
285. Brown CJ, et al. Nat. Rev. Cancer (2009) PMID: 19935675
286. Joerger AC, et al. Annu. Rev. Biochem. (2008) PMID: 18410249
287. Kato S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12826609
288. Kamada R, et al. J. Biol. Chem. (2011) PMID: 20978130
289. Zerdouni Y, et al. Hum. Mol. Genet. (2017) PMID: 28472496
290. Yamada H, et al. Carcinogenesis (2007) PMID: 17690113
291. Bougeard G, et al. J. Clin. Oncol. (2015) PMID: 26014290
292. Sorrell AD, et al. Mol Diagn Ther (2013) PMID: 23351000
293. Nichols KE, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11219776
294. Kleihues P, et al. Am. J. Pathol. (1997) PMID: 9006316
295. Gonzalez KD, et al. J. Clin. Oncol. (2009) PMID: 19204208
296. Lalloo F, et al. Lancet (2003) PMID: 12672316
297. Mandelker D, et al. Ann. Oncol. (2019) PMID: 31050713
298. Herman SE, et al. Blood (2014) PMID: 24659631
299. Herman SE, et al. Blood (2011) PMID: 21422473
300. Rushworth SA, et al. Cell. Signal. (2013) PMID: 22975686
301. Rahal R, et al. Nat. Med. (2014) PMID: 24362935
302. Xie P, et al. Cytokine Growth Factor Rev. () PMID: 18499506
303. Kayagaki N, et al. Science (2007) PMID: 17991829
304. Keats JJ, et al. Cancer Cell (2007) PMID: 17692805
305. Liao G, et al. J. Biol. Chem. (2004) PMID: 15084608
306. Annunziata CM, et al. Cancer Cell (2007) PMID: 17692804
307. Nagel I, et al. Leukemia (2009) PMID: 19693093
308. Gardam S, et al. Blood (2011) PMID: 21300983
309. Marabelle et al., 2019; ESMO Abstract 11920
310. Andre et al., 2021; ASCO GI Abstract 9
311. Oaknin A, et al. JAMA Oncol (2020) PMID: 33001143
312. Berton et al., 2021; ASCO Abstract 2564
313. Andre et al., 2021; ESMO GI Abstract SO-9
314. Muro K, et al. Lancet Oncol. (2016) PMID: 27157491
315. Doi et al., 2015; ASCO Abstract 4010
316. Ott et al., 2016; ASCO Abstract 5581

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor. Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX **References**

317. Herbst RS, et al. Lancet (2016) PMID: 26712084
318. Pilmack et al., 2014; ESMO Abstract LBA23
319. Chow et al., 2014; ESMO Abstract LBA31
320. Hsu et al., 2015; ESMO Abstract 2801
321. Cohen et al., 2016; ASCO Abstract 6017
322. Seiwert TY, et al. Lancet Oncol. (2016) PMID: 27247226
323. Nanda et al., 2014; SABCS Abstract S1-09
324. Nanda R, et al. J. Clin. Oncol. (2016) PMID: 27138582
325. Frenel et al., 2016; ASCO Abstract 5515
326. Mehnert et al., 2016; ASCO Abstract 6091
327. Alley et al., 2015; AACR Abstract CT103
328. Georger B, et al. Lancet Oncol. (2020) PMID: 31812554
329. Patnaik et al., 2015; ASCO Abstract 8011
330. Gubens et al., 2016; ASCO Abstract 9027
331. Schaff et al., 2015; SNO Abstract IMCT-18
332. Papadimitrakopoulou et al., 2015; ASCO Abstract 8031
333. Gangadhar et al., 2015; SITC Abstract 511
334. Reardon et al., 2016; ASCO Abstract 2010
335. Makker V, et al. Lancet Oncol. (2019) PMID: 30922731
336. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
337. Smith et al., 2016; ASCO Abstract 9028
338. Mazieres et al., 2016; ASCO Abstract 9032
339. Besse et al., 2015; ECC Abstract 16LBA
340. Spigel et al., 2015; ASCO Abstract 8028
341. Fehrenbacher L, et al. Lancet (2016) PMID: 26970723
342. Herbst RS, et al. Nature (2014) PMID: 25428504
343. Balar et al., 2016; ASCO Abstract LBA4500
344. Dreicer et al., 2016; ASCO Abstract 4515
345. Rosenberg JE, et al. Lancet (2016) PMID: 26952546
346. Powles T, et al. Nature (2014) PMID: 25428503
347. Georger B, et al. Lancet Oncol. (2020) PMID: 31780255
348. McDermott DF, et al. J. Clin. Oncol. (2016) PMID: 26755520
349. Adams et al., 2016; ASCO Abstract 1009
350. Hellmann MD, et al. Ann. Oncol. (2019) PMID: 30918950
351. Bendell et al., 2016; ASCO Abstract 3502
352. Verschraegen et al., 2016; ASCO Abstract 9036
353. Chung et al., 2016; ASCO Abstract 4009
354. Patel et al., 2016; ESMO Abstract 777PD
355. Hellmann et al., 2017; ASCO Abstract 8503
356. Disis et al., 2016; ASCO Abstract 5533
357. Dirix et al., 2016; SABCS Abstract S1-04
358. Larkin et al., 2016; ESMO Abstract 775PD
359. Le Tourneau et al., 2016; ASCO Abstract 4516
360. Fakhrejahani et al., 2017; ASCO GU Abstract 159
361. Rajan et al., 2016; ASCO Abstract e20106
362. Paik PK, et al. Cancer Discov (2015) PMID: 25971939
363. Klemptner SJ, et al. J Thorac Oncol (2017) PMID: 27693535
364. Yakes FM, et al. Mol. Cancer Ther. (2011) PMID: 21926191
365. Weber H, et al. J Biomol Screen (2014) PMID: 25260782
366. Navis AC, et al. PLoS ONE (2013) PMID: 23484006
367. Yeh I, et al. Nat Commun (2015) PMID: 26013381
368. Lee YH, et al. Cancers (Basel) (2014) PMID: 25534569
369. Torres KE, et al. Clin. Cancer Res. (2011) PMID: 21540237
370. Sameni M, et al. Clin. Cancer Res. (2016) PMID: 26432786
371. Schöffski P, et al. Eur. J. Cancer (2017) PMID: 29059635
372. Hellerstedt BA, et al. Clin Lung Cancer (2019) PMID: 30528315
373. Nokihara H, et al. Clin Lung Cancer (2019) PMID: 30718102
374. Elisei R, et al. J. Clin. Oncol. (2013) PMID: 24002501
375. Abou-Alfa GK, et al. N. Engl. J. Med. (2018) PMID: 29972759
376. Choueiri TK, et al. J. Clin. Oncol. (2017) PMID: 28199818
377. Italiano A, et al. Lancet Oncol. (2020) PMID: 32078813
378. Wu et al., 2018; WLCL Abstract P1.01-97
379. Wolf et al., 2020; ASCO Abstract 9509
380. Schuler M, et al. Ann. Oncol. (2020) PMID: 32240796
381. Wolf J, et al. N Engl J Med (2020) PMID: 32877583
382. van den Bent M, et al. J. Neurooncol. (2020) PMID: 31776899
383. Bang YJ, et al. Cancer Sci. (2020) PMID: 31778267
384. Esaki T, et al. Cancer Sci. (2019) PMID: 30724423
385. Migden MR, et al. N. Engl. J. Med. (2018) PMID: 29863979
386. Stratigos et al., 2020; ESMO Abstract LBA47
387. Lewis et al. 2020; DOI: 10.1136/jitc-2020-SITC2020.0428
388. Sezer et al., 2020; ESMO Abstract LBA52
389. Vassal et al., 2015; ASCO Abstract 2595
390. Li et al., 2015; ASCO Abstract 8090
391. Frampton GM, et al. Cancer Discov (2015) PMID: 25971938
392. Benderra MA, et al. J Thorac Oncol (2016) PMID: 26845121
393. Waqar SN, et al. J Thorac Oncol (2015) PMID: 25898962
394. Mendenhall MA, et al. J Thorac Oncol (2015) PMID: 25898965
395. Jenkins RW, et al. Clin Lung Cancer (2015) PMID: 25769807
396. Stein MN, et al. Eur. Urol. (2015) PMID: 25457019
397. Awad et al., 2017; ASCO Abstract 8511
398. Shaw et al., 2016; ASCO Abstract 9066
399. Lu et al., 2016; ASCO Abstract 9058
400. Yoshida T, et al. J. Clin. Oncol. (2016) PMID: 27354483
401. Solomon BJ, et al. N. Engl. J. Med. (2014) PMID: 25470694
402. Shaw AT, et al. N. Engl. J. Med. (2013) PMID: 23724913
403. Moro-Sibilot et al., 2015; ASCO Abstract 8065
404. Goto et al., 2016; ASCO Abstract 9022
405. Shaw AT, et al. N. Engl. J. Med. (2014) PMID: 25264305
406. Mazières J, et al. J. Clin. Oncol. (2015) PMID: 25667280
407. Scheffler M, et al. Oncotarget (2015) PMID: 25868855
408. Vaishnavi A, et al. Nat. Med. (2013) PMID: 24162815
409. Drilon et al., 2016; ASCO Abstract 108
410. Camidge et al., 2014; ASCO Abstract 8001
411. Schrock AB, et al. J Thorac Oncol (2016) PMID: 27343443
412. Jorge SE, et al. Lung Cancer (2015) PMID: 26791794
413. Mahjoubi L, et al. Invest New Drugs (2016) PMID: 26892698
414. Awad MM, et al. J. Clin. Oncol. (2016) PMID: 26729443
415. Zhang Y, et al. J Thorac Oncol (2016) PMID: 26724472
416. Diamond JR, et al. J. Clin. Oncol. (2013) PMID: 23610116
417. Gambacorti-Passerini et al., 2017; ASH Abstract 4128
418. Mossé YP, et al. Lancet Oncol. (2013) PMID: 23598171
419. Bais et al., 2017; AACR Abstract 3720/5
420. Garassino et al., 2016; IASLC Abstract PLO4a.03
421. Segal et al., 2016; ESMO Abstract 9490
422. Segal et al., 2015; ASCO Abstract 3011
423. Lutzky et al., 2014; ASCO Abstract 3001
424. Iguchi et al., 2015; ASCO Abstract 3039
425. Ribas et al., 2015; ASCO Abstract 3003
426. Karzai et al., 2017; ASCO Genitourinary Abstract 162
427. Lee et al., 2016; ASCO Abstract 3015
428. Necchi A, et al. Eur. Urol. (2019) PMID: 30243800
429. Hamid et al., 2016; ESMO Abstract 1050PD
430. Hong et al., 2016; ESMO 2016 Abstract 1049PD
431. Yap et al., 2016; EORTC-NCI-AACR Abstract 1LBA
432. Levy A, et al. Eur. J. Cancer (2016) PMID: 27764686
433. Postow MA, et al. N. Engl. J. Med. (2015) PMID: 25891304
434. Larkin J, et al. N. Engl. J. Med. (2015) PMID: 26027431
435. Robert C, et al. N. Engl. J. Med. (2015) PMID: 25399552
436. Topalian SL, et al. J. Clin. Oncol. (2014) PMID: 24590637
437. Weber JS, et al. Lancet Oncol. (2015) PMID: 25795410
438. Brahmer et al., 2014; ASCO Abstract 8112
439. Gettinger et al., 2014; ASCO Abstract 8024
440. Antonia et al., 2014; ASCO Abstract 8023
441. Rizvi et al., 2014; ASCO Abstract 8022
442. Antonia et al., 2014; ASCO Abstract 8113
443. Brahmer J, et al. N. Engl. J. Med. (2015) PMID: 26028407
444. Borghaei H, et al. N. Engl. J. Med. (2015) PMID: 26412456
445. Sharma P, et al. J. Clin. Oncol. (2019) PMID: 31100038
446. Sharma P, et al. Lancet Oncol. (2017) PMID: 28131785
447. Amin et al., 2014; ASCO Abstract 5010
448. McDermott et al., 2016; ASCO Abstract 4507
449. Motzer RJ, et al. J. Clin. Oncol. (2015) PMID: 25452452
450. Topalian SL, et al. N. Engl. J. Med. (2012) PMID: 22658127
451. Brahmer JR, et al. J. Clin. Oncol. (2010) PMID: 20516446
452. Motzer RJ, et al. N. Engl. J. Med. (2015) PMID: 26406148
453. Overman MJ, et al. Lancet Oncol. (2017) PMID: 28734759
454. Overman MJ, et al. J. Clin. Oncol. (2018) PMID: 29355075
455. Overman et al., 2019; ASCO GI Abstract 481
456. Gillison et al., 2016; AACR Abstract CT099
457. Ferris et al., 2016; ASCO Abstract 6009
458. Ferris et al., 2019; 31239321
459. Brahmer JR, et al. N. Engl. J. Med. (2012) PMID: 22658128
460. Hamanishi J, et al. J. Clin. Oncol. (2015) PMID: 26351349
461. Normann MC, et al. J Gynecol Oncol (2019) PMID: 31074244
462. Calvo et al., 2015; ECC Abstract 3098
463. Antonia SJ, et al. Lancet Oncol. (2016) PMID: 27269741
464. Le et al., 2016; ASCO GI Cancers Abstract 06
465. Kojima et al., 2016; ASCO GI Cancers Abstract TPS175
466. Janjigian YY, et al. J. Clin. Oncol. (2018) PMID: 30110194
467. Tanizaki J, et al. Ann Oncol (2021) PMID: 34843940
468. Ansell et al., 2015; ASH Abstract 583
469. Ansell SM, et al. N. Engl. J. Med. (2015) PMID: 25482239
470. Younes A, et al. Lancet Oncol. (2016) PMID: 27451390
471. Hodi FS, et al. Lancet Oncol. (2018) PMID: 30361170
472. Hodi FS, et al. Lancet Oncol. (2016) PMID: 27622997
473. Larkin J, et al. N. Engl. J. Med. (2019) PMID: 31562797
474. Hellmann MD, et al. N. Engl. J. Med. (2019) PMID: 31562796
475. Lenz et al., 2020; ASCO GI Abstract 11
476. Motzer RJ, et al. Lancet Oncol. (2019) PMID: 31427204
477. Motzer RJ, et al. N. Engl. J. Med. (2018) PMID: 29562145
478. Paoluzzi et al., 2016; ASCO Abstract 11047
479. Broto et al., 2019; ESMO Abstract 16690
480. Nadal et al., 2018; ASCO Abstract 4528
481. Kollipara R, et al. Oncologist (2017) PMID: 28778959
482. Hodi et al., 2019; AACR Abstract CT037
483. Paik PK, et al. N. Engl. J. Med. (2020) PMID: 32469185
484. Mazieres et al., 2020; ESMO Abstract 1283P
485. Ryoo et al., 2018; ESMO Abstract 186P
486. Ryoo et al., 2018; ESMO Abstract 621PD
487. Decaens et al., 2019; ESMO Abstract 698P
488. Falchook GS, et al. Clin. Cancer Res. (2020) PMID: 31822497

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 • CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1336448-02

APPENDIX

References

489. Shitara K, et al. Jpn. J. Clin. Oncol. (2020) PMID:

32328660

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2022 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Leah Commander, M.D. | 19 April 2022
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | 1.888.988.3639

Sample Preparation: 7010 Kit Creek Road, Morrisville, NC 27560 · CLIA: 34D2044309
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531