

Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

**Date:** 27 Jun 2023 1 of 3

# **Sample Information**

Patient Name: 蔡況治 Gender: Female ID No.: W200235640 History No.: 36397292

**Age**: 66

Ordering Doctor: DOC1697J 蔡淳光 Ordering REQ.: 0CMGZEE Signing in Date: 2023/06/27

**Path No.:** M112-00154 **MP No.:** MY23035

**Assay:** Oncomine Myeloid Assay **Sample Type:** Bone Marrow

Bone Marrow Aspirating Date: 2023/06/16

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

## Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page
Variants (Exclude variant in Taiwan	2
BioBank with >1% allele frequency)	
Biomarker Descriptions	2

# **Report Highlights**

0 Relevant Biomarkers0 Therapies Available0 Clinical Trials

#### **Relevant Acute Myeloid Leukemia Variants**

Gene	Finding	Gene	Finding
ABL1	None detected	MECOM	None detected
ASXL1	None detected	MLLT3	None detected
CEBPA	None detected	MYH11	None detected
CREBBP	None detected	NPM1	None detected
FLT3	None detected	NUP214	None detected
IDH1	None detected	RARA	None detected
IDH2	None detected	RUNX1	None detected
KMT2A	None detected	TP53	None detected

# **Relevant Biomarkers**

No clinically significant biomarkers found in this sample.

Prevalent cancer biomarkers without relevant evidence based on included data sources

JAK2 p.(V617F) c.1849G>T

**Date**: 27 Jun 2023 2 of 3

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

#### **DNA Sequence Variants** Allele Gene Amino Acid Change Codina Variant ID Locus Frequency Transcript Variant Effect Coverage JAK2 p.(V617F) c.1849G>T COSM12600 chr9:5073770 79.71% NM 004972.4 missense 1986 TP53 p.(I232T) c.695T>C chr17:7577586 62.28% NM\_000546.5 missense 1999 U2AF1 p.(P40=)c.120G>A chr21:44524437 52.13% NM\_006758.2 synonymous 1997

### **Biomarker Descriptions**

#### JAK2 (Janus kinase 2)

Background: The JAK2 gene encodes a non-receptor, membrane associated protein tyrosine kinase (PTK). JAK2 is a member of the Janus kinase (JAK) family that includes JAK1, JAK2, JAK3, and TYK2. Janus kinases are characterized by the presence of a second phosphotransferase-related or pseudokinase domain immediately N-terminal to the PTK domain¹. JAK kinases function with signal transducer and activator of transcription (STAT) proteins to facilitate intracellular signal transduction required for cytokine receptor and interferon-alpha/beta/gamma signaling¹,2,3. Since JAK2 functions in interferon receptor signaling, inactivation of JAK2 is proposed to inhibit presentation of tumor antigens and contribute to immune evasion⁴,5.

Alterations and prevalence: Clonal expansion of hematopoietic cells in myeloproliferative neoplasms (MPNs) has been associated with loss of heterozygosity on chromosome 9p and subsequently to the acquisition of a dominant somatic gain-of-function V617F mutation in the pseudokinase domain of JAK2<sup>6,7</sup>. The JAK2 V617F mutation has been observed rarely in acute myeloid leukemia (AML)<sup>8,9</sup>. Mutations in the pseudokinase domain of JAK2 including R683G have been detected in 8% of ALL<sup>10,11</sup>. JAK2 fusions are observed in myeloid and lymphoid leukemias with partner genes including TEL, PCM1, and BCR genes<sup>12,13,14,15</sup>. JAK2 fusions are infrequently observed in solid tumors<sup>16</sup>. As with JAK1, truncating mutations in JAK2 are common in solid tumors and particularly enriched in uterine cancers<sup>16</sup>.

Potential relevance: Currently, no therapies are approved for JAK2 aberrations. JAK2 V617F and JAK2 exon 12 mutations are considered major diagnostic criteria of PV<sup>17,18</sup>. Ruxolitinib<sup>19</sup> (2011) is a JAK1/2 inhibitor FDA approved for PMF and PV, although specific JAK2 alterations are not indicated. Other JAK inhibitors including tofacitinib (2012) and baricitinib (2018) are approved for the treatment of rheumatoid arthritis. Clinical cases associated with high tumor mutational burden (TMB) but failure to respond to anti-PD1 therapy were associated with loss of function mutations in JAK1/2<sup>20</sup>. Some case studies report efficacy with ruxolitinib in myeloid and lymphoid leukemias, although duration of complete response was limited<sup>12,13,14,15</sup>.

Date: 27 Jun 2023

#### References

- 1. Babon et al. The molecular regulation of Janus kinase (JAK) activation. Biochem. J. 2014 Aug 15;462(1):1-13. PMID: 25057888
- 2. Müller et al. The protein tyrosine kinase JAK1 complements defects in interferon-alpha/beta and -gamma signal transduction. Nature. 1993 Nov 11;366(6451):129-35. PMID: 8232552
- 3. Ren et al. JAK1 truncating mutations in gynecologic cancer define new role of cancer-associated protein tyrosine kinase aberrations. Sci Rep. 2013 Oct 24;3:3042. PMID: 24154688
- 4. Zaretsky et al. Mutations Associated with Acquired Resistance to PD-1 Blockade in Melanoma. N. Engl. J. Med. 2016 Sep 1;375(9):819-29. PMID: 27433843
- 5. Garcia-Diaz et al. Interferon Receptor Signaling Pathways Regulating PD-L1 and PD-L2 Expression. Cell Rep. 2017 May 9;19(6):1189-1201. PMID: 28494868
- Baxter et al. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. Lancet. 2005 Mar 19-25;365(9464):1054-61. PMID: 15781101
- 7. Kralovics et al. A gain-of-function mutation of JAK2 in myeloproliferative disorders. N. Engl. J. Med. 2005 Apr 28;352(17):1779-90. PMID: 15858187
- 8. Hidalgo-López et al. Morphologic and Molecular Characteristics of De Novo AML With JAK2 V617F Mutation. J Natl Compr Canc Netw. 2017 Jun;15(6):790-796. PMID: 28596259
- 9. Aynardi et al. JAK2 V617F-positive acute myeloid leukaemia (AML): a comparison between de novo AML and secondary AML transformed from an underlying myeloproliferative neoplasm. A study from the Bone Marrow Pathology Group. Br. J. Haematol. 2018 Jul;182(1):78-85. PMID: 29767839
- 10. Mullighan et al. JAK mutations in high-risk childhood acute lymphoblastic leukemia. Proc. Natl. Acad. Sci. U.S.A. 2009 Jun 9;106(23):9414-8. PMID: 19470474
- 11. Scott. Lymphoid malignancies: Another face to the Janus kinases. Blood Rev. 2013 Mar;27(2):63-70. PMID: 23340138
- 12. Chase et al. Ruxolitinib as potential targeted therapy for patients with JAK2 rearrangements. Haematologica. 2013 Mar;98(3):404-8. PMID: 22875628
- 13. Rumi et al. Efficacy of ruxolitinib in chronic eosinophilic leukemia associated with a PCM1-JAK2 fusion gene. J. Clin. Oncol. 2013 Jun 10;31(17):e269-71. PMID: 23630205
- 14. Schwaab et al. Limited duration of complete remission on ruxolitinib in myeloid neoplasms with PCM1-JAK2 and BCR-JAK2 fusion genes. Ann. Hematol. 2015 Feb;94(2):233-8. PMID: 25260694
- 15. Rumi et al. Efficacy of ruxolitinib in myeloid neoplasms with PCM1-JAK2 fusion gene. Ann. Hematol. 2015 Nov;94(11):1927-8. PMID: 26202607
- 16. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
- 17. NCCN Guidelines® NCCN-Myeloproliferative Neoplasms [Version 3.2022]
- 18. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia. 2022 Jul;36(7):1703-1719. PMID: 35732831
- 19. https://www.accessdata.fda.gov/drugsatfda\_docs/label/2023/202192s028lbl.pdf
- 20. Shin et al. Primary Resistance to PD-1 Blockade Mediated by JAK1/2 Mutations. Cancer Discov. 2017 Feb;7(2):188-201. PMID: 27903500