



Sample Information

Patient Name: 曾魏茂蘭

Gender: Female

ID No.: J200477333

History No.: 2314694

Age: 79

Ordering Doctor: DOC3064F 陳育民

Ordering REQ.: C24MHJN

Signing in Date: 2021/10/15

Path No.: S110-99779

MP No.: F21083

Assay: Oncomine Focus Assay

Sample Type: FFPE

Block No.: S110-28279A

Percentage of tumor cells: 50%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page
Variant Details	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	4
Clinical Trials Summary	8
Alert Details	8

Report Highlights

- 1 Relevant Biomarkers
- 3 Therapies Available
- 4 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	MET exon 14 skipping		

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	MET exon 14 skipping MET proto-oncogene, receptor tyrosine kinase Prognostic significance: None Diagnostic significance: None	capmatinib ¹ tepotinib ¹ crizotinib	None	4

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ALK	p.(D1529E)	c.4587C>G	.	chr2:29416366	48.35%	NM_004304.5	missense	1998
PDGFRA	p.(G313=)	c.939T>G	.	chr4:55133726	49.62%	NM_006206.6	synonymous	1997
PDGFRA	p.(V824=)	c.2472C>T	.	chr4:55152040	48.07%	NM_006206.6	synonymous	1999
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.55%	NM_213647.3	missense	1999
EGFR	p.(Q787=)	c.2361G>A	.	chr7:55249063	50.90%	NM_005228.5	synonymous	1998

Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
MET-MET	MET-MET.M13M15	chr7:116411708 - chr7:116414935	33890

Biomarker Descriptions

MET (MET proto-oncogene, receptor tyrosine kinase)

Background: The MET proto-oncogene encodes a receptor tyrosine kinase for the hepatocyte growth factor (HGF) protein, which is expressed by mesenchymal cells. Ubiquitin-dependent proteolysis regulates the steady state level of the MET protein via recognition of the tyrosine phosphorylation site Y1003 in the MET Cbl-binding domain within the juxtamembrane region^{1,2,3}. Growth factor signaling leads to MET dimerization and subsequent initiation of downstream effectors including those involved in the RAS/RAF/MEK/ERK and PI3K/AKT signaling pathways, which regulate cell migration, proliferation, and survival^{4,5}.

Alterations and prevalence: Recurrent somatic MET alterations include activating mutations, gene amplification, and translocations generating MET gene fusions. Recurrent somatic mutations fall into two classes, mutations in the MET kinase domain, which are uncommon, and splice-site mutations affecting exon 14. Recurrent kinase domain mutations are observed in papillary renal cell carcinoma (PRCC) (1-2%) and include M1250T, H1094Y, and V1070E. Mutation of the Y1003 phosphorylation site is reported in lung cancer but is uncommon (<1%)^{6,7}. In contrast, splice-site mutations flanking exon 14 are observed in 4% of non-small cell lung cancer (NSCLC). These mutations include canonical splice site mutations affecting exon 14 and deletions that extend into the splicing motifs within intron 13^{8,9}. Such mutations disrupt splicing leading to the formation of an alternative transcript that joins exon 13 directly to exon 15 and skips exon 14 entirely. The MET exon 14 skipping transcript lacks the juxtamembrane domain that contains the recognition motif for ubiquitin-dependent proteolysis and thus leads to a marked increase in steady-state level of the MET protein¹⁰. MET exon 14 skipping mutations act as oncogenic drivers in lung cancer mutually exclusive to activating mutations in EGFR and KRAS and other oncogenic fusions such as ALK and ROS1^{8,11,12}. MET is amplified in 2-5% of ovarian cancer, esophageal adenocarcinoma, stomach adenocarcinoma, glioblastoma, and lung adenocarcinoma^{6,7,13}. Recurrent MET fusions, although infrequent, are observed in

Biomarker Descriptions (continued)

adult and pediatric glioblastoma, papillary renal cell carcinoma, lung cancer, liver cancer, thyroid cancer, and melanoma^{14,15,16}. MET alterations are believed to be enriched in late-stage cancers where they drive tumor progression and metastasis^{17,18,19}.

Potential relevance: In 2020, the FDA granted accelerated approval to capmatinib²⁰ for NSCLC harboring MET exon 14 skipping positive as detected by an FDA-approved test²¹. The kinase inhibitor, tepotinib²², is also approved (2021) for MET exon 14 skipping mutations in NSCLC²². MET exon 14 skipping mutations confer sensitivity to approved kinase inhibitors including crizotinib (2011), which is recommended for MET amplifications and exon 14 skipping mutations^{8,11,12,21}. The FDA also granted breakthrough therapy designation (2018) to crizotinib for metastatic non-small cell lung cancer (NSCLC) with MET exon 14 alterations with disease progression on or after platinum-based chemotherapy²³. Conversely, amplification of MET has been observed to mediate resistance to EGFR tyrosine kinase inhibitors (TKIs)^{24,25,26,27,28}. In a phase II trial testing the MET inhibitor savolitinib, patients with advanced PRCC exhibited median progression free survival (PFS) of 6.2 and 1.4 months for MET-driven and MET-independent PRCC, respectively²⁹.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ✕ No evidence

MET exon 14 skipping

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
capmatinib	●	●	✕	●	✕
tepotinib	●	●	✕	●	✕
crizotinib	✕	●	✕	●	✕
datopotamab deruxtecan	✕	✕	✕	✕	● (II)
bozitinib	✕	✕	✕	✕	● (I/II)
amivantamab	✕	✕	✕	✕	● (I)
HLX55	✕	✕	✕	✕	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

FDA information is current as of 2021-08-18. For the most up-to-date information, search www.fda.gov.

MET exon 14 skipping

● capmatinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2020-05-06

Variant class: MET exon 14 skipping

Indications and usage:

TABRECTA™ is a kinase inhibitor indicated for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trial(s).

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/213591s000lbl.pdf

● tepotinib

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2021-02-03

Variant class: MET exon 14 skipping

Indications and usage:

TEPMETKO® is a kinase inhibitor indicated for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) harboring mesenchymal-epithelial transition (MET) exon 14 skipping alterations.

This indication is approved under accelerated approval based on overall response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214096s000lbl.pdf

Current NCCN Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

NCCN information is current as of 2021-08-02. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

MET exon 14 skipping

● capmatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Brain Metastases (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 1.2021]

● capmatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Useful in certain circumstances
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy)
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression (Subsequent therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

MET exon 14 skipping (continued)

● tepotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (First-line therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic (Subsequent therapy); Preferred intervention
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]

Current ESMO Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

ESMO information is current as of 2021-08-02. For the most up-to-date information, search www.esmo.org.

MET exon 14 skipping

● capmatinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

● crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

● tepotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Non-Small-Cell Lung Cancer [Online Guideline (15SEP2020 - <https://www.esmo.org/guidelines/lung-and-chest-tumours/clinical-practice-living-guidelines-metastatic-non-small-cell-lung-cancer>); Ann Oncol (2018) 29 (suppl 4): iv192–iv237.]

Clinical Trials in Taiwan region:

Clinical Trials Summary

MET exon 14 skipping

NCT ID	Title	Phase
NCT04484142	Phase II, Single-arm, Open-label Study of DS-1062a in Advanced or Metastatic Non-small Cell Lung Cancer With Actionable Genomic Alterations and Progressed on or After Kinase Inhibitor Therapy and Platinum Based Chemotherapy (TROPION-Lung05)	II
NCT03175224	Phase I/II Multicenter Study of the Safety, Pharmacokinetics, and Preliminary Efficacy of APL-101 in Subjects With Non-Small Cell Lung Cancer With c-Met EXON 14 Skip Mutations and c-Met Dysregulation Advanced Solid Tumors.	I/II
NCT02609776	A Phase I, First-in-Human, Open-Label, Dose Escalation Study of JNJ-61186372, a Human Bispecific EGFR and cMet Antibody, in Subjects With Advanced Non-Small Cell Lung Cancer.	I
NCT04169178	A Phase I Dose Finding/Expansion Study of HLX55, A Monoclonal Antibody Targeting Tyrosine-Protein Kinase MET (C-MET) in Patients With Advanced Solide Tumors Refractory to Standard Therapy	I

Alerts Informed By Public Data Sources

Current FDA Information

 Contraindicated
  Not recommended
  Resistance
  Breakthrough
  Fast Track

FDA information is current as of 2021-08-18. For the most up-to-date information, search www.fda.gov.

MET exon 14 skipping

crizotinib

Cancer type: Non-Small Cell Lung Cancer

Variant class: MET exon 14 skipping

Supporting Statement:

The FDA has granted Breakthrough Therapy Designation to the tyrosine kinase inhibitor, crizotinib, for metastatic non-small cell lung cancer (NSCLC) with MET exon 14 alterations with disease progression on or after platinum-based chemotherapy.

Reference:

https://www.pfizer.com/news/press-release/press-release-detail/pfizer_s_xalkori_crizotinib_receives_fda_breakthrough_therapy_designation_in_two_new_indications-0

Signatures

Testing Personnel:

Laboratory Supervisor:

Pathologist:

References

1. Peschard et al. A conserved DpYR motif in the juxtamembrane domain of the Met receptor family forms an atypical c-Cbl/Cbl-b tyrosine kinase binding domain binding site required for suppression of oncogenic activation. *J. Biol. Chem.* 2004 Jul 9;279(28):29565-71. PMID: 15123609
2. Peschard et al. Mutation of the c-Cbl TKB domain binding site on the Met receptor tyrosine kinase converts it into a transforming protein. *Mol. Cell.* 2001 Nov;8(5):995-1004. PMID: 11741535
3. Abella et al. Met/Hepatocyte growth factor receptor ubiquitination suppresses transformation and is required for Hrs phosphorylation. *Mol. Cell. Biol.* 2005 Nov;25(21):9632-45. PMID: 16227611
4. Sierra et al. c-MET as a potential therapeutic target and biomarker in cancer. *Ther Adv Med Oncol.* 2011 Nov;3(1 Suppl):S21-35. PMID: 22128285
5. Mo et al. Targeting MET in cancer therapy. *Chronic Dis Transl Med.* 2017 Sep;3(3):148-153. PMID: 29063069
6. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
7. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
8. Frampton et al. Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov.* 2015 Aug;5(8):850-9. PMID: 25971938
9. Schrock et al. Characterization of 298 Patients with Lung Cancer Harboring MET Exon 14 Skipping Alterations. *J Thorac Oncol.* 2016 Sep;11(9):1493-502. PMID: 27343443
10. Pilotto et al. MET exon 14 juxtamembrane splicing mutations: clinical and therapeutical perspectives for cancer therapy. *Ann Transl Med.* 2017 Jan;5(1):2. doi: 10.21037/atm.2016.12.33. PMID: 28164087
11. Reungwetwattana et al. The race to target MET exon 14 skipping alterations in non-small cell lung cancer: The Why, the How, the Who, the Unknown, and the Inevitable. *Lung Cancer.* 2017 Jan;103:27-37. PMID: 28024693
12. Saffroy et al. MET exon 14 mutations as targets in routine molecular analysis of primary sarcomatoid carcinoma of the lung. *Oncotarget.* 2017 Jun 27;8(26):42428-42437. PMID: 28418914
13. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
14. Yeh et al. Activating MET kinase rearrangements in melanoma and Spitz tumours. *Nat Commun.* 2015 May 27;6:7174. doi: 10.1038/ncomms8174. PMID: 26013381
15. Bao et al. RNA-seq of 272 gliomas revealed a novel, recurrent PTPRZ1-MET fusion transcript in secondary glioblastomas. *Genome Res.* 2014 Nov;24(11):1765-73. PMID: 25135958
16. Sebastian et al. Recurrent MET fusion genes represent a drug target in pediatric glioblastoma. *Nat. Med.* 2016 Nov;22(11):1314-1320. PMID: 27748748
17. Zeng et al. c-Met gene amplification is associated with advanced stage colorectal cancer and liver metastases. *Cancer Lett.* 2008 Jul 8;265(2):258-69. PMID: 18395971
18. Tsugawa et al. Amplification of the c-met, c-erbB-2 and epidermal growth factor receptor gene in human gastric cancers: correlation to clinical features. *Oncology.* 1998 Sep-Oct;55(5):475-81. PMID: 9732228
19. Di et al. Overexpression and amplification of the met/HGF receptor gene during the progression of colorectal cancer. *Clin. Cancer Res.* 1995 Feb;1(2):147-54. PMID: 9815967
20. https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/213591s000lbl.pdf
21. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 5.2021]
22. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214096s000lbl.pdf
23. https://www.pfizer.com/news/press-release/press-release-detail/pfizer_s_xalkori_crizotinib_receives_fda_breakthrough_therapy_designation_in_two_new_indications-0
24. Bean et al. MET amplification occurs with or without T790M mutations in EGFR mutant lung tumors with acquired resistance to gefitinib or erlotinib. *Proc. Natl. Acad. Sci. U.S.A.* 2007 Dec 26;104(52):20932-7. PMID: 18093943
25. Chen et al. Clinicopathologic and molecular features of epidermal growth factor receptor T790M mutation and c-MET amplification in tyrosine kinase inhibitor-resistant Chinese non-small cell lung cancer. *Pathol Oncol Res.* 2009 Dec;15(4):651-8. doi: 10.1007/s12253-009-9167-8. Epub 2009 Apr 21. PMID: 19381876
26. Suda et al. Reciprocal and complementary role of MET amplification and EGFR T790M mutation in acquired resistance to kinase inhibitors in lung cancer. *Clin. Cancer Res.* 2010 Nov 15;16(22):5489-98. PMID: 21062933

References (continued)

27. Zhang et al. Current mechanism of acquired resistance to epidermal growth factor receptor-tyrosine kinase inhibitors and updated therapy strategies in human nonsmall cell lung cancer. *J Cancer Res Ther.* 2016 Dec;12(Supplement):C131-C137. PMID: 28230005
28. Nguyen et al. Acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancers dependent on the epidermal growth factor receptor pathway. *Clin Lung Cancer.* 2009 Jul;10(4):281-9. PMID: 19632948
29. Choueiri et al. Biomarker-Based Phase II Trial of Savolitinib in Patients With Advanced Papillary Renal Cell Cancer. *J. Clin. Oncol.* 2017 Sep 10;35(26):2993-3001. PMID: 28644771