Taipei Veterans General Hospital

Department of Pathology and Laboratory Medicine
No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan
11217, R.O.C.
Tel: 02-2875-7449

Sample Information

Patient Name: 鄧淦
Gender: Male
ID No.: A103621986
History No.: 26488540
Age: 77

Ordering Doctor: DOC2591E 許文虎
Ordering REQ.: 0BRDMBJ
Signing in Date: 2022/01/19

Path No.: S111-98208
MP No.: F22010
Assay: Oncomine Focus Assay
Sample Type: FFPE
Block No.: S111-01370E
Percentage of tumor cells: 80%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents

	Page
Variant Details	2
Biomarker Descriptions	2
Relevant Therapy Summary	3
Relevant Therapy Details	3
Clinical Trials Summary	7

Report Highlights

1 Relevant Biomarkers
1 Therapies Available
1 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).The content of this report has not been evaluated or approved by the FDA or other regulatory agencies.

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	*PIK3CA p.(H1047R) c.3140A>G* phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha Allele Frequency: 57.20% **Prognostic significance:** None **Diagnostic significance:** None	None	**alpelisib + hormone therapy** [1, 2]	1

Public data sources included in relevant therapies: FDA[1], NCCN, **EMA**[2], ESMO
Public data sources included in prognostic and diagnostic significance: NCCN, ESMO
Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
PIK3CA	p.(H1047R)	c.3140A>G	COSM775	chr3:178952085	57.20%	NM_006218.4	missense	2000
JAK1	p.(P733=)	c.2199A>G	.	chr1:65310489	79.03%	NM_002227.4	synonymous	1998
ALK	p.(A1200=)	c.3600G>C	.	chr2:29443617	41.50%	NM_004304.5	synonymous	1971
PDGFRA	p.(G313=)	c.939T>G	.	chr4:55133726	76.98%	NM_006206.6	synonymous	1998
PDGFRA	p.(V824=)	c.2472C>T	.	chr4:55152040	74.89%	NM_006206.6	synonymous	1999
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.30%	NM_213647.3	missense	2000

Biomarker Descriptions

PIK3CA (phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha)

Background: The PIK3CA gene encodes the phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha of the class I phosphatidylinositol 3-kinase (PI3K) enzyme[1]. PI3K is a heterodimer that contains a p85 regulatory subunit, which couples one of four p110 catalytic subunits to activated tyrosine protein kinases[2,3]. The p110 catalytic subunits include p110α, β, δ, γ and are encoded by genes PIK3CA, PIK3CB, PIK3CD, and PIK3CG, respectively[2]. PI3K catalyzes the conversion of phosphatidylinositol (4,5)-bisphosphate (PI(4,5)P2) into phosphatidylinositol (3,4,5)-trisphosphate (PI(3,4,5)P3) while the phosphatase and tensin homolog (PTEN) catalyzes the reverse reaction[4,5]. The reversible phosphorylation of inositol lipids regulates diverse aspects of cell growth and metabolism[4,5,6,7]. Recurrent somatic alterations in PIK3CA are frequent in cancer and result in the activation of PI3K/AKT/MTOR pathway, which can influence several hallmarks of cancer including cell proliferation, apoptosis, cancer cell metabolism and invasion, and genetic instability[8,9,10].

Alterations and prevalence: Recurrent somatic activating mutations in PIK3CA are common in diverse cancers and are observed in 20-30% of breast, cervical, and uterine cancers and 10-20% of bladder, gastric, head and neck, and colorectal cancers[11,12]. Activating mutations in PIK3CA commonly cluster in two regions corresponding to the exon 9 helical (codons E542/E545) and exon 20 kinase (codon H1047) domains, each having distinct mechanisms of activation[13,14,15]. PIK3CA resides in the 3q26 cytoband, a region frequently amplified (10-30%) in diverse cancers including squamous carcinomas of the lung, cervix, head and neck, and esophagus, and in serous ovarian and uterine cancers[11,12].

Potential relevance: The PI3K inhibitor, alpelisib[16], is FDA approved (2019) in combination with fulvestrant for the treatment of patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, advanced or metastatic breast cancer. Additionally, a phase Ib study of alpelisib with letrozole in patients with metastatic estrogen receptor (ER)-positive breast cancer, the clinical benefit rate, defined as lack of disease progression ≥ 6 months, was 44% (7/16) in PIK3CA-mutated tumors and 20% (2/20) in PIK3CA wild-type tumors[17]. Specifically, exon 20 H1047R mutations were associated with more durable clinical responses in comparison to exon 9 E545K mutations[17]. However, alpelisib did not improve response when administered

Biomarker Descriptions (continued)

with letrozole in patients with ER+ early breast cancer with PIK3CA mutations[18]. Case studies with MTOR inhibitors sirolimus and temsirolimus report isolated cases of clinical response in PIK3CA mutated refractory cancers[19,20].

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

PIK3CA p.(H1047R) c.3140A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
alpelisib + fulvestrant	○	○	○	○	✖
inavolisib	✖	✖	✖	✖	● (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

FDA information is current as of 2021-11-17. For the most up-to-date information, search www.fda.gov.

PIK3CA p.(H1047R) c.3140A>G

○ alpelisib + fulvestrant

Cancer type: Breast Cancer **Label as of:** 2021-07-20 **Variant class:** PIK3CA H1047R mutation

Other criteria: ERBB2 negative, Hormone receptor positive

Indications and usage:
PIQRAY® is a kinase inhibitor indicated in combination with fulvestrant for the treatment of postmenopausal women, and men, with hormone receptor (HR)- positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated, advanced or metastatic breast cancer as detected by an FDA-approved test following progression on or after an endocrine-based regimen.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212526Orig1s004lbl.pdf

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

Current NCCN Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

NCCN information is current as of 2021-11-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

PIK3CA p.(H1047R) c.3140A>G

○ alpelisib + fulvestrant

Cancer type: Breast Cancer **Variant class:** PIK3CA activating mutation

Other criteria: ERBB2 negative, Hormone receptor positive

NCCN Recommendation category: 1

Population segment (Line of therapy):
- Stage IV; Invasive, Recurrent, Unresectable, Local, Regional (Second-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Breast Cancer [Version 8.2021]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

Current EMA Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

EMA information is current as of 2021-11-17. For the most up-to-date information, search www.ema.europa.eu/ema.

PIK3CA p.(H1047R) c.3140A>G

○ alpelisib + fulvestrant

Cancer type: Breast Cancer **Label as of:** 2021-10-11 **Variant class:** PIK3CA H1047R mutation

Other criteria: ERBB2 negative, Hormone receptor positive

Reference:

https://www.ema.europa.eu/en/documents/product-information/piqray-epar-product-information_en.pdf

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

Current ESMO Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

ESMO information is current as of 2021-11-01. For the most up-to-date information, search www.esmo.org.

PIK3CA p.(H1047R) c.3140A>G

○ alpelisib + fulvestrant

Cancer type: Breast Cancer

Variant class: PIK3CA exon 20 mutation

Other criteria: ERBB2 negative, ER positive

ESMO Level of Evidence/Grade of Recommendation: I / B

Population segment (Line of therapy):

- Luminal A; Advanced, Metastatic (Second-line therapy); ESMO-MCBS v1.1 score: 2

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Breast Cancer [Ann Oncol (2021); DOI:https://doi.org/10.1016/j.annonc.2021.09.019]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

Clinical Trials in Taiwan region:

Clinical Trials Summary

PIK3CA p.(H1047R) c.3140A>G

NCT ID	Title	Phase
NCT04589845	Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial	II

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

Signatures

Testing Personnel:

Shu-Ying Wang 2022.01.20 11:45:12 +08'00'

Laboratory Supervisor:

2022.01.20 11:45:35 +08'00'

Pathologist:

Yi-Che Yeh 2022.01.20 11:45:54 +08'00'

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).

References

1. Volinia et al. Molecular cloning, cDNA sequence, and chromosomal localization of the human phosphatidylinositol 3-kinase p110 alpha (PIK3CA) gene. Genomics. 1994 Dec;24(3):472-7. PMID: 7713498
2. Whale et al. Functional characterization of a novel somatic oncogenic mutation of PIK3CB. Signal Transduct Target Ther. 2017;2:17063. PMID: 29279775
3. Osaki et al. PI3K-Akt pathway: its functions and alterations in human cancer. Apoptosis. 2004 Nov;9(6):667-76. PMID: 15505410
4. Cantley. The phosphoinositide 3-kinase pathway. Science. 2002 May 31;296(5573):1655-7. PMID: 12040186
5. Fruman et al. The PI3K Pathway in Human Disease. Cell. 2017 Aug 10;170(4):605-635. PMID: 28802037
6. Engelman et al. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. Nat. Rev. Genet. 2006 Aug;7(8):606-19. PMID: 16847462
7. Vanhaesebroeck et al. PI3K signalling: the path to discovery and understanding. Nat. Rev. Mol. Cell Biol. 2012 Feb 23;13(3):195-203. PMID: 22358332
8. Yuan et al. PI3K pathway alterations in cancer: variations on a theme. Oncogene. 2008 Sep 18;27(41):5497-510. PMID: 18794884
9. Liu et al. Targeting the phosphoinositide 3-kinase pathway in cancer. Nat Rev Drug Discov. 2009 Aug;8(8):627-44. PMID: 19644473
10. Hanahan et al. Hallmarks of cancer: the next generation. Cell. 2011 Mar 4;144(5):646-74. PMID: 21376230
11. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4. PMID: 22588877
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
13. Miled et al. Mechanism of two classes of cancer mutations in the phosphoinositide 3-kinase catalytic subunit. Science. 2007 Jul 13;317(5835):239-42. PMID: 17626883
14. Burke et al. Synergy in activating class I PI3Ks. Trends Biochem. Sci. 2015 Feb;40(2):88-100. PMID: 25573003
15. Burke et al. Oncogenic mutations mimic and enhance dynamic events in the natural activation of phosphoinositide 3-kinase p110α (PIK3CA). Proc. Natl. Acad. Sci. U.S.A. 2012 Sep 18;109(38):15259-64. PMID: 22949682
16. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/212526Orig1s004lbl.pdf
17. Mayer et al. A Phase Ib Study of Alpelisib (BYL719), a PI3Kα-Specific Inhibitor, with Letrozole in ER+/HER2- Metastatic Breast Cancer. Clin. Cancer Res. 2017 Jan 1;23(1):26-34. PMID: 27126994
18. Mayer et al. A Phase II Randomized Study of Neoadjuvant Letrozole Plus Alpelisib for Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Breast Cancer (NEO-ORB). Clin. Cancer Res. 2019 Feb 5. PMID: 30723140
19. Jung et al. Pilot study of sirolimus in patients with PIK3CA mutant/amplified refractory solid cancer. Mol Clin Oncol. 2017 Jul;7(1):27-31. PMID: 28685070
20. Janku et al. PIK3CA mutations in patients with advanced cancers treated with PI3K/AKT/mTOR axis inhibitors. Mol. Cancer Ther. 2011 Mar;10(3):558-65. PMID: 21216929

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2021.12(004).