Taipei Veterans General Hospital

Department of Pathology and Laboratory Medicine
No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan
11217, R.O.C.
Tel: 02-2875-7449

Sample Information

Patient Name: 柯芳庭
Gender: Male
ID No.: C120530922
History No.: 47492546
Age: 58

Ordering Doctor: DOC6278K 陳逸安
Ordering REQ.: 0BSYWUX
Signing in Date: 2022/03/18

Path No.: S111-98718
MP No.: F22020
Assay: Oncomine Focus Assay
Sample Type: FFPE
Block No.: S110-23675G
Percentage of tumor cells: 60%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents

	Page
Variant Details	2
Biomarker Descriptions	3
Relevant Therapy Summary	3
Relevant Therapy Details	4
Clinical Trials Summary	7
Alert Details	8

Report Highlights

2 Relevant Biomarkers
2 Therapies Available
3 Clinical Trials

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	***KRAS p.(G12C) c.34G>T, KRAS amplification***	ROS1	None detected
MET	None detected		

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).The content of this report has not been evaluated or approved by the FDA or other regulatory agencies.

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	*KRAS p.(G12C) c.34G>T* KRAS proto-oncogene, GTPase Allele Frequency: 70.31% **Prognostic significance:** None **Diagnostic significance:** None	**sotorasib** [1]	cabozantinib	3
IIC	*KRAS amplification* KRAS proto-oncogene, GTPase **Prognostic significance:** None **Diagnostic significance:** None	None	None	1

Public data sources included in relevant therapies: FDA1, NCCN, **EMA**2, ESMO
Public data sources included in prognostic and diagnostic significance: NCCN, ESMO
Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
KRAS	p.(G12C)	c.34G>T	COSM516	chr12:25398285	70.31%	NM_033360.4	missense	1994
JAK1	p.(P733=)	c.2199A>G	.	chr1:65310489	51.55%	NM_002227.4	synonymous	1998
ALK	p.(D1529E)	c.4587C>G	.	chr2:29416366	49.45%	NM_004304.5	missense	1998
ALK	p.(E1158Q)	c.3472G>C	.	chr2:29445253	4.40%	NM_004304.5	missense	2000
ALK	p.(G1125=)	c.3375C>A	.	chr2:29445458	48.35%	NM_004304.5	synonymous	1998
PDGFRA	p.(G313=)	c.939T>G	.	chr4:55133726	48.92%	NM_006206.6	synonymous	1997
KIT	p.(M541L)	c.1621A>C	.	chr4:55593464	48.27%	NM_000222.3	missense	1999
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.35%	NM_213647.3	missense	2000
MET	p.(N375S)	c.1124A>G	.	chr7:116340262	49.82%	NM_001127500.3	missense	1999
RET	p.(L769=)	c.2307G>T	.	chr10:43613843	100.00%	NM_020975.6	synonymous	1999
RET	p.(S904=)	c.2712C>G	.	chr10:43615633	46.72%	NM_020975.6	synonymous	1999
HRAS	p.(G12=)	c.36C>T	.	chr11:534287	46.40%	NM_001130442.2	synonymous	2000

Copy Number Variations

Gene	Locus	Copy Number
KRAS	chr12:25364761	7.23

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Biomarker Descriptions

KRAS (KRAS proto-oncogene, GTPase)

Background: The KRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes NRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival[1,2,3].

Alterations and prevalence: Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. KRAS mutations are observed in up to 10-20% of uterine cancer, 30-35% of lung adenocarcinoma and colorectal cancer, and about 60% of pancreatic cancer[4]. The majority of KRAS mutations consist of point mutations occurring at G12, G13, and Q61[4,5,6]. Mutations at A59, K117, and A146 have also been observed but are less frequent[7,8].

Potential relevance: The KRAS inhibitor, sotorasib[9], is approved (2021) for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC). The FDA has granted breakthrough therapy designation (2021) to the small molecule inhibitor, adagrasib, for KRAS G12C positive in non-small cell lung cancer following prior systemic therapy[10].The small molecular inhibitor, RO-5126766, was also granted breakthrough designation (2021) alone for KRAS G12V mutant non-small cell lung cancer or in combination with defactinib, for KRAS mutant endometrial carcinoma and KRAS G12V mutant non-small cell lung cancer[11]. Additionally, onvansertib[12] was granted fast track designation (2020) for second-line treatment of patients with KRAS-mutated metastatic colorectal cancer (mCRC). The EGFR antagonists, cetuximab[13] and panitumumab[14], are contraindicated for treatment of colorectal cancer patients with KRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)[8]. Additionally, KRAS mutations are associated with poor prognosis in NSCLC[15].

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

KRAS p.(G12C) c.34G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
sotorasib	●	●	✖	✖	✖
cabozantinib	✖	✖	✖	○	✖
D-1553	✖	✖	✖	✖	● (I/II)
RMC-4630, pembrolizumab	✖	✖	✖	✖	● (I/II)
mRNA-5671, pembrolizumab	✖	✖	✖	✖	● (I)

KRAS amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
RMC-4630, pembrolizumab	✖	✖	✖	✖	● (I/II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Relevant Therapy Details

Current FDA Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

FDA information is current as of 2022-01-19. For the most up-to-date information, search www.fda.gov.

KRAS p.(G12C) c.34G>T

● sotorasib

Cancer type: Non-Small Cell Lung Cancer **Label as of:** 2021-05-28 **Variant class:** KRAS G12C mutation

Indications and usage:
LUMAKRAS™ is an inhibitor of the RAS GTPase family indicated for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC), as determined by an FDA-approved test, who have received at least one prior systemic therapy. This indication is approved under accelerated approval based on overall response rate (ORR) and duration of response (DOR). Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214665s000lbl.pdf

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Current NCCN Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

NCCN information is current as of 2022-01-04. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

KRAS p.(G12C) c.34G>T

● sotorasib

Cancer type: Non-Small Cell Lung Cancer **Variant class:** KRAS G12C mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- Adenocarcinoma, Large Cell, Squamous Cell, Not otherwise specified (NOS); Advanced, Metastatic, Progression (Subsequent therapy)

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 1.2022]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Current ESMO Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

ESMO information is current as of 2022-01-04. For the most up-to-date information, search www.esmo.org.

KRAS p.(G12C) c.34G>T

○ cabozantinib

Cancer type: Thyroid Gland Medullary Carcinoma **Variant class:** RAS mutation

ESMO Level of Evidence/Grade of Recommendation: II / C

Population segment (Line of therapy):
- Metastatic (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Thyroid Cancer [Annals of Oncology 30: 1856–1883, 2019 doi:10.1093/annonc/mdz400]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Clinical Trials in Taiwan region:

Clinical Trials Summary

KRAS p.(G12C) c.34G>T

NCT ID	Title	Phase
NCT03948763	A Phase I, Open-Label, Multicenter Study to Assess the Safety and Tolerability of mRNA-5671/V941 as a Monotherapy and in Combination With Pembrolizumab in Participants With KRAS Mutant Advanced or Metastatic Non-Small Cell Lung Cancer, Colorectal Cancer or Pancreatic Adenocarcinoma	I
NCT04418661	A Phase 1/2, Open-label, Multicenter, Dose Escalation and Dose Expansion Study of SAR442720 in Combination With Pembrolizumab in Patients With Advanced Malignancies	I/II
NCT04585035	A Phase I/II, Open Label Study to Evaluate the Safety, Tolerability, Pharmacokinetics and Efficacy of D-1553 in Subjects With Advanced or Metastatic Solid Tumors With KRasG12C Mutation	I/II

KRAS amplification

NCT ID	Title	Phase
NCT04418661	A Phase 1/2, Open-label, Multicenter, Dose Escalation and Dose Expansion Study of SAR442720 in Combination With Pembrolizumab in Patients With Advanced Malignancies	I/II

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Alerts Informed By Public Data Sources

Current FDA Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

FDA information is current as of 2022-01-19. For the most up-to-date information, search www.fda.gov.

KRAS p.(G12C) c.34G>T

cetuximab

Cancer type: Colorectal Cancer **Label as of:** 2021-09-24 **Variant class:** KRAS G12 mutation

Indications and usage:

Erbitux® is an epidermal growth factor receptor (EGFR) antagonist indicated for treatment of:

Head and Neck Cancer

- Locally or regionally advanced squamous cell carcinoma of the head and neck in combination with radiation therapy.
- Recurrent locoregional disease or metastatic squamous cell carcinoma of the head and neck in combination with platinum-based therapy with fluorouracil.
- Recurrent or metastatic squamous cell carcinoma of the head and neck progressing after platinum-based therapy.

Colorectal Cancer

K-Ras wild-type, EGFR-expressing, metastatic colorectal cancer as determined by FDA-approved test

- in combination with FOLFIRI for first-line treatment,
- in combination with irinotecan in patients who are refractory to irinotecan-based chemotherapy,
- as a single agent in patients who have failed oxaliplatin- and irinotecan-based chemotherapy or who are intolerant to irinotecan.

Limitations of Use: Erbitux® is not indicated for treatment of Ras-mutant colorectal cancer or when the results of the Ras mutation tests are unknown.

BRAF V600E Mutation-Positive Metastatic Colorectal Cancer (CRC)

- in combination with encorafenib, for the treatment of adult patients with metastatic colorectal cancer (CRC) with a BRAF V600E mutation, as detected by an FDA-approved test, after prior therapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf

panitumumab

Cancer type: Colorectal Cancer **Label as of:** 2021-08-25 **Variant class:** KRAS G12 mutation

Indications and usage:

VECTIBIX® is an epidermal growth factor receptor (EGFR) antagonist indicated for the treatment of wild-type RAS (defined as wild-type in both KRAS and NRAS as determined by an FDA-approved test for this use) metastatic colorectal cancer (mCRC):

- In combination with FOLFOX for first-line treatment.
- As monotherapy following disease progression after prior treatment with fluoropyrimidine, oxaliplatin, and irinotecan-containing chemotherapy.
- Limitation of Use: VECTIBIX® is not indicated for the treatment of patients with RAS-mutant mCRC or for whom RAS mutation status is unknown.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

KRAS p.(G12C) c.34G>T (continued)

adagrasib

Cancer type: Non-Small Cell Lung Cancer **Variant class:** KRAS G12C mutation

Supporting Statement:
The FDA has granted Breakthrough Designation to the small molecule inhibitor, adagrasib, for KRAS G12C positive in non-small cell lung cancer following prior systemic therapy.

Reference:

https://ir.mirati.com/press-releases/press-release-details/2021/Mirati-Therapeutics-Adagrasib-Receives-Breakthrough-Therapy-Designation-from-U.S.-Food-and-Drug-Administration-for-Patients-with-Advanced-Non-Small-Cell-Lung-Cancer-Harboring-the-KRAS-G12C-Mutation/default.aspx

---

defactinib + RO-5126766

Cancer type: Endometrial Carcinoma **Variant class:** KRAS mutation

Supporting Statement:
The FDA has granted Breakthrough Designation to the small molecule inhibitor, RO-5126766 alone for KRAS G12V mutant non-small cell lung cancer or in combination with defactinib, for KRAS mutant endometrial carcinoma and KRAS G12V mutant non-small cell lung cancer.

Reference:

https://investor.verastem.com//news-releases/news-release-details/verastem-oncology-receives-breakthrough-therapy-designation-vs

---

bevacizumab + onvansertib + FOLFIRI

Cancer type: Colorectal Cancer **Variant class:** KRAS mutation

Supporting Statement:
The FDA has granted Fast Track Designation to the Polo-like Kinase 1 (PLK1) inhibitor, onvansertib, in combination with FOLFIRI and bevacizumab, for KRAS mutations in metastatic colorectal cancer in the second line.

Reference:

https://cardiffoncology.investorroom.com/2020-05-28-Cardiff-Oncology-Announces-Fast-Track-Designation-Granted-by-the-FDA-to-Onvansertib-for-Second-Line-Treatment-of-KRAS-Mutated-Colorectal-Cancer

---

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Current NCCN Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

NCCN information is current as of 2022-01-04. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/international_adaptations.aspx.

KRAS p.(G12C) c.34G>T

cetuximab

Cancer type: Colon Cancer **Variant class:** KRAS exon 2 mutation

Summary:
NCCN Guidelines® include the following supporting statement(s):
- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab."

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2021]

cetuximab

Cancer type: Rectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
NCCN Guidelines® include the following supporting statement(s):
- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab."

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2021]

panitumumab

Cancer type: Colon Cancer **Variant class:** KRAS exon 2 mutation

Summary:
NCCN Guidelines® include the following supporting statement(s):
- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab."

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2021]

panitumumab

Cancer type: Rectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
NCCN Guidelines® include the following supporting statement(s):
- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab."

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2021]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Current EMA Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

EMA information is current as of 2022-01-19. For the most up-to-date information, search www.ema.europa.eu/ema.

KRAS p.(G12C) c.34G>T

cetuximab, cetuximab + oxaliplatin

Cancer type: Colorectal Cancer **Label as of:** 2020-01-30 **Variant class:** KRAS exon 2 mutation

Reference:

https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information_en.pdf

panitumumab + oxaliplatin

Cancer type: Colorectal Cancer **Label as of:** 2021-12-07 **Variant class:** KRAS exon 2 mutation

Reference:

https://www.ema.europa.eu/en/documents/product-information/vectibix-epar-product-information_en.pdf

Current ESMO Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

ESMO information is current as of 2022-01-04. For the most up-to-date information, search www.esmo.org.

KRAS p.(G12C) c.34G>T

cetuximab

Cancer type: Colorectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "It has been demonstrated that the (potential) benefit of anti-EGFR antibodies in all treatment lines and either as a single agent or in combination with any chemotherapy regimen is limited to patients in whom a RAS mutation is excluded. It was shown that the 'expanded RAS' analysis (also including the detection of mutations in exons 3 and 4 of the KRAS gene as well as mutations in the NRAS [exons 2-4] gene) is superior to the KRAS (exon 2) analysis in predicting both more efficacy in the expanded RAS wild-type (WT) patients and a potential detrimental effect in patients harbouring any RAS mutation in their tumour genome [II/A]."

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2014) 25 (suppl 3): iii1-iii9. (eUpdate: 20 September 2016; Corrigendum: 21 July 2015)]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

KRAS p.(G12C) c.34G>T (continued)

cetuximab + chemotherapy

Cancer type: Colorectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "It has been demonstrated that the (potential) benefit of anti-EGFR antibodies in all treatment lines and either as a single agent or in combination with any chemotherapy regimen is limited to patients in whom a RAS mutation is excluded. It was shown that the 'expanded RAS' analysis (also including the detection of mutations in exons 3 and 4 of the KRAS gene as well as mutations in the NRAS [exons 2-4] gene) is superior to the KRAS (exon 2) analysis in predicting both more efficacy in the expanded RAS wild-type (WT) patients and a potential detrimental effect in patients harbouring any RAS mutation in their tumour genome [II/A]."
- "Thus, the activity of the anti-EGFR antibodies is confined to RAS WT tumours (and not only KRAS WT tumours). This is true for the combinations of cetuximab or panitumumab alone or with irinotecan- and oxaliplatin-based regimens. Treatment with anti-EGFR antibodies may even harm patients with a RAS mutation, especially when combined with oxaliplatin [I/A]."

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2014) 25 (suppl 3): iii1-iii9. (eUpdate: 20 September 2016; Corrigendum: 21 July 2015)]

panitumumab

Cancer type: Colorectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "It has been demonstrated that the (potential) benefit of anti-EGFR antibodies in all treatment lines and either as a single agent or in combination with any chemotherapy regimen is limited to patients in whom a RAS mutation is excluded. It was shown that the 'expanded RAS' analysis (also including the detection of mutations in exons 3 and 4 of the KRAS gene as well as mutations in the NRAS [exons 2-4] gene) is superior to the KRAS (exon 2) analysis in predicting both more efficacy in the expanded RAS wild-type (WT) patients and a potential detrimental effect in patients harbouring any RAS mutation in their tumour genome [II/A]."

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2014) 25 (suppl 3): iii1-iii9. (eUpdate: 20 September 2016; Corrigendum: 21 July 2015)]

panitumumab + chemotherapy

Cancer type: Colorectal Cancer **Variant class:** KRAS exon 2 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "It has been demonstrated that the (potential) benefit of anti-EGFR antibodies in all treatment lines and either as a single agent or in combination with any chemotherapy regimen is limited to patients in whom a RAS mutation is excluded. It was shown that the 'expanded RAS' analysis (also including the detection of mutations in exons 3 and 4 of the KRAS gene as well as mutations in the NRAS [exons 2-4] gene) is superior to the KRAS (exon 2) analysis in predicting both more efficacy in the expanded RAS wild-type (WT) patients and a potential detrimental effect in patients harbouring any RAS mutation in their tumour genome [II/A]."
- "Thus, the activity of the anti-EGFR antibodies is confined to RAS WT tumours (and not only KRAS WT tumours). This is true for the combinations of cetuximab or panitumumab alone or with irinotecan- and oxaliplatin-based regimens. Treatment with anti-EGFR antibodies may even harm patients with a RAS mutation, especially when combined with oxaliplatin [I/A]."

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2014) 25 (suppl 3): iii1-iii9. (eUpdate: 20 September 2016; Corrigendum: 21 July 2015)]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

Signatures

Testing Personnel:

Shu-Ying Wang 2022.03.21 11:49:38 +08'00'

Laboratory Supervisor:

2022.03.21 11:50:17 +08'00'

Pathologist:

Yi-Chu Yeh 2022.03.21 11:50:29 +08'00'

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).

References

1. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. Nat. Rev. Cancer. 2011 Oct 13;11(11):761-74. PMID: 21993244
2. Karnoub et al. Ras oncogenes: split personalities. Nat. Rev. Mol. Cell Biol. 2008 Jul;9(7):517-31. PMID: 18568040
3. Scott et al. Therapeutic Approaches to RAS Mutation. Cancer J. 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.0000000000000187. PMID: 27341593
4. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
5. Román et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. Mol Cancer. 2018 Feb 19;17(1):33. doi: 10.1186/s12943-018-0789-x. PMID: 29455666
6. Dinu et al. Prognostic significance of KRAS gene mutations in colorectal cancer--preliminary study. J Med Life. 2014 Oct-Dec;7(4):581-7. PMID: 25713627
7. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4. PMID: 22588877
8. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. J. Clin. Oncol. 2016 Jan 10;34(2):179-85. PMID: 26438111
9. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/214665s000lbl.pdf
10. https://ir.mirati.com/press-releases/press-release-details/2021/Mirati-Therapeutics-Adagrasib-Receives-Breakthrough-Therapy-Designation-from-U.S.-Food-and-Drug-Administration-for-Patients-with-Advanced-Non-Small-Cell-Lung-Cancer-Harboring-the-KRAS-G12C-Mutation/default.aspx
11. https://investor.verastem.com//news-releases/news-release-details/verastem-oncology-receives-breakthrough-therapy-designation-vs
12. https://cardiffoncology.investorroom.com/2020-05-28-Cardiff-Oncology-Announces-Fast-Track-Designation-Granted-by-the-FDA-to-Onvansertib-for-Second-Line-Treatment-of-KRAS-Mutated-Colorectal-Cancer
13. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf
14. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf
15. Slebos et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. N. Engl. J. Med. 1990 Aug 30;323(9):561-5. PMID: 2199829

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2022.02(004).