

ACT Onco[®] + Report

| PATIENT | | |
|------------------------------|---------------------------|----------------------|
| Identifier: 梁怡秋 | | Patient ID: 28591200 |
| Date of Birth: Oct 26, 1967 | | Gender: Female |
| Diagnosis: Pancreatic cancer | | |
| ORDERING PHYSICIAN | | |
| Name: 姜乃榕醫師 | | Tel: 886-228712121 |
| Facility: 臺北榮總 | | |
| Address: 臺北市北投區石牌路二段 201 號 | | |
| SPECIMEN | | |
| Specimen ID: S11232826B | Collection site: Pancreas | Type: FFPE tissue |
| Date received: Aug 02, 2023 | Lab ID: AA-23-04980 | D/ID: NA |

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

| Genomic Alterations/Biomarkers | Probable Effects in Patient's Cancer Type | | Probable Sensitive in Other Cancer Types |
|--------------------------------|---|-----------|--|
| | Sensitive | Resistant | |
| Not detected | | | |

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

| Genomic Alterations/Biomarkers | Possibly Sensitive | Possibly Resistant |
|--------------------------------|--------------------|------------------------|
| KRAS G12V | - | Cetuximab, Panitumumab |

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACTOnco® + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

| Gene | Amino Acid Change | Allele Frequency |
|--------|-------------------|------------------|
| CDKN2A | Splice donor | 14.4% |
| KRAS | G12V | 9.5% |
| TP53 | R273C | 11.8% |

- Copy Number Alterations

| Chromosome | Gene | Variation | Copy Number |
|--|------|---------------|-------------|
| Chr8 | MYC | Amplification | 9 |
| Copy number loss cannot be determined because of low tumor purity (<30%) | | | |

- Fusions

| Fusion Gene & Exon | Transcript ID |
|--|---------------|
| No fusion gene detected in this sample | |

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

| Biomarker | Results |
|----------------------------------|----------------------|
| Tumor Mutational Burden (TMB) | Cannot be determined |
| Microsatellite Instability (MSI) | Cannot be determined |

Note:

- Variant(s) enlisted in the SNV table may currently exhibit no relevance to treatment response prediction. Please refer to INTERPRETATION for more biological information and/or potential clinical impacts of the variants.
- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on <30% tumor purity.
- TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is < 30%.
- This specimen did not meet acceptance criteria due to low tumor purity. The test was performed with client's consent.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS

TARGETED THERAPIES

| Genomic Alterations | Therapies | Effect |
|---------------------|------------------------|-----------|
| Level 3A | | |
| KRAS G12V | Cetuximab, Panitumumab | resistant |

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

| Level | Description |
|-------|--|
| 1 | FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication |
| 2 | Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication |
| 3A | Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type |
| 3B | Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required) |
| 4 | Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies |

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

| Genomic Alterations | Potential Clinical Effects |
|---------------------|----------------------------|
| Not detected | |

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

| Genomic Alterations | Therapies | Effect | Level of Evidence | Cancer Type |
|-----------------------------|-------------------------------------|-----------------------|-------------------|----------------|
| MYC Amplification | FAC, CMF, and P-FEC regimens | Sensitive | Clinical | Breast cancer |
| | Platinum-based regimens | Sensitive | Clinical | Ovarian cancer |
| TP53 R273C | Platinum- and taxane-based regimens | Less sensitive | Clinical | Ovarian cancer |

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

CDKN2A Splice donor

Biological Impact

The Cyclin-Dependent Kinase Inhibitor 2A (CDKN2A) gene encodes the p16 (p16INK4a) and p14 (ARF) proteins. p16INK4a binds to CDK4 and CDK6, inhibiting these CDKs from binding D-type cyclins and phosphorylating the retinoblastoma (RB) protein whereas p14 (ARF) blocks the oncogenic activity of MDM2 by inhibiting MDM2-induced degradation of p53^{[1][2][3]}. CDKN2A has been reported as a haploinsufficient tumor suppressor with one copy loss that may lead to weak protein expression and is insufficient to execute its original physiological functions^[4]. Loss of CDKN2A has been frequently found in human tumors that result in uncontrolled cell proliferation^{[5][6]}.

CDKN2A c.150+1G>T is a variant located at the splice donor region, which may result in the exon skipping.

Therapeutic and prognostic relevance

Intact p16-Cdk4-Rb axis is known to be associated with sensitivity to cyclin-dependent kinase inhibitors^{[7][8]}. Several case reports also revealed that patients with CDKN2A-deleted tumors respond to the CDK4/6-specific inhibitor treatments^{[9][10][11]}. However, there are clinical studies that demonstrated CDKN2A nuclear expression, CDKN2A/CDKN2B co-deletion, or CDKN2A inactivating mutation was not associated with clinical benefit from CDK4/6 inhibitors, such as palbociclib and ribociclib, in RB-positive patients^{[12][13][14]}. CDKN2A loss or mutation has been determined as an inclusion criterion for the trial evaluating CDK4/6 inhibitors efficacy in different types of solid tumors (NCT02693535, NCT02187783).

The phase II TAPUR trial demonstrated clinical benefits to palbociclib monotherapy in advanced NSCLC or head and neck cancer harboring a CDKN2A mutation or copy number loss. However, pancreatic and biliary cancer patients harboring a CDKN2A mutation or copy number loss did not demonstrate an objective response or stable disease when treated with palbociclib monotherapy for 16 weeks (DOI: 10.1200/JCO.2021.39.15_suppl.6043)^{[15][16]}.

Notably, the addition of several CDK4/6 inhibitors to hormone therapies, including palbociclib in combination with letrozole, ribociclib plus letrozole, and abemaciclib combines with fulvestrant, have been approved by the U.S. FDA for the treatment of ER+ and HER2- breast cancer^{[8][17][18]}.

In a Phase I trial, a KRAS wild-type squamous non-small cell lung cancer (NSCLC) patient with CDKN2A loss had a partial response when treated with CDK4/6 inhibitor abemaciclib^[10]. Administration of combined palbociclib and MEK inhibitor PD-0325901 yield promising progression-free survival among patients with KRAS mutant non-small cell lung cancer (NSCLC) (AACR 2017, Abstract CT046). Moreover, MEK inhibitor in combination with CDK4/6 inhibitor demonstrates significant anti-KRAS-mutant NSCLC activity and radiosensitizing effect in preclinical models^[19].

A retrospective analysis demonstrated that concurrent deletion of CDKN2A with EGFR mutation in patients with non-small cell lung cancer (NSCLC), predicts worse overall survival after EGFR-TKI treatment^[20].

KRAS G12V

Biological Impact

The V-Ki-Ras2 Kirsten Rat Sarcoma 2 Viral Oncogene Homolog (KRAS) gene encodes a small GTPase protein, a member of the RAS family of small GTPases, which catalyze the hydrolysis of GTP to GDP. RAS proteins cycle between an active (GTP-bound) and an inactive (GDP-bound) state, to activate the downstream oncogenic pathways, including the PI3K/AKT/mTOR and MAPK pathways^[21]. KRAS mutations occur primarily in three hotspots G12, G13 and Q61, and less frequently in codon A146^{[21][22]}. These are activating mutations that lead to constitutive activation and persistent stimulation of the downstream signaling pathways^{[23][24]}.

ACT Onco[®] + Report

Mutations in KRAS have been reported in a diverse spectrum of human malignancies, including pancreatic carcinomas (>80%)^{[21][25]}, colon carcinomas (40-50%)^{[26][27]}, and lung carcinomas (30-50%)^{[28][29]}, but are also present in biliary tract malignancies, endometrial cancer, cervical cancer, bladder cancer, liver cancer, myeloid leukemia and breast cancer^[22].

KRAS G12V is a hotspot mutation that has been shown to result in the increased activation of downstream signaling pathways^[30].

Therapeutic and prognostic relevance

Cetuximab and panitumumab are FDA-approved for treating RAS wild-type metastatic colorectal cancer. The NCCN for CRC recommends that patients with any known KRAS or NRAS mutation (exons 2, 3, and 4) should not be treated with either cetuximab or panitumumab.

KRAS mutation has been determined as an inclusion criterion for the trials evaluating MEK inhibitors efficacies in various types of solid tumors (NCT03704688, NCT02399943, NCT02285439, NCT03637491, NCT04214418).

KRAS mutations are associated with a lack of efficacy of EGFR TKIs^{[31][32][33]}. Some case reports suggest that MEK inhibitors may benefit patients with KRAS mutations, as shown in cervical and ovarian cancer cases (Am J Clin Exp Obstet Gynecol 2015;2(3):140-143)^{[34][35]}. However, a randomized Phase II study did not find trametinib to be superior to docetaxel in KRAS-mutant non-small cell lung cancer patients^[36]. MEK inhibitors as a monotherapy have limited response^[37].

Combining MEK and mTOR inhibitors is being evaluated as a potential strategy in RAS-mutant CRC^{[38][39]}. The combination of trametinib and palbociclib has resulted in objective responses in KRAS mutant models^[40].

Sorafenib has been shown to be beneficial in KRAS-mutant CRC/NSCLC, and KRAS-amplified melanoma^{[41][42][43]}. KRAS mutations in exon 2 (codon 12 or 13) and codon 61 have been associated with poor prognosis in CRC^[44].

Patients with KRAS or BRAF mutations in low-grade serous carcinoma of the ovary or peritoneum had better overall survival than those with wild-type genes^[45]. In ovarian serous borderline tumor, KRAS G12V mutation was linked to shorter survival time^[46].

In patients with metastatic colorectal cancer treated with bevacizumab, the shortest survival was observed in patients with tumors harboring G12V or G12A KRAS mutation, and the PFS and OS for patients with G12V/A KRAS mutation was 6.6 and 16.8 compared to 11.6 and 23.6 months for patients with tumors harboring other KRAS mutation type^[47]. In another retrospective study, Patients with KRAS G12V exhibited worse OS and higher recurrence incidences compared with the entire cohort (OS: 26 months vs 60 months; DFS: 15 months vs 24 months) in lung adenocarcinoma^[48].

TP53 R273C

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[49]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[50].

R273C is a hotspot mutation occurred at the DNA-binding domain (DBD) of the p53 protein^[51]. This is a gain-of-function mutation that has been shown to cause aberrant activation of gene expression, increased cell proliferation, migration and increase the HER2 promoter activity and mRNA expression in vitro^{[52][53][54]}.

ACT Onco[®] + Report

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[55].

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[56]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[57].

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[58][59][60]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[61]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[62][63]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[64].

TP53 oncomorphic mutations, including P151S, Y163C, R175H, L194R, Y220C, R248Q, R248W, R273C, R273H, R273L, and R282W have been shown to predict resistance to platinum- and taxane-based chemotherapy in advanced serous ovarian carcinoma patients^[65].

MYC Amplification

Biological Impact

The v-myc avian myelocytomatosis viral oncogene homolog, also known as c-myc (MYC) gene encodes a transcription factor involved in cellular proliferation, inhibiting exit from the cell cycle, stimulating vascularization and enhancing genomic instability^{[66][67][68]}. Dysregulated MYC expression is implicated in a wide range of human cancers^[69].

Therapeutic and prognostic relevance

MYC amplification was associated with better clinical outcome in breast cancer patients treated with FAC (5-fluorouracil, doxorubicin, and cyclophosphamide), CMF (cyclophosphamide, methotrexate and 5-fluorouracil) and P-FEC (paclitaxel followed by 5-fluorouracil, epirubicin and cyclophosphamide) and higher expression of MYC was also associated with a better response rate in platinum-treated ovarian cancer patients^{[70][71][72]}.

CDK inhibition using the dinaciclib, a CDK1, 2, 5 and 9 inhibitors, exerted antitumor activity in triple-negative breast cancer (TNBC) tumor xenograft and cell lines with increased activity of the MYC pathway^{[73][74]}.

Overexpression of MYC has been reported as a favorable prognostic biomarker in colorectal carcinoma (CRC)^{[75][76]}. However, the favorable prognostic value of MYC in CRC is abrogated by the TP53 mutation^[76].

MYC amplification with the loss of tumor suppressor pathways such as p53 and RB has been shown to be associated with poor outcomes and was correlated with shortened disease-free survival in breast cancer with BRCA1 deficiency in TNBC^{[73][77]}.

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Not Applicable.

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

| Gene | Amino Acid Change | Exon | cDNA Change | Accession Number | COSMIC ID | Allele Frequency | Coverage |
|--------|-------------------|------|-------------|------------------|-----------|------------------|----------|
| CDKN2A | Splice donor | - | c.150+1G>T | NM_000077 | - | 14.4% | 1349 |
| KRAS | G12V | 2 | c.35G>T | NM_004985 | COSM520 | 9.5% | 3141 |
| TP53 | R273C | 8 | c.817C>T | NM_000546 | COSM10659 | 11.8% | 1323 |

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

| Gene | Amino Acid Change | Exon | cDNA Change | Accession Number | COSMIC ID | Allele Frequency | Coverage |
|--------|-------------------|------|-------------|------------------|-------------|------------------|----------|
| ATR | N744D | 10 | c.2230A>G | NM_001184 | - | 11.2% | 1594 |
| BRCA2 | Q3036E | 23 | c.9106C>G | NM_000059 | COSM6925801 | 51.6% | 190 |
| CREBBP | V1924M | 31 | c.5770G>A | NM_004380 | - | 34.7% | 98 |
| EPHA2 | P63L | 3 | c.188C>T | NM_004431 | COSM1205442 | 50.5% | 3460 |
| GNAQ | D130N | 3 | c.388G>A | NM_002072 | - | 51.4% | 889 |
| KDR | T48A | 2 | c.142A>G | NM_002253 | - | 65.0% | 391 |
| MUC16 | V8883A | 3 | c.26648T>C | NM_024690 | - | 45.8% | 1077 |
| UBE2A | V141fs | 6 | c.420dup | NM_003336 | - | 10.0% | 2379 |



Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Jul 08, 2023
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11232826B
- Collection site: Pancreas
- Examined by: Dr. Yun-An Chen
- 1. The percentage of viable tumor cells in total cells in the whole slide (%): 10%
- 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 30%
- 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
- 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
- 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 1252x
- Target Base Coverage at 100x: 95%

RNA test

- Average unique RNA Start Sites per control GSP2: 142

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®] to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

ACT Onco[®] + Report

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師黃靖婷 博士
Ching-Ting Huang Ph.D.
檢字第 016511 號

CT Huang

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

ACT Onco[®] + Report

GENE LIST SNV & CNV

| | | | | | | | | | | | |
|----------|---------|---------|----------|----------|---------|-----------|-----------|----------|----------|----------|----------|
| ABCB1* | ABCC2* | ABCG2* | ABL1 | ABL2 | ADAMTS1 | ADAMTS13 | ADAMTS15 | ADAMTS16 | ADAMTS18 | ADAMTS6 | ADAMTS9 |
| ADAMTSL1 | ADGRA2 | ADH1C* | AKT1 | AKT2 | AKT3 | ALDH1A1* | ALK | AMER1 | APC | AR | ARAF |
| ARID1A | ARID1B | ARID2 | ASXL1 | ATM | ATR | ATRX | AURKA | AURKB | AXIN1 | AXIN2 | AXL |
| B2M | BAP1 | BARD1 | BCL10 | BCL2* | BCL2L1 | BCL2L2* | BCL6 | BCL9 | BCOR | BIRC2 | BIRC3 |
| BLM | BMP1R1A | BRAF | BRCA1 | BRCA2 | BRD4 | BRIP1 | BTG1 | BTG2* | BTB | BUB1B | CALR |
| CANX | CARD11 | CASP8 | CBFB | CBL | CCNA1 | CCNA | CCNB1 | CCNB2 | CCNB3 | CCND1 | CCND2 |
| CCND3 | CCNE1 | CCNE2 | CCNH | CD19 | CD274 | CD58 | CD70* | CD79A | CD79B | CDC73 | CDH1 |
| CDK1 | CDK12 | CDK2 | CDK4 | CDK5 | CDK6 | CDK7 | CDK8 | CDK9 | CDKN1A | CDKN1B | CDKN2A |
| CDKN2B | CDKN2C | CEBPA* | CHEK1 | CHEK2 | CIC | CREBBP | CRKL | CRLF2 | CSF1R | CTCF | CTLA4 |
| CTNNA1 | CTNNB1 | CUL3 | CYLD | CYP1A1* | CYP2B6* | CYP2C19* | CYP2C8* | CYP2D6 | CYP2E1* | CYP3A4* | CYP3A5* |
| DAXX | DCUN1D1 | DDR2 | DICER1 | DNMT3A | DOT1L | DPYD | DTX1 | E2F3 | EGFR | EP300 | EPCAM |
| EPHA2 | EPHA3 | EPHA5 | EPHA7 | EPHB1 | ERBB2 | ERBB3 | ERBB4 | ERCC1 | ERCC2 | ERCC3 | ERCC4 |
| ERCC5 | ERG | ESR1 | ESR2 | ETV1 | ETV4 | EZH2 | FAM46C | FANCA | FANCC | FANCD2 | FANCE |
| FANCF | FANCG | FANCL | FAS | FAT1 | FBXW7 | FCGR2B | FGF1* | FGF10 | FGF14 | FGF19* | FGF23 |
| FGF3 | FGF4* | FGF6 | FGFR1 | FGFR2 | FGFR3 | FGFR4 | FH | FLCN | FLT1 | FLT3 | FLT4 |
| FOXL2* | FOXP1 | FRG1 | FUBP1 | GATA1 | GATA2 | GATA3 | GNA11 | GNA13 | GNAQ | GNAS | GREM1 |
| GRIN2A | GSK3B | GSTP1* | GSTT1* | HGF | HIF1A | HIST1H1C* | HIST1H1E* | HNF1A | HR | HRAS* | HSP90AA1 |
| HSP90AB1 | HSPA4 | HSPA5 | IDH1 | IDH2 | IFNL3* | IGF1 | IGF1R | IGF2 | IKBKB | IKBKE | IKZF1 |
| IL6 | IL7R | INPP4B | INSR | IRF4 | IRS1 | IRS2* | JAK1 | JAK2 | JAK3 | JUN* | KAT6A |
| KDM5A | KDM5C | KDM6A | KDR | KEAP1 | KIT | KMT2A | KMT2C | KMT2D | KRAS | LCK | LIG1 |
| LIG3 | LMO1 | LRP1B | LYN | MALT1 | MAP2K1 | MAP2K2 | MAP2K4 | MAP3K1 | MAP3K7 | MAPK1 | MAPK3 |
| MAX | MCL1 | MDM2 | MDM4 | MED12 | MEF2B | MEN1 | MET | MITF | MLH1 | MPL | MRE11 |
| MSH2 | MSH6 | MTHFR* | MTOR | MUC16 | MUC4 | MUC6 | MUTYH | MYC | MYCL | MYCN | MYD88 |
| NAT2* | NBN | NEFH | NF1 | NF2 | NFE2L2 | NFKB1 | NFKBIA | NKX2-1* | NOTCH1 | NOTCH2 | NOTCH3 |
| NOTCH4 | NPM1 | NQO1* | NRAS | NSD1 | NTRK1 | NTRK2 | NTRK3 | PAK3 | PALB2 | PARP1 | PAX5 |
| PAX8 | PBRM1 | PDCD1 | PDCD1LG2 | PDGFRA | PDGFRB | PDIA3 | PGF | PHOX2B* | PIK3C2B | PIK3C2G | PIK3C3 |
| PIK3CA | PIK3CB | PIK3CD | PIK3CG | PIK3R1 | PIK3R2 | PIK3R3 | PIM1 | PMS1 | PMS2 | POLB | POLD1 |
| POLE | PPARG | PPP2R1A | PRDM1 | PRKAR1A | PRKCA | PRKCB | PRKCG | PRKCI | PRKCQ | PRKDC | PRKN |
| PSMB8 | PSMB9 | PSME1 | PSME2 | PSME3 | PTCH1 | PTEN | PTGS2 | PTPN11 | PTPRD | PTPRT | RAC1 |
| RAD50 | RAD51 | RAD51B | RAD51C | RAD51D | RAD52 | RAD54L | RAF1 | RARA | RB1 | RBM10 | RECQL4 |
| REL | RET | RHOA | RICTOR | RNF43 | ROS1 | RPPH1 | RPTOR | RUNX1 | RUNX1T1 | RXRA | SDHA |
| SDHB | SDHC | SDHD | SERPINB3 | SERPINB4 | SETD2 | SF3B1 | SGK1 | SH2D1A* | SLC19A1* | SLC22A2* | SLC1B1* |
| SLC1B3* | SMAD2 | SMAD3 | SMAD4 | SMARCA4 | SMARCB1 | SMO | SOCS1* | SOX2* | SOX9 | SPEN | SPOP |
| SRC | STAG2 | STAT3 | STK11 | SUFU | SYK | SYNE1 | TAF1 | TAP1 | TAP2 | TAPBP | TBX3 |
| TEK | TERT | TET1 | TET2 | TGFBR2 | TMSB4X* | TNF | TNFAIP3 | TNFRSF14 | TNFSF11 | TOP1 | TP53 |
| TPMT* | TSC1 | TSC2 | TSHR | TYMS | U2AF1 | UBE2A* | UBE2K | UBR5 | UGT1A1* | USH2A | VDR* |
| VEGFA | VEGFB | VHL | WT1 | XIAP | XPO1 | XRCC2 | ZNF217 | | | | |

*Analysis of copy number alterations NOT available.

FUSION

| | | | | | | | | | | | | |
|-----|------|------|-------|-------|-------|-----|------|-------|-------|-------|-----|------|
| ALK | BRAF | EGFR | FGFR1 | FGFR2 | FGFR3 | MET | NRG1 | NTRK1 | NTRK2 | NTRK3 | RET | ROS1 |
|-----|------|------|-------|-------|-------|-----|------|-------|-------|-------|-----|------|

ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Not Applicable.

SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS

Not Applicable.

ACT Onco[®] + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 17055429; 2006, Cell;127(2):265-75
The regulation of INK4/ARF in cancer and aging.
2. PMID: 8521522; 1995, Cell;83(6):993-1000
Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest.
3. PMID: 9529249; 1998, Cell;92(6):725-34
ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppression pathways.
4. PMID: 16115911; 2005, Clin Cancer Res;11(16):5740-7
Comprehensive analysis of CDKN2A status in microdissected urothelial cell carcinoma reveals potential haploinsufficiency, a high frequency of homozygous co-deletion and associations with clinical phenotype.
5. PMID: 7550353; 1995, Nat Genet;11(2):210-2
Frequency of homozygous deletion at p16/CDKN2 in primary human tumours.
6. PMID: 24089445; 2013, Clin Cancer Res;19(19):5320-8
The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma.
7. PMID: 27849562; 2017, Gut;66(7):1286-1296
Palbociclib (PD-0332991), a selective CDK4/6 inhibitor, restricts tumour growth in preclinical models of hepatocellular carcinoma.
8. PMID: 25524798; 2015, Lancet Oncol;16(1):25-35
The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study.
9. PMID: 28283584; 2017, Oncologist;22(4):416-421
Clinical Benefit in Response to Palbociclib Treatment in Refractory Uterine Leiomyosarcomas with a Common CDKN2A Alteration.
10. PMID: 27217383; 2016, Cancer Discov;6(7):740-53
Efficacy and Safety of Abemaciclib, an Inhibitor of CDK4 and CDK6, for Patients with Breast Cancer, Non-Small Cell Lung Cancer, and Other Solid Tumors.
11. PMID: 26715889; 2015, Curr Oncol;22(6):e498-501
Does CDKN2A loss predict palbociclib benefit?
12. PMID: 25501126; 2015, Clin Cancer Res;21(5):995-1001
CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment.
13. PMID: 27542767; 2016, Clin Cancer Res;22(23):5696-5705
A Phase I Study of the Cyclin-Dependent Kinase 4/6 Inhibitor Ribociclib (LEE011) in Patients with Advanced Solid Tumors and Lymphomas.
14. PMID: 24797823; 2014, Oncologist;19(6):616-22
Enabling a genetically informed approach to cancer medicine: a retrospective evaluation of the impact of comprehensive tumor profiling using a targeted next-generation sequencing panel.
15. PMID: 35050752; 2020, JCO Precis Oncol;4():757-766
Palbociclib in Patients With Non-Small-Cell Lung Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
16. PMID: 35100714; 2019, JCO Precis Oncol;3():1-8
Palbociclib in Patients With Pancreatic and Biliary Cancer With CDKN2A Alterations: Results From the Targeted Agent and Profiling Utilization Registry Study.
17. PMID: 27717303; 2016, N Engl J Med;375(18):1738-1748
Ribociclib as First-Line Therapy for HR-Positive, Advanced Breast Cancer.

ACT Onco[®] + Report

18. PMID: 28580882; 2017, J Clin Oncol;35(25):2875-2884
MONARCH 2: Abemaciclib in Combination With Fulvestrant in Women With HR+/HER2- Advanced Breast Cancer Who Had Progressed While Receiving Endocrine Therapy.
19. PMID: 26728409; 2016, Clin Cancer Res;22(1):122-33
Coadministration of Trametinib and Palbociclib Radiosensitizes KRAS-Mutant Non-Small Cell Lung Cancers In Vitro and In Vivo.
20. PMID: 31401335; 2019, Transl Oncol;12(11):1425-1431
Concomitant Genetic Alterations are Associated with Worse Clinical Outcome in EGFR Mutant NSCLC Patients Treated with Tyrosine Kinase Inhibitors.
21. PMID: 2453289; 1988, Cell;53(4):549-54
Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes.
22. PMID: 2114981; 1990, Eur J Clin Invest;20(3):225-35
ras oncogenes: their role in neoplasia.
23. PMID: 20617134; 2010, J Biomed Biotechnol;2010():150960
Clinical relevance of KRAS in human cancers.
24. PMID: 21993244; 2011, Nat Rev Cancer;11(11):761-74
RAS oncogenes: weaving a tumorigenic web.
25. PMID: 3047672; 1988, Nucleic Acids Res;16(16):7773-82
KRAS codon 12 mutations occur very frequently in pancreatic adenocarcinomas.
26. PMID: 3587348; 1987, Nature;327(6120):293-7
Prevalence of ras gene mutations in human colorectal cancers.
27. PMID: 1942608; 1991, Nihon Shokakibyo Gakkai Zasshi;88(8):1539-44
[Prevalence of K-ras gene mutations in human colorectal cancers].
28. PMID: 2252272; 1990, Am Rev Respir Dis;142(6 Pt 2):S27-30
The ras oncogenes in human lung cancer.
29. PMID: 1486840; 1992, Environ Health Perspect;98():13-24
Role of proto-oncogene activation in carcinogenesis.
30. PMID: 23455880; 2013, J Cancer Res Clin Oncol;139(6):953-61
KRAS allele-specific activity of sunitinib in an isogenic disease model of colorectal cancer.
31. PMID: 18349398; 2008, J Clin Oncol;26(9):1472-8
Molecular characteristics of bronchioloalveolar carcinoma and adenocarcinoma, bronchioloalveolar carcinoma subtype, predict response to erlotinib.
32. PMID: 23401440; 2013, J Clin Oncol;31(8):1112-21
KRAS mutation: should we test for it, and does it matter?
33. PMID: 18024870; 2007, J Clin Oncol;25(33):5240-7
Prognostic and predictive importance of p53 and RAS for adjuvant chemotherapy in non small-cell lung cancer.
34. PMID: 29946554; 2018, Gynecol Oncol Rep;25():41-44
Binimetinib (MEK162) in recurrent low-grade serous ovarian cancer resistant to chemotherapy and hormonal treatment.
35. PMID: 26075998; 2014, Gynecol Oncol Rep;10():28-9
Response to MEK inhibitor in small cell neuroendocrine carcinoma of the cervix with a KRAS mutation.
36. PMID: 25722381; 2015, Ann Oncol;26(5):894-901
A randomized phase II study of the MEK1/MEK2 inhibitor trametinib (GSK1120212) compared with docetaxel in KRAS-mutant advanced non-small-cell lung cancer (NSCLC)†.

ACT Onco[®] + Report

37. PMID: 24947927; 2014, Clin Cancer Res;20(16):4251-61
Phase I expansion and pharmacodynamic study of the oral MEK inhibitor RO4987655 (CH4987655) in selected patients with advanced cancer with RAS-RAF mutations.
38. PMID: 27340376; 2016, Curr Colorectal Cancer Rep;12():141-150
Molecular Subtypes and Personalized Therapy in Metastatic Colorectal Cancer.
39. PMID: 22392911; 2012, Clin Cancer Res;18(9):2515-25
Inhibition of MEK and PI3K/mTOR suppresses tumor growth but does not cause tumor regression in patient-derived xenografts of RAS-mutant colorectal carcinomas.
40. PMID: 26369631; 2016, Clin Cancer Res;22(2):405-14
Sensitivity of KRAS-Mutant Colorectal Cancers to Combination Therapy That Cotargets MEK and CDK4/6.
41. PMID: 24407191; 2014, Br J Cancer;110(5):1148-54
Sorafenib and irinotecan (NEXIRI) as second- or later-line treatment for patients with metastatic colorectal cancer and KRAS-mutated tumours: a multicentre Phase I/II trial.
42. PMID: 23224737; 2013, Clin Cancer Res;19(3):743-51
A phase II study of sorafenib in patients with platinum-pretreated, advanced (Stage IIIB or IV) non-small cell lung cancer with a KRAS mutation.
43. PMID: 26307133; 2016, Clin Cancer Res;22(2):374-82
Copy Number Changes Are Associated with Response to Treatment with Carboplatin, Paclitaxel, and Sorafenib in Melanoma.
44. PMID: 15923428; 2005, Ann Oncol;16 Suppl 4():iv44-49
Prognostic and predictive factors in colorectal cancer: Kirsten Ras in CRC (RASCAL) and TP53CRC collaborative studies.
45. PMID: 26484411; 2015, Br J Cancer;113(9):1254-8
Impact of mutational status on survival in low-grade serous carcinoma of the ovary or peritoneum.
46. PMID: 24549645; 2013, J Pathol;231(4):449-56
KRAS (but not BRAF) mutations in ovarian serous borderline tumour are associated with recurrent low-grade serous carcinoma.
47. PMID: 26662311; 2016, Tumour Biol;37(5):6823-30
G12V and G12A KRAS mutations are associated with poor outcome in patients with metastatic colorectal cancer treated with bevacizumab.
48. PMID: 26372703; 2015, Br J Cancer;113(8):1206-15
Prognostic value of the KRAS G12V mutation in 841 surgically resected Caucasian lung adenocarcinoma cases.
49. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
50. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
51. PMID: 22713868; 2012, Genes Dev;26(12):1268-86
Mutant p53: one name, many proteins.
52. PMID: 23264849; 2012, Genes Cancer;3(7-8):491-502
Gain-of-Function Activity of Mutant p53 in Lung Cancer through Up-Regulation of Receptor Protein Tyrosine Kinase Axl.
53. PMID: 23612969; 2013, J Biol Chem;288(23):16704-16714
A novel p53 mutant found in iatrogenic urothelial cancers is dysfunctional and can be rescued by a second-site global suppressor mutation.
54. PMID: 29970031; 2018, BMC Cancer;18(1):709
Mutant p53 gain of function induces HER2 over-expression in cancer cells.
55. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-

ACT Onco[®] + Report

Line Therapy Within 3 Months.

56. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
57. PMID: 25669829; 2015, Ann Oncol;26(5):1012-1018
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
58. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
59. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
60. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
61. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
62. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
63. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
64. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
65. PMID: 25385265; 2015, Int J Oncol;46(2):607-18
TP53 oncomorphic mutations predict resistance to platinum and taxane-based standard chemotherapy in patients diagnosed with advanced serous ovarian carcinoma.
66. PMID: 19029958; 2008, Nat Rev Cancer;8(12):976-90
Reflecting on 25 years with MYC.
67. PMID: 22464321; 2012, Cell;149(1):22-35
MYC on the path to cancer.
68. PMID: 10378696; 1999, Oncogene;18(19):3004-16
MYC oncogenes and human neoplastic disease.
69. PMID: 16934487; 2006, Semin Cancer Biol;16(4):318-30
The Myc oncoprotein as a therapeutic target for human cancer.
70. PMID: 22113465; 2012, Clin Exp Med;12(4):217-23
C-myc as a predictive marker for chemotherapy in metastatic breast cancer.
71. PMID: 21741827; 2011, Eur J Cancer;47(12):1779-88
Association between c-myc amplification and pathological complete response to neoadjuvant chemotherapy in breast cancer.
72. PMID: 15132769; 2004, Cancer Sci;95(5):418-23
Expression of the c-myc gene as a predictor of chemotherapy response and a prognostic factor in patients with ovarian cancer.
73. PMID: 22430491; 2012, J Exp Med;209(4):679-96
MYC pathway activation in triple-negative breast cancer is synthetic lethal with CDK inhibition.
74. PMID: 27486754; 2016, Oncotarget;7(35):56864-56875

ACT Onco[®] + Report

Inhibition of cyclin dependent kinase 9 by dinaciclib suppresses cyclin B1 expression and tumor growth in triple negative breast cancer.

75. PMID: 24503701; 2014, PLoS One;9(2):e87456
Immunohistochemistry for myc predicts survival in colorectal cancer.
76. PMID: 9816266; 1996, Clin Cancer Res;2(6):1049-53
Overexpression of the c-myc proto-oncogene in colorectal carcinoma is associated with a reduced mortality that is abrogated by point mutation of the p53 tumor suppressor gene.
77. PMID: 23860775; 2013, Tumour Biol;34(6):3945-58
MYC overexpression and poor prognosis in sporadic breast cancer with BRCA1 deficiency.