



Sample Information

Patient Name: 阮氏賢
Gender: Female
ID No.: F260146776
History No.: 49088406
Age: 38

Ordering Doctor: DOC1483K 王浩元
Ordering REQ.: 0CUCNHU
Signing in Date: 2023/11/30

Path No.: M112-00310
MP No.: MY23081
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/11/30

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Lymphoblastic Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	18 Therapies Available
Relevant Therapy Summary	3	0 Clinical Trials
Relevant Therapy Details	4	
Diagnostic Details	13	
Alert Details	13	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	TP53 p.(R282W) c.844C>T tumor protein p53 Allele Frequency: 63.19%	None	idelalisib + rituximab ² acalabrutinib allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO
Public data sources included in diagnostic significance: NCCN, ESMO

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.10(004). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
			decitabine ibrutinib liposomal cytarabine-daunorubicin CPX-351 obinutuzumab + venetoclax rituximab + venetoclax venetoclax venetoclax + chemotherapy	
	Diagnostic significance: Acute Lymphoblastic Leukemia			

Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

Public data sources included in diagnostic significance: NCCN, ESMO

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
TP53	p.(R282W)	c.844C>T	COSM10704	chr17:7577094	63.19%	NM_000546.5	missense	1967
EZH2	p.(T86=)	c.258C>T	.	chr7:148529831	48.49%	NM_004456.5	synonymous	1825

Gene Fusions (RNA)			
Genes	Variant ID	Locus	Read Count
SET-NUP214	SET-NUP214.S7N18.Non-Targeted	chr9:131456321 - chr9:134034770	4960

Biomarker Descriptions

TP53 (tumor protein p53)

Background: The TP53 gene encodes the p53 tumor suppressor protein that binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis. Alterations in TP53 is required for oncogenesis as they result in loss of protein function and gain of transforming potential¹. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers^{2,3}.

Alterations and prevalence: TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)^{4,5,6,7,8,9}. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common including substitutions at codons R158, R175, Y220, R248, R273, and R282^{4,5}. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes^{10,11,12,13}.

Potential relevance: The small molecule p53 reactivator, PC14586, received a fast track designation (2020) by the FDA for advanced tumors harboring a TP53 Y220C mutation¹⁴. The FDA has granted fast track designation (2019) to the p53 reactivator, eprentapopt,¹⁵ and breakthrough designation¹⁶ (2020) in combination with azacitidine or azacitidine and venetoclax for acute myeloid leukemia patients (AML) and myelodysplastic syndrome (MDS) harboring a TP53 mutation, respectively. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation^{17,18}. TP53 mutations confer poor prognosis and poor risk in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), and chronic lymphocytic leukemia (CLL), and acute lymphoblastic leukemia (ALL)^{19,20,21,22,23,24}. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant²⁵. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system²⁶.

Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

TP53 p.(R282W) c.844C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	×	○	×	×	×
azacitidine	×	○	×	×	×
cytarabine	×	○	×	×	×
cytarabine + daunorubicin	×	○	×	×	×
cytarabine + daunorubicin + etoposide	×	○	×	×	×
cytarabine + etoposide + idarubicin	×	○	×	×	×
cytarabine + fludarabine + idarubicin + filgrastim	×	○	×	×	×
cytarabine + idarubicin	×	○	×	×	×
cytarabine + mitoxantrone	×	○	×	×	×
decitabine	×	○	×	×	×
liposomal cytarabine-daunorubicin CPX-351	×	○	×	×	×
venetoclax + azacitidine	×	○	×	×	×
venetoclax + cytarabine	×	○	×	×	×
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	×	○	×	×	×
venetoclax + decitabine	×	○	×	×	×
idelalisib + rituximab	×	×	○	○	×
acalabrutinib	×	×	×	○	×
ibrutinib	×	×	×	○	×
obinutuzumab + venetoclax	×	×	×	○	×
rituximab + venetoclax	×	×	×	○	×
venetoclax	×	×	×	○	×

Relevant Therapy Details

Current NCCN Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-09-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

TP53 p.(R282W) c.844C>T

☐ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

☐ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

☐ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

☐ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

TP53 p.(R282W) c.844C>T (continued)**○ cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

TP53 p.(R282W) c.844C>T (continued)**○ decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

TP53 p.(R282W) c.844C>T (continued)**○ venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

TP53 p.(R282W) c.844C>T (continued)**○ cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

○ venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

Current EMA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

EMA information is current as of 2023-09-13. For the most up-to-date information, search www.ema.europa.eu/ema.

TP53 p.(R282W) c.844C>T

☐ idelalisib + rituximab

Cancer type: Chronic Lymphocytic Leukemia Label as of: 2023-06-23

Variant class: TP53 mutation

Reference:

https://www.ema.europa.eu/en/documents/product-information/zydelig-epar-product-information_en.pdf

Current ESMO Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

ESMO information is current as of 2023-09-01. For the most up-to-date information, search www.esmo.org.

TP53 p.(R282W) c.844C>T

☐ acalabrutinib

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma **Variant class:** TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Symptomatic, Relapsed (Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

☐ ibrutinib

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma **Variant class:** TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Symptomatic, Relapsed (Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

☐ rituximab + venetoclax

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma **Variant class:** TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- Symptomatic, Relapsed (Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

☐ idelalisib + rituximab

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma **Variant class:** TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: II / B

Population segment (Line of therapy):

- Symptomatic, Relapsed (Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

TP53 p.(R282W) c.844C>T (continued)

○ acalabrutinib

Cancer type: Chronic Lymphocytic Leukemia, Small **Variant class:** TP53 mutation
Lymphocytic Lymphoma

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Early-stage; Symptomatic (First-line therapy)
- Advanced-stage (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

○ ibrutinib

Cancer type: Chronic Lymphocytic Leukemia, Small **Variant class:** TP53 mutation
Lymphocytic Lymphoma

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Early-stage; Symptomatic (First-line therapy)
- Advanced-stage (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

○ idelalisib + rituximab

Cancer type: Chronic Lymphocytic Leukemia, Small **Variant class:** TP53 mutation
Lymphocytic Lymphoma

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Early-stage; Symptomatic (First-line therapy)
- Advanced-stage (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

○ obinutuzumab + venetoclax

Cancer type: Chronic Lymphocytic Leukemia, Small **Variant class:** TP53 mutation
Lymphocytic Lymphoma

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Early-stage; Symptomatic (First-line therapy)
- Advanced-stage (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

TP53 p.(R282W) c.844C>T (continued)

○ venetoclax

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma
Variant class: TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Early-stage; Symptomatic (First-line therapy)
- Advanced-stage (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

○ venetoclax

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma
Variant class: TP53 mutation

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Symptomatic, Relapsed (Subsequent therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Lymphocytic Leukaemia [Annals of Oncology (2020), <https://doi.org/10.1016/j.annonc.2020.09.019>.(published)]

Diagnostic Details

Current ESMO Information

ESMO information is current as of 2023-09-01. For the most up-to-date information, search www.esmo.org.

TP53 p.(R282W) c.844C>T

Diagnostic significance: Acute Lymphoblastic Leukemia

Variant class: TP53 aberration

Diagnostic notes:

- ALL with adverse clinico-biological features (Recommended for new clinical trials)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2016) 27 (suppl 5): v69-v82.]

Alerts Informed By Public Data Sources

Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2023-09-13. For the most up-to-date information, search www.fda.gov.

TP53 p.(R282W) c.844C>T

eprenetapopt + azacitidine, eprenetapopt + venetoclax + azacitidine

Cancer type: Myelodysplastic Syndrome

Variant class: TP53 mutation

Supporting Statement:

The FDA has granted Breakthrough Designation to the small molecule inhibitor, eprenetapopt, in combination with azacitidine or azacitidine and venetoclax for TP53 mutant myelodysplastic syndrome (MDS).

Reference:

<http://vp280.alertir.com/en/pressreleases/karolinska-development%27s-portfolio-company-aprea-therapeutics-receives-fda-breakthrough-therapy-designation-1769167>

eprenetapopt + azacitidine, eprenetapopt + venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: TP53 mutation

Supporting Statement:

The FDA has granted Fast Track Designation to the small molecule inhibitor, eprenetapopt, in combination with azacitidine or azacitidine and venetoclax for TP53 mutant acute myeloid leukemia (AML).

Reference:

<https://ir.aprea.com//news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>

Current NCCN Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

NCCN information is current as of 2023-09-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

TP53 p.(R282W) c.844C>T

chemoimmunotherapy

Cancer type: Chronic Lymphocytic Leukemia, Small Lymphocytic Lymphoma
Variant class: TP53 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Chemoimmunotherapy is not recommended since del(17p)/TP53 mutation is associated with low response rates."

Reference: NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2023]

Current ESMO Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

ESMO information is current as of 2023-09-01. For the most up-to-date information, search www.esmo.org.

TP53 p.(R282W) c.844C>T

lenalidomide

Cancer type: Myelodysplastic Syndrome **Variant class:** TP53 mutation

Summary:

ESMO Clinical Practice Guidelines include the following supporting statement:

- "TP53 gene mutations, found in ~20% of lower-risk MDS with del(5q), confer resistance to LEN and a higher risk of AML progression."

Reference: ESMO Clinical Practice Guidelines - ESMO-Myelodysplastic Syndromes [Ann of Oncol (2020), <https://doi.org/10.1016/j.annonc.2020.11.002>]

References

1. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell*. 2014 Mar 17;25(3):304-17. PMID: 24651012
2. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol*. 2010 Jan;2(1):a001008. PMID: 20182602
3. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med*. 2017 Apr 3;7(4). PMID: 28270529
4. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. PMID: 22588877
5. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet*. 2013 Oct;45(10):1113-20. PMID: 24071849
6. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature*. 2012 Sep 27;489(7417):519-25. PMID: 22960745
7. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature*. 2015 Jan 29;517(7536):576-82. PMID: 25631445
8. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet*. 2016 Jun;48(6):607-16. PMID: 27158780
9. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature*. 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
10. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat*. 2002 Jun;19(6):607-14. PMID: 12007217
11. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer*. 2011 Apr;2(4):466-74. PMID: 21779514
12. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene*. 2007 Apr 2;26(15):2157-65. PMID: 17401424
13. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat*. 2014 Jun;35(6):766-78. PMID: 24729566
14. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
15. <https://ir.aprea.com//news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>
16. <http://vp280.alertir.com/en/pressreleases/karolinska-development%27s-portfolio-company-aprea-therapeutics-receives-fda-breakthrough-therapy-designation-1769167>
17. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol*. 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
18. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci*. 2017 Nov;74(22):4171-4187. PMID: 28643165
19. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]
20. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
21. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
22. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2023]
23. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2023]
24. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 2.2023]
25. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 5.2023]
26. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med*. 2020 Aug 3. PMID: 32747829