



Sample Information

**Patient Name:** 張予欣  
**Gender:** Female  
**ID No.:** J223599592  
**History No.:** 44280286  
**Age:** 7

**Ordering Doctor:** DOC3939K 侯明欣  
**Ordering REQ.:** 0CUSBV  
**Signing in Date:** 2023/12/14

**Path No.:** M112-00327  
**MP No.:** MY23085  
**Assay:** Oncomine Myeloid Assay  
**Sample Type:** Bone Marrow  
**Bone Marrow Aspirating Date:** 2023/12/14

**Reporting Doctor:** DOC5444B 楊靜芬 (Phone: 8#5444)

**Note:**

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	1 Relevant Biomarkers
Biomarker Descriptions	2	12 Therapies Available
Relevant Therapy Summary	4	0 Clinical Trials
Relevant Therapy Details	5	
Diagnostic Details	10	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>RUNX1::MECOM fusion</i> RUNX family transcription factor 1 - MDS1 and EVI1 complex locus	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine liposomal cytarabine-daunorubicin CPX-351	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO  
Public data sources included in diagnostic significance: NCCN, ESMO

## Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
		venetoclax + chemotherapy		
	<b>Diagnostic significance:</b> Acute Myeloid Leukemia			
Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO				
Public data sources included in diagnostic significance: NCCN, ESMO				

### Prevalent cancer biomarkers without relevant evidence based on included data sources

*CBL p.(C416R) c.1246T>C*

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
CBL	p.(C416R)	c.1246T>C	COSM1724993	chr11:119149238	51.53%	NM_005188.4	missense	1999
RB1	p.(T58I)	c.173C>T	.	chr13:48881451	52.50%	NM_000321.2	missense	1998
CEBPA	p.(H195_P196dup)	c.589_590insACCCG C	.	chr19:33792731	37.04%	NM_004364.4	nonframeshift Insertion	945
BCOR	p.(A1527=)	c.4581C>T	.	chrX:39916422	49.40%	NM_001123385.2	synonymous	2000

### Gene Fusions (RNA)

Genes	Variant ID	Locus	Read Count
RUNX1-MECOM	RUNX1-MECOM.R4M2	chr21:36206707 - chr3:169099312	7751
RUNX1-MECOM	RUNX1-MECOM.R3M2	chr21:36231771 - chr3:169099312	2061

## Biomarker Descriptions

### CBL (Cbl proto-oncogene)

**Background:** The CBL gene encodes the casitas B-lineage lymphoma (CBL) ubiquitin ligase, a member of the ubiquitin ligase (E3) protein family that also includes CBL-b and CBL-c<sup>1</sup>. CBL proteins are characterized by their highly conserved N-terminal tyrosine kinase binding (TKB) domain and RING finger (RF) catalytic domain which are directly involved in the regulation of receptor tyrosine kinase (RTK) signaling<sup>1,2</sup>. Upon recognition of an activated RTK via its TKB domain, CBL mediates the transfer of ubiquitin from the ubiquitin-conjugating enzyme (E2) via its RF domain, consequently targeting the RTK for proteasome degradation. CBL can also function as an adaptor protein via recruitment of signaling molecules to active RTKs<sup>2</sup>. CBL is the target of genetic aberrations, including missense mutations and translocations, which can lead to oncogenic transformation in hematological malignancies as well as solid tumors<sup>2,3,4,5</sup>. Mutations in CBL often result in a loss of E3 ligase activity, thereby preventing proteasome-mediated RTK degradation, which supports the role of CBL as a tumor suppressor gene<sup>3</sup>. However, CBL mutants often maintain their adapter function, contributing to their transforming potential and suggesting a simultaneous oncogenic role for CBL in cancer<sup>2</sup>. Hereditary mutations in CBL lead to constitutive activation of RAS and MAPK pathways resulting in genetic disorders known as RASopathies which can lead to increased cancer risk<sup>6</sup>.

**Alterations and prevalence:** Genetic alterations in CBL were first recognized in acute myeloid leukemia (AML) as a result of an interstitial deletion leading to MLL-CBL fusion<sup>7,8</sup>. However, fusions involving CBL are relatively rare. Aberrations in CBL most often involve missense mutations which commonly cluster in the linker region or RF domain corresponding to exons 8 and 9<sup>2,3</sup>. Such mutations lead to disruption of E3 ligase activity and have been reported in systemic mastocytosis (SM), 1-3% of de novo AML, 10% of secondary AML, 8% of atypical AML, and 10-15% of juvenile myelomonocytic leukemia (JMML) and chronic myelomonocytic leukemia (CMML)<sup>2,9,10,11,12,13,14</sup>. Mutations in CBL have also been reported in 1-6% of melanomas, lung, stomach, colorectal, esophageal, and uterine cancers<sup>5,11</sup>.

## Biomarker Descriptions (continued)

Potential relevance: Mutations in CBL confer adverse prognosis in SM and have been shown to be independently predictive of inferior survival<sup>10,15</sup>.

### MECOM (MDS1 and EVI1 complex locus)

**Background:** The MECOM gene encodes the MDS1 and EVI1 complex locus (MECOM), a zinc-finger transcriptional factor that regulates hematopoietic cell differentiation<sup>16</sup>. The MECOM locus encodes multiple alternative splice variants that result in MDS1-EVI1, MDS1, and EVI1 protein isoforms<sup>17</sup>. The EVI1 isoform is the most abundant and oncogenic form of MECOM that is expressed in various cancers including acute myeloid leukemia (AML)<sup>17,18</sup>. MECOM is a frequent target of chromosomal translocation which can lead to MECOM overexpression and leukemogenesis<sup>19</sup>.

**Alterations and prevalence:** Somatic mutations MECOM are observed in up to 22% of malignant melanoma; 75% of these mutations are missense and the remaining 25% are truncating mutations<sup>11,20,21</sup>. MECOM amplifications are observed in up to 35% of lung squamous cell carcinoma, 30% of ovarian serous cystadenocarcinoma, and 20% of esophageal adenocarcinoma, uterine carcinosarcoma, and cervical squamous cell carcinoma<sup>11,21</sup>. MECOM rearrangements occur with various partner genes including ETV6, RUNX1, and H2AFY<sup>22</sup>. The t(3;21)(q26;q22) translocation that results in the MECOM-RUNX1 fusion is most commonly observed in chronic myeloid leukemia (CML) in blast crisis. The t(3;3)(q21.3;q26.2)/ inv(3)(q21.3;q26.3) translocation, also referred to as inv(3)/t(3;3), results in a GATA2-MECOM fusion and is observed in AML, primary myelofibrosis (PMF), and myelodysplastic syndrome (MDS)<sup>6,23,24</sup>. The inv(3)/t(3;3) translocation repositions the distal GATA enhancer element and activates MECOM expression while simultaneously causing GATA2 haploinsufficiency<sup>25</sup>.

**Potential relevance:** AML with MECOM rearrangement is considered a distinct molecular subtype of AML as defined by the World Health Organization (WHO)<sup>26</sup>. MECOM rearrangements, including GATA2-MECOM fusions, are associated with poor/adverse risk in AML<sup>23,27</sup>. Inv(3) is associated with poor cytogenetic risk in MDS as defined by the revised international prognostic scoring system (IPSS-R) scoring system<sup>6</sup>. In PMF, inv(3) is considered an unfavorable karyotype associated with intermediate risk as defined by the dynamic international prognostic scoring system (DIPSS)-Plus scoring system<sup>24</sup>. MECOM overexpression is observed in 10% of de novo AML associated with poor prognosis, and is commonly found in MLL-rearranged cases<sup>28,29</sup>. Amplification of MECOM is associated with favorable prognosis in ovarian cancer<sup>30</sup>.

### RUNX1 (RUNX family transcription factor 1)

**Background:** The RUNX1 gene encodes the runt-related transcription factor (RUNX) 1, part of the RUNX family of transcription factors which also includes RUNX2 and RUNX3<sup>31</sup>. All RUNX proteins share several conserved regions with similar functionality including a highly conserved N-terminal 'runt' domain responsible for binding DNA, a C-terminal region composed of an activation domain, inhibitory domain, protein interacting motifs, and a nuclear matrix targeting signal<sup>32</sup>. Each of these proteins are capable of interacting with core binding factor beta (CBFβ) to form the core binding factor (CBF) complex. Consequently, RUNX1, RUNX2, and RUNX3 are collectively known as core binding factor alpha (CBFα) since they can each function as the alpha subunit of CBF. Specifically, CBFβ binds to the 'runt' domain of RUNX1 leading to RUNX1 stabilization and increased affinity of the CBF complex for promoters involved in hematopoietic differentiation and cell cycle regulation<sup>33,34</sup>. RUNX1 is frequently mutated in various hematological malignancies<sup>34</sup>. Germline mutations in RUNX1 result in a rare autosomal dominant condition known as familial platelet disorder, with predisposition to acute myeloid leukemia (FPD/AML)<sup>35,36</sup>. Somatic mutations and chromosomal translocations in RUNX1 are often observed in myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), and chronic myelomonocytic leukemia (CMML)<sup>34</sup>.

**Alterations and prevalence:** RUNX1 is frequently rearranged in hematological malignancies with over 50 different observed translocations<sup>37</sup>. The most recurrent translocation, t(12;21)(q34;q11), results in ETV6-RUNX1 fusion and is observed in 20-25% of childhood ALL<sup>38,39,40</sup>. This translocation is also observed in adult ALL at a lower frequency (2%)<sup>39,40</sup>. Another recurrent translocation, t(8;21)(q22;q22), results in RUNX1-RUNX1T1 fusion and is observed in 5-10% of AML<sup>41</sup>. The RUNX1-RUNX1T1 fusion, consists of the RHD domain of RUNX1 and the majority of RUNX1T1, which promotes oncogenesis by altering transcriptional regulation of RUNX1 target genes<sup>34,41</sup>. Somatic mutations in RUNX1 include missense, nonsense, and frameshift mutations resulting in loss of function or dominant negative effects<sup>34</sup>. RUNX1 mutations are reported in approximately 10% of de novo AML as well as 10-15% of MDS<sup>6,11,23,34</sup>.

**Potential relevance:** AML with RUNX1-RUNX1T1 fusions is considered a distinct molecular subtype by the World Health Organization (WHO)<sup>23,26</sup>. Translocations involving RUNX1, specifically t(8;21)(q22;q22)/RUNX1-RUNX1T1 in AML and t(12;21)(q34;q11)/ETV6-RUNX1 in ALL, are associated with favorable risk<sup>27,42</sup>. On the other hand, mutations in RUNX1 confer poor prognosis in AML, MDS, and systemic mastocytosis (SM)<sup>6,15,27</sup>.

## Relevant Therapy Summary

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### RUNX1::MECOM fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✕	●	✕	✕	✕
azacitidine	✕	●	✕	✕	✕
cytarabine	✕	●	✕	✕	✕
cytarabine + daunorubicin	✕	●	✕	✕	✕
cytarabine + daunorubicin + etoposide	✕	●	✕	✕	✕
cytarabine + etoposide + idarubicin	✕	●	✕	✕	✕
cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
cytarabine + idarubicin	✕	●	✕	✕	✕
cytarabine + mitoxantrone	✕	●	✕	✕	✕
decitabine	✕	●	✕	✕	✕
liposomal cytarabine-daunorubicin CPX-351	✕	●	✕	✕	✕
venetoclax + azacitidine	✕	●	✕	✕	✕
venetoclax + cytarabine	✕	●	✕	✕	✕
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
venetoclax + decitabine	✕	●	✕	✕	✕

## Relevant Therapy Details

### Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-09-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org).  
For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

### RUNX1::MECOM fusion

#### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ● cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ● cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ● cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

## RUNX1::MECOM fusion (continued)

### ● cytarabine + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)
- (Consolidation therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

## RUNX1::MECOM fusion (continued)

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**RUNX1::MECOM fusion (continued)****● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]



## RUNX1::MECOM fusion (continued)

### ● cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

### ● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: MECOM fusion

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

## Diagnostic Details

### Current ESMO Information

ESMO information is current as of 2023-09-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

#### RUNX1::MECOM fusion

### Diagnostic significance: Acute Myeloid Leukemia

**Variant class:** t(3;21)(q26.2;q22.1)

**Diagnostic notes:**

- AML with myelodysplasia-related changes; WHO classification of AML

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## References

1. Ryan et al. The N terminus of Cbl-c regulates ubiquitin ligase activity by modulating affinity for the ubiquitin-conjugating enzyme. *J. Biol. Chem.* 2010 Jul 30;285(31):23687-98. PMID: 20525694
2. Liyasova et al. Molecular pathways: cbl proteins in tumorigenesis and antitumor immunity-opportunities for cancer treatment. *Clin. Cancer Res.* 2015 Apr 15;21(8):1789-94. PMID: 25477533
3. Katzav et al. Mutations of c-Cbl in myeloid malignancies. *Oncotarget.* 2015 May 10;6(13):10689-96. PMID: 26028666
4. Kales et al. Cbl and human myeloid neoplasms: the Cbl oncogene comes of age. *Cancer Res.* 2010 Jun 15;70(12):4789-94. PMID: 20501843
5. Tan et al. CBL is frequently altered in lung cancers: its relationship to mutations in MET and EGFR tyrosine kinases. *PLoS ONE.* 2010 Jan 29;5(1):e8972. PMID: 20126411
6. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
7. Tefferi. Novel mutations and their functional and clinical relevance in myeloproliferative neoplasms: JAK2, MPL, TET2, ASXL1, CBL, IDH and IKZF1. *Leukemia.* 2010 Jun;24(6):1128-38. PMID: 20428194
8. Fu et al. Identification of CBL, a proto-oncogene at 11q23.3, as a novel MLL fusion partner in a patient with de novo acute myeloid leukemia. *Genes Chromosomes Cancer.* 2003 Jun;37(2):214-9. PMID: 12696071
9. Traina et al. Single nucleotide polymorphism array lesions, TET2, DNMT3A, ASXL1 and CBL mutations are present in systemic mastocytosis. *PLoS ONE.* 2012;7(8):e43090. PMID: 22905207
10. Pardanani et al. ASXL1 and CBL mutations are independently predictive of inferior survival in advanced systemic mastocytosis. *Br. J. Haematol.* 2016 Nov;175(3):534-536. PMID: 26628266
11. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
12. Loh et al. Mutations in CBL occur frequently in juvenile myelomonocytic leukemia. *Blood.* 2009 Aug 27;114(9):1859-63. PMID: 19571318
13. Jankowska et al. Mutational spectrum analysis of chronic myelomonocytic leukemia includes genes associated with epigenetic regulation: UTX, EZH2, and DNMT3A. *Blood.* 2011 Oct 6;118(14):3932-41. PMID: 21828135
14. Itzykson et al. Prognostic score including gene mutations in chronic myelomonocytic leukemia. *J. Clin. Oncol.* 2013 Jul 1;31(19):2428-36. PMID: 23690417
15. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
16. Hinai et al. Review: Aberrant EVI1 expression in acute myeloid leukaemia. *Br. J. Haematol.* 2016 Mar;172(6):870-8. PMID: 26729571
17. Bard-Chapeau et al. EVI1 oncoprotein interacts with a large and complex network of proteins and integrates signals through protein phosphorylation. *Proc. Natl. Acad. Sci. U.S.A.* 2013 Jul 30;110(31):E2885-94. PMID: 23858473
18. Ogawa et al. Abnormal expression of Evi-1 gene in human leukemias. *Hum. Cell.* 1996 Dec;9(4):323-32. PMID: 9183665
19. Choi et al. Intratumoral Heterogeneity of Frameshift Mutations in MECOM Gene is Frequent in Colorectal Cancers with High Microsatellite Instability. *Pathol. Oncol. Res.* 2017 Jan;23(1):145-149. PMID: 27620344
20. Lee et al. Targeted next-generation sequencing reveals high frequency of mutations in epigenetic regulators across treatment-naïve patient melanomas. 2015 Jun 9;7:59. PMID: 26221190
21. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
22. Han et al. H2AFY is a novel fusion partner of MECOM in acute myeloid leukemia. *Cancer Genet.* 2018 Apr;222-223:9-12. PMID: 29666008
23. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]
24. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2023]
25. Gröschel et al. A single oncogenic enhancer rearrangement causes concomitant EVI1 and GATA2 deregulation in leukemia. *Cell.* 2014 Apr 10;157(2):369-381. PMID: 24703711
26. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia.* 2022 Jul;36(7):1703-1719. PMID: 35732831
27. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
28. Barjesteh et al. High EVI1 expression predicts poor survival in acute myeloid leukemia: a study of 319 de novo AML patients. *Blood.* 2003 Feb 1;101(3):837-45. PMID: 12393383
29. Stevens et al. EVI1 expression in childhood acute lymphoblastic leukaemia is not restricted to MLL and BCR/ABL rearrangements and is influenced by age. *Blood Cancer J.* 2014 Jan 24;4:e179. PMID: 24464103

## References (continued)

30. Nanjundan et al. Amplification of MDS1/EVI1 and EVI1, located in the 3q26.2 amplicon, is associated with favorable patient prognosis in ovarian cancer. *Cancer Res.* 2007 Apr 1;67(7):3074-84. PMID: 17409414
31. de et al. Runx transcription factors in the development and function of the definitive hematopoietic system. *Blood.* 2017 Apr 13;129(15):2061-2069. PMID: 28179276
32. Chuang et al. RUNX family: Regulation and diversification of roles through interacting proteins. *Int. J. Cancer.* 2013 Mar 15;132(6):1260-71. PMID: 23180629
33. Jung et al. Prognostic factor analysis in core-binding factor-positive acute myeloid leukemia. *Anticancer Res.* 2014 Feb;34(2):1037-45. PMID: 24511052
34. Sood et al. Role of RUNX1 in hematological malignancies. *Blood.* 2017 Apr 13;129(15):2070-2082. PMID: 28179279
35. Béri-Dexheimer et al. Clinical phenotype of germline RUNX1 haploinsufficiency: from point mutations to large genomic deletions. *Eur. J. Hum. Genet.* 2008 Aug;16(8):1014-8. PMID: 18478040
36. Hayashi et al. Myeloid neoplasms with germ line RUNX1 mutation. *Int. J. Hematol.* 2017 Aug;106(2):183-188. PMID: 28534116
37. De et al. RUNX1 translocations and fusion genes in malignant hemopathies. *Future Oncol.* 2011 Jan;7(1):77-91. PMID: 21174539
38. De et al. ETV6 fusion genes in hematological malignancies: a review. *Leuk. Res.* 2012 Aug;36(8):945-61. PMID: 22578774
39. Pui et al. Acute lymphoblastic leukemia. *N. Engl. J. Med.* 2004 Apr 8;350(15):1535-48. PMID: 15071128
40. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
41. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D920-4. PMID: 23161685
42. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 2.2023]