


REPORTING	PHYSICIAN	 Complete Tumor Response Map on page 3
Report Date: APR-08-2024	Chih-Hsueh Chen	
Receipt Date: APR-03-2024	Account: Genconn Biotech Co., LTD	
Collection Date: APR-02-2024	Address: F15., No 207-5 Sec 3, Beixin Rd, Xindian	
Specimen: Blood	Dist, New Taipei City, 23143, Taiwan	
Status: FINAL	Ph: +886 963 820 633 Fax: N/A	
	Additional Recipient: N/A	

Summary of Detected Somatic Alterations, Immunotherapy Biomarkers & Associated Treatment Options

KEY ✔ Approved in indication ⚡ Approved in other indication ✖ Lack of response

Detected Alteration(s) / Biomarker(s)	Associated FDA-approved therapies	Clinical trial availability (see page 5)	% cfDNA or Amplification
EGFR T790M	✔ Osimertinib ✖ Afatinib, Dacomitinib, Erlotinib, Erlotinib+ramucirumab, Gefitinib, Neratinib	Yes	2.1%
EGFR S768I	⚡ Osimertinib	Yes	3.0%
EGFR L747_T751del (Exon 19 deletion)	✔ Osimertinib	Yes	4.7%
TP53 P92fs	None	Yes	1.9%

Variants of Uncertain Clinical Significance
PDGFRA G898S (0.9%), MET K324N (0.2%), AR E804D (0.1%)
The functional consequences and/or clinical significance of alterations are unknown. Relevance of therapies targeting these alterations is uncertain.

Synonymous Alterations
MTOR A849A (1.8%), RHOA R5R (0.1%)
This sequence change does not alter the amino acid at this position and is unlikely to be a therapeutic target. Clinical correlation is advised.

Comments
Reported by: JV4


Biomarker	Additional Details
MSI-High	NOT DETECTED

We evaluated this sample for 74 genes, including the following guideline-recommended genes for NSCLC







EGFR(T790M and others)	ALK	ROS1	BRAF	MET	ERBB2(HER2)	RET	NTRK	KRAS
------------------------	-----	------	------	-----	-------------	-----	------	------

Guardant360 Tumor Response Map

The Guardant360 Tumor Response Map illustrates the variant allele fraction (% cfDNA) of observed somatic variants at each sample submission. Amplifications are not plotted, and only the first and last five test dates are plotted. Please see the Physician Portal (portal.guardanthealth.com) for the Tumor Response Map with all test dates.



Detected Alteration(s) / Biomarker(s)	% cfDNA or Amp	Alteration Trend	
EGFR L747_T751del (Exon 19 deletion)	4.7%	17.1% 4.7%	
EGFR S768I	3.0%	12.6% 3%	
EGFR T790M	2.1%	ND 2.1%	
TP53 P92fs	1.9%	11.3% 1.9%	
MTOR A849A	1.8%	8.5% 1.8%	Synonymous Alteration §
PDGFRA G898S	0.9%	0.4% 0.9%	Variants of Uncertain Clinical Significance §

Detected Alteration(s) / Biomarker(s)	% cfDNA or Amp	Alteration Trend	
MET K324N	0.2%		Variants of Uncertain Clinical Significance §
AR E804D	0.1%		Variants of Uncertain Clinical Significance §
RHOA R5R	0.1%		Synonymous Alteration §
TERT A651D	ND		
ATM R1619R	ND		
ATM Y2755C	ND		

The table above annotates the variant allele fraction (% cfDNA) detected in this sample, listed in descending order.
§ See definitions section for more detail

Available Clinical Trials (within the same state as the ordering physician)

There may be additional trials not listed here. Visit: portal.guardanthealth.com or email clientservices@guardanthealth.com with A1008957 in the subject line of the email, for additional trials.

Alteration	Trial ID / Contact	Title	Phase	Site(s)
EGFR T790M	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Kaohsiung, Taiwan Taipei City, Taiwan Tainan, Taiwan Taichung, Taiwan
	NCT05120349 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Taipei City, Taiwan Tainan, Taiwan Taoyuan, Taiwan Taipei, Taiwan (3) Additional trial sites available
	NCT05526755 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study of 5 Years of Adjuvant Osimertinib in Completely Resected Epidermal Growth Factor Receptor Mutation (EGFRm) Non-small Cell Lung Carcinoma (NSCLC)	Phase 2	Kaohsiung City, Taiwan Kaohsiung, Taiwan Hualien, Taiwan Taipei, Taiwan (2) Additional trial sites available
	NCT05647122 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	First in Human Study of AZD9592 in Solid Tumors	Phase 1	Taipei City, Taiwan Taipei, Taiwan Taoyuan, Taiwan Taichung, Taiwan
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Yunlin, Taiwan Taipei City, Taiwan Kaohsiung, Taiwan (2) Taipei, Taiwan (2) Additional trial sites available
Visit portal.guardanthealth.com for trials not within the same state as the physician's office				
EGFR S768I	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Kaohsiung, Taiwan Taipei City, Taiwan Tainan, Taiwan Taichung, Taiwan
	NCT05120349 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Taipei City, Taiwan Tainan, Taiwan Taoyuan, Taiwan Taipei, Taiwan (3) Additional trial sites available
	NCT05663866 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Premedication to Reduce Amivantamab Associated Infusion Related Reactions	Phase 2	ChangHua, Taiwan Kaohsiung, Taiwan Taipei City, Taiwan Tainan City, Taiwan Additional trial sites available
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Yunlin, Taiwan Taipei City, Taiwan Kaohsiung, Taiwan (2) Taipei, Taiwan (2) Additional trial sites available
	NCT06120140 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Taipei, Taiwan Taoyuan City, Taiwan Hsin Chu, Taiwan
Visit portal.guardanthealth.com for trials not within the same state as the physician's office				

Alteration	Trial ID / Contact	Title	Phase	Site(s)
EGFR L747_T751del	NCT04077463 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Kaohsiung, Taiwan Taipei City, Taiwan Tainan, Taiwan Taichung, Taiwan
	NCT05120349 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Taipei City, Taiwan Tainan, Taiwan Taoyuan, Taiwan Taipei, Taiwan (3) Additional trial sites available
	NCT05663866 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Premedication to Reduce Amivantamab Associated Infusion Related Reactions	Phase 2	ChangHua, Taiwan Kaohsiung, Taiwan Taipei City, Taiwan Tainan City, Taiwan Additional trial sites available
	NCT05801029 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Yunlin, Taiwan Taipei City, Taiwan Kaohsiung, Taiwan (2) Taipei, Taiwan (2) Additional trial sites available
	NCT06120140 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Taipei, Taiwan Taoyuan City, Taiwan Hsin Chu, Taiwan
Visit portal.guardanthhealth.com for trials not within the same state as the physician's office				
TP53 P92fs	NCT04768868 Jian Wang, Jian. wang@impacttherapeutics.com, +86 18613056501	The Safety and Pharmacokinetics Preliminary Efficacy of IMP7068 in Patients With Advanced Solid Tumors	Phase 1	Taipei, Taiwan Taoyuan, Taiwan Taichung, Taiwan Tainan, Taiwan (2)
Visit portal.guardanthhealth.com for trials not within the same state as the physician's office				

More clinical trial options available at portal.guardanthhealth.com

Definitions

Somatic Alterations Not Detected (ND): Somatic alterations may be present that are below the limit of detection of this test. Certain sample or variant characteristics may result in reduced analytic sensitivity. The absence of detectable somatic alterations in circulating cell-free DNA does not preclude the presence of somatic alterations in the tumor.

Variants of Uncertain Clinical Significance: The functional consequences and/or clinical significance of alterations are unknown. Relevance of therapies targeting these alterations is uncertain.

Synonymous Alteration: This sequence change does not alter the amino acid at this position and is unlikely to be a therapeutic target. Clinical correlation is advised.

Deletion (Del): The following alteration was detected in this patient: *EGFR* L747_T751del; *TP53* P92fs. Guardant360 detects short deletions in exons of certain genes (see Table 1), including potential splice site-disrupting events.

Interpretation

Somatic alterations were detected in the circulating cell-free DNA isolated from this patient's blood specimen. These genomic alterations are cancer-associated somatic variants, some of which have been associated with either increased or reduced clinical response to specific treatments. The percentage of altered cell-free DNA circulating (% cfDNA) in blood is related to the unique tumor biology of each patient. Factors that may affect the % cfDNA of detected somatic alterations include tumor growth, turn over, size, heterogeneity, vascularization, disease progression, and treatment.

Method and Limitations

Guardant360 sequences 74 cancer-associated genes to identify somatic alterations. Cell-free DNA (cfDNA) is extracted from plasma, enriched for targeted regions, and sequenced using the Illumina platform and hg19 as the reference genome. All exons are sequenced in some genes; only clinically significant exons are sequenced in other genes. The types of genomic alterations detected by Guardant360 include single nucleotide variants, gene amplifications, fusions, short insertions/deletions (longest detected, 70 base pairs), and splice site disrupting events (see Table 1). Microsatellite Instability (MSI) is assessed for all cancer types by evaluating somatic changes in the length of repetitive sequences on the Guardant360 panel. A "Not Detected" result in samples where the highest % cfDNA is < 0.2% is an inconclusive result because it does not preclude MSI-High status in tissue. MSI status is currently not reported for earlier panel versions. This version of the Guardant360 test is not validated for the detection of other types of genomic alterations, such as complex rearrangements or gene deletions. Certain sample or variant characteristics, such as low cfDNA concentration, may result in reduced analytic sensitivity. Guardant360 cannot discern the source of circulating cfDNA, and for some variants in the range of ~40 to 60% cfDNA, the test cannot easily distinguish germline variants from somatic alterations. Guardant360 is not validated for the detection of germline or de novo variants that are associated with hereditary cancer risk. Tissue genotyping should be considered when plasma genotyping is negative, if clinically appropriate.

Table 1: Genes on the Guardant360 Panel

Guardant360 reports single nucleotide variants, splice site mutations, and insertion and deletion variants (indels) in all clinically relevant exons in 74 genes and reports other variant types in select genes as indicated below.

AKT1	ALK #	APC	AR †	ARAF	ARID1A	ATM	BRAF †	BRCA1
BRCA2	CCND1 †	CCND2 †	CCNE1 †	CDH1	CDK12	CDK4 †	CDK6 †	CDKN2A
CTNNB1	DDR2	EGFR †	ERBB2 †	ESR1	EZH2	FBXW7	FGFR1 †	FGFR2 # †
FGFR3 #	GATA3	GNA11	GNAQ	GNAS	HNF1A	HRAS	IDH1	IDH2
JAK2	JAK3	KIT †	KRAS †	MAP2K1	MAP2K2	MAPK1	MAPK3	MET †
MLH1	MPL	MTOR	MYC †	NF1	NFE2L2	NOTCH1	NPM1	NRAS
NTRK1 #	NTRK3	PDGFRA †	PIK3CA †	PTEN	PTPN11	RAF1 †	RB1	RET #
RHEB	RHOA	RIT1	ROS1 #	SMAD4	SMO	STK11	TERT ‡	TP53
TSC1	VHL							

‡ Guardant360 reports alterations in the promoter region of this gene.
Guardant360 reports fusion events involving this gene.
† Guardant360 reports amplifications of this gene.

About the Test

The Guardant360 assay was developed and its performance characteristics were determined by Guardant Health, Inc. This test has not been cleared or approved by the U.S. Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. This test may be used for clinical purposes and should not be regarded as investigational or for research only. Guardant Health's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high complexity clinical laboratory testing. The laboratory report should be interpreted in the context of other clinical information and laboratory, pathology, and imaging studies by a qualified medical professional prior to initiating or changing a patient's treatment plan. The selection of any, all, or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) is entirely at the discretion of the treating medical professional. Drug and trial information are based on the diagnosis written on the submitted test request form; this information is not based on any supplemental information provided by the requesting medical professional, including pathology reports or other molecular studies. Some drugs listed in this report may not be approved or cleared by the FDA for the indicated use. Guardant Health makes no endorsement, express or implied, of any product, physician, or procedure contained in this report. This report makes no promises or guarantees that a particular medication will affect (or not affect) the clinical outcome of any patient.

Testing Performed at: Guardant Health

Laboratory Director: Martina Lefterova, MD PhD | CLIA ID: 05D2070300 | CAP #: 8765297 | 505 Penobscot Drive, Redwood City, CA, 94063, USA

Additional information is available

Any therapeutic annotations are based on publicly available information. This information is described in the "Detailed Therapy Results" and "Relevance of Detected Alterations" sections.

Visit portal.guardanthealth.com or email clientservices@guardanthealth.com with A1008957 in the subject line of the email for:

- Additional clinical trials
- Detailed Therapy Results
- Relevance of Detected Alterations
- References

If you would like to receive this additional information with every Guardant360 report, please call client services at [855.698.8887](tel:855.698.8887) to opt-in.

Additional information begins on the next page.

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
EGFR T790M	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Seattle, WA; Detroit, MI; Saint Louis, MO; Philadelphia, PA; Portland, OR; Salt Lake City, UT; Tampa, FL; Fairfax, VA; Boston, MA (3); New York, NY (2); CA (5); Puerto Rico; Japan (7); China (13); Taiwan (4); Korea, Republic of (4); Italy (5); France (7); Germany (8); Spain (8)
	NCT05120349 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Louisville, KY; Fort Belvoir, VA; Newark, DE; Chicago, IL; Orange, CA; Morristown, NJ; San Francisco, CA; Los Angeles, CA; Houston, TX; Frederick, MD; Atlanta, GA; Grand Junction, CO; Lexington, KY; Anchorage, AK; NY (6); Argentina (8); Singapore (2); Romania (5); United Kingdom (7); Malaysia (4); Spain (5); Canada (3); Vietnam (3); Turkey (5); China (23); Taiwan (8); Poland (4); Brazil (7); Italy (10); Germany (7)
	NCT05401110 Clinical Trial Recruitment Navigator, cancer.trial.info@cshs.org,310-423-2133	Study of Osimertinib With Carotuximab in Advanced, EGFR-mutated Non-Small Cell Lung Cancer	Phase 1	CA (5)
	NCT05469022 In Ae Kim, MD. PhD.,20180618@kuh.ac.kr,+821035438353	Neoadjuvant Lazertinib Therapy in EGFR-Mutation Positive Lung Adenocarcinoma Detected by BALF Liquid Biopsy	Phase 2	Korea, Republic of
	NCT05507606 See https://clinicaltrials.gov/ct2/show/NCT05507606	Study of Osimertinib+Bevacizumab+Chemotherapy for EGFR+ Advanced Non-Small Cell Lung Cancer With Concurrent Mutations	Phase 2	China
	NCT05526755 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study of 5 Years of Adjuvant Osimertinib in Completely Resected Epidermal Growth Factor Receptor Mutation (EGFRm) Non-small Cell Lung Carcinoma (NSCLC)	Phase 2	Yuma, AZ; Rockville, MD; Las Vegas, NV; Santa Rosa, CA; White Plains, NY; Singapore; Hong Kong (2); Philippines (4); Taiwan (6); Korea, Republic of (15); United Kingdom (3); Italy (13); Malaysia (4); Thailand (2); Spain (8)
	NCT05647122 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	First in Human Study of AZD9592 in Solid Tumors	Phase 1	Providence, RI; Houston, TX; Duarte, CA; Mineola, NY; Milford, MA; Philadelphia, PA; Irvine, CA; Baltimore, MD; Fairfax, VA; North Haven, CT; New York, NY (3); Malaysia; Canada (2); Japan (2); China (4); Taiwan (4); Korea, Republic of (4); Italy (4); France (2); Australia (2); Spain (3)
	NCT05686434 chen chen,chen_checn@tmu.edu.cn, 13920761627	Osimertinib Therapy After Resection in High-risk Stage I EGFRm NSCLC (OSTAR)	Phase 2	China
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Canada (3); Singapore (3); Hong Kong (3); Taiwan (9); Korea, Republic of (7); Malaysia (6); Thailand (5)
	NCT06195189 Li Li, BA,tracy.li_2010@hotmail.com,	Sunvozertinib Combined With Chemotherapy for EGFRm After EGFR-TKI Treatment Failure#	Phase 1 /Phase 2	China

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
	113880343287 x+86	Phase I/II		
EGFR S768I	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Seattle, WA; Detroit, MI; Saint Louis, MO; Philadelphia, PA; Portland, OR; Salt Lake City, UT; Tampa, FL; Fairfax, VA; Boston, MA (3); New York, NY (2); CA (5); Puerto Rico; Japan (7); China (13); Taiwan (4); Korea, Republic of (4); Italy (5); France (7); Germany (8); Spain (8)
	NCT04197934 Clinical Trials Referral Office, mayocliniccancerstudies@mayo.edu,855-776-0015	WSD0922-FU for the Treatment of Glioblastoma, Anaplastic Astrocytoma, or Non-small Cell Lung Cancer With Central Nervous System Metastases	Phase 1	Rochester, MN; Scottsdale, AZ; Jacksonville, FL
	NCT05068024 Jo-Han Wang, Project Manager,jo-han.wang@wuxiapptec.com,+1-647-649-2850	A Study of FWD1509 in Adults With Non-Small Cell Lung Cancer	Phase 1 /Phase 2	Houston, TX; Canton, OH
	NCT05120349 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Louisville, KY; Fort Belvoir, VA; Newark, DE; Chicago, IL; Orange, CA; Morristown, NJ; San Francisco, CA; Los Angeles, CA; Houston, TX; Frederick, MD; Atlanta, GA; Grand Junction, CO; Lexington, KY; Anchorage, AK; NY (6); Argentina (8); Singapore (2); Romania (5); United Kingdom (7); Malaysia (4); Spain (5); Canada (3); Vietnam (3); Turkey (5); China (23); Taiwan (8); Poland (4); Brazil (7); Italy (10); Germany (7)
	NCT05326425 Jin Hyoung Kang, oncologykang@naver.com,82-2-2258-6043	Lazertinib in Patients With NSCLC With Asymptomatic or Mild Symptomatic Brain Metastases After Failure of EGFR TKI.	Phase 2	Korea, Republic of (6)
	NCT05469022 In Ae Kim, MD. PhD.,20180618@kuh.ac.kr,+821035438353	Neoadjuvant Lazertinib Therapy in EGFR-Mutation Positive Lung Adenocarcinoma Detected by BALF Liquid Biopsy	Phase 2	Korea, Republic of
	NCT05663866 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Premedication to Reduce Amivantamab Associated Infusion Related Reactions	Phase 2	Renton, WA; Fountain Valley, CA; Fairfax, VA; Taiwan (7); Korea, Republic of (6); France (6); Spain (14)
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Canada (3); Singapore (3); Hong Kong (3); Taiwan (9); Korea, Republic of (7); Malaysia (6); Thailand (5)
	NCT06120140 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Hinsdale, IL; Westbury, NY; W. Salem, WI; Reno, NV; Flemington, NJ; CA (5); Argentina; Turkey; China (3); Taiwan (3); Korea, Republic of (2); Malaysia (4); Spain (2)
	NCT06195189 Li Li, BA,tracy.li_2010@hotmail.com, 113880343287 x+86	Sunvozertinib Combined With Chemotherapy for EGFRm After EGFR-TKI Treatment Failure# Phase I/II	Phase 1 /Phase 2	China
EGFR L747_T751del	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Seattle, WA; Detroit, MI; Saint Louis, MO; Philadelphia, PA; Portland, OR; Salt Lake City, UT; Tampa, FL; Fairfax, VA; Boston, MA (3); New York, NY (2); CA (5); Puerto Rico; Japan (7); China (13); Taiwan (4);

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
				Korea, Republic of (4); Italy (5); France (7); Germany (8); Spain (8)
	NCT05120349 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Global Study to Assess the Effects of Osimertinib in Participants With EGFRm Stage IA2-IA3 NSCLC Following Complete Tumour Resection	Phase 3	Louisville, KY; Fort Belvoir, VA; Newark, DE; Chicago, IL; Orange, CA; Morristown, NJ; San Francisco, CA; Los Angeles, CA; Houston, TX; Frederick, MD; Atlanta, GA; Grand Junction, CO; Lexington, KY; Anchorage, AK; NY (6); Argentina (8); Singapore (2); Romania (5); United Kingdom (7); Malaysia (4); Spain (5); Canada (3); Vietnam (3); Turkey (5); China (23); Taiwan (8); Poland (4); Brazil (7); Italy (10); Germany (7)
	NCT05326425 Jin Hyoung Kang, oncologykang@naver.com, 82-2-2258-6043	Lazertinib in Patients With NSCLC With Asymptomatic or Mild Symptomatic Brain Metastases After Failure of EGFR TKI.	Phase 2	Korea, Republic of (6)
	NCT05469022 In Ae Kim, MD, PhD., 20180618@kuh.ac.kr, +821035438353	Neoadjuvant Lazertinib Therapy in EGFR-Mutation Positive Lung Adenocarcinoma Detected by BALF Liquid Biopsy	Phase 2	Korea, Republic of
	NCT05663866 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Premedication to Reduce Amivantamab Associated Infusion Related Reactions	Phase 2	Renton, WA; Fountain Valley, CA; Fairfax, VA; Taiwan (7); Korea, Republic of (6); France (6); Spain (14)
	NCT05801029 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Canada (3); Singapore (3); Hong Kong (3); Taiwan (9); Korea, Republic of (7); Malaysia (6); Thailand (5)
	NCT05826483 Zhou Chengzhi, MD, doctorzcz@163.com, 13560351186	Almonertinib in the First-line Treatment of Patients of NSCLC With Poor Performance Status	Phase 1	China
	NCT06043973 Degan Lu, Professor, deganlu@126.com, 18753157623	Almonertinib Combined With Anlotinib as First-line Treatment for Advanced Non-small Cell Lung Cancer	Phase 3	China
	NCT06120140 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Hinsdale, IL; Westbury, NY; W. Salem, WI; Reno, NV; Flemington, NJ; CA (5); Argentina; Turkey; China (3); Taiwan (3); Korea, Republic of (2); Malaysia (4); Spain (2)
	NCT06195189 Li Li, BA, tracy.li_2010@hotmail.com, 113880343287 x+86	Sunvozertinib Combined With Chemotherapy for EGFRm After EGFR-TKI Treatment Failure# Phase I/II	Phase 1 /Phase 2	China
TP53 P92fs	NCT02769962 Danielle F Pinkiert, R.N., danielle.pinkiert@nih.gov, (240) 858-7566	Trial of EP0057, a Nanoparticle Camptothecin With Olaparib in People With Relapsed /Refractory Small Cell Lung Cancer	Phase 1 /Phase 2	Bethesda, MD
	NCT03968653 Debiopharm International S.A, clinicaltrials@debiopharm.com, +41 21 321 01 11	Study of Oral Debio 0123 in Combination With Carboplatin in Participants With Advanced Solid Tumors	Phase 1	Spain; Netherlands (3)
	NCT04768868 Jian Wang, Jian, wang@impacttherapeutics.com, +86 18613056501	The Safety and Pharmacokinetics Preliminary Efficacy of IMP7068 in Patients With Advanced Solid Tumors	Phase 1	Louisville, KY; Boston, MA; Atlanta, GA; Dallas, TX; Fairway, KS; San Antonio, TX; China (4); Taiwan (5)
	NCT04869475 Min Shi, MD & Ph. D, sm11998@rjh.com.	Arsenic Trioxide in Refractory Solid Tumors With Rescuable p53 Mutation	Phase 2	China

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
	cn,+86-21-64370045			
	NCT05109975 Debiopharm International S.A, clinicaltrials@debiopharm.com,+41 21 321 01 11	A Study to Evaluate Safety and Preliminary Anti-tumor Activity of Debio 0123 as Monotherapy in Adult Participants With Advanced Solid Tumors	Phase 1	Grand Rapids, MI; San Antonio, TX; Switzerland; Spain (7)
	NCT05489731 li zhang, professor,zhangli6@mail.sysu.edu.cn,13902282893	VIC-1911 Combined With Osimertinib for EGFR -Mutant Non-small Cell Lung Cancer	Phase 1	China

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
EGFR S768I T790M L747_T751del (Exon 19 deletion)	ABT-101		Egfr/Her2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Head and neck squamous cell carcinoma (HNSCC))
	Amivantamab	Rybrevant	Bispecific anti-Met/Egfr antibody.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 20 insertion)
	Avitinib		Irreversible mutation-specific Egfr kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Non-Hodgkin lymphoma (NHL))
	AZD3759		Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	BAY2927088		Egfr/Her2 kinase inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	BBP-398		Shp-2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	BBT-176		Fourth generation Egfr inhibitor targeting exon 19del/L858R, T790M, and C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Befotertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	BLU-451		Egfr inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	BLU-945		Fourth generation Egfr inhibitor targeting T790M and T790M /C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	BPI-361175		Fourth generation Egfr inhibitor targeting T790M and T790M /C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	BPI-7711		Egfr T790M inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	CLN-081		Covalent mutation-specific (L858R, T790M, exon 19 deletion, exon 20 insertion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	ERAS-601		Shp-2 inhibitor.	Phase 2 (Solid Tumor)
	ET0038		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Furmonertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	FWD1509		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	H002		Fourth generation Egfr inhibitor targeting exon 19del/L858R, T790M, and C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	HBI-2376		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Hemay022		Egfr tyrosine kinase inhibitor.	Phase 1 (Breast carcinoma (HER2+))
	JIN-A02		Fourth generation Egfr inhibitor targeting T790M and T790M /C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Lazertinib		Third generation mutation-specific	Phase 2 (Non-small cell lung carcinoma (NSCLC))

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
TP53 P92fs			Egfr tyrosine kinase inhibitor.	
	Lifirafenib		Dual Braf/Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	MCLA-129		Anti-EGFR/c-Met bispecific antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Esophageal squamous cell carcinoma)
	Mobocertinib	Exkivity	Mutation-specific Egfr/Her2 inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (NSCLC with EGFR exon 20 insertion, Lung cancer)
	Modotuximab		Anti-EGFR antibody.	Phase 1 (Gastric carcinoma, Colorectal carcinoma (CRC))
	Naquotinib		EGFR mutant-specific inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	NX-019		Egfr inhibitor.	Phase 1 (Solid Tumor)
	Osimertinib	Tagrisso	Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)
	Pirotinib		ErbB family inhibitor.	Phase 1 (Solid Tumor)
	Pozotinib		Egfr/Her2/ErbB4 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Breast carcinoma, Esophageal squamous cell carcinoma, Colorectal carcinoma (CRC))
	Pyrotinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Breast carcinoma)
	SKLB1028		Egfr/Flt3/c-Abl inhibitor.	Phase 2 (Acute myeloid leukemia (AML))
	Sunvozertinib		Bispecific anti-Egfr/Her2 monoclonal antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Non-Hodgkin lymphoma (NHL))
	TAS2940		Egfr/Her2 kinase inhibitor.	Phase 1 (Solid Tumor)
	TAS3351		Fourth generation Egfr inhibitor targeting T790M and T790M /C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	TAVO412		Anti-c-Met/anti-EGFR/anti-VEGF trispecific antibody.	Phase 1 (Solid Tumor)
	Varlitinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Gastric carcinoma, Hepatocellular carcinoma (HCC), Pancreatic carcinoma, Cholangiocarcinoma)
	ZN-e4		Egfr T790M inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	Adavosertib		Wee1 tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Lymphoma, Embryonal tumor with multi-layered rosettes (ETMR), Medulloblastoma, Small cell lung carcinoma (SCLC), Solid Tumor, Primary myelofibrosis (PMF), Ovarian carcinosarcoma, Acute myeloid leukemia (AML), MDS/MPN, unclassifiable, Chronic myelomonocytic leukemia)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
				(CMML), Peritoneal papillary serous carcinoma, Myelodysplastic Syndrome (MDS))
	AL8326		Aurora kinase B/VEGFRs/Fgfr multi-kinase inhibitor.	Phase 1 (Solid Tumor) Phase 3 (Small cell lung carcinoma (SCLC))
	Alisertib		Aurora kinase A inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Peripheral T-cell lymphoma (PTCL))
	ATO	Trisenox	PML-RARA inhibitor. Inhibits multiple signaling pathways, including the Hedgehog pathway.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Acute myeloid leukemia (AML), Acute promyelocytic leukemia (APL))
	AZD2811		Nanoparticle formulation of Aurora kinase B inhibitor barasertib (AZD1152).	Phase 1 (Solid Tumor) Phase 2 (Acute myeloid leukemia (AML), Myelodysplastic Syndrome (MDS))
	Azenosertib		Wee1 tyrosine kinase inhibitor.	Phase 1 (Solid Tumor) Phase 2 (High-grade serous ovarian carcinoma, Uterine serous/clear cell carcinoma, Osteosarcoma, Ovarian epithelial carcinoma, Colorectal adenocarcinoma, Acute myeloid leukemia (AML), Fallopian tube carcinoma, Peritoneal carcinoma, Pancreatic adenocarcinoma)
	Debio 0123		Wee1 tyrosine kinase inhibitor.	Phase 1 (Solid Tumor)
	EP0042		Aurora kinase A/B and Flt3 inhibitor.	Phase 2 (Acute myeloid leukemia (AML), Chronic myelomonocytic leukemia (CMML), Myelodysplastic Syndrome (MDS))
	IMP7068		Wee1 tyrosine kinase inhibitor.	Phase 1 (Solid Tumor)
	JAB-2485		Aurora kinase A inhibitor.	Phase 2 (Solid Tumor)
	LY3295668		Aurora kinase A inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Head and neck squamous cell carcinoma (HNSCC), Small cell lung carcinoma (SCLC), Breast carcinoma (triple negative), Breast carcinoma (hormone receptor +, HER2-))
	SGT-53		TP53 gene therapy delivered via transferrin-targeted nanoparticles.	Phase 1 (Solid Tumor) Phase 2 (Glioblastoma, Glioma, Pancreatic carcinoma)
	SY-4835		Wee1 tyrosine kinase inhibitor.	Phase 1 (Solid Tumor)
	TAS-119		Aurora kinase A inhibitor.	Phase 1 (Solid Tumor)
	Tinengotinib		Aurora kinase A/B inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Breast carcinoma (triple negative))
EGFR T790M L747_T751del (Exon 19 deletion)	Aumolertinib		Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	CM93		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 1 (Glioblastoma)
	Nazartinib		Third generation EGFR mutant-specific (T790M, L858R, exon 19 deletion) tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Olafertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	Olmutinib		Egfr inhibitor.	Phase 2 (Non-small cell lung carcinoma)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
EGFR S768I L747_T751del (Exon 19 deletion)				(NSCLC))
	PF-06747775		Egfr T790M-specific inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	BDTX-1535		Irreversible brain-penetrant fourth generation Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioma)
	BDTX-189		Irreversible Egfr/Her2 inhibitor.	Phase 2 (Solid Tumor)
	BLU-701		Fourth generation Egfr inhibitor targeting exon 19del, L858R, and C797X resistance mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Icotinib	Conmana	Egfr inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Esophageal carcinoma)
	WSD0922-FU		Blood-brain barrier penetrable EGFR /EGFRvIII inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioblastoma, Anaplastic astrocytoma)

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
EGFR T790M	The presence of an EGFR abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Egfr protein, which can lead to excessive proliferation. ⁽¹⁾ The EGFR T790M mutation has typically been reported as a secondary resistance mutation to the Egfr inhibitors erlotinib, gefitinib and afatinib, but has also been reported as a rare germline variant in de novo non-small cell lung cancer, particularly in lung adenocarcinoma. ^(2,3) Several studies have reported the presence of the T790M mutation in the germline in 0-1% of NSCLC cases, although one study did detect this mutation in 50% (5/10) of cases; in addition, T790M was associated with a 31% risk of lung cancer in never smokers in one study. ⁽⁴⁻⁷⁾	The presence of a sensitizing EGFR mutation in a tumor is the strongest biological predictor of sensitivity to an Egfr tyrosine kinase inhibitor (TKI). Compared with conventional chemotherapy, Egfr TKIs have been shown to improve progression-free survival in non-small cell lung cancer patients whose tumors harbor EGFR mutations. ⁽⁸⁻¹¹⁾ The EGFR T790M mutation reported here has been described as a "gatekeeper" mutation that confers resistance to the tyrosine kinase inhibitors erlotinib and gefitinib. ^(12,13) Studies have also reported the emergence of EGFR T790M upon resistance to afatinib monotherapy. ^(3,15) Third generation irreversible Egfr TKIs that target the EGFR T790M mutation have shown efficacy in T790M-mutant NSCLC and are under clinical investigation. Osimertinib has received approval by the FDA, EMA, and PMDA for the treatment of EGFR T790M-mutant metastatic NSCLC as well as EGFR-mutated NSCLC that has not been exposed to previous TKI treatment. ⁽¹⁶⁻²¹⁾ However, this EGFR mutation has been associated with resistance to erlotinib; thus, the relevance of ramucirumab plus erlotinib is uncertain.	A Phase 2 study has reported no clinical responses in 12 non-small cell lung cancer (NSCLC) patients who harbored EGFR T790M and were treated with neratinib. ⁽²²⁾ In addition, Phase 2 studies have reported limited clinical activity with dacomitinib in NSCLC patients who harbored EGFR T790M, with 2/32 patients having shown a partial response. ⁽²³⁻²⁵⁾ Some patients with EGFR-mutant NSCLC exhibit resistance to Egfr inhibition; resistance has been associated with insertions in EGFR exon 20, the T790M mutation in EGFR, and amplification of the gene MET. ⁽²⁶⁻²⁸⁾ Several studies have reported that resistance to Egfr TKIs in NSCLC is mediated by the transformation of NSCLC cell types to those of SCLC with neuroendocrine features. ⁽²⁹⁻³²⁾
EGFR S768I	The presence of an EGFR abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Egfr protein, which can lead to excessive proliferation. ⁽¹⁾	The presence of a sensitizing EGFR mutation in a tumor is the strongest biological predictor of sensitivity to an Egfr tyrosine kinase inhibitor (TKI). Compared with conventional chemotherapy, Egfr TKIs have been shown to improve progression-free survival in non-small cell lung cancer patients whose tumors harbor EGFR mutations. ⁽⁸⁻¹¹⁾ The Egfr TKIs erlotinib, afatinib, gefitinib, osimertinib, and dacomitinib have been approved by the FDA for the treatment of non-small cell lung cancer (NSCLC) with exon 19 deletion or L858R EGFR mutations; osimertinib has additionally been approved for the treatment of NSCLC with EGFR T790M. ^(8,11,16,33-37) Afatinib has additionally been FDA-approved for the treatment of NSCLC with S768I, L861Q, and/or G719X mutations. ⁽³⁾ The combination of erlotinib and ramucirumab as well as osimertinib plus platinum-based chemotherapy have been FDA-approved for the treatment of metastatic NSCLC patients with	Some patients with EGFR-mutant NSCLC exhibit resistance to Egfr inhibition; resistance has been associated with insertions in EGFR exon 20, the T790M mutation in EGFR, and amplification of either MET or ERBB2. ^(26-28,49) Third generation irreversible Egfr TKIs that target the EGFR T790M mutation have shown efficacy in T790M-mutant NSCLC, including osimertinib, which has received approval by the FDA, EMA, and PMDA for the treatment of EGFR T790M-mutant metastatic NSCLC. ⁽¹⁷⁻²¹⁾ Several studies have reported that resistance to Egfr TKIs in NSCLC is mediated by the transformation of NSCLC cell types to those of SCLC with neuroendocrine features. ⁽²⁹⁻³²⁾ Preclinical studies have reported increased Smo expression in NSCLC cell lines resistant to first, second, and third generation Egfr inhibitors as compared with sensitive ones; treatment with Smo inhibitors was observed to restore sensitivity in the resistant cell lines. ⁽⁵⁰⁻⁵²⁾

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
		tumors harboring an EGFR exon 19 deletion or the exon 21 L858R mutation. (38-40). Amivantamab has been approved by the FDA for NSCLC patients with EGFR exon 20 insertions, whose disease has progressed on or after platinum-based chemotherapy and as frontline therapy in combination with carboplatin and pemetrexed. The accelerated FDA approval of mobocertinib for NSCLC patients with EGFR exon 20 insertions has been withdrawn due to lack of progression-free survival benefit in the confirmatory Phase 3 trial. (41-44). Studies have reported non-squamous NSCLC patients with metastatic disease and tumors harboring an EGFR exon 19 deletion or L858R mutation to be sensitive to osimertinib, erlotinib, afatinib, gefitinib, dacomitinib, and the combination of erlotinib plus ramucirumab. (8,11,16,33,38,45). Less common activating EGFR mutations have variable sensitivity to EGFR tyrosine kinase inhibitors. (46). However, EGFR T790M, also detected in this tumor, has been associated with resistance to the Egfr tyrosine kinase inhibitors erlotinib, gefitinib, afatinib, and dacomitinib. (3,12,15,15-25,27,47).	
EGFR L747_T751del	The presence of an EGFR abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Egfr protein, which can lead to excessive proliferation. (1).	The presence of a sensitizing EGFR mutation in a tumor is the strongest biological predictor of sensitivity to an Egfr tyrosine kinase inhibitor (TKI). Compared with conventional chemotherapy, Egfr TKIs have been shown to improve progression-free survival in non-small cell lung cancer patients whose tumors harbor EGFR mutations. (8-11). The Egfr TKIs erlotinib, afatinib, gefitinib, osimertinib, and dacomitinib have been approved by the FDA for the treatment of non-small cell lung cancer (NSCLC) with exon 19 deletion or L858R EGFR mutations; osimertinib has additionally been approved for the treatment of NSCLC with EGFR T790M. (8,11,16,33-37). Afatinib has additionally been FDA-approved for the treatment of NSCLC with S768I, L861Q, and/or G719X mutations. (3). The combination of erlotinib and ramucirumab as well as osimertinib plus platinum-based chemotherapy have been FDA-approved for the treatment of metastatic NSCLC patients with tumors harboring an EGFR exon 19 deletion or the exon 21 L858R	Some patients with EGFR-mutant NSCLC exhibit resistance to Egfr inhibition; resistance has been associated with insertions in EGFR exon 20, the T790M mutation in EGFR, and amplification of either MET or ERBB2. (26-28,49). Third generation irreversible Egfr TKIs that target the EGFR T790M mutation have shown efficacy in T790M-mutant NSCLC, including osimertinib, which has received approval by the FDA, EMA, and PMDA for the treatment of EGFR T790M-mutant metastatic NSCLC. (17-21). Several studies have reported that resistance to Egfr TKIs in NSCLC is mediated by the transformation of NSCLC cell types to those of SCLC with neuroendocrine features. (29-32). Preclinical studies have reported increased Smo expression in NSCLC cell lines resistant to first, second, and third generation Egfr inhibitors as compared with sensitive ones; treatment with Smo inhibitors was observed to restore sensitivity in the resistant cell lines. (50-52).

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
		<p>mutation. (38-40). Amivantamab has been approved by the FDA for NSCLC patients with EGFR exon 20 insertions, whose disease has progressed on or after platinum-based chemotherapy and as frontline therapy in combination with carboplatin and pemetrexed. The accelerated FDA approval of mobocertinib for NSCLC patients with EGFR exon 20 insertions has been withdrawn due to lack of progression-free survival benefit in the confirmatory Phase 3 trial. (41-44). Studies have reported non-squamous NSCLC patients with metastatic disease and tumors harboring an EGFR exon 19 deletion or L858R mutation to be sensitive to osimertinib, erlotinib, afatinib, gefitinib, dacomitinib, and the combination of erlotinib plus ramucirumab. (8,11,16,33,38,45). Less common activating EGFR mutations have variable sensitivity to EGFR tyrosine kinase inhibitors. (46). However, one of the EGFR mutations in the report has been associated with resistance to erlotinib; thus, the relevance of ramucirumab plus erlotinib is uncertain. However, EGFR T790M, also detected in this tumor, has been associated with resistance to the Egfr tyrosine kinase inhibitors erlotinib, gefitinib, afatinib, and dacomitinib. (3,12,15,15-25,27,47).</p>	
TP53 P92fs	Loss of tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers. (53). Carriers of a germline mutation in TP53 have Li-Fraumeni Syndrome, an inherited cancer syndrome resulting in multiple tumors in early adulthood, including breast cancer, brain tumors, and leukemias. (54-56). Expression of p53 in normal cells is low; however, TP53 alterations, including those that result in loss of p53 tumor suppressor function, may lead to stabilization and increased expression of p53, particularly in the nucleus, and several studies have shown that it may have oncogenic gain-of-function effects. (57-61). TP53 alterations are believed to be early events in NSCLC, preceding lymph node metastasis. (62). TP53 mutation and expression of p53 have been correlated with the lung squamous cell carcinoma subtype, and p53 expression in lung squamous cell carcinoma has also been associated	At present, there are no approved therapies targeting TP53 alterations, despite their high prevalence in cancer. Therapeutic approaches under investigation include gene therapy for TP53 and (dendritic cell-based) TP53 vaccines. (73-75). Inhibition of components of the DNA damage checkpoint, including Wee1, has been reported to enhance the activity of DNA-damaging agents in preclinical cancer models with deficiency of p53 function. (76-78). Clinical trials of the Wee1 inhibitor adavosertib (MK-1775) are currently underway for patients with solid tumors and hematologic malignancies. Studies have reported Aurora kinase A to be activated in cells harboring TP53 mutation, and Aurora kinase A and B inhibitors have been reported to activate wild-type p53 in cellular assays; thus, tumors retaining a wild-type TP53 allele may benefit from Aurora kinase inhibitors. (79-84).	Mutations in TP53 may increase resistance to ionizing radiation therapy. (85,86).

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
	with disease stage and higher grade tumors. (63-66). TP53 mutation has been associated with PD-L1 expression and T-cell infiltration in lung adenocarcinoma samples. (67-71). TP53 mutations have been significantly associated with the development of distant metastases after diagnosis in early-stage NSCLC in a cohort of 759 patients. (72).		

References

1. Ciardiello F, Tortora G "EGFR antagonists in cancer treatment." *The New England journal of medicine*(2008): 1160-74
2. Wu J, Yu C, Chang Y, Yang C, Shih J, Yang P "Effectiveness of tyrosine kinase inhibitors on "uncommon" epidermal growth factor receptor mutations of unknown clinical significance in non-small cell lung cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 3812-21
3. Yang J, Sequist L, Geater S, Tsai C, Mok T, Schuler M, Yamamoto N, Yu C, Ou S, Zhou C, Massey D, Zazulina V, Wu Y "Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6." *The Lancet. Oncology*(2015): 830-8
4. Gazdar A, Robinson L, Oliver D, Xing C, Travis W, Soh J, Toyooka S, Watumull L, Xie Y, Kernstine K, Schiller J "Hereditary lung cancer syndrome targets never smokers with germline EGFR gene T790M mutations." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2014): 456-63
5. Girard N, Lou E, Azzoli C, Reddy R, Robson M, Harlan M, Orlow I, Yatabe Y, Nafa K, Ladanyi M, Viale A, Kris M, Riely G, Miller V, Klein R, Matsuo K, Pao W "Analysis of genetic variants in never-smokers with lung cancer facilitated by an Internet-based blood collection protocol: a preliminary report." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2010): 755-63
6. Vikis H, Sato M, James M, Wang D, Wang Y, Wang M, Jia D, Liu Y, Bailey-Wilson J, Amos C, Pinney S, Petersen G, de Andrade M, Yang P, Wiest J, Fain P, Schwartz A, Gazdar A, Gaba C, Rothschild H, Mandal D, Kupert E, Seminara D, Viswanathan A, Govindan R, Minna J, Anderson M, You M "EGFR-T790M is a rare lung cancer susceptibility allele with enhanced kinase activity." *Cancer research*(2007): 4665-70
7. Rosell R, Molina M, Costa C, Simonetti S, Gimenez-Capitan A, Bertran-Alamillo J, Mayo C, Moran T, Mendez P, Cardenal F, Isla D, Provencio M, Cobo M, Insa A, Garcia-Campelo R, Reguart N, Majem M, Viteri S, Carcereny E, Porta R, Massuti B, Queralt C, de Aguirre I, Sanchez J, Sanchez-Ronco M, Mate J, Ariza A, Benlloch S, Sanchez J, Bivona T, Sawyers C, Taron M "Pretreatment EGFR T790M mutation and BRCA1 mRNA expression in erlotinib-treated advanced non-small-cell lung cancer patients with EGFR mutations." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 1160-8
8. Mok T, Wu Y, Thongprasert S, Yang C, Chu D, Saijo N, Sunpaweravong P, Han B, Margono B, Ichinose Y, Nishiwaki Y, Ohe Y, Yang J, Chewaskulyong B, Jiang H, Duffield E, Watkins C, Armour A, Fukuoka M "Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma." *The New England journal of medicine*(2009): 947-57
9. Rosell R, Moran T, Queralt C, Porta R, Cardenal F, Camps C, Majem M, Lopez-Vivanco G, Isla D, Provencio M, Insa A, Massuti B, Gonzalez-Larriba J, Paz-Ares L, Bover I, Garcia-Campelo R, Moreno M, Catot S, Rolfo C, Reguart N, Palmero R, Sánchez J, Bastus R, Mayo C, Bertran-Alamillo J, Molina M, Sanchez J, Taron M "Screening for epidermal growth factor receptor mutations in lung cancer." *The New England journal of medicine*(2009): 958-67
10. Tsao M, Sakurada A, Cutz J, Zhu C, Kamel-Reid S, Squire J, Lorimer I, Zhang T, Liu N, Daneshmand M, Marrano P, da Cunha Santos G, Lagarde A, Richardson F, Seymour L, Whitehead M, Ding K, Pater J, Shepherd F "Erlotinib in lung cancer - molecular and clinical predictors of outcome." *The New England journal of medicine* (2005): 133-44
11. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia-Gomez R, Pallares C, Sanchez J, Porta R, Cobo M, Garrido P, Longo F, Moran T, Insa A, De Marinis F, Corre R, Bover I, Illiano A, Dansin E, de Castro J, Milella M, Reguart N, Altavilla G, Jimenez U, Provencio M, Moreno M, Terrasa J, Muñoz-Langa J, Valdivia J, Isla D, Domine M, Molinier O, Mazieres J, Baize N, Garcia-Campelo R, Robinet G, Rodriguez-Abreu D, Lopez-Vivanco G, Gebbia V, Ferrera-Delgado L, Bombardieri P, Bernabe R, Bearz A, Arta A, Cortesi E, Rolfo C, Sanchez-Ronco M, Drozdowskyj A, Queralt C, de Aguirre I, Ramirez J, Sanchez J, Molina M, Taron M, Paz-Ares L "Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTC): a multicentre, open-label, randomised phase 3 trial." *The Lancet. Oncology*(2012): 239-46
12. Suda K, Onozato R, Yatabe Y, Mitsudomi T "EGFR T790M mutation: a double role in lung cancer cell survival?" *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2009): 1-4
13. Yun C, Mengwasser K, Toms A, Woo M, Greulich H, Wong K, Meyerson M, Eck M "The T790M mutation in EGFR kinase causes drug resistance by increasing the affinity for ATP." *Proceedings of the National Academy of Sciences of the United States of America*(2008): 2070-5
14. Wu S, Liu Y, Tsai M, Chang Y, Yu C, Yang P, Yang J, Wen Y, Shih J "The mechanism of acquired resistance to irreversible EGFR tyrosine kinase inhibitor-afatinib in lung adenocarcinoma patients." *Oncotarget*(2016): 12404-13
15. Kim Y, Ko J, Cui Z, Abolhoda A, Ahn J, Ou S, Ahn M, Park K "The EGFR T790M mutation in acquired resistance to an irreversible second-generation EGFR inhibitor." *Molecular cancer therapeutics*(2012): 784-91
16. Soria J, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee K, Dechaphunkul A, Imamura F, Nogami N, Kurata T, Okamoto I, Zhou C, Cho B, Cheng Y, Cho E, Voon P, Planchard D, Su W, Gray J, Lee S, Hodge R, Marotti M, Rukazekov Y, Ramalingam S "Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer." *The New England journal of medicine*(2018): 113-125
17. Jänne P, Boss D, Camidge D, Britten C, Engelman J, Garon E, Guo F, Wong S, Liang J, Letrent S, Millham R, Taylor I, Eckhardt S, Schellens J "Phase I dose-escalation study of the pan-HER inhibitor, PF299804, in patients with advanced malignant solid tumors." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 1131-9
18. Jänne P, Yang J, Kim D, Planchard D, Ohe Y, Ramalingam S, Ahn M, Kim S, Su W, Horn L, Haggstrom D, Felip E, Kim J, Frewer P, Cantarini M, Brown K, Dickinson P, Ghiorghiu S, Ranson M "AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer." *The New England journal of medicine*(2015): 1689-99
19. Greig S "Osimertinib: First Global Approval." *Drugs*(2016): 263-73
20. Yang J, Ahn M, Kim D, Ramalingam S, Sequist L, Su W, Kim S, Kim J, Planchard D, Felip E, Blackhall F, Haggstrom D, Yoh K, Novello S, Gold K, Hirashima T, Lin C, Mann H, Cantarini M, Ghiorghiu S, Jänne P "Osimertinib in Pretreated T790M-Positive Advanced Non-Small-Cell Lung Cancer: AURA Study Phase II Extension Component." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2017): 1288-1296
21. Wang S, Cang S, Liu D "Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer." *Journal of hematology & oncology*(2016): 34
22. Sequist L, Besse B, Lynch T, Miller V, Wong K, Gitlitz B, Eaton K, Zacharchuk C, Freyman A, Powell C, Ananthakrishnan R, Quinn S, Soria J "Neratinib, an irreversible pan-ErbB receptor tyrosine kinase inhibitor: results of a phase II trial in patients with advanced non-small-cell lung cancer." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2010): 3076-83
23. Reckamp K, Giaccone G, Camidge D, Gadgeel S, Khuri F, Engelman J, Koczywas M, Rajan A, Campbell A, Gernhardt D, Ruiz-Garcia A, Letrent S, Liang J, Taylor I, O'Connell J, Jänne P "A phase 2 trial of dacomitinib (PF-00299804), an oral, irreversible pan-HER (human epidermal growth factor receptor) inhibitor, in patients with advanced non-small cell lung cancer after failure of prior chemotherapy and erlotinib." *Cancer*(2014): 1145-54

References

24. Jänne P, Shaw A, Camidge D, Giaccone G, Shreeve S, Tang Y, Goldberg Z, Martini J, Xu H, James L, Solomon B "Combined Pan-HER and ALK/ROS1/MET Inhibition with Dacomitinib and Crizotinib in Advanced Non-Small Cell Lung Cancer: Results of a Phase I Study." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2016): 737-747
25. Yu H, Ahn M, Cho B, Gerber D, Natale R, Socinski M, Giri N, Quinn S, Sbar E, Zhang H, Giaccone G "Phase 2 study of intermittent pulse dacomitinib in patients with advanced non-small cell lung cancers." *Lung cancer (Amsterdam, Netherlands)*(2017): 195-199
26. Engelman J, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park J, Lindeman N, Gale C, Zhao X, Christensen J, Kosaka T, Holmes A, Rogers A, Cappuzzo F, Mok T, Lee C, Johnson B, Cantley L, Jänne P "MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling." *Science (New York, N.Y.)*(2007): 1039-43
27. Greulich H, Chen T, Feng W, Jänne P, Alvarez J, Zappaterra M, Bulmer S, Frank D, Hahn W, Sellers W, Meyerson M "Oncogenic transformation by inhibitor-sensitive and -resistant EGFR mutants." *PLoS medicine*(2005): e313
28. Kwak E, Sordella R, Bell D, Godin-Heymann N, Okimoto R, Brannigan B, Harris P, Driscoll D, Fidias P, Lynch T, Rabindran S, McGinnis J, Wissner A, Sharma S, Isselbacher K, Settleman J, Haber D "Irreversible inhibitors of the EGF receptor may circumvent acquired resistance to gefitinib." *Proceedings of the National Academy of Sciences of the United States of America*(2005): 7665-70
29. Watanabe S, Sone T, Matsui T, Yamamura K, Tani M, Okazaki A, Kurokawa K, Tambo Y, Takato H, Ohkura N, Waseda Y, Katayama N, Kasahara K "Transformation to small-cell lung cancer following treatment with EGFR tyrosine kinase inhibitors in a patient with lung adenocarcinoma." *Lung cancer (Amsterdam, Netherlands)*(2013): 370-2
30. Chang Y, Kim S, Choi Y, So K, Rho J, Kim W, Lee J, Chung J, Choi C "Neuroendocrine differentiation in acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitor." *Tuberculosis and respiratory diseases*(2013): 95-103
31. Popat S, Wotherspoon A, Nutting C, Gonzalez D, Nicholson A, O'Brien M "Transformation to "high grade" neuroendocrine carcinoma as an acquired drug resistance mechanism in EGFR-mutant lung adenocarcinoma." *Lung cancer (Amsterdam, Netherlands)*(2013): 1-4
32. Sequist L, Waltman B, Dias-Santagata D, Digumarthy S, Turke A, Fidias P, Bergethon K, Shaw A, Gettinger S, Cosper A, Akhavanfar S, Heist R, Temel J, Christensen J, Wain J, Lynch T, Vernovsky K, Mark E, Lanuti M, Iafrate A, Mino-Kenudson M, Engelman J "Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors." *Science translational medicine*(2011): 75a26
33. Wu Y, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Tsuji F, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Nadanaciva S, Sandin R, Mok T "Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial." *The Lancet. Oncology*(2017): 1454-1466
34. Mok T, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Lee M, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Wu Y "Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2018): 2244-2250
35. Shepherd F, Rodrigues Pereira J, Ciuleanu T, Tan E, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, Findlay B, Tu D, Johnston D, Bezjak A, Clark G, Santabárbara P, Seymour L "Erlotinib in previously treated non-small-cell lung cancer." *The New England journal of medicine*(2005): 123-32
36. Sequist L, Yang J, Yamamoto N, O'Byrne K, Hirsh V, Mok T, Geater S, Orlov S, Tsai C, Boyer M, Su W, Bennouna J, Kato T, Gorbunova V, Lee K, Shah R, Massey D, Zazulina V, Shahidi M, Schuler M "Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2013): 3327-34
37. Douillard J, Ostoros G, Cobo M, Ciuleanu T, McCormack R, Webster A, Milenkova T "First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study." *British journal of cancer*(2014): 55-62
38. Nakagawa K, Garon E, Seto T, Nishio M, Ponce Aix S, Paz-Ares L, Chiu C, Park K, Novello S, Nadal E, Imamura F, Yoh K, Shih J, Au K, Moro-Sibilot D, Enatsu S, Zimmermann A, Fridolf-Moller B, Visseren-Grul C, Reck M "Ramucicromab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial." *The Lancet. Oncology*(2019): 1655-1669
39. Planchard D, Jänne P, Cheng Y, Yang J, Yanagitani N, Kim S, Sugawara S, Yu Y, Fan Y, Geater S, Laktionov K, Lee C, Valdiviezo N, Ahmed S, Maurel J, Andrasina I, Goldman J, Ghiorghiu D, Rukazkenov Y, Todd A, Kobayashi K "Osimertinib with or without Chemotherapy in EGFR-Mutated Advanced NSCLC." *The New England journal of medicine*(2023): 1935-1948
40. "A Phase III, Open-label, Randomized Study of Osimertinib With or Without Platinum Plus Pemetrexed Chemo, as First-line Treatment in Patients With Epidermal Growth Factor Receptor (EGFR) Mutation Positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer (FLAURA2)." (2024)
41. Cho B, Lee K, Cho E, et al. "Amivantamab (JNJ-61186372), an EGFR-MET bispecific antibody, in combination with lazertinib, a 3rd-generation tyrosine kinase inhibitor (TKI), in advanced EGFR NSCLC" *Annals of Oncology*(2020): Abstract 12580
42. Park K, John T, Kim S, et al. "Amivantamab (JNJ-61186372), an anti-EGFR-MET bispecific antibody, in patients with EGFR exon 20 insertion (exon20ins)-mutated non-small cell lung cancer (NSCLC)." *Journal of Clinical Oncology*(2020)
43. Riely G, Neal J, Camidge D, Spira A, Piotrowska Z, Costa D, Tsao A, Patel J, Gadgil S, Bazhenova L, Zhu V, West H, Mekhail T, Gentzler R, Nguyen D, Vincent S, Zhang S, Lin J, Bunn V, Jin S, Li S, Jänne P "Activity and Safety of Mobocertinib (TAK-788) in Previously Treated Non-Small Cell Lung Cancer with EGFR Exon 20 Insertion Mutations from a Phase I/II Trial." *Cancer discovery*(2021): 1688-1699
44. Zhou C, Ramalingam S, Li B, et al. "Mobocertinib in NSCLC With EGFR Exon 20 Insertions: Results From EXCLAIM and Pooled Platinum-Pretreated Patient Populations" *Journal of Thoracic Oncology*(2021)
45. Yang J, Wu Y, Schuler M, Sebastian M, Popat S, Yamamoto N, Zhou C, Hu C, O'Byrne K, Feng J, Lu S, Huang Y, Geater S, Lee K, Tsai C, Gorbunova V, Hirsh V, Bennouna J, Orlov S, Mok T, Boyer M, Su W, Lee K, Kato T, Massey D, Shahidi M, Zazulina V, Sequist L "Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials." *The Lancet. Oncology*(2015): 141-51
46. O'Kane G, Bradbury P, Feld R, Leigh N, Liu G, Pisters K, Kamel-Reid S, Tsao M, Shepherd F "Uncommon EGFR mutations in advanced non-small cell lung cancer." *Lung cancer (Amsterdam, Netherlands)*(2017): 137-144
47. Denis M, Vallée A, Théolyre S "EGFR T790M resistance mutation in non small-cell lung carcinoma." *Clinica chimica acta; international journal of clinical chemistry*(2015): 81-5

References

48. Takezawa K, Pirazzoli V, Arcila M, Nebhan C, Song X, de Stanchina E, Ohashi K, Janjigian Y, Spitzler P, Melnick M, Riely G, Kris M, Miller V, Ladanyi M, Politi K, Pao W "HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation." *Cancer discovery*(2012): 922-33
49. Yu H, Arcila M, Rekhtman N, Sima C, Zakowski M, Pao W, Kris M, Miller V, Ladanyi M, Riely G "Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2013): 2240-7
50. Della Corte C, Bellicic C, Vicidomini G, Vitagliano D, Malapelle U, Accardo M, Fabozzi A, Fiorelli A, Fasano M, Papaccio F, Martinelli E, Troiani T, Troncone G, Santini M, Bianco R, Ciardiello F, Morgillo F "SMO Gene Amplification and Activation of the Hedgehog Pathway as Novel Mechanisms of Resistance to Anti-Epidermal Growth Factor Receptor Drugs in Human Lung Cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2015): 4686-97
51. Bai X, Zhang X, Yang S, An S, Chen Z, Su J, Xie Z, Gou L, Wu Y "Blockade of Hedgehog Signaling Synergistically Increases Sensitivity to Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitors in Non-Small-Cell Lung Cancer Cell Lines." *PLoS one*(2016): e0149370
52. Della Corte C, Malapelle U, Viglieri E, Pepe F, Troncone G, Ciaramella V, Troiani T, Martinelli E, Belli V, Ciardiello F, Morgillo F "Efficacy of continuous EGFR-inhibition and role of Hedgehog in EGFR acquired resistance in human lung cancer cells with activating mutation of EGFR." *Oncotarget*(2017): 23020-23032
53. Brown C, Lain S, Verma C, Fersht A, Lane D "Awakening guardian angels: drugging the p53 pathway." *Nature reviews. Cancer*(2009): 862-73
54. Malkin D, Li F, Strong L, Fraumeni J, Nelson C, Kim D, Kassel J, Gryka M, Bischoff F, Tainsky M "Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms." *Science (New York, N.Y.)*(1990): 1233-8
55. Srivastava S, Zou Z, Pirolo K, Blattner W, Chang E "Germ-line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome." *Nature*(1991): 747-9
56. Santibáñez-Koref M, Birch J, Hartley A, Jones P, Craft A, Eden T, Crowther D, Kelsey A, Harris M "p53 germline mutations in Li-Fraumeni syndrome." *Lancet (London, England)*(1991): 1490-1
57. Wang Y, Lin R, Tan Y, Chen J, Chen C, Wang Y "Wild-type p53 overexpression and its correlation with MDM2 and p14ARF alterations: an alternative pathway to non-small-cell lung cancer." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2005): 154-64
58. Koga T, Hashimoto S, Sugio K, Yoshino I, Nakagawa K, Yonemitsu Y, Sugimachi K, Sueishi K "Heterogeneous distribution of P53 immunoreactivity in human lung adenocarcinoma correlates with MDM2 protein expression, rather than with P53 gene mutation." *International journal of cancer*(2001): 232-9
59. Kato S, Han S, Liu W, Otsuka K, Shibata H, Kanamaru R, Ishioka C "Understanding the function-structure and function-mutation relationships of p53 tumor suppressor protein by high-resolution missense mutation analysis." *Proceedings of the National Academy of Sciences of the United States of America*(2003): 8424-9
60. Houben R, Hesbacher S, Schmid C, Kauczok C, Flohr U, Haferkamp S, Müller C, Schrama D, Wischhusen J, Becker J "High-level expression of wild-type p53 in melanoma cells is frequently associated with inactivity in p53 reporter gene assays." *PLoS one*(2011): e22096
61. Olivier M, Petitjean A, Marcel V, Pétré A, Mounawar M, Plymoth A, de Fromental C, Hainaut P "Recent advances in p53 research: an interdisciplinary perspective." *Cancer gene therapy*(2009): 1-12
62. Chang Y, Wu C, Shih J, Lee Y "Comparison of p53 and epidermal growth factor receptor gene status between primary tumors and lymph node metastases in non-small cell lung cancers." *Annals of surgical oncology*(2011): 543-50
63. Jiang R, Zhang B, Teng X, Hu P, Xu S, Zheng Z, Liu R, Tang T, Ye F "Validating a targeted next-generation sequencing assay and profiling somatic variants in Chinese non-small cell lung cancer patients." *Scientific reports*(2020): 2070
64. Mattioli M, Soddu S, Prodosmo A, Visca P, Conti S, Alessandrini G, Facciolo F, Strigari L "Prognostic role of serum p53 antibodies in lung cancer." *BMC cancer*(2015): 148
65. Bircan A, Bircan S, Kapucuoglu N, Songur N, Ozturk O, Akkaya A "Maspin, VEGF and p53 expression in small biopsies of primary advanced lung cancer and relationship with clinicopathologic parameters." *Pathology oncology research : POR*(2010): 553-61
66. Kim Y, Hammerman P, Kim J, Yoon J, Lee Y, Sun J, Wilkerson M, Pedamallu C, Cibulskis K, Yoo Y, Lawrence M, Stojanov P, Carter S, McKenna A, Stewart C, Sivachenko A, Oh I, Kim H, Choi Y, Kim K, Shim Y, Kim K, Song S, Na K, Choi Y, Hayes D, Kim J, Cho S, Kim Y, Ahn J, Ahn M, Getz G, Meyerson M, Park K "Integrative and comparative genomic analysis of lung squamous cell carcinomas in East Asian patients." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2014): 121-8
67. Dong Z, Zhong W, Zhang X, Su J, Xie Z, Liu S, Tu H, Chen H, Sun Y, Zhou Q, Yang J, Yang X, Lin J, Yan H, Zhai H, Yan L, Liao R, Wu S, Wu Y "Potential Predictive Value of TP53 and KRAS Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2017): 3012-3024
68. Scheel A, Ansén S, Schultheis A, Scheffler M, Fischer R, Michels S, Hellmich M, George J, Zander T, Brockmann M, Stoelben E, Groen H, Timens W, Perner S, von Bergwelt-Baildon M, Büttner R, Wolf J "PD-L1 expression in non-small cell lung cancer: Correlations with genetic alterations." *Oncotarget*(2016): e1131379
69. Albitar M, Sudarsanam S, Ma W, Jiang S, Chen W, Funari V, Blocker F, Agersborg S "Correlation of MET gene amplification and TP53 mutation with PD-L1 expression in non-small cell lung cancer." *Oncotarget*(2018): 13682-13693
70. Mansuet-Lupo A, Alifano M, Pécuchet N, Biton J, Becht E, Goc J, Germain C, Ouakrim H, Régnard J, Cremer I, Laurent-Puig P, Dieu-Nosjean M, Blons H, Damotte D "Intratumoral Immune Cell Densities Are Associated with Lung Adenocarcinoma Gene Alterations." *American journal of respiratory and critical care medicine*(2016): 1403-1412
71. Kadara H, Choi M, Zhang J, Parra E, Rodriguez-Canales J, Gaffney S, Zhao Z, Behrens C, Fujimoto J, Chow C, Yoo Y, Kalhor N, Moran C, Rimm D, Swisher S, Gibbons D, Heymach J, Kaftan E, Townsend J, Lynch T, Schlessinger J, Lee J, Lifton R, Wistuba I, Herbst R "Whole-exome sequencing and immune profiling of early-stage lung adenocarcinoma with fully annotated clinical follow-up." *Annals of oncology : official journal of the European Society for Medical Oncology*(2017): 75-82
72. Van Egeren D, Kohli K, Warner J, Bedard P, Riely G, Lepisto E, Schrag D, LeNoue-Newton M, Catalano P, Kehl K, Michor F "Genomic analysis of early-stage lung cancer reveals a role for TP53 mutations in distant metastasis." *Scientific reports*(2022): 19055
73. Schuler P, Harasymczuk M, Visus C, Deleo A, Trivedi S, Lei Y, Argiris A, Gooding W, Butterfield L, Whiteside T, Ferris R "Phase I dendritic cell p53 peptide vaccine for head and neck cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2014): 2433-44
74. Vermeij R, Leffers N, van der Burg S, Melief C, Daemen T, Nijman H "Immunological and clinical effects of vaccines targeting p53-overexpressing malignancies." *Journal of biomedicine & biotechnology*(2011): 702146

References

75. Saito H, Ando S, Morishita N, Lee K, Dator D, Dy D, Shigemura K, Adhim Z, Nibu K, Fujisawa M, Shirakawa T "A combined lymphokine-activated killer (LAK) cell immunotherapy and adenovirus-p53 gene therapy for head and neck squamous cell carcinoma." *Anticancer research*(2014): 3365-70
76. Ma C, Janetka J, Piwnica-Worms H "Death by releasing the breaks: CHK1 inhibitors as cancer therapeutics." *Trends in molecular medicine*(2011): 88-96
77. Hirai H, Arai T, Okada M, Nishibata T, Kobayashi M, Sakai N, Imagaki K, Ohtani J, Sakai T, Yoshizumi T, Mizuarai S, Iwasawa Y, Kotani H "MK-1775, a small molecule Wee1 inhibitor, enhances anti-tumor efficacy of various DNA-damaging agents, including 5-fluorouracil." *Cancer biology & therapy*(2010): 514-22
78. Bridges K, Hirai H, Buser C, Brooks C, Liu H, Buchholz T, Molkenkine J, Mason K, Meyn R "MK-1775, a novel Wee1 kinase inhibitor, radiosensitizes p53-defective human tumor cells." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 5638-48
79. Vilgelm A, Pawlikowski J, Liu Y, Hawkins O, Davis T, Smith J, Weller K, Horton L, McClain C, Ayers G, Turner D, Essaka D, Stewart C, Sosman J, Kelley M, Ecsedy J, Johnston J, Richmond A "Mdm2 and aurora kinase a inhibitors synergize to block melanoma growth by driving apoptosis and immune clearance of tumor cells." *Cancer research*(2015): 181-93
80. Li Z, Sun Y, Chen X, Squires J, Nowroozizadeh B, Liang C, Huang J "p53 Mutation Directs AURKA Overexpression via miR-25 and FBXW7 in Prostatic Small Cell Neuroendocrine Carcinoma." *Molecular cancer research : MCR*(2015): 584-91
81. Katayama H, Sen S "Functional significance of Aurora kinase A regulatory interactions with p53-ERα complex in human breast cancer cells." *Hormones & cancer*(2011): 117-24
82. Tentler J, Ionkina A, Tan A, Newton T, Pitts T, Glogowska M, Kabos P, Sartorius C, Sullivan K, Espinosa J, Eckhardt S, Diamond J "p53 Family Members Regulate Phenotypic Response to Aurora Kinase A Inhibition in Triple-Negative Breast Cancer." *Molecular cancer therapeutics*(2015): 1117-29
83. Gully C, Velazquez-Torres G, Shin J, Fuentes-Mattei E, Wang E, Carlock C, Chen J, Rothenberg D, Adams H, Choi H, Guma S, Phan L, Chou P, Su C, Zhang F, Chen J, Yang T, Yeung S, Lee M "Aurora B kinase phosphorylates and instigates degradation of p53." *Proceedings of the National Academy of Sciences of the United States of America*(2012): E1513-22
84. Marxer M, Ma H, Man W, Poon R "p53 deficiency enhances mitotic arrest and slippage induced by pharmacological inhibition of Aurora kinases." *Oncogene*(2014): 3550-60
85. El-Deiry W "The role of p53 in chemosensitivity and radiosensitivity." *Oncogene*(2003): 7486-95
86. Miyasaka A, Oda K, Ikeda Y, Sone K, Fukuda T, Inaba K, Makii C, Enomoto A, Hosoya N, Tanikawa M, Uehara Y, Arimoto T, Kuramoto H, Wada-Hiraike O, Miyagawa K, Yano T, Kawana K, Osuga Y, Fujii T "PI3K/mTOR pathway inhibition overcomes radioresistance via suppression of the HIF1-α/VEGF pathway in endometrial cancer." *Gynecologic oncology*(2015): 174-80