

ACT Onco[®] + Report

PATIENT		
Identifier: 洪睿承		Patient ID: 45543042
Date of Birth: Feb 21, 2007		Gender: Male
Diagnosis: Osteosarcoma		
ORDERING PHYSICIAN		
Name: 顏秀如醫師		Tel: 886-228712121
Facility: 臺北榮總		
Address: 臺北市北投區石牌路二段 201 號		
SPECIMEN		
Specimen ID: S11226789B	Collection site: Lung	Type: FFPE tissue
Date received: Jun 26, 2023	Lab ID: AA-23-04122	D/ID: NA

ABOUT ACT Onco[®]+

The test is a next-generation sequencing (NGS)-based assay developed for efficient and comprehensive genomic profiling of cancers. This test interrogates coding regions of 440 genes associated with cancer treatment, prognosis and diagnosis. Genetic mutations detected by this test include small-scale mutations like single nucleotide variants (SNVs), small insertions and deletions (InDels) (≤ 15 nucleotides) and large-scale genomic alterations like copy number alterations (CNAs). The test also includes an RNA test, detecting fusion transcripts of 13 genes.

SUMMARY FOR ACTIONABLE VARIANTS

VARIANTS/BIOMARKERS WITH EVIDENCE OF CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Probable Effects in Patient's Cancer Type		Probable Sensitive in Other Cancer Types
	Sensitive	Resistant	
Not detected			

VARIANTS/BIOMARKERS WITH POTENTIAL CLINICAL SIGNIFICANCE

Genomic Alterations/Biomarkers	Possibly Sensitive	Possibly Resistant
CCNE1 Amplification	-	Palbociclib, Trastuzumab
KDR Amplification	Pazopanib, Sunitinib	-
KIT Amplification	-	Imatinib, Nilotinib, Sunitinib
PDGFRA Amplification	Imatinib, Pazopanib, Sorafenib, Sunitinib	-

Note:

- The above summary tables present genomic variants and biomarkers based on the three-tiered approach proposed by US FDA for reporting tumor profiling NGS testing. "Variants/biomarkers with evidence of clinical significance" refers to mutations that are widely recognized as standard-of-care biomarkers (FDA level 2/AMP tier 1). "Variants/biomarkers with potential clinical significance" refers to mutations that are not included in the standard of care but are informational for clinicians, which are commonly biomarkers used as inclusion criteria for clinical trials (FDA level 3/AMP tier 2).
- The therapeutic agents and possible effects to a given drug are based on mapping the variants/biomarkers with ACT Genomics clinical knowledge database. The mapping results only provide information for reference, but not medical recommendation.
- Please refer to corresponding sections for more detailed information about genomic alteration and clinical relevance listed above.

ACTOnco® + Report

TESTING RESULTS

VARIANT(S) WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Allele Frequency
Not detected		

- Copy Number Alterations

Chromosome	Gene	Variation	Copy Number
Chr17	TP53	Homozygous deletion	0
Chr13	BRCA2	Heterozygous deletion	1
Chr16	PALB2	Heterozygous deletion	1
Chr17	NF1	Heterozygous deletion	1
Chr19	STK11	Heterozygous deletion	1
Chr3	ATR	Heterozygous deletion	1
Chr5	RAD50	Heterozygous deletion	1
Chr4	KDR, KIT	Amplification	7
Chr19	CCNE1	Amplification	11
Chr4	PDGFRA	Amplification	16

- Fusions

Fusion Gene & Exon	Transcript ID
No fusion gene detected in this sample	

- Immune Checkpoint Inhibitor (ICI) Related Biomarkers

Biomarker	Results
Tumor Mutational Burden (TMB)	2.6 muts/Mb
Microsatellite Instability (MSI)	Microsatellite stable (MSS)

Note:

- Loss of heterozygosity (LOH) information was used to infer tumor cellularity. Copy number alteration in the tumor was determined based on 53% tumor purity.
- For more therapeutic agents which are possibly respond to heterozygous deletion of genes listed above, please refer to APPENDIX for more information.
- TMB was calculated by using the sequenced regions of ACTOnco®+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The threshold for high mutation load is set at ≥ 7.5 mutations per megabase. TMB, microsatellite status and gene copy number deletion cannot be determined if calculated tumor purity is $< 30\%$.

ACT Onco[®] + Report

THERAPEUTIC IMPLICATIONS TARGETED THERAPIES

Genomic Alterations	Therapies	Effect
Level 3A		
<i>KIT</i> Amplification	Imatinib, Nilotinib, Sunitinib	resistant
Level 4		
<i>KDR</i> Amplification	Pazopanib, Sunitinib	sensitive
<i>PDGFRA</i> Amplification	Imatinib, Pazopanib, Sorafenib, Sunitinib	sensitive
<i>CCNE1</i> Amplification	Palbociclib, Trastuzumab	resistant

Therapies associated with benefit or lack of benefit are based on biomarkers detected in this tumor and published evidence in professional guidelines or peer-reviewed journals.

Level	Description
1	FDA-recognized biomarkers predictive of response or resistance to FDA approved drugs in this indication
2	Standard care biomarkers (recommended by the NCCN guideline) predictive of response or resistance to FDA approved drugs in this indication
3A	Biomarkers predictive of response or resistance to therapies approved by the FDA or NCCN guideline in a different cancer type
3B	Biomarkers that serve as inclusion criteria for clinical trials (minimal supportive data required)
4	Biomarkers that show plausible therapeutic significance based on small studies, few case reports, or preclinical studies

ACT Onco[®] + Report

IMMUNE CHECKPOINT INHIBITORS (ICIs)

No genomic alterations detected to confer sensitivity or lack of benefit to immune checkpoint therapies.

- Other Biomarkers with Potential Clinical Effects for ICIs

Genomic Alterations	Potential Clinical Effects
Not detected	

Note: Tumor non-genomic factors, such as patient germline genetics, PDL1 expression, tumor microenvironment, epigenetic alterations or other factors not provided by this test may affect ICI response.

CHEMOTHERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to chemotherapies.

HORMONAL THERAPIES

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to hormonal therapies.

OTHERS

No genomic alterations detected in this tumor predicted to confer sensitivity or lack of benefit to other therapies.

Note:

Therapeutic implications provided in the test are based solely on the panel of 440 genes sequenced. Therefore, alterations in genes not covered in this panel, epigenetic and post-transcriptional and post-translational factors may also determine a patient's response to therapies. In addition, several other patient-associated clinical factors, including but not limited to, prior lines of therapies received, dosage and combinations with other therapeutic agents, patient's cancer types, sub-types, and/or stages, may also determine the patient's clinical response to therapies.

ACT Onco[®] + Report

VARIANT INTERPRETATION

ATR Heterozygous deletion

Biological Impact

Ataxia Telangiectasia and Rad3-related protein (ATR) gene encodes a serine/threonine kinase that is involved in the DNA damage response. ATR plays as a central coordinator of the DNA damage response (DDR) by responding to single-stranded regions of the DNA^{[1][2]} and the maintenance of genome stability^[3]. ATR has also been implicated as a haploinsufficient tumor suppressor with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[4][5]}. Germline mutation of ATR is associated with cancer predisposition and Seckel syndrome, a condition associated with CNS disorders^{[6][7]}. Somatic mutations of ATR are associated with microsatellite instability and are found in colorectal cancer^[8], urothelial cancer^[9], gastric cancer^[10], endometrial cancer^[11] and myelomas^[12].

Therapeutic and prognostic relevance

Talazoparib is FDA-approved for treating mCRPC patients harboring mutations in homologous recombination repair (HRR) genes, including ATR.

In a clinical study, a metastatic castration-resistant prostate cancer patient harboring deleterious mutation in the ATR gene (K2106fs) had a PSA remission of 62% and PSA-PFS of 13 months by olaparib treatment^[13].

ATR has been determined as an inclusion criterion for the trials evaluating olaparib efficacy in ovarian cancer^[14] and advanced solid tumors (NCT03297606; CAPTUR trial), rucaparib efficacy in ovarian cancer^[15], niraparib efficacy in pancreatic cancer (NCT03553004), and any malignancy, except prostate (NCT03207347), and talazoparib efficacy in HER2-negative solid tumors (NCT02401347), prostate cancer (NCT03148795), and lung cancer (NCT03377556), respectively.

BRCA2 Heterozygous deletion

Biological Impact

The BRCA2 gene encodes a tumor suppressor involved in the homologous recombination pathway for double-strand DNA repair^[16]. BRCA2 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[17]. BRCA2 germline mutations confer an increased lifetime risk of developing breast, ovarian, prostate and pancreatic cancer, limited reports of related gastric cancer, and Fanconi anemia subtype D1-associated risk of brain cancer, medulloblastoma, pharyngeal cancer, chronic lymphocytic leukemia and acute myeloid leukemia^[18]. Somatic mutations in BRCA2 are highest in colorectal, non-small cell lung cancer (NSCLC), and ovarian cancers^[19].

Therapeutic and prognostic relevance

Multiple PARP inhibitors, including olaparib, rucaparib, niraparib, and talazoparib, have been approved by the U.S. FDA for the treatment of cancer. Olaparib is approved for multiple settings in advanced ovarian cancer, metastatic breast cancer with BRCA mutations, metastatic pancreatic cancer, and mCRPC with BRCA mutation or HRR gene mutations, including BRCA. Rucaparib is approved for maintenance treatment of recurrent ovarian cancer with BRCA mutations and mCRPC with BRCA mutations. Niraparib is approved for maintenance treatment of advanced ovarian cancer and recurrent ovarian cancer with BRCA mutations. Talazoparib is approved for locally advanced or metastatic breast cancer with BRCA mutations and mCRPC with HRR gene mutations, including BRCA.

According to the NCCN guidelines, rucaparib is recommended as recurrence therapy for patients with BRCA-mutated ovarian cancer who have been treated with multiple lines of chemotherapy. It is also recommended as maintenance therapy for patients with metastatic pancreatic cancer who have undergone prior platinum-based therapy and harbor germline or somatic BRCA mutations. Additionally, niraparib is recommended as maintenance therapy for ovarian

ACT Onco[®] + Report

cancer patients with BRCA mutations.

CCNE1 Amplification

Biological Impact

The CCNE1 gene encodes the cyclin E1 protein, a regulator of the cell cycle that activates the cyclin-dependent protein kinase 2 (CDK2) and plays a role in regulating cells' transition from G1 to S phase and the maintenance of genomic stability^[20]. Increasing in cyclin E1 level, either by gene amplification or overexpression, is found in a diverse range of cancers and can be indicative of poor prognosis^[21].

Therapeutic and prognostic relevance

There are no FDA-approved therapies targeting cyclin E1 currently available^[22]. Dinaciclib, a CDK1/2 specific inhibitor, is currently under clinical evaluation^[23]. A combination of dinaciclib, a small molecule CDK2 inhibitor, and AKT inhibitors that may selectively target patients with CCNE1-amplified high-grade serous ovarian cancer (HGSC) in preclinical setting^[24]. A preclinical study in breast cancer cell lines showed that amplification of CCNE1 is associated with acquired resistance to CDK4/6 inhibition by palbociclib^[25]. A study of HER2-amplified breast cancer patients indicated that amplification of CCNE1 was associated with trastuzumab resistance and shorter progression-free survival^[26].

There are retrospective study and meta-analysis demonstrated that amplification and overexpression of CCNE1 are associated with poor survival in cancer patients^{[27][28]}. From the result of PALOMA-3 phase III trial, pre-treated hormone receptor-positive/HER2-negative metastatic breast cancer patients were resistant to palbociclib treatment when CCNE1 was highly expressed (median PFS: CCNE1 high, 7.6 months; CCNE1 low, 14.1 months)^[29]. CCNE1 amplification has been selected as an inclusion criteria for the trial examining palbociclib in malignant solid tumor (NCT02896335, NCT03155620, NCT01037790, NCT03526250).

KDR Amplification

Biological Impact

KDR (kinase domain receptor), also known as VEGFR2 or Flk-1, is a tyrosine kinase receptor for the vascular endothelial growth factor (VEGF) and involves in angiogenesis pathway^[30]. Binding of VEGF to KDR results in activation of phospholipase C (PLC-gamma) and downstream signaling via protein kinase C (PKC) and RAF/MEK/ERK^[31]. Mutations of KDR are rare in tumors, and alterations of KDR activity typically occur via KDR amplification and subsequent overexpression^{[32][33]}.

Therapeutic and prognostic relevance

To date, there are four VEGF inhibitors (sorafenib, sunitinib, pazopanib, bevacizumab) and one VEGFR2 inhibitor cabozantinib that are FDA-approved for the treatment of cancers^{[34][35][36]}. Notably, a case report showed that an angiosarcoma patient with concurrent KDR and FLT4 amplification developed a progressive disease when treated with sorafenib, but experienced a potent antitumor response and achieved clinically stable disease for 6 months after receiving pazopanib therapy^[37]. Besides, an angiosarcoma patient with upregulated VEGFR2 responded to sunitinib treatment^[38]. VEGFR2 inhibitors like apatinib and vandetanib are in early clinical phase trial^{[39][40][41]}.

KDR amplification and/or mutation has been selected as an inclusion criteria for the trial examining cabozantinib in metastatic castrate resistant prostate cancer (mCRPC) (NCT04631744) and sunitinib in malignant solid tumors (NCT03297606). The increased copy number of KIT or KDR significantly correlated with a worse 5-year breast cancer-specific survival (BCSS) in triple-negative breast cancer (TNBC) patients^[42].

ACT Onco[®] + Report

KIT Amplification

Biological Impact

KIT is a proto-oncogene that encodes a type 3 transmembrane receptor tyrosine kinase. Activation of KIT through dimerization and autophosphorylation upon binding by its ligand results in increased intracellular PI3K/AKT/mTOR, MAPK/ERK and JAK/STAT signaling pathways to promote cell proliferation and survival^[43]. KIT activating mutations are frequently found in 80 - 90% of gastrointestinal stromal tumors (GISTs) which distributed over multiple exons with different frequencies (exons 11 (66.1%), exon 9 (13%), exon 13 (1.2%), and exon 17 (0.6%))^{[44][45]}.

Therapeutic and prognostic relevance

NCCN guidelines recommend using KIT inhibitors (imatinib, sunitinib, nilotinib) for cutaneous melanoma patients with specific KIT hotspot mutations in exon 11 and exon 13, but not for those with KIT exon 17 mutations or amplification. Dasatinib and ripretinib are recommended for metastatic or unresectable cutaneous melanoma patients with KIT activating mutations. Also, imatinib is a preferred regimen for neoadjuvant therapy of resectable GIST with significant morbidity patient harboring KIT mutation, and ponatinib is recommended for advanced GIST patients with KIT exon 11 mutations.

KIT mutation has been determined as an inclusion criterion for the trials evaluating dasatinib, avapritinib, sunitinib, ponatinib, regorafenib, ripretinib, imatinib, and cabozantinib efficacies in various types of solid tumors (NCT03297606, NCT03353753, NCT03465722, NCT02272998, NCT02501551, NCT03353753, NCT02461849, NCT02712112, NCT04631744, NCT04116541).

The efficacy of U.S. FDA-approved KIT TKIs such as imatinib, sunitinib, regorafenib, and ripretinib for GIST strongly depends on the location of the activating KIT mutations^{[46][47][48][49][50][51]}. Patients with GIST harboring KIT exon 9 mutations have intermediate sensitivity to imatinib and better survival than those with KIT exon 11 mutations^[47]. Newly developing agents, including avapritinib, show potential to be better inhibitors for clinically relevant KIT/PDGFR mutations in GIST^[52]. Ponatinib and dasatinib show promise in GIST patients with KIT exon 11 mutations, with a disease control rate of 67% and partial control rate of 32%(DOI:10.1200/jco.2011.29.15_suppl.10006)^[53]. Meanwhile, a Phase II trial involving melanoma showed a 38.5% response rate to nilotinib in patients with KIT exon 11 mutations^[54].

Both KIT and PDGFRA overexpression were associated with high tumor grade, high proliferation index, and poor outcome in patients with the serous type of ovarian carcinoma^[55].

A phase II trial of imatinib in melanoma showed that patients with KIT amplification had lower disease control rate compared with patients carrying KIT mutations (18% amplified vs. 77% mutated)^[56].

NF1 Heterozygous deletion

Biological Impact

The neurofibromin 1 (NF1) gene encodes a GTPase activating protein (GAP) which is an important negative regulator of the Ras cellular proliferation pathways^{[57][58][59][60]}. Besides, NF1 also physically interacts with the N-terminal domain of focal adhesion kinase (FAK) and involves in the regulation of cell adhesion, growth, and other pathways^{[61][62]}. NF1 is considered a classical haploinsufficient tumor suppressor gene with loss of one allele through inherited or acquired mutation may lead to reduced protein expression and is insufficient to execute normal cellular functions contributing to tumor development^{[63][64][65][66][67]}. NF1 syndrome is a germline condition resulting in a predisposition to several types of cancer such as neurofibromas, melanoma, lung cancer, ovarian cancer, breast cancer, colorectal cancer, hematological malignancies^{[68][69][70]}. Meanwhile, sporadic NF1 mutations have been observed in multiple cancer types^[71], including myelodysplastic syndromes, melanomas, colon cancer^[72], glioblastomas^[73], lung cancer^[74], ovarian cancer, and breast cancer^[68].

ACT Onco® + Report

Therapeutic and prognostic relevance

Selumetinib is FDA-approved for treating pediatric patients 2 years of age and older with neurofibromatosis type 1 (NF1) who have symptomatic, inoperable plexiform neurofibromas (PN).

In the NCCN guidelines for CNS cancers, selumetinib is recommended as a treatment option for recurrent or progressive NF-1 mutated glioma patient.

NF1 mutation/ loss has been determined as an inclusion criterion for the trials evaluating selumetinib efficacy in solid tumors (NCT02664935, NCT03155620)^[75].

NF1 depletion is associated with drug resistance to various inhibitors, such as RAF, EGFR, tamoxifen, and retinoic acid^{[71][76]}. Loss of NF1 in lung adenocarcinomas, colorectal cancer, and BRAF-mutated melanomas is associated with resistance to anti-EGFR and BRAF inhibitors^{[77][78][79][80][81][82]}. NF1 loss contributes to trastuzumab resistance in HER2-positive metastatic gastric cancer, but a combination of HER2 and MEK/ERK inhibitors may overcome this resistance^[83]. Trametinib is effective in treating neurofibromatosis type I-associated glioblastoma^[84]. Patients with mutations in the mTOR pathway, including NF1, have responded to everolimus^[85]. However, a patient with metastatic lung cancer harboring CCDC6-ROS1 and NF1 truncating mutation treated with crizotinib had a short overall survival of one month^[86].

NF1 depletion has been linked to drug resistance to several inhibitors in vitro, including RAF, EGFR, and trastuzumab. However, adding MEK inhibitors could restore sensitivity to erlotinib^[77], and MEK and mTOR inhibitors showed promise in NF1-deficient tumors^{[87][88]}. Knockdown of NF1 also led to resistance to crizotinib and cabozantinib treatment in ROS1 fusion-positive cells^[86].

PALB2 Heterozygous deletion

Biological Impact

The partner and localizer of BRCA2 (PALB2) gene encodes a protein that plays a critical role in homologous recombination repair (HRR) through its ability to interact with BRCA2 in nuclear foci, promoting its localization and stability in key nuclear structures^[89]. The Fanconi anemia complementation group (FANC) which includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM and FANCN (also called PALB2) are involved in the repair of DNA double-strand breaks (DSBs) by homologous recombination (HR)^{[90][91][92]}. PALB2 is a haploinsufficient tumor suppressor gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological function^[93]. Biallelic germline loss-of-function mutations in PALB2 cause Fanconi anemia, whereas monoallelic loss-of-function mutations are associated with an increased risk of breast cancer and pancreatic cancer^[94]. Fanconi Anemia is an autosomal recessive disease characterized by hematological abnormalities, bone marrow failure, limb deformities, skin hyperpigmentation, and susceptibility to hematologic and solid malignancies, such as acute myeloid leukemia and head and neck carcinoma^{[95][96]}.

Therapeutic and prognostic relevance

Olaparib and talazoparib are FDA-approved for treating mCRPC patients harboring mutations in homologous recombination repair (HRR) genes, including PALB2.

The NCCN guidelines recommend rucaparib as maintenance therapy for pancreatic adenocarcinoma patients with PALB2 mutations after platinum-based therapy. For breast cancer patients with stage IV disease and germline PALB2 alterations, olaparib treatment is recommended by the NCCN guidelines regardless of subtype.

PALB2 mutation has been determined as an inclusion criterion for the trials evaluating rucaparib, niraparib, and talazoparib efficacies in various types of solid tumors (NCT02401347, NCT03553004, NCT02952534)^{[15][97]}.

ACT Onco[®] + Report

A case report demonstrated an exceptional response to mitomycin C and cisplatin treatment in a gemcitabine-resistant pancreatic cancer patient with biallelic inactivation of PALB2^[98].

PDGFRA Amplification

Biological Impact

The PDGFRA gene encodes for the protein platelet-derived growth factor alpha (PDGFRA). The Ligand binding to the extracellular domain of PDGFRA induces receptor dimerization, enabling autophosphorylation of specific tyrosine residues and subsequently results in the activation of downstream pathways such as RAS-MAPK, PI3K and PLC-γ that are involved in developmental and cellular responses^{[99][100]}. Mutations, insertions, deletions, fusions and genomic amplification of PDGFRA lead to its activation in several tumor types: ~7% of gastrointestinal stromal tumors (GISTs) have PDGFRA activating mutations and these mutations are mutually exclusive from KIT mutations^[101]; activating mutations in PDGFRA have been observed in ~5% of Chinese melanoma patients^[102]; amplification of PDGFRA is the second most frequent receptor tyrosine kinase amplification in glioblastoma (GBM)^{[103][104][105][73][106]}, intimal sarcomas^[107], malignant peripheral nerve sheath tumors^[108], non-small cell lung adenocarcinomas and non-small cell lung squamous cell carcinomas^[109].

Therapeutic and prognostic relevance

A retrospective study showed that either KIT, PDGFRA, or EGFR amplification in glioma at the time of the first diagnosis was associated with an unfavorable overall survival^[110].

In a preclinical study, a PDGFRA-amplified cell line was sensitive to imatinib, sunitinib and sorafenib treatment, demonstrated by disruption of downstream signaling and reduced cell viability in vitro^[111]. Another study also showed that pazopanib could inhibit tumor growth in the PDGFRA-amplified pleomorphic liposarcoma xenograft mouse model^[112].

RAD50 Heterozygous deletion

Biological Impact

The RAD50 gene encodes a highly-conserved DNA double-strand break (DSB) repair factor. It forms MRN complex with NBS1 and MRE11 protein and is involved in sensing and early processing of DSB, cell cycle checkpoints, DNA recombination and maintenance of telomeres^{[113][114]}. Mutations in the components of the MRN complex could increase susceptibility to familial breast cancer^{[115][116]}, gastric cancer^[117], colorectal cancer^[118], and urothelial cancer^[119]. RAD50 has been implicated as a haploinsufficient gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^[120]. Besides, RAD50 deletion was also suggested as a marker of BRCAness, a phenotype shared between non-BRCA1/2-mutated ovarian cancers and BRCA1/2-mutated ovarian cancers^[121].

Therapeutic and prognostic relevance

Preclinical data showed that knockdown of the RAD50 gene in ovarian cancer cell lines was significantly associated with better responses to two PARP inhibitors, olaparib and rucaparib^[121]. RAD50 has been selected as an inclusion criterion for the trials examining talazoparib efficacy in HER2-negative breast cancer, olaparib efficacy in breast cancer, rucaparib efficacy in metastatic prostate cancer and niraparib efficacy in any malignancy (except prostate) (NCT02401347, NCT03207347, NCT03344965, NCT03413995).

STK11 Heterozygous deletion

Biological Impact

The serine/threonine kinase 11 (STK11, also known as LKB1) gene encodes the multifunctional serine/threonine kinase, a tumor suppressor that functions as an inhibitor for the mTOR signaling pathway^{[122][123]}. STK11 is a haploinsufficient

ACT Onco[®] + Report

gene with one copy loss may lead to weak protein expression and is insufficient to execute its original physiological functions^{[124][125]}. In the mouse model, loss of STK11 promotes aggressive endometrial and squamous cell carcinomas^{[126][127]}. Mutations in STK11 have been found in lung, breast, cervical, testicular, and liver cancers, as well as malignant melanoma, pancreatic and biliary carcinoma^[128]. Germline mutations in STK11 are found in 30-70% of Peutz-Jeghers syndrome^[129].

Therapeutic and prognostic relevance

A clinical study in a pancreatic cancer patient with Peutz-Jeghers syndrome whose tumor harboring an STK11 D194E mutation coupled with the loss of heterozygosity of the other STK11 allele displayed partial response to the everolimus treatment^[130]. In another clinical case study, an adrenocorticotrophic pituitary carcinoma patient whose tumor bearing an STK11 inactivating mutation responded to a combination of everolimus and radiotherapy^[131].

Preclinical data suggested that lung cancer cell lines with STK11 inactivating mutations may confer increased sensitivity to the MEK-1 and MEK-2 inhibitor, trametinib^[132].

Inactivating mutations of STK11 was shown to be associated with resistance to immune checkpoint blockade in KRAS-mutant lung adenocarcinoma (LUAC) and NSCLC (DOI: 10.1200/JCO.2017.35.15_suppl.9016)^{[133][134][135]}. It was proposed that loss of STK11 negatively impacts the number and function of tumor-infiltrating T cells (TILs) and PD-L1 expression on tumor cells and therefore results in an ineffective response to PD-1-targeting antibodies^[136].

TP53 Homozygous deletion

Biological Impact

TP53 encodes the p53 protein, a crucial tumor suppressor that orchestrates essential cellular processes including cell cycle arrest, senescence and apoptosis^[137]. TP53 is a proto-typical haploinsufficient gene, such that loss of a single copy of TP53 can result in tumor formation^[138].

Therapeutic and prognostic relevance

Despite having a high mutation rate in cancers, there are currently no approved targeted therapies for TP53 mutations. A phase II trial demonstrated that Wee1 inhibitor (AZD1775) in combination with carboplatin was well tolerated and showed promising anti-tumor activity in TP53-mutated ovarian cancer refractory or resistant (< 3 months) to standard first-line therapy (NCT01164995)^[139].

In a retrospective study (n=19), advanced sarcoma patients with TP53 loss-of-function mutations displayed improved progression-free survival (208 days versus 136 days) relative to patients with wild-type TP53 when treated with pazopanib^[140]. Results from another Phase I trial of advanced solid tumors (n=78) demonstrated that TP53 hotspot mutations are associated with better clinical response to the combination of pazopanib and vorinostat^[141].

Advanced solid tumor and colorectal cancer patients harboring a TP53 mutation have been shown to be more sensitive to bevacizumab when compared with patients harboring wild-type TP53^{[142][143][144]}. In a pilot trial (n=21), TP53-negative breast cancer patients demonstrated increased survival following treatment with bevacizumab in combination with chemotherapy agents, Adriamycin (doxorubicin) and Taxotere (docetaxel)^[145]. TP53 mutations were correlated with poor survival of advanced breast cancer patients receiving tamoxifen or primary chemotherapy^{[146][147]}. In a retrospective study of non-small cell lung cancer (NSCLC), TP53 mutations were associated with high expression of VEGF-A, the primary target of bevacizumab, offering a mechanistic explanation for why patients exhibit improved outcomes after bevacizumab treatment when their tumors harbor mutant TP53 versus wild-type TP53^[148].

ACT Onco[®] + Report

US FDA-APPROVED DRUG(S)

Binimetinib (MEKTOVI)

Binimetinib is an oral kinase inhibitor that targets MEK. Binimetinib is developed and marketed by Array BioPharma under the trade name MEKTOVI.

- FDA Approval Summary of Binimetinib (MEKTOVI)

MEKTOVI ^[149] NCT01909453	Melanoma (Approved on 2018/06/27)
	BRAF V600E/K
	Encorafenib + binimetinib vs. Vemurafenib [PFS(M): 14.9 vs. 7.3]

Cobimetinib (COTELLIC)

Cobimetinib is a reversible inhibitor which targets MEK1 and MEK2. Cobimetinib is developed by Exelixis and Genentech, and marketed by Genentech under the trade name COTELLIC.

- FDA Approval Summary of Cobimetinib (COTELLIC)

coBRIM ^[150] NCT01689519	Melanoma (Approved on 2015/11/10)
	BRAF V600E/K
	Cobimetinib + vemurafenib vs. Placebo + vemurafenib [PFS(M): 12.3 vs. 7.2]

Everolimus (AFINITOR)

Everolimus, a derivative of sirolimus, works as an inhibitor of mammalian target of rapamycin complex 1 (mTORC1) and blocks mTORC1-mediated downstream signals for cell growth, proliferation, and survival. Everolimus is developed and marketed by Novartis under the trade name AFINITOR.

- FDA Approval Summary of Everolimus (AFINITOR)

RADIANT-4 ^[151] NCT01524783	Lung or gastrointestinal neuroendocrine tumor (Approved on 2016/02/26)
	-
	Everolimus vs. Placebo [PFS(M): 11 vs. 3.9]
BOLERO-2 ^[152] NCT00863655	Breast cancer (Approved on 2012/07/20)
	ER+/HER2-
	Everolimus + exemestane vs. Placebo + exemestane [PFS(M): 7.8 vs. 3.2]
EXIST-2 NCT00790400	Tuberous sclerosis complex (tsc)-associated renal angiomyolipoma (Approved on 2012/04/26)
	-
	Everolimus vs. Placebo [ORR(%): 41.8 vs. 0]
RADIANT-3 ^[153] NCT00510068	Pancreatic neuroendocrine tumor (Approved on 2011/05/05)
	-
	Everolimus vs. Placebo [PFS(M): 11 vs. 4.6]
EXIST-1 ^[154] NCT00789828	Subependymal giant cell astrocytoma (Approved on 2010/10/29)
	-
	Everolimus vs. Placebo [ORR(%): 35.0]

ACT Onco[®] + Report

RECORD-1 ^[155] NCT00410124	Renal cell carcinoma (Approved on 2009/05/30)
	-
	Everolimus vs. Placebo [PFS(M): 4.9 vs. 1.9]

Imatinib (GLEEVEC)

Imatinib is an oral, small molecule inhibitor of tyrosine kinase enzymes, namely, the Abelson proto-oncogene (ABL), c-KIT, and platelet-derived growth factor receptor (PDGFR). Imatinib is developed and marketed by Novartis under the trade name GLEEVEC.

- FDA Approval Summary of Imatinib (GLEEVEC)

[156] NCT00022737	Acute lymphocytic leukemia (Approved on 2013/01/25)
	-
	Imatinib [EFS(%): 70]
	Gastrointestinal stromal tumor (Approved on 2012/01/31)
	KIT positive
	Imatinib [RFS(%): 42 (imatinib for 12) 25 (imatinib for 36)]
	Gastrointestinal stromal tumor (Approved on 2009/02/10)
	KIT+
	Imatinib vs. Placebo [RFS(%): 21 vs. 28]
	Myelodysplastic myeloproliferative cancer (Approved on 2006/10/19)
	-
	Imatinib [MCyR(%): 39, CHR(%): 45]
[157]	Acute lymphocytic leukemia (Approved on 2006/10/19)
	Ph+
	Imatinib [MCyR(%): 35, CHR(%): 19]
	Dermatofibrosarcoma protuberans (Approved on 2006/10/19)
	-
	Imatinib [ORR(%): 83.0]
	Systemic mastocytosis (Approved on 2006/10/19)
	-
	Imatinib [CHR(%): 29]
	Chronic eosinophilic leukemia (Approved on 2006/10/19)
	-
	Imatinib [CHR(%): 61]
[158] NCT00471497	Chronic myeloid leukemia (Approved on 2003/05/20)
	Ph+
	Imatinib vs. Nilotinib [MMR(%): 22 vs. 44]
[159] NCT00333840	Chronic myeloid leukemia (Approved on 2003/04/18)
	-
	Imatinib vs. Interferon-α+ cytarabine [PFS(%): 81.2 vs. 60.6]
[160] NCT00009906	Gastrointestinal stromal tumor (Approved on 2002/02/01)
	-
	Imatinib [PFS(M): 18.9 (imatinib 400 mg) 23.2 (imatinib 800 mg)]

ACT Onco® + Report

Niraparib (ZEJULA)

Niraparib is an oral, small molecule inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1 and -2 (PARP-1, -2). Niraparib is developed and marketed by Tesaro under the trade name ZEJULA.

- FDA Approval Summary of Niraparib (ZEJULA)

PRIMA NCT02655016	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2020/04/29)
	- Niraparib vs. Placebo [PFS (overall population)(M): 13.8 vs. 8.2]
NOVA^[161] NCT01847274	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/03/27)
	- Niraparib vs. Placebo [PFS (overall population)(M): 11.3 vs. 4.7]

Olaparib (LYNPARZA)

Olaparib is an oral, small molecule inhibitor of poly (ADP-ribose) polymerase-1, -2, and -3 (PARP-1, -2, -3). Olaparib is developed by KuDOS Pharmaceuticals and marketed by AstraZeneca under the trade name LYNPARZA.

- FDA Approval Summary of Olaparib (LYNPARZA)

PROpel NCT03732820	Prostate cancer (Approved on 2023/05/31)
	BRCA mutation Olaparib + abiraterone + prednisone vs. Placebo + abiraterone + prednisone [rPFS(M): not reached vs. 8]
OlympiA NCT02032823	HER2-negative high-risk early breast cancer (Approved on 2022/03/11)
	HER2-/gBRCA mutation Olaparib vs. Placebo [invasive disease-free survival (IDFS)(M):]
PROfound^[162] NCT02987543	Prostate cancer (Approved on 2020/05/19)
	HRR genes mutation Olaparib vs. Enzalutamide or abiraterone acetate [PFS(M): 5.8 vs. 3.5]
PAOLA-1^[163] NCT02477644	Ovarian cancer (Approved on 2020/05/08)
	HRD+ Olaparib + bevacizumab vs. Placebo + bevacizumab [PFS(M): 37.2 vs. 17.7]
POLO^[164] NCT02184195	Pancreatic adenocarcinoma (Approved on 2019/12/27)
	gBRCA mutation Olaparib vs. Placebo [ORR(%): 23.0 vs. 12.0, PFS(M): 7.4 vs. 3.8]
SOLO-1^[165] NCT01844986	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/12/19)
	gBRCA mutation or sBRCA mutation Olaparib vs. Placebo [PFS(M): NR vs. 13.8]
OlympiAD^[166] NCT02000622	Breast cancer (Approved on 2018/02/06)
	HER2-/gBRCA mutation Olaparib vs. Chemotherapy [PFS(M): 7 vs. 4.2]
SOLO-2/ENGOT-Ov21^[167] NCT01874353	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	gBRCA mutation Olaparib vs. Placebo [PFS(M): 19.1 vs. 5.5]
Study19^[168] NCT00753545	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2017/08/17)
	- Olaparib vs. Placebo [PFS(M): 8.4 vs. 4.8]

ACT Onco[®] + Report

Pazopanib (VOTRIENT)

Pazopanib is an oral, small molecule, multi-kinase inhibitor that targets receptor tyrosine kinase including vascular endothelial growth factor receptor-1, -2, -3 (VEGFR-1, -2, -3), platelet-derived growth factor receptor- α , - β (PDGFR- α , - β), c-kit, fibroblast growth factor-1 and -3 (FGFR-1, -3), thereby inhibiting angiogenesis. Pazopanib is developed and marketed by GlaxoSmithKline under the trade name VOTRIENT.

- FDA Approval Summary of Pazopanib (VOTRIENT)

PALETTE ^[169] NCT00753688	Sarcoma (Approved on 2016/04/26)
	-
VEG105192 ^[170] NCT00334282	Pazopanib vs. Placebo [PFS(M): 4.6 vs. 1.6]
	Renal cell carcinoma (Approved on 2009/10/19)
	-
	Pazopanib vs. Placebo [PFS(M): 9.2 vs. 4.2]

Rucaparib (RUBRACA)

Rucaparib is an inhibitor of the DNA repair enzyme poly (ADP-ribose) polymerase-1, -2 and -3 (PARP-1, -2, -3). Rucaparib is developed and marketed by Clovis Oncology under the trade name RUBRACA.

- FDA Approval Summary of Rucaparib (RUBRACA)

TRITON2 NCT02952534	Prostate cancer (Approved on 2020/05/15)
	gBRCA mutation or sBRCA mutation
ARIEL3 ^[15] NCT01968213	Rucaparib [ORR(%): 44.0, DOR(M): NE]
	Ovarian cancer, Fallopian tube cancer, Peritoneal carcinoma (Approved on 2018/04/06)
	-
	Rucaparib vs. Placebo [PFS (All)(M): 10.8 vs. 5.4, PFS (HRD)(M): 13.6 vs. 5.4, PFS (tBRCA)(M): 16.6 vs. 5.4]

Selumetinib (KOSELUGO)

Selumetinib is a kinase inhibitor. Selumetinib is developed and marketed by AstraZeneca under the trade name KOSELUGO.

- FDA Approval Summary of Selumetinib (KOSELUGO)

SPRINT NCT01362803	Plexiform neurofibromas (Approved on 2020/04/10)
	-
	Selumetinib [ORR(%): 66.0]

ACT Onco[®] + Report

Sorafenib (NEXAVAR)

Sorafenib is a small molecule multi-kinase inhibitor that targets multiple kinase families including VEGFR, PDGFRB, and the RAF family kinases. Sorafenib is co-developed and co-marketed by Bayer HealthCare Pharmaceuticals and Onyx Pharmaceuticals under the trade name NEXAVAR.

- FDA Approval Summary of Sorafenib (NEXAVAR)

DECISION ^[171] NCT00984282	Differentiated thyroid carcinoma (Approved on 2013/11/22)
	- Sorafenib vs. Placebo [PFS(M): 10.8 vs. 5.8]
SHARP ^[172] NCT00105443	Hepatocellular carcinoma (Approved on 2007/11/16)
	- Sorafenib vs. Placebo [OS(M): 10.7 vs. 7.9]
TARGET ^[173] NCT00073307	Renal cell carcinoma (Approved on 2005/12/20)
	- Sorafenib vs. Placebo [PFS(D): 167 vs. 84]

Sunitinib (SUTENT)

Sunitinib is an oral, small molecule, multi-kinase inhibitor that targets receptor tyrosine kinase including platelet-derived growth factor receptor- α , - β (PDGFR- α , - β), vascular endothelial growth factor receptors-1, -2, -3 (VEGFR-1, -2, -3), c-kit, Fms-like tyrosine kinase-3 (FLT3), colony stimulating factor receptor type 1 (CSF-1R), and the glial cell-line derived neurotrophic factor receptor (RET), thereby inhibiting angiogenesis. Sunitinib is developed and marketed by Pfizer under the trade name SUTENT.

- FDA Approval Summary of Sunitinib (SUTENT)

^{[174][175][176]} NCT00428597	Pancreatic cancer (Approved on 2011/05/20)
	- Sunitinib vs. Placebo [PFS(M): 10.2 vs. 5.4]
^{[177][178]} NCT00083889	Renal cell carcinoma (Approved on 2007/02/02)
	- Sunitinib vs. Ifn- α [PFS(W): 47.3 vs. 22]
^{[179][180][178]} NCT00077974	Renal cell carcinoma (Approved on 2007/02/02)
	- Sunitinib [ORR(%): 34.0]
^{[180][178]} NCT00054886	Renal cell carcinoma (Approved on 2007/02/02)
	- Sunitinib [ORR(%): 36.5]
^[181] NCT00075218	Gastrointestinal stromal tumor (Approved on 2006/01/26)
	- Sunitinib vs. Placebo [TTP(W): 27.3 vs. 6.4]

ACT Onco[®] + Report

Talazoparib (TALZENNA)

Talazoparib is an inhibitor of poly (ADP-ribose) polymerase (PARP) enzymes, including PARP1 and PARP2. Talazoparib is developed and marketed by Pfizer under the trade name TALZENNA.

- FDA Approval Summary of Talazoparib (TALZENNA)

TALAPRO-2 NCT03395197	Prostate cancer (Approved on 2023/06/20)
	HRR genes mutation Talazoparib + enzalutamide vs. Placebo + enzalutamide [rPFS(M): Not reached vs. 13.8]
EMBRACA ^[182] NCT01945775	Breast cancer (Approved on 2018/10/16)
	HER2-/gBRCA mutation Talazoparib vs. Chemotherapy [PFS(M): 8.6 vs. 5.6]

Temsirolimus (TORISEL)

Temsirolimus is a soluble ester of sirolimus (rapamycin, brand-name drug Rapamune) and functions as an inhibitor of mammalian target of rapamycin complex (mTORC). The inhibitory molecular mechanism is similar to Everolimus. Temsirolimus is developed by Wyeth Pharmaceuticals and marketed by Pfizer under the trade name TORISEL.

- FDA Approval Summary of Temsirolimus (TORISEL)

[183] NCT00065468	Renal cell carcinoma (Approved on 2007/05/30)
	- Temsirolimus vs. Ifn-α [OS(M): 10.9 vs. 7.3]

Trametinib (MEKINIST)

Trametinib is an anti-cancer inhibitor which targets MEK1 and MEK2. Trametinib is developed and marketed by GlaxoSmithKline (GSK) under the trade name MEKINIST.

- FDA Approval Summary of Trametinib (MEKINIST)

CDRB436G2201 NCT02684058	Low-grade glioma (Approved on 2023/03/09)
	BRAF V600E Dabrafenib + trametinib vs. Carboplatin + vincristine [ORR(%): 46.6 vs. 10.8]
BRF117019, NCI-MATCH, CTMT212X2101 NCT02034110, NCT02465060, NCT02124772	Cancer (Approved on 2022/06/22)
	BRAF V600E Dabrafenib + trametinib [ORR(adult patients)(%): 41.0, ORR(pediatric patients)(%): 25.0]
BRF117019 ^[184] NCT02034110	Anaplastic thyroid cancer (Approved on 2018/05/04)
	BRAF V600E Dabrafenib + trametinib [ORR(%): 61.0]
BRF113928 ^[185] NCT01336634	Non-small cell lung cancer (Approved on 2017/06/22)
	BRAF V600E Trametinib + dabrafenib vs. Dabrafenib [ORR(%): 63.0 vs. 27.0, DOR(M): 12.6 vs. 9.9]

ACT Onco[®] + Report

COMBI-d ^[186] NCT01584648	Melanoma (Approved on 2014/01/10)
	BRAF V600E/K
	Trametinib + dabrafenib vs. Dabrafenib + placebo [PFS(M): 9.3 vs. 8.8]
METRIC ^[187] NCT01245062	Melanoma (Approved on 2013/05/29)
	BRAF V600E/K
	Trametinib vs. Dacarbazine or paclitaxel [PFS(M): 4.8 vs. 1.5]

D=day; W=week; M=month

ACT Onco[®] + Report

ONGOING CLINICAL TRIALS

Trials were searched by applying filters: study status, patient's diagnosis, intervention, location and/or biomarker(s). Please visit <https://clinicaltrials.gov> to search and view for a complete list of open available and updated matched trials.

No trial has been found.

ACT Onco[®] + Report


SUPPLEMENTARY INFORMATION OF TESTING RESULTS DETAILED INFORMATION OF VARIANTS WITH CLINICAL RELEVANCE

- Single Nucleotide and Small InDel Variants

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
Not Detected							

- Copy Number Alterations

Observed copy number (CN) for each evaluated position is shown on the y-axis. Regions referred to as amplification or deletion are shown in color. Regions without significant changes are represented in gray.



ACT Onco[®] + Report

OTHER DETECTED VARIANTS

Gene	Amino Acid Change	Exon	cDNA Change	Accession Number	COSMIC ID	Allele Frequency	Coverage
ASXL1	G587C	13	c.1759G>T	NM_015338	-	14.3%	1224
ATM	E1353K	27	c.4057G>A	NM_000051	-	43.6%	518
AXL	R4W	1	c.10C>T	NM_021913	-	98.2%	283
BRD4	T221M	5	c.662C>T	NM_058243	-	28.0%	858
EPHA2	Splice region	-	c.1738+8G>A	NM_004431	-	42.7%	881
KMT2A	Q2337R	27	c.7010A>G	NM_001197104	-	51.2%	823
MAP3K1	G616W	10	c.1846G>T	NM_005921	-	28.1%	1415
MSH2	I632V	12	c.1894A>G	NM_000251	-	41.5%	518
PALB2	T1012I	10	c.3035C>T	NM_024675	-	69.4%	607
PALB2	V425M	4	c.1273G>A	NM_024675	COSM1286951	67.3%	928
POLD1	D644E	16	c.1932C>G	NM_001256849	-	50.3%	922
PRKN	R396G	11	c.1186A>G	NM_004562	-	34.3%	1339
PTCH1	Splice region	-	c.2560+7C>T	NM_000264	-	46.8%	2490
RXRA	E243G	5	c.728A>G	NM_002957	-	16.1%	801
TSC2	I195V	6	c.583A>G	NM_000548	-	71.6%	559


Note:

- This table enlists variants detected by the panel other than those with clinical relevance (reported in Testing Result section). The clinical impact of a genetic variant is determined according to ACT Genomics in-house clinical knowledge database. A negative result does not necessarily indicate absence of biological effect on the tumor. Some variants listed here may possibly have preclinical data or may show potential clinical relevance in the future.

ACT Onco[®] + Report

TEST DETAILS

SPECIMEN RECEIVED AND PATHOLOGY REVIEW



- Collection date: Jun 08, 2023
- Facility retrieved: 臺北榮總
- H&E-stained section No.: S11226789B
- Collection site: Lung
- Examined by: Dr. Yun-An Chen
- 1. The percentage of viable tumor cells in total cells in the whole slide (%): 45%
- 2. The percentage of viable tumor cells in total cells in the encircled areas in the whole slide (%): 85%
- 3. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the whole slide (%): 0%
- 4. The percentage of necrotic cells (including necrotic tumor cells) in total cells in the encircled areas in the whole slide (%): 0%
- 5. Additional comment: NA
- Manual macrodissection: Performed on the highlighted region
- The outline highlights the area of malignant neoplasm annotated by a pathologist.

RUN QC

- Panel: ACTOnco[®]+

DNA test

- Mean Depth: 1072x
- Target Base Coverage at 100x: 95%

RNA test

- Average unique RNA Start Sites per control GSP2: 132

ACT Onco[®] + Report

LIMITATIONS

1. This test does not provide information of variant causality and does not detect variants in non-coding regions that could affect gene expression. This report does not report polymorphisms and we do not classify whether a mutation is germline or somatic. Variants identified by this assay were not subject to validation by Sanger or other technologies.
2. The possibility cannot be excluded that certain pathogenic variants detected by other sequencing tools may not be reported in the test because of technical limitation of bioinformatics algorithm or the NGS sequencing platform, e.g. low coverage.
3. This test has been designed to detect fusions in 13 genes sequenced. Therefore, fusion in genes not covered by this test would not be reported. For novel fusions detected in this test, Sanger sequencing confirmation is recommended if residue specimen is available.

NEXT-GENERATION SEQUENCING (NGS) METHODS

DNA test

Extracted genomic DNA was amplified using primers targeting coding exons of analyzed genes and subjected to library construction. Barcoded libraries were subsequently conjugated with sequencing beads by emulsion PCR and enriched using Ion Chef system. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific).

Raw reads generated by the sequencer were mapped to the hg19 reference genome using the Ion Torrent Suite. Coverage depth was calculated using Torrent Coverage Analysis plug-in. Single nucleotide variants (SNVs) and short insertions/deletions (InDels) were identified using the Torrent Variant Caller plug-in. VEP (Variant Effect Predictor) was used to annotate every variant using databases from Clinvar, COSMIC and Genome Aggregation database. Variants with coverage ≥ 20 , allele frequency $\geq 5\%$ and actionable variants with allele frequency $\geq 2\%$ were retained. This test provides uniform coverage of the targeted regions, enabling target base coverage at $100\times \geq 85\%$ with a mean coverage $\geq 500\times$.

Variants reported in Genome Aggregation database with $> 1\%$ minor allele frequency (MAF) were considered as polymorphisms. ACT Genomics in-house database was used to determine technical errors. Clinically actionable and biologically significant variants were determined based on the published medical literature.

The copy number alterations (CNAs) were predicted as described below:

Amplicons with read counts in the lowest 5th percentile of all detectable amplicons and amplicons with a coefficient of variation ≥ 0.3 were removed. The remaining amplicons were normalized to correct the pool design bias. ONCOCNV (an established method for calculating copy number aberrations in amplicon sequencing data by Boeva et al., 2014) was applied for the normalization of total amplicon number, amplicon GC content, amplicon length, and technology-related biases, followed by segmenting the sample with a gene-aware model. The method was used as well for establishing the baseline of copy number variations.

Tumor mutational burden (TMB) was calculated by using the sequenced regions of ACTOnco[®]+ to estimate the number of somatic nonsynonymous mutations per megabase of all protein-coding genes (whole exome). The TMB calculation predicted somatic variants and applied a machine learning model with a cancer hotspot correction. TMB may be reported as "TMB-High", "TMB-Low" or "Cannot Be Determined". TMB-High corresponds to ≥ 7.5 mutations per megabase (Muts/Mb); TMB-Low corresponds to < 7.5 Muts/Mb. TMB is reported as "Cannot Be Determined" if the tumor purity of the sample is $< 30\%$.

Classification of microsatellite instability (MSI) status is determined by a machine learning prediction algorithm. The change of a number of repeats of different lengths from a pooled microsatellite stable (MSS) baseline in > 400 genomic loci are used as the features for the algorithm. The final output of the results is either microsatellite Stable (MSS) or microsatellite instability high (MSI-H).

RNA test

Extracted RNA was reverse-transcribed and subjected to library construction. Sequencing was performed according to Ion Proton or Ion S5 sequencer protocol (Thermo Fisher Scientific). To ensure sequencing quality for fusion variant analysis, the average unique RNA Start Sites (SS) per control Gene Specific Primer 2 (GSP 2) should be ≥ 10 .

ACT Onco[®] + Report

The fusion analysis pipeline aligned sequenced reads to the human reference genome, identified regions that map to noncontiguous regions of the genome, applied filters to exclude probable false-positive events and, annotated previously characterized fusion events according to Quiver Gene Fusion Database, a curated database owned and maintained by ArcherDX. In general, samples with detectable fusions need to meet the following criteria: (1) Number of unique start sites (SS) for the GSP2 ≥ 3 ; (2) Number of supporting reads spanning the fusion junction ≥ 5 ; (3) Percentage of supporting reads spanning the fusion junction $\geq 10\%$; (4) Fusions annotated in Quiver Gene Fusion Database.

DATABASE USED

- Reference genome: Human genome sequence hg19
- COSMIC v.92
- Genome Aggregation database r2.1.1
- ClinVar (version 20210404)
- ACT Genomics in-house database
- Quiver Gene Fusion Database version 5.1.18

Variant Analysis:

醫檢師陳韻鈺 博士
Yun-Yu Chen Ph.D.
檢字第 015647 號

Yun Yu Chen

Sign Off

解剖病理專科醫師王業翰
Yeh-Han Wang M.D.
病解字第 000545 號

Yeh

ACT Onco[®] + Report

GENE LIST SNV & CNV

ABCB1*	ABCC2*	ABCG2*	ABL1	ABL2	ADAMTS1	ADAMTS13	ADAMTS15	ADAMTS16	ADAMTS18	ADAMTS6	ADAMTS9
ADAMTSL1	ADGRA2	ADH1C*	AKT1	AKT2	AKT3	ALDH1A1*	ALK	AMER1	APC	AR	ARAF
ARID1A	ARID1B	ARID2	ASXL1	ATM	ATR	ATRX	AURKA	AURKB	AXIN1	AXIN2	AXL
B2M	BAP1	BARD1	BCL10	BCL2*	BCL2L1	BCL2L2*	BCL6	BCL9	BCOR	BIRC2	BIRC3
BLM	BMPR1A	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1	BTG2*	BTB	BUB1B	CALR
CANX	CARD11	CASP8	CBFB	CBL	CCNA1	CCNA	CCNB1	CCNB2	CCNB3	CCND1	CCND2
CCND3	CCNE1	CCNE2	CCNH	CD19	CD274	CD58	CD70*	CD79A	CD79B	CDC73	CDH1
CDK1	CDK12	CDK2	CDK4	CDK5	CDK6	CDK7	CDK8	CDK9	CDKN1A	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA*	CHEK1	CHEK2	CIC	CREBBP	CRKL	CRLF2	CSF1R	CTCF	CTLA4
CTNNA1	CTNNB1	CUL3	CYLD	CYP1A1*	CYP2B6*	CYP2C19*	CYP2C8*	CYP2D6	CYP2E1*	CYP3A4*	CYP3A5*
DAXX	DCUN1D1	DDR2	DICER1	DNMT3A	DOT1L	DPYD	DTX1	E2F3	EGFR	EP300	EPCAM
EPHA2	EPHA3	EPHA5	EPHA7	EPHB1	ERBB2	ERBB3	ERBB4	ERCC1	ERCC2	ERCC3	ERCC4
ERCC5	ERG	ESR1	ESR2	ETV1	ETV4	EZH2	FAM46C	FANCA	FANCC	FANCD2	FANCE
FANCF	FANCG	FANCL	FAS	FAT1	FBXW7	FCGR2B	FGF1*	FGF10	FGF14	FGF19*	FGF23
FGF3	FGF4*	FGF6	FGFR1	FGFR2	FGFR3	FGFR4	FH	FLCN	FLT1	FLT3	FLT4
FOXL2*	FOXP1	FRG1	FUBP1	GATA1	GATA2	GATA3	GNA11	GNA13	GNAQ	GNAS	GREM1
GRIN2A	GSK3B	GSTP1*	GSTT1*	HGF	HIF1A	HIST1H1C*	HIST1H1E*	HNF1A	HR	HRA5*	HSP90AA1
HSP90AB1	HSPA4	HSPA5	IDH1	IDH2	IFNL3*	IGF1	IGF1R	IGF2	IKBK	IKBE	IKZF1
IL6	IL7R	INPP4B	INSR	IRF4	IRS1	IRS2*	JAK1	JAK2	JAK3	JUN*	KAT6A
KDM5A	KDM5C	KDM6A	KDR	KEAP1	KIT	KMT2A	KMT2C	KMT2D	KRAS	LCK	LIG1
LIG3	LMO1	LRP1B	LYN	MALT1	MAP2K1	MAP2K2	MAP2K4	MAP3K1	MAP3K7	MAPK1	MAPK3
MAX	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MET	MITF	MLH1	MPL	MRE11
MSH2	MSH6	MTHFR*	MTOR	MUC16	MUC4	MUC6	MUTYH	MYC	MYCL	MYCN	MYD88
NAT2*	NBN	NEFH	NF1	NF2	NFE2L2	NFKB1	NFKBIA	NKX2-1*	NOTCH1	NOTCH2	NOTCH3
NOTCH4	NPM1	NQO1*	NRAS	NSD1	NTRK1	NTRK2	NTRK3	PAK3	PALB2	PARP1	PAX5
PAX8	PBRM1	PDCD1	PDCD1LG2	PDGFRA	PDGFRB	PDIA3	PGF	PHOX2B*	PIK3C2B	PIK3C2G	PIK3C3
PIK3CA	PIK3CB	PIK3CD	PIK3CG	PIK3R1	PIK3R2	PIK3R3	PIM1	PMS1	PMS2	POLB	POLD1
POLE	PPARG	PPP2R1A	PRDM1	PRKAR1A	PRKCA	PRKCB	PRKCG	PRKCI	PRKCQ	PRKDC	PRKN
PSMB8	PSMB9	PSME1	PSME2	PSME3	PTCH1	PTEN	PTGS2	PTPN11	PTPRD	PTPRT	RAC1
RAD50	RAD51	RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10	RECQL4
REL	RET	RHOA	RICTOR	RNF43	ROS1	RPPH1	RPTOR	RUNX1	RUNX1T1	RXRA	SDHA
SDHB	SDHC	SDHD	SERPINB3	SERPINB4	SETD2	SF3B1	SGK1	SH2D1A*	SLC19A1*	SLC22A2*	SLC1B1*
SLC1B3*	SMAD2	SMAD3	SMAD4	SMARCA4	SMARCB1	SMO	SOC1*	SOX2*	SOX9	SPEN	SPOP
SRC	STAG2	STAT3	STK11	SUFU	SYK	SYNE1	TAF1	TAP1	TAP2	TAPBP	TBX3
TEK	TERT	TET1	TET2	TGFBR2	TMSB4X*	TNF	TNFAIP3	TNFRSF14	TNFSF11	TOP1	TP53
TPMT*	TSC1	TSC2	TSHR	TYMS	U2AF1	UBE2A*	UBE2K	UBR5	UGT1A1*	USH2A	VDR*
VEGFA	VEGFB	VHL	WT1	XIAP	XPO1	XRCC2	ZNF217				

*Analysis of copy number alterations NOT available.

FUSION

ALK	BRAF	EGFR	FGFR1	FGFR2	FGFR3	MET	NRG1	NTRK1	NTRK2	NTRK3	RET	ROS1
-----	------	------	-------	-------	-------	-----	------	-------	-------	-------	-----	------


ACT Onco[®] + Report

APPENDIX

POSSIBLE THERAPEUTIC IMPLICATIONS FOR HETEROZYGOUS DELETION

Gene	Therapies	Possible effect
NF1	Binimetinib, Cobimetinib, Trametinib, Selumetinib, Everolimus, Temsirolimus	sensitive
STK11	Binimetinib, Cobimetinib, Trametinib, Everolimus, Temsirolimus	sensitive
ATR	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
BRCA2	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
PALB2	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
RAD50	Niraparib, Olaparib, Rucaparib, Talazoparib	sensitive
NF1	Afatinib, Cabozantinib, Crizotinib, Erlotinib, Gefitinib, Lapatinib, Cetuximab, Trastuzumab, Vemurafenib	resistant


SIGNALING PATHWAYS AND MOLECULAR-TARGETED AGENTS




1: Olaparib, Niraparib, Rucaparib, Talazoparib

ACT Onco[®] + Report

Receptor Tyrosine Kinase/Growth Factor Signalling



1: Pazopanib; 2: Imatinib, Sunitinib, Pazopanib; 3: Sunitinib, Pazopanib; 4: Sunitinib, Pazopanib; 5: Sunitinib; 6: Imatinib, Sunitinib, Pazopanib, Sorafenib



1: Everolimus, Temsirolimus; 2: Sorafenib; 3: Trametinib, Selumetinib, Binimetinib, Cobimetinib

ACT Onco® + Report

DISCLAIMER

法律聲明

本檢驗報告僅提供專業醫療參考，結果需經專業醫師解釋及判讀。基因突變資訊非必具備藥物或治療有效性指標，反之亦然。本檢驗報告提供之用藥指引不聲明或保證其臨床有效性，反之亦然。本基因檢測方法係由本公司研究開發，已經過有效性測試。

本檢驗報告非經本公司許可，不得私自變造、塗改，或以任何方式作為廣告及其他宣傳之用途。

本公司於提供檢驗報告後，即已完成本次契約義務，後續之報告解釋、判讀及用藥、治療，應自行尋求相關專業醫師協助，若需將報告移件其他醫師，本人應取得該醫師同意並填寫移件申請書，主動告知行動基因，行動基因僅能配合該醫師意願與時間提供醫師解說。

醫療決策需由醫師決定

任何治療與用藥需經由醫師在考慮病患所有健康狀況相關資訊包含健檢、其他檢測報告和病患意願後，依照該地區醫療照護標準由醫師獨立判斷。醫師不應僅依據單一報告結果(例如本檢測或本報告書內容)做決策。

基因突變與用藥資訊並非依照有效性排序

本報告中列出之生物標記變異與藥物資訊並非依照潛在治療有效性排序。

證據等級

藥物潛在臨床效益(或缺乏潛在臨床效益)的實證證據是依據至少一篇臨床療效個案報告或臨床前試驗做為評估。本公司盡力提供適時及準確之資料，但由於醫學科技之發展日新月異，本公司不就本報告提供的資料是否為準確、適宜或最新作保證。

責任

本檢驗報告僅提供專業醫療參考，本公司及其員工不對任何由使用本報告之內容引起的直接、間接、特殊、連帶或衍生的損失或損害承擔責任。

ACT Onco[®] + Report

REFERENCE

1. PMID: 11544175; 2001, Genes Dev;15(17):2177-96
Cell cycle checkpoint signaling through the ATM and ATR kinases.
2. PMID: 11163154; 2001, Curr Opin Genet Dev;11(1):71-7
ATM and ATR: networking cellular responses to DNA damage.
3. PMID: 12526805; 2002, Cell;111(6):779-89
ATR regulates fragile site stability.
4. PMID: 10097108; 1999, Proc Natl Acad Sci U S A;96(7):3745-50
A human Cds1-related kinase that functions downstream of ATM protein in the cellular response to DNA damage.
5. PMID: 15282542; 2004, EMBO J;23(15):3164-74
ATR functions as a gene dosage-dependent tumor suppressor on a mismatch repair-deficient background.
6. PMID: 22341969; 2012, Am J Hum Genet;90(3):511-7
Germline mutation in ATR in autosomal- dominant oropharyngeal cancer syndrome.
7. PMID: 12640452; 2003, Nat Genet;33(4):497-501
A splicing mutation affecting expression of ataxia-telangiectasia and Rad3-related protein (ATR) results in Seckel syndrome.
8. PMID: 17879369; 2007, Genes Chromosomes Cancer;46(12):1061-8
Mutations in the ataxia telangiectasia and rad3-related-checkpoint kinase 1 DNA damage response axis in colon cancers.
9. PMID: 16288216; 2006, Oncogene;25(14):2113-8
Microsatellite instability and mutation analysis of candidate genes in urothelial cell carcinomas of upper urinary tract.
10. PMID: 11691784; 2001, Cancer Res;61(21):7727-30
Somatic mutations in the DNA damage-response genes ATR and CHK1 in sporadic stomach tumors with microsatellite instability.
11. PMID: 19470935; 2009, J Clin Oncol;27(19):3091-6
ATR mutation in endometrioid endometrial cancer is associated with poor clinical outcomes.
12. PMID: 26282654; 2015, J Clin Oncol;33(33):3911-20
Mutational Spectrum, Copy Number Changes, and Outcome: Results of a Sequencing Study of Patients With Newly Diagnosed Myeloma.
13. PMID: 35599270; 2022, Med Oncol;39(5):96
Olaparib for Chinese metastatic castration-resistant prostate cancer: A real-world study of efficacy and gene predictive analysis.
14. PMID: 30353044; 2018, Br J Cancer;119(11):1401-1409
Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the BRCA genes.
15. PMID: 28916367; 2017, Lancet;390(10106):1949-1961
Rucaparib maintenance treatment for recurrent ovarian carcinoma after response to platinum therapy (ARIEL3): a randomised, double-blind, placebo-controlled, phase 3 trial.
16. PMID: 11239455; 2001, Mol Cell;7(2):263-72
BRCA2 is required for homology-directed repair of chromosomal breaks.
17. PMID: 17597348; 2007, Ann Surg Oncol;14(9):2510-8
Heterogenic loss of the wild-type BRCA allele in human breast tumorigenesis.
18. PMID: 22193408; 2011, Nat Rev Cancer;12(1):68-78
BRCA1 and BRCA2: different roles in a common pathway of genome protection.
19. PMID: 27283171; 2016, J Natl Compr Canc Netw;14(6):795-806
The Relevance of Hereditary Cancer Risks to Precision Oncology: What Should Providers Consider When Conducting Tumor Genomic

ACT Onco[®] + Report

Profiling?

20. PMID: 1833068; 1991, Cell;66(6):1217-28
Human cyclin E, a new cyclin that interacts with two members of the CDC2 gene family.
21. PMID: 22186781; 2012, Cell Cycle;11(1):57-64
An integrated view of cyclin E function and regulation.
22. PMID: 20336084; 2010, J Invest Dermatol;130(7):1914-21
UVR-induced regulatory T cells switch antigen-presenting cells from a stimulatory to a regulatory phenotype.
23. PMID: 25395429; 2015, Blood;125(3):443-8
Dinaciclib, a novel CDK inhibitor, demonstrates encouraging single-agent activity in patients with relapsed multiple myeloma.
24. PMID: 27663592; 2017, Clin Cancer Res;23(7):1862-1874
Selective Targeting of Cyclin E1-Amplified High-Grade Serous Ovarian Cancer by Cyclin-Dependent Kinase 2 and AKT Inhibition.
25. PMID: 27020857; 2016, Cancer Res;76(8):2301-13
Early Adaptation and Acquired Resistance to CDK4/6 Inhibition in Estrogen Receptor-Positive Breast Cancer.
26. PMID: 21321214; 2011, Proc Natl Acad Sci U S A;108(9):3761-6
Cyclin E amplification/overexpression is a mechanism of trastuzumab resistance in HER2+ breast cancer patients.
27. PMID: 20336784; 2010, Cancer;116(11):2621-34
Gene amplification CCNE1 is related to poor survival and potential therapeutic target in ovarian cancer.
28. PMID: 30026836; 2018, J Cancer;9(13):2397-2407
Prognostic Values of CCNE1 Amplification and Overexpression in Cancer Patients: A Systematic Review and Meta-analysis.
29. PMID: 30807234; 2019, J Clin Oncol;37(14):1169-1178
Cyclin E1 Expression and Palbociclib Efficacy in Previously Treated Hormone Receptor-Positive Metastatic Breast Cancer.
30. PMID: 23172303; 2013, J Biochem;153(1):13-9
Vascular endothelial growth factor and its receptor system: physiological functions in angiogenesis and pathological roles in various diseases.
31. PMID: 12778165; 2003, Nat Med;9(6):669-76
The biology of VEGF and its receptors.
32. PMID: 21382095; 2011, Neuropathology;31(6):583-8
Amplification of the PDGFRA, KIT and KDR genes in glioblastoma: a population-based study.
33. PMID: 31604903; 2020, Oncologist;25(1):e39-e47
The Pan-Cancer Landscape of Coamplification of the Tyrosine Kinases KIT, KDR, and PDGFRA.
34. PMID: 23028128; 2012, Cold Spring Harb Perspect Med;2(10):
Anti-VEGF therapies in the clinic.
35. PMID: 27718781; 2016, N Engl J Med;375(23):2246-2254
Adjuvant Sunitinib in High-Risk Renal-Cell Carcinoma after Nephrectomy.
36. PMID: 29972759; 2018, N Engl J Med;379(1):54-63
Cabozantinib in Patients with Advanced and Progressing Hepatocellular Carcinoma.
37. PMID: 27160228; 2016, J Natl Compr Canc Netw;14(5):499-502
Antitumor Response of VEGFR2- and VEGFR3-Amplified Angiosarcoma to Pazopanib.
38. PMID: 25639617; 2015, Breast J;21(2):205-7
Refractory angiosarcoma of the breast with VEGFR2 upregulation successfully treated with sunitinib.
39. PMID: 26578684; 2016, Clin Cancer Res;22(8):1940-50

ACT Onco[®] + Report

KDR Amplification Is Associated with VEGF-Induced Activation of the mTOR and Invasion Pathways but does not Predict Clinical Benefit to the VEGFR TKI Vandetanib.

40. PMID: 29855279; 2018, BMC Cancer;18(1):618
Apatinib treatment for KIT- and KDR-amplified angiosarcoma: a case report.
41. PMID: 31786466; 2020, Biomed Pharmacother;122():109587
Efficacy and safety of the VEGFR2 inhibitor Apatinib for metastatic soft tissue sarcoma: Chinese cohort data from NCT03121846.
42. PMID: 22170730; 2012, Genes Chromosomes Cancer;51(4):375-83
Increased gene copy number of KIT and VEGFR2 at 4q12 in primary breast cancer is related to an aggressive phenotype and impaired prognosis.
43. PMID: 17546049; 2007, Oncogene;26(54):7560-8
KIT oncogenic signaling mechanisms in imatinib-resistant gastrointestinal stromal tumor: PI3-kinase/AKT is a crucial survival pathway.
44. PMID: 15365079; 2004, J Clin Oncol;22(18):3813-25
Biology of gastrointestinal stromal tumors.
45. PMID: 22089421; 2011, Nat Rev Cancer;11(12):865-78
Gastrointestinal stromal tumours: origin and molecular oncology.
46. PMID: 12181401; 2002, N Engl J Med;347(7):472-80
Efficacy and safety of imatinib mesylate in advanced gastrointestinal stromal tumors.
47. PMID: 18955458; 2008, J Clin Oncol;26(33):5352-9
Primary and secondary kinase genotypes correlate with the biological and clinical activity of sunitinib in imatinib-resistant gastrointestinal stromal tumor.
48. PMID: 23177515; 2013, Lancet;381(9863):295-302
Efficacy and safety of regorafenib for advanced gastrointestinal stromal tumours after failure of imatinib and sunitinib (GRID): an international, multicentre, randomised, placebo-controlled, phase 3 trial.
49. PMID: 25641662; 2015, Cancer;121(9):1405-13
Clinical outcomes of patients with advanced gastrointestinal stromal tumors: safety and efficacy in a worldwide treatment-use trial of sunitinib.
50. PMID: 19282169; 2009, Eur J Cancer;45(11):1959-68
Clinical evaluation of continuous daily dosing of sunitinib malate in patients with advanced gastrointestinal stromal tumour after imatinib failure.
51. PMID: 32511981; 2020, Lancet Oncol;21(7):923-934
Ripretinib in patients with advanced gastrointestinal stromal tumours (INVICTUS): a double-blind, randomised, placebo-controlled, phase 3 trial.
52. PMID: 32350132; 2020, Sci Transl Med;12(541):
Discovery and pharmacological characterization of AZD3229, a potent KIT/PDGFR α inhibitor for treatment of gastrointestinal stromal tumors.
53. PMID: 35091442; 2022, Clin Cancer Res;28(7):1268-1276
Phase II Study of Ponatinib in Advanced Gastrointestinal Stromal Tumors: Efficacy, Safety, and Impact of Liquid Biopsy and Other Biomarkers.
54. PMID: 28327988; 2017, Ann Oncol;28(6):1380-1387
Efficacy and safety of nilotinib in patients with KIT-mutated metastatic or inoperable melanoma: final results from the global, single-arm, phase II TEAM trial.
55. PMID: 15583695; 2004, Br J Cancer;91(12):2048-55
Genetic alterations and protein expression of KIT and PDGFRA in serous ovarian carcinoma.
56. PMID: 23775962; 2013, J Clin Oncol;31(26):3182-90
Imatinib for melanomas harboring mutationally activated or amplified KIT arising on mucosal, acral, and chronically sun-damaged skin.
57. PMID: 8563751; 1996, Nat Genet;12(2):144-8

ACT Onco[®] + Report

Loss of NF1 results in activation of the Ras signaling pathway and leads to aberrant growth in haematopoietic cells.

58. PMID: 1946382; 1991, Proc Natl Acad Sci U S A;88(21):9658-62
Identification of the neurofibromatosis type 1 gene product.
59. PMID: 2116237; 1990, Cell;62(3):599-608
The neurofibromatosis type 1 gene encodes a protein related to GAP.
60. PMID: 2121370; 1990, Cell;63(4):843-9
The GAP-related domain of the neurofibromatosis type 1 gene product interacts with ras p21.
61. PMID: 14502561; 2003, J Cell Physiol;197(2):214-24
NF1 modulates the effects of Ras oncogenes: evidence of other NF1 function besides its GAP activity.
62. PMID: 19479903; 2009, Mol Carcinog;48(11):1005-17
Neurofibromin physically interacts with the N-terminal domain of focal adhesion kinase.
63. PMID: 28680740; 2017, Adv Med Biol;118():83-122
Haploinsufficient tumor suppressor genes.
64. PMID: 10442636; 1999, Oncogene;18(31):4450-9
Haploinsufficiency for the neurofibromatosis 1 (NF1) tumor suppressor results in increased astrocyte proliferation.
65. PMID: 16288202; 2006, Oncogene;25(16):2297-303
NF1 haploinsufficiency augments angiogenesis.
66. PMID: 18089636; 2008, Hum Mol Genet;17(7):936-48
Rac1 mediates the osteoclast gains-in-function induced by haploinsufficiency of Nf1.
67. PMID: 7920653; 1994, Nat Genet;7(3):353-61
Tumour predisposition in mice heterozygous for a targeted mutation in Nf1.
68. PMID: 25026295; 2014, Oncotarget;5(15):5873-92
The NF1 gene revisited - from bench to bedside.
69. PMID: 29892687; 2018, Gynecol Oncol Rep;23():41-44
Clonal lineage of high grade serous ovarian cancer in a patient with neurofibromatosis type 1.
70. PMID: 29926297; 2018, Breast Cancer Res Treat;171(3):719-735
Breast cancer in women with neurofibromatosis type 1 (NF1): a comprehensive case series with molecular insights into its aggressive phenotype.
71. PMID: 28637487; 2017, Hum Genomics;11(1):13
The NF1 somatic mutational landscape in sporadic human cancers.
72. PMID: 15840687; 2005, Gut;54(8):1129-35
NF1 gene loss of heterozygosity and expression analysis in sporadic colon cancer.
73. PMID: 20129251; 2010, Cancer Cell;17(1):98-110
Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1.
74. PMID: 27158780; 2016, Nat Genet;48(6):607-16
Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas.
75. PMID: 32669708; 2020, Nature;583(7818):807-812
The National Lung Matrix Trial of personalized therapy in lung cancer.
76. PMID: 21482774; 2012, Proc Natl Acad Sci U S A;109(8):2730-5
Genome-wide functional screen identifies a compendium of genes affecting sensitivity to tamoxifen.

ACT Onco[®] + Report

77. PMID: 24535670; 2014, Cancer Discov;4(5):606-19
Reduced NF1 expression confers resistance to EGFR inhibition in lung cancer.
78. PMID: 29703253; 2018, BMC Cancer;18(1):479
SMAD4 and NF1 mutations as potential biomarkers for poor prognosis to cetuximab-based therapy in Chinese metastatic colorectal cancer patients.
79. PMID: 30858928; 2019, Oncotarget;10(14):1440-1457
CRISPR-induced RASGAP deficiencies in colorectal cancer organoids reveal that only loss of NF1 promotes resistance to EGFR inhibition.
80. PMID: 24576830; 2014, Cancer Res;74(8):2340-50
Loss of NF1 in cutaneous melanoma is associated with RAS activation and MEK dependence.
81. PMID: 23288408; 2013, Cancer Discov;3(3):350-62
A genome-scale RNA interference screen implicates NF1 loss in resistance to RAF inhibition.
82. PMID: 24265153; 2014, Cancer Discov;4(1):94-109
The genetic landscape of clinical resistance to RAF inhibition in metastatic melanoma.
83. PMID: 30269082; 2019, Gut;68(7):1152-1161
Liquid biopsies to track trastuzumab resistance in metastatic HER2-positive gastric cancer.
84. PMID: 26936308; 2016, J Clin Pharm Ther;41(3):357-359
Prolonged disease control with MEK inhibitor in neurofibromatosis type I-associated glioblastoma.
85. PMID: 26859683; 2016, Oncotarget;7(9):10547-56
Next-generation sequencing reveals somatic mutations that confer exceptional response to everolimus.
86. PMID: 32122926; 2020, Clin Cancer Res;26(12):2932-2945
MAPK Pathway Alterations Correlate with Poor Survival and Drive Resistance to Therapy in Patients with Lung Cancers Driven by ROS1 Fusions.
87. PMID: 22573716; 2012, Cancer Res;72(13):3350-9
Sensitivity of glioblastomas to clinically available MEK inhibitors is defined by neurofibromin 1 deficiency.
88. PMID: 23209032; 2013, Clin Cancer Res;19(2):450-61
Prognostic significance of AKT/mTOR and MAPK pathways and antitumor effect of mTOR inhibitor in NF1-related and sporadic malignant peripheral nerve sheath tumors.
89. PMID: 16793542; 2006, Mol Cell;22(6):719-729
Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2.
90. PMID: 23325218; 2013, Nature;493(7432):356-63
Fanconi anaemia and the repair of Watson and Crick DNA crosslinks.
91. PMID: 15905196; 2005, Carcinogenesis;26(10):1731-40
The Fanconi anemia group A protein modulates homologous repair of DNA double-strand breaks in mammalian cells.
92. PMID: 27037238; 2016, EMBO J;35(9):909-23
Interplay between Fanconi anemia and homologous recombination pathways in genome integrity.
93. PMID: 24153426; 2013, Nat Commun;4(1):2578
Heterozygous mutations in PALB2 cause DNA replication and damage response defects.
94. PMID: 20858716; 2010, Cancer Res;70(19):7353-9
PALB2/FANCN: recombining cancer and Fanconi anemia.
95. PMID: 25754594; 2015, Hum Mutat;36(5):562-8
Loss-of-Function FANCL Mutations Associate with Severe Fanconi Anemia Overlapping the VACTERL Association.

ACT Onco[®] + Report

96. PMID: 28678401; 2017, Cancer;123(20):3943-3954
Assessing the spectrum of germline variation in Fanconi anemia genes among patients with head and neck carcinoma before age 50.
97. PMID: 32086346; 2020, Clin Cancer Res;26(11):2487-2496
Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study.
98. PMID: 21135251; 2011, Mol Cancer Ther;10(1):3-8
Personalizing cancer treatment in the age of global genomic analyses: PALB2 gene mutations and the response to DNA damaging agents in pancreatic cancer.
99. PMID: 18483217; 2008, Genes Dev;22(10):1276-312
Role of platelet-derived growth factors in physiology and medicine.
100. PMID: 3013421; 1986, Cell;46(2):155-69
The biology of platelet-derived growth factor.
101. PMID: 20023271; 2010, Am J Clin Pathol;133(1):149-55
Detection of c-KIT and PDGFRA gene mutations in gastrointestinal stromal tumors: comparison of DHPLC and DNA sequencing methods using a single population-based cohort.
102. PMID: 24132921; 2013, Clin Cancer Res;19(24):6935-42
Large-scale analysis of PDGFRA mutations in melanomas and evaluation of their sensitivity to tyrosine kinase inhibitors imatinib and crenolanib.
103. PMID: 18772890; 2008, Nature;455(7216):1061-8
Comprehensive genomic characterization defines human glioblastoma genes and core pathways.
104. PMID: 19915670; 2009, PLoS One;4(11):e7752
Glioblastoma subclasses can be defined by activity among signal transduction pathways and associated genomic alterations.
105. PMID: 22323597; 2012, Proc Natl Acad Sci U S A;109(8):3041-6
Intratumoral heterogeneity of receptor tyrosine kinases EGFR and PDGFRA amplification in glioblastoma defines subpopulations with distinct growth factor response.
106. PMID: 20889717; 2010, Genes Dev;24(19):2205-18
PDGFRA gene rearrangements are frequent genetic events in PDGFRA-amplified glioblastomas.
107. PMID: 20685895; 2010, Cancer Res;70(18):7304-14
Coactivated platelet-derived growth factor receptor {alpha} and epidermal growth factor receptor are potential therapeutic targets in intimal sarcoma.
108. PMID: 16357008; 2006, Carcinogenesis;27(3):664-71
Mutation and expression of PDGFRA and KIT in malignant peripheral nerve sheath tumors, and its implications for imatinib sensitivity.
109. PMID: 19755855; 2009, Cancer Biol Ther;8(21):2042-50
Amplification of chromosomal segment 4q12 in non-small cell lung cancer.
110. PMID: 17189383; 2006, Mol Cancer Res;4(12):927-34
Amplification of KIT, PDGFRA, VEGFR2, and EGFR in gliomas.
111. PMID: 19366796; 2009, Cancer Res;69(9):3937-46
Ligand-dependent platelet-derived growth factor receptor (PDGFR)-alpha activation sensitizes rare lung cancer and sarcoma cells to PDGFR kinase inhibitors.
112. PMID: 30060824; 2018, Tissue Cell;53():30-36
Doxorubicin-resistant pleomorphic liposarcoma with PDGFRA gene amplification is targeted and regressed by pazopanib in a patient-derived orthotopic xenograft mouse model.

ACT Onco[®] + Report

113. PMID: 9315668; 1997, Mol Cell Biol;17(10):6087-96
hMre11 and hRad50 nuclear foci are induced during the normal cellular response to DNA double-strand breaks.
114. PMID: 16467875; 2006, Cell Res;16(1):45-54
The role of NBS1 in DNA double strand break repair, telomere stability, and cell cycle checkpoint control.
115. PMID: 16385572; 2006, Int J Cancer;118(11):2911-6
Evaluation of RAD50 in familial breast cancer predisposition.
116. PMID: 24894818; 2014, Breast Cancer Res;16(3):R58
Rare key functional domain missense substitutions in MRE11A, RAD50, and NBN contribute to breast cancer susceptibility: results from a Breast Cancer Family Registry case-control mutation-screening study.
117. PMID: 18440592; 2008, Hum Pathol;39(6):925-32
Gastric cancer with high-level microsatellite instability: target gene mutations, clinicopathologic features, and long-term survival.
118. PMID: 11196187; 2001, Cancer Res;61(1):36-8
Frameshift mutations at coding mononucleotide repeats of the hRAD50 gene in gastrointestinal carcinomas with microsatellite instability.
119. PMID: 24934408; 2014, Cancer Discov;4(9):1014-21
Synthetic lethality in ATM-deficient RAD50-mutant tumors underlies outlier response to cancer therapy.
120. PMID: 16474176; 2006, Carcinogenesis;27(8):1593-9
RAD50 and NBS1 are breast cancer susceptibility genes associated with genomic instability.
121. PMID: 27016230; 2016, Gynecol Oncol;141(1):57-64
Copy number deletion of RAD50 as predictive marker of BRCAness and PARP inhibitor response in BRCA wild type ovarian cancer.
122. PMID: 19029933; 2008, Oncogene;27(55):6908-19
LKB1; linking cell structure and tumor suppression.
123. PMID: 19584313; 2009, Physiol Rev;89(3):777-98
LKB1 and AMPK family signaling: the intimate link between cell polarity and energy metabolism.
124. PMID: 20142330; 2010, Dis Model Mech;3(3-4):181-93
Lkb1 inactivation is sufficient to drive endometrial cancers that are aggressive yet highly responsive to mTOR inhibitor monotherapy.
125. PMID: 17676035; 2007, Nature;448(7155):807-10
LKB1 modulates lung cancer differentiation and metastasis.
126. PMID: 18245476; 2008, Cancer Res;68(3):759-66
Loss of Lkb1 provokes highly invasive endometrial adenocarcinomas.
127. PMID: 18172296; 2008, Cancer Res;68(1):55-63
LKB1 deficiency sensitizes mice to carcinogen-induced tumorigenesis.
128. PMID: 25244018; 2014, Int J Mol Sci;15(9):16698-718
Recent progress on liver kinase B1 (LKB1): expression, regulation, downstream signaling and cancer suppressive function.
129. PMID: 9425897; 1998, Nat Genet;18(1):38-43
Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase.
130. PMID: 21189378; 2011, J Clin Oncol;29(6):e150-3
mTOR inhibitor treatment of pancreatic cancer in a patient With Peutz-Jeghers syndrome.
131. PMID: 27615706; 2016, CNS Oncol;5(4):203-9
Widely metastatic atypical pituitary adenoma with mTOR pathway STK11(F298L) mutation treated with everolimus therapy.
132. PMID: 27821489; 2017, Cancer Res;77(1):153-163
A Transcriptional Signature Identifies LKB1 Functional Status as a Novel Determinant of MEK Sensitivity in Lung Adenocarcinoma.

ACT Onco[®] + Report

133. PMID: 29764856; 2018, Clin Cancer Res;24(22):5710-5723
TP53, STK11, and EGFR Mutations Predict Tumor Immune Profile and the Response to Anti-PD-1 in Lung Adenocarcinoma.
134. PMID: 29773717; 2018, Cancer Discov;8(7):822-835
STK11/LKB1 Mutations and PD-1 Inhibitor Resistance in KRAS-Mutant Lung Adenocarcinoma.
135. PMID: 29337640; 2018, J Clin Oncol;36(7):633-641
Molecular Determinants of Response to Anti-Programmed Cell Death (PD)-1 and Anti-Programmed Death-Ligand 1 (PD-L1) Blockade in Patients With Non-Small-Cell Lung Cancer Profiled With Targeted Next-Generation Sequencing.
136. PMID: 26833127; 2016, Cancer Res;76(5):999-1008
STK11/LKB1 Deficiency Promotes Neutrophil Recruitment and Proinflammatory Cytokine Production to Suppress T-cell Activity in the Lung Tumor Microenvironment.
137. PMID: 24739573; 2014, Nat Rev Cancer;14(5):359-70
Unravelling mechanisms of p53-mediated tumour suppression.
138. PMID: 21125671; 2011, J Pathol;223(2):137-46
Haplo-insufficiency: a driving force in cancer.
139. PMID: 27998224; 2016, J Clin Oncol;34(36):4354-4361
Phase II Study of WEE1 Inhibitor AZD1775 Plus Carboplatin in Patients With TP53-Mutated Ovarian Cancer Refractory or Resistant to First-Line Therapy Within 3 Months.
140. PMID: 26646755; 2016, Ann Oncol;27(3):539-43
TP53 mutational status is predictive of pazopanib response in advanced sarcomas.
141. PMID: 25669829; 2015, Ann Oncol;26(5):1012-1018
Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation.
142. PMID: 27466356; 2016, Mol Cancer Ther;15(10):2475-2485
TP53 Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics.
143. PMID: 23670029; 2013, Oncotarget;4(5):705-14
P53 mutations in advanced cancers: clinical characteristics, outcomes, and correlation between progression-free survival and bevacizumab-containing therapy.
144. PMID: 17145525; 2006, Semin Oncol;33(5 Suppl 10):S8-14
Bevacizumab in combination with chemotherapy: first-line treatment of patients with metastatic colorectal cancer.
145. PMID: 21399868; 2011, Int J Oncol;38(5):1445-52
p53, HER2 and tumor cell apoptosis correlate with clinical outcome after neoadjuvant bevacizumab plus chemotherapy in breast cancer.
146. PMID: 20549698; 2011, Int J Cancer;128(8):1813-21
p53 status influences response to tamoxifen but not to fulvestrant in breast cancer cell lines.
147. PMID: 10786679; 2000, Cancer Res;60(8):2155-62
Complete sequencing of TP53 predicts poor response to systemic therapy of advanced breast cancer.
148. PMID: 25672981; 2015, Cancer Res;75(7):1187-90
VEGF-A Expression Correlates with TP53 Mutations in Non-Small Cell Lung Cancer: Implications for Antiangiogenesis Therapy.
149. PMID: 29573941; 2018, Lancet Oncol;19(5):603-615
Encorafenib plus binimetinib versus vemurafenib or encorafenib in patients with BRAF-mutant melanoma (COLUMBUS): a multicentre, open-label, randomised phase 3 trial.
150. PMID: 27480103; 2016, Lancet Oncol;17(9):1248-60
Cobimetinib combined with vemurafenib in advanced BRAF(V600)-mutant melanoma (coBRIM): updated efficacy results from a randomised,

ACT Onco[®] + Report

double-blind, phase 3 trial.

151. PMID: 26703889; 2016, Lancet;387(10022):968-977
Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study.
152. PMID: 22149876; 2012, N Engl J Med;366(6):520-9
Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer.
153. PMID: 21306238; 2011, N Engl J Med;364(6):514-23
Everolimus for advanced pancreatic neuroendocrine tumors.
154. PMID: 23158522; 2013, Lancet;381(9861):125-32
Efficacy and safety of everolimus for subependymal giant cell astrocytomas associated with tuberous sclerosis complex (EXIST-1): a multicentre, randomised, placebo-controlled phase 3 trial.
155. PMID: 18653228; 2008, Lancet;372(9637):449-56
Efficacy of everolimus in advanced renal cell carcinoma: a double-blind, randomised, placebo-controlled phase III trial.
156. PMID: 19805687; 2009, J Clin Oncol;27(31):5175-81
Improved early event-free survival with imatinib in Philadelphia chromosome-positive acute lymphoblastic leukemia: a children's oncology group study.
157. PMID: 12200353; 2002, Blood;100(6):1965-71
A phase 2 study of imatinib in patients with relapsed or refractory Philadelphia chromosome-positive acute lymphoid leukemias.
158. PMID: 21856226; 2011, Lancet Oncol;12(9):841-51
Nilotinib versus imatinib for the treatment of patients with newly diagnosed chronic phase, Philadelphia chromosome-positive, chronic myeloid leukaemia: 24-month minimum follow-up of the phase 3 randomised ENESTnd trial.
159. PMID: 18256322; 2008, Blood;111(8):4022-8
Imatinib pharmacokinetics and its correlation with response and safety in chronic-phase chronic myeloid leukemia: a subanalysis of the IRIS study.
160. PMID: 28196207; 2017, JAMA Oncol;3(7):944-952
Correlation of Long-term Results of Imatinib in Advanced Gastrointestinal Stromal Tumors With Next-Generation Sequencing Results: Analysis of Phase 3 SWOG Intergroup Trial S0033.
161. PMID: 27717299; 2016, N Engl J Med;375(22):2154-2164
Niraparib Maintenance Therapy in Platinum-Sensitive, Recurrent Ovarian Cancer.
162. PMID: 32343890; 2020, N Engl J Med;382(22):2091-2102
Olaparib for Metastatic Castration-Resistant Prostate Cancer.
163. PMID: 31851799; 2019, N Engl J Med;381(25):2416-2428
Olaparib plus Bevacizumab as First-Line Maintenance in Ovarian Cancer.
164. PMID: 31157963; 2019, N Engl J Med;381(4):317-327
Maintenance Olaparib for Germline BRCA-Mutated Metastatic Pancreatic Cancer.
165. PMID: 30345884; 2018, N Engl J Med;379(26):2495-2505
Maintenance Olaparib in Patients with Newly Diagnosed Advanced Ovarian Cancer.
166. PMID: 28578601; 2017, N Engl J Med;377(6):523-533
Olaparib for Metastatic Breast Cancer in Patients with a Germline BRCA Mutation.
167. PMID: 28754483; 2017, Lancet Oncol;18(9):1274-1284
Olaparib tablets as maintenance therapy in patients with platinum-sensitive, relapsed ovarian cancer and a BRCA1/2 mutation (SOLO2/ENGOT-Ov21): a double-blind, randomised, placebo-controlled, phase 3 trial.

ACT Onco[®] + Report

168. PMID: 27617661; 2016, Lancet Oncol;17(11):1579-1589
Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: an updated analysis from a randomised, placebo-controlled, double-blind, phase 2 trial.
169. PMID: 22595799; 2012, Lancet;379(9829):1879-86
Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial.
170. PMID: 20100962; 2010, J Clin Oncol;28(6):1061-8
Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial.
171. PMID: 24768112; 2014, Lancet;384(9940):319-28
Sorafenib in radioactive iodine-refractory, locally advanced or metastatic differentiated thyroid cancer: a randomised, double-blind, phase 3 trial.
172. PMID: 18650514; 2008, N Engl J Med;359(4):378-90
Sorafenib in advanced hepatocellular carcinoma.
173. PMID: 17189398; 2006, Clin Cancer Res;12(24):7271-8
Sorafenib for the treatment of advanced renal cell carcinoma.
174. PMID: 27924459; 2016, Target Oncol;11(6):815-824
Patient-Reported Outcomes and Quality of Life with Sunitinib Versus Placebo for Pancreatic Neuroendocrine Tumors: Results From an International Phase III Trial.
175. PMID: 27836885; 2017, Ann Oncol;28(2):339-343
Sunitinib in pancreatic neuroendocrine tumors: updated progression-free survival and final overall survival from a phase III randomized study.
176. PMID: 21306237; 2011, N Engl J Med;364(6):501-13
Sunitinib malate for the treatment of pancreatic neuroendocrine tumors.
177. PMID: 17227905; 2007, Oncologist;12(1):107-13
Food and Drug Administration drug approval summary: Sunitinib malate for the treatment of gastrointestinal stromal tumor and advanced renal cell carcinoma.
178. PMID: 27238653; 2016, Eur Urol;70(6):1006-1015
Early Tumour Shrinkage: A Tool for the Detection of Early Clinical Activity in Metastatic Renal Cell Carcinoma.
179. PMID: 16757724; 2006, JAMA;295(21):2516-24
Sunitinib in patients with metastatic renal cell carcinoma.
180. PMID: 25577718; 2015, Eur Urol;67(5):952-8
Depth of remission is a prognostic factor for survival in patients with metastatic renal cell carcinoma.
181. PMID: 17046465; 2006, Lancet;368(9544):1329-38
Efficacy and safety of sunitinib in patients with advanced gastrointestinal stromal tumour after failure of imatinib: a randomised controlled trial.
182. PMID: 30110579; 2018, N Engl J Med;379(8):753-763
Talzoparib in Patients with Advanced Breast Cancer and a Germline BRCA Mutation.
183. PMID: 17538086; 2007, N Engl J Med;356(22):2271-81
Temsilolimus, interferon alfa, or both for advanced renal-cell carcinoma.
184. PMID: 29072975; 2018, J Clin Oncol;36(1):7-13
Dabrafenib and Trametinib Treatment in Patients With Locally Advanced or Metastatic BRAF V600-Mutant Anaplastic Thyroid Cancer.
185. PMID: 27080216; 2016, Lancet Oncol;17(5):642-50
Dabrafenib in patients with BRAF(V600E)-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, phase 2 trial.
186. PMID: 25265492; 2014, N Engl J Med;371(20):1877-88
Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma.

ACT Onco[®] + Report

187. PMID: 22663011; 2012, N Engl J Med;367(2):107-14
Improved survival with MEK inhibition in BRAF-mutated melanoma.