

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Prostate acinar adenocarcinoma	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Prostate
	NAME Sun, I Li		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S112-01461 E (PF23064)
	DATE OF BIRTH 03 October 1949		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Male		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 11 January 2023
	MEDICAL RECORD # 44360125		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 25 May 2023

Biomarker Findings

Microsatellite status - MSI-High

Tumor Mutational Burden - 350 Muts/Mb

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

BARD1 R406*	ERRF1 F411fs*41
BRCA2 D946fs*14	FH splice site 1237-1G>A
BRIP1 splice site 918+1G>A - subclonal†	FUBP1 I301fs*4
CDK12 R1048*, N474fs*8 - subclonal†	IRF2 N167fs*7
CHEK1 T226fs*14	KMT2D (MLL2) A221fs*40
FANCL G119*	MAPK1 R191H
EGFR G724S	MLH1 R127I
APC R876*	MSH2 loss
PIK3CA E970K	MSH3 N385fs*19
PTEN P246L, K267fs*9, R173C	MSH6 loss
RNF43 G659fs*41, H683fs*17	NOTCH3 G463*
TSC2 R57H	NTRK2 A11V
ASXL1 G646fs*12	PMS2 Y268fs*39
ATR F222fs*11	PRKAR1A Q275*
CASP8 R250Q, Q308*	QKI K134fs*14, R124*
CBL splice site 1007+2T>C	SETD2 R1407fs*5
CD79A V69I - subclonal†	SMAD4 G386S - subclonal†
CDK8 Q331H - subclonal†	SPEN K1999fs*23
CTCF T204fs*26	TGFB2 R497* - subclonal†
CTNNA1 K695fs*27	TP53 R196*, Y205H - subclonal, R175C†
EPHB4 V420I	XRCC2 L117fs*17

8 Disease relevant genes with no reportable alterations: **ATM**, **BRCA1**, **CHEK2**, **PALB2**, **RAD51B**, **RAD51C**, **RAD51D**, **RAD54L**

† See About the Test in appendix for details.

Report Highlights

- Targeted therapies with **NCCN categories of evidence** in this tumor type: Olaparib (p. 33), Pembrolizumab (p. 35), Rucaparib (p. 36)
- Variants that may inform **nontargeted treatment approaches** (e.g., chemotherapy) in this tumor type: **BRCA2**D946fs*14 (p. 8), **CDK12**N474fs*8, R1048* (p. 10)
- Evidence-matched **clinical trial options** based on this patient's genomic findings: (p. 45)
- Variants with **prognostic implications** for this tumor type that may impact treatment decisions: **BRCA2**D946fs*14 (p. 8), **PTEN**K267fs*9, P246L, R173C (p. 14), **TP53**R175C, R196*, Y205H (p. 30)
- Variants in select cancer susceptibility genes to consider for possible **follow-up germline testing** in the appropriate clinical context: **BRCA2**D946fs*14 (p. 8), **MLH1**R127I (p. 23), **PMS2**Y268fs*39 (p. 26)
- Variants that may represent **clonal hematopoiesis** and may originate from non-tumor sources: **ASXL1**G646fs*12 (p. 16), **KMT2D (MLL2)**A221fs*40 (p. 22)

PATHOLOGIST COMMENTS

Brennan Decker, M.D., Ph.D. 04-Jun-2023

The very high TMB in this case is likely attributable to the observed microsatellite instability - high (MSI-H) phenotype. In the setting of MSI-H associated hypermutation, co-occurring mutations are often subclonal passengers resulting from the large number of mutations, rather than acting as traditional drivers. This may impact

potential for therapy response (e.g., for the homologous recombination pathway alterations and PARPi, in this case; PMID: 35772050). High TMB resulting from MSI-H has been associated with response to immunotherapy. Please see the Professional Services section below for additional information.

BIOMARKER FINDINGS

Microsatellite status - MSI-High

10 Trials see p. 45

Tumor Mutational Burden - 350 Muts/Mb

10 Trials see p. 47

GENOMIC FINDINGS

BARD1 - R406*

10 Trials see p. 50

BRCA2 - D946fs*14

10 Trials see p. 52

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

Pembrolizumab 2A

Dostarlimab

Pembrolizumab 2A

Dostarlimab

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

Olaparib 1

Rucaparib

Olaparib 1

Rucaparib 2A

Olaparib + Abiraterone

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Atezolizumab

Avelumab

Cemiplimab

Durvalumab

Durvalumab + Tremelimumab

Nivolumab

Nivolumab + Ipilimumab

Retifanlimab

Atezolizumab

Avelumab

Cemiplimab

Durvalumab

Nivolumab

Nivolumab + Ipilimumab

Retifanlimab

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

Niraparib

Talazoparib

Niraparib

Talazoparib

☐ NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

GENOMIC FINDINGS	THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)	THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
BRIP1 - splice site 918+1G>A - subclonal	Olaparib 1	Niraparib
10 Trials see p. 54	Rucaparib	Talazoparib
CDK12 - R1048*, N474fs*8 - subclonal	Olaparib 1	Cemiplimab
	Dostarlimab	Niraparib
	Pembrolizumab	Nivolumab
	Rucaparib	Retifanlimab
		Talazoparib
10 Trials see p. 56		
CHEK1 - T226fs*14	Olaparib 1	Niraparib
10 Trials see p. 58	Rucaparib	Talazoparib
FANCL - G119*	Olaparib 1	Niraparib
10 Trials see p. 61	Rucaparib	Talazoparib
EGFR - G724S	none	Erlotinib
5 Trials see p. 60		Gefitinib
APC - R876*	none	none
2 Trials see p. 49		
PIK3CA - E970K	none	none
10 Trials see p. 63		
PTEN - P246L, K267fs*9, R173C	none	none
10 Trials see p. 65		
RNF43 - G659fs*41, H683fs*17	none	none
2 Trials see p. 67		
TSC2 - R57H	none	none
10 Trials see p. 68		

1 NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING IN SELECT CANCER SUSCEPTIBILITY GENES

Findings below have been previously reported as pathogenic germline in the ClinVar genomic database and were detected at an allele frequency of >10%. See appendix for details.

BRCA2 - D946fs*14 p. 8 **PMS2** - Y268fs*39 p. 26
MLH1 - R127I p. 23

This report does not indicate whether variants listed above are germline or somatic in this patient. In the appropriate clinical context, follow-up germline testing would be needed to determine whether a finding is germline or somatic.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS (CH)

Genomic findings below may include nontumor somatic alterations, such as CH. The efficacy of targeting such nontumor somatic alterations is unknown. This content should be interpreted based on clinical context. Refer to appendix for additional information on CH.

ASXL1 - G646fs*12 p. 16 **KMT2D (MLL2)** - A221fs*40 p. 22

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

ASXL1 - G646fs*12 p. 16	MLH1 - R127I p. 23
ATR - F222fs*11 p. 16	MSH2 - loss p. 24
CASP8 - R250Q, Q308* p. 17	MSH3 - N385fs*19 p. 24
CBL - splice site 1007+2T>C p. 17	MSH6 - loss p. 25
CD79A - V69I - subclonal p. 18	NOTCH3 - G463* p. 25
CDK8 - Q331H - subclonal p. 18	NTRK2 - A11V p. 26
CTCF - T204fs*26 p. 19	PMS2 - Y268fs*39 p. 26
CTNNA1 - K695fs*27 p. 19	PRKAR1A - Q275* p. 27
EPHB4 - V420I p. 20	QKI - K134fs*14, R124* p. 27
ERF1 - F411fs*41 p. 20	SETD2 - R1407fs*5 p. 27
FH - splice site 1237-1G>A p. 21	SMAD4 - G386S - subclonal p. 28
FUBP1 - I301fs*4 p. 21	SPEN - K1999fs*23 p. 28
IRF2 - N167fs*7 p. 22	TGFBR2 - R497* - subclonal p. 29
KMT2D (MLL2) - A221fs*40 p. 22	TP53 - R196*, Y205H - subclonal, R175C p. 30
MAPK1 - R191H p. 23	XRCC2 - L117fs*17 p. 31

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

MSI-High

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in multiple solid tumor types, microsatellite instability (MSI) and associated increased tumor mutational burden (TMB)¹⁻² may predict sensitivity to immune checkpoint inhibitors, including the approved PD-1-targeting agents cemiplimab, dostarlimab, nivolumab (alone or in combination with ipilimumab), retifanlimab, and pembrolizumab³⁻⁹, as well as PD-L1-targeting agents atezolizumab, avelumab, and durvalumab (alone or in combination with tremelimumab)¹⁰⁻¹¹.

FREQUENCY & PROGNOSIS

MSI has been reported in 3.1-14.6% of prostate cancer samples¹²⁻¹⁶. A study of prostate cancer in hereditary nonpolyposis colorectal cancer (HNPCC) families reported MSI-H in 4-50% of cases¹⁷⁻¹⁹. For patients with advanced prostate cancer, dMMR/MSI status was associated with shorter median OS compared with patients with proficient MMR (3.8 vs. 7.0 years) by univariate and multivariate analysis (adjusted HR=4.09; P=0.005)²⁰.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor²¹. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2²¹⁻²³. This sample has a high

level of MSI, equivalent to the clinical definition of an MSI-high (MSI-H) tumor: one with mutations in >30% of microsatellite markers²⁴⁻²⁶. MSI-H status indicates high-level deficiency in MMR and typically correlates with loss of expression of at least one, and often two, MMR family proteins^{21,23,25-26}.

POTENTIAL GERMLINE IMPLICATIONS

While approximately 80% of MSI-H tumors arise due to somatic inactivation of an MMR pathway protein, about 20% arise due to germline mutations in one of the MMR genes²¹, which are associated with a condition known as Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC)²⁷. Lynch syndrome leads to an increased risk of colorectal, endometrial, gastric, and other cancers²⁷⁻²⁹ and has an estimated prevalence in the general population ranging from 1:600 to 1:2000³⁰⁻³². Therefore, in the appropriate clinical context, germline testing of MLH1, MSH2, MSH6, and PMS2 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

350 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1³³⁻³⁵, anti-PD-1 therapies³³⁻³⁶, and combination nivolumab and ipilimumab³⁷⁻⁴². In multiple pan-tumor studies, increased tissue tumor mutational burden (TMB) was associated with sensitivity to immune checkpoint inhibitors^{33-36,43-47}. In the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors, significant improvement in ORR was observed for patients with TMB ≥ 10 Muts/Mb (as measured by this assay) compared with those with TMB < 10 Muts/Mb in a large cohort that included multiple tumor types⁴³; similar findings were observed in the KEYNOTE 028 and 012 trials³⁶. At the same TMB cutpoint, retrospective analysis of patients with solid tumors treated with any checkpoint inhibitor identified that tissue TMB scores ≥ 10 Muts/Mb were associated with prolonged time to treatment failure compared with scores < 10 muts/Mb (HR=0.68)⁴⁷. For patients with solid tumors treated with nivolumab plus

ipilimumab in the CheckMate 848 trial, improved responses were observed in patients with a tissue TMB ≥ 10 Muts/Mb independent of blood TMB at any cutpoint in matched samples⁴⁸. However, support for higher TMB thresholds and efficacy was observed in the prospective Phase 2 MyPathway trial of atezolizumab for patients with pan-solid tumors, where improved ORR and DCR was seen in patients with TMB ≥ 16 Muts/Mb than those with TMB ≥ 10 and < 16 Muts/Mb⁴⁶. Similarly, analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy compared with patients with higher TMB treated with chemotherapy⁴⁹ or those with lower TMB treated with PD-1 or PD-L1-targeting agents³⁴. The Phase 2 CheckMate 650 trial of nivolumab and ipilimumab treatment for patients with metastatic castration-resistant prostate cancer reported that patients harboring above the median study TMB experienced increased ORR and PSA responses⁴². A real-world study for patients with pretreated metastatic castration-resistant prostate cancer (mCRPC) reported longer time to next therapy (TTNT) for patients with high tumor mutational burden (TMB; ≥ 10 Muts/Mb) treated with immune checkpoint inhibitors (ICIs) compared with those treated with taxane chemotherapies, whereas for patients with low TMB (< 10 Muts/Mb), treatment with ICIs resulted in worse TTNT compared with taxanes⁵⁰.

FREQUENCY & PROGNOSIS

Prostate acinar adenocarcinoma harbors a median

TMB of 2.7 mutations per megabase (muts/Mb), and 3.4% of cases have high TMB (> 20 muts/Mb)⁵¹. Prostate cancer has been reported to harbor a relatively low TMB among solid tumors⁵²⁻⁵³, with approximately 0.5-1.5 (muts/Mb) in localized tumor samples⁵⁴⁻⁵⁶, and a higher but still low TMB of 2-5 muts/Mb in metastatic, castration-resistant prostate cancer (mCRPC) samples⁵⁷⁻⁵⁹. One study reported that 4 of 150 (2.7%) mCRPC cases harbored high TMB (nearly 50 muts/Mb), which was due to defects in mismatch repair genes MLH1 and MSH2 in 3 of the 4 cases⁵⁹. Published data investigating the prognostic implications of tumor mutational burden in prostate cancer are limited (PubMed, Feb 2023).

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁶⁰⁻⁶¹ and cigarette smoke in lung cancer^{8,62}, treatment with temozolomide-based chemotherapy in glioma⁶³⁻⁶⁴, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes⁶⁵⁻⁶⁹, and microsatellite instability (MSI)^{65,68-69}. This sample harbors a TMB level that may be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors in multiple solid tumor types^{34-36,43}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

BARD1

ALTERATION

R406*

HGVS VARIANT

NM_000465.2:c.1216C>T (p.R406*)

VARIANT CHROMOSOMAL POSITION

chr2:215645382

VARIANT ALLELE FREQUENCY (% VAF)

40.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

Clinical benefit from rucaparib has been observed in a patient with BARD1-mutated ovarian cancer⁷⁰. On the basis of preclinical evidence, tumors with BARD1 inactivation may be sensitive to PARP inhibitors⁷¹⁻⁷⁴.

FREQUENCY & PROGNOSIS

BARD1 mutations have been reported in 0-2% of prostate adenocarcinoma samples^{54-55,57,75}. Published data investigating the prognostic implications of BARD1 alterations in prostate

carcinoma are limited (PubMed, Mar 2023).

FINDING SUMMARY

BARD1 encodes the BRCA1-associated RING domain 1 protein, which is required for stabilization and nuclear localization of BRCA1 as well as formation of the E3 ubiquitin ligase⁷⁶. The BARD1 ANK repeats and BRCT motifs play important roles in chromosome stability, and both these regions and the RING domain are necessary for homology-directed repair^{71,77-78}. Alterations such as seen here may disrupt BARD1 function or expression.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

BRCA2

ALTERATION

D946fs*14

HGVS VARIANT

NM_000059.3:c.2835del (p.D946fs*14)

VARIANT CHROMOSOMAL POSITION

chr13:32911321-32911322

VARIANT ALLELE FREQUENCY (% VAF)

37.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Alterations that inactivate BRCA1 or BRCA2 may confer sensitivity to PARP inhibitors⁷⁹⁻⁹⁶ or ATR inhibitors⁹⁷⁻⁹⁹. Clinical responses to PARP inhibitors have been reported for patients with either germline or somatic BRCA1/2 mutations^{80,85,88,95-96} and for patients with platinum-resistant or -refractory disease^{79,84,91,94}. The WEE1 inhibitor adavosertib has been evaluated as a monotherapy and in combination with PARP-inhibitor, olaparib. In a Phase 2 study for patients with PARP-resistant ovarian cancer, the combination of olaparib and adavosertib elicited improved clinical benefit (ORR: 29%; DCR: 89%) compared to adavosertib alone (ORR: 23%; DCR: 63%); however, in the BRCA-mutated cohort, no significant difference in clinical benefit was observed between the combination (ORR: 19%) and monotherapy (ORR: 20%) treatments¹⁰⁰. In a Phase 1 monotherapy trial of adavosertib that included 9 patients with BRCA1/2-mutated solid tumors, 2 patients with BRCA1-mutated cancers (1 with ovarian serous carcinoma and 1 with oral squamous cell carcinoma) achieved PRs, and a third patient with ovarian serous carcinoma harboring mutations in BRCA1 and TP53 experienced 14% tumor shrinkage prior to disease progression¹⁰¹. In a case study, a patient with therapy-induced neuroendocrine prostate cancer and an inactivating BRCA2 rearrangement experienced a CR ongoing for 20 months to the ATR inhibitor berzosertib⁹⁹. Preclinical studies of BRCA1/2 inactivation in T-cell acute lymphoblastic leukemia (T-ALL)¹⁰², ovarian carcinoma¹⁰³, and triple-negative breast cancer (TNBC)¹⁰⁴ showing reduced cell viability and increased DNA damage during ATR treatment further support the sensitivity of BRCA2-deficient cells to ATR inhibitors. The Phase 3 PROfound study for patients with metastatic castration-

resistant prostate cancer (CRPC) who had progressed on a new hormonal agent reported improved radiographic PFS with olaparib compared with physician's choice of abiraterone/prednisone or enzalutamide for patients with BRCA1/2 or ATM alterations (7.4 vs. 3.6 mo., HR=0.34)¹⁰⁵. The Phase 1/2 PETRA study of PARP1 inhibitor AZD5305 observed a 25% (10/40) ORR (10 PR), for patients with BRCA1/2, PALB2, or RAD51C-mutated breast, ovarian, pancreatic, or prostate cancer, including 5 patients with prior PARP inhibitor treatment¹⁰⁶.

— Nontargeted Approaches —

Alterations in DNA repair genes such as BRCA1, BRCA2, ATM, PALB2, FANCA, RAD51D, CHEK2, and CDK12 have been reported to be predictive for sensitivity to platinum agents in castration resistant prostate cancer (CRPC) (NCCN Prostate Cancer Guidelines, v1.2023)¹⁰⁷⁻¹¹⁰. Clinical data from small retrospective studies¹¹¹ and case reports¹¹²⁻¹¹⁶ are conflicting as to whether alterations in DNA repair genes such as BRCA1, BRCA2, and ATM are associated with outcomes for patients with CRPC treated with PSMA-targeted radionuclide-based therapies such as lutetium Lu 177 vipivotide tetraxetan (177Lu-PSMA-617). Germline BRCA2 mutations in mCRPC were associated with relative benefit from first-line abiraterone or enzalutamide compared with taxanes (CSS 24.0 vs. 17.0 months, PFS on the second systemic therapy 18.9 vs. 8.6 months) in a large prospective cohort study¹¹⁷. Three patients with non-neuroendocrine prostate cancer harboring BRCA2 mutations derived clinical benefit from treatment with platinum-based chemotherapy^{107,118}. Inactivation of BRCA2 may also predict sensitivity to DNA-damaging drugs such as trabectedin, lurbinectedin, and the platinum chemotherapies cisplatin and carboplatin¹¹⁹⁻¹²⁹.

FREQUENCY & PROGNOSIS

BRCA2 mutations have been identified in 3-6% of primary and 6-7% of metastatic prostate cancer specimens^{56,59,130}, with deleterious germline BRCA2 mutations present for 5% of patients with metastatic prostate cancer¹³¹. BRCA2 homozygous deletion has been reported in 3-6% of prostate adenocarcinoma cases^{75,132-133}. The positive predictive value of prostate specific antigen (PSA) levels was found to be higher in patients with BRCA1/2 mutations than in the general population¹³⁴. BRCA2 germline mutations predict

poor prognosis for patients with prostate cancer (NCCN Prostate Cancer Guidelines, v1.2023)¹³⁵⁻¹⁴¹ and have been associated with attributes of aggressive prostate cancer at diagnosis, including high Gleason score, nodal involvement, advanced tumor stage, and metastatic spread¹³⁵. An exome sequencing study of aggressive and non-aggressive prostate cancer cases reported that patients with pathogenic, likely pathogenic, or deleterious BRCA2 mutations had statistically higher odds of aggressive disease (4.3-fold), prostate cancer death (4.7-fold), and metastatic disease (5.7-fold)¹⁴². In another study, germline BRCA2 mutation carriers had a significantly shorter cause-specific survival (CSS, 8.6 vs. 15.7 years) than noncarriers¹³⁵. Following radical conventional treatment for localized prostate cancer, patients with germline BRCA1/2 mutations experienced significantly shorter metastasis-free survival (HR=2.36) and CSS (HR=2.17) than noncarriers¹⁴³. Patients with germline or somatic BRCA2 mutations have been reported to experience significantly shorter time to castration-resistant disease progression on first-line abiraterone or enzalutamide treatment (9.4 vs. 18.4 months, HR=1.58) and significantly shorter OS (16.9 vs. 21.8 months, HR=1.64)¹⁴⁴. For patients with metastatic castration-resistant prostate cancer (mCRPC), germline BRCA2 mutations were an independent marker of poor prognosis (CSS 17.4 vs. 33.2 months, HR=2.11) in 1 study¹¹⁷.

FINDING SUMMARY

The BRCA2 tumor suppressor gene encodes a protein that regulates the response to DNA damage¹⁴⁵. Inactivating mutations in BRCA2 can lead to the inability to repair DNA damage and loss of cell cycle checkpoints, which can lead to tumorigenesis¹⁴⁶. Alterations such as seen here may disrupt BRCA2 function or expression^{145,147-162}.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the BRCA2 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with hereditary breast and ovarian cancer syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Inactivating germline mutations in BRCA1 or BRCA2 are associated with autosomal dominant hereditary breast and ovarian cancer¹⁶⁴⁻¹⁶⁵, and the lifetime risk of breast and ovarian cancer in BRCA2 mutation carriers has

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

been estimated to be as high as >80% and 23%, respectively¹⁶⁶. Elevated risk for other cancer types, including gastric, pancreatic, prostate, and colorectal, has also been identified, with an

increase in risk ranging from 20 to 60%¹⁶⁷. The estimated prevalence of deleterious germline BRCA1/2 mutations in the general population is between 1:400 and 1:800, with an approximately

10-fold higher prevalence in the Ashkenazi Jewish population^{166,168-173}. In the appropriate clinical context, germline testing of BRCA2 is recommended.

GENE

BRIP1

ALTERATION

splice site 918+1G>A - subclonal

HGVS VARIANT

NM_032043.2:c.918+1G>A (p.?)

VARIANT CHROMOSOMAL POSITION

chr17:59885827

VARIANT ALLELE FREQUENCY (% VAF)

6.8%

FREQUENCY & PROGNOSIS

BRIP1 mutation has been reported in <1% of prostate cancer samples (cBioPortal, Apr 2023)¹⁷⁶⁻¹⁷⁷. Published data investigating the prognostic implications of BRIP1 alterations in prostate cancer are limited (PubMed, Aug 2022).

FINDING SUMMARY

BRIP1, also known as FANCI (Fanconi Anemia complementation group J) and BACH1, encodes a DNA helicase required for DNA repair and the maintenance of chromosomal stability¹⁷⁸⁻¹⁸⁰. Alterations such as seen here may disrupt BRIP1 function or expression^{178,181-183}.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the BRIP1 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters)

associated with familial breast cancer (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in BRIP1 are associated with increased risk of breast, ovarian, and cervical cancers¹⁸⁴⁻¹⁸⁸. Germline mutations in BRIP1 are also associated with Fanconi anemia (FA), a rare autosomal disorder that predisposes patients to a subset of cancers, including acute myeloid leukemia (AML), myelodysplastic syndrome (MDS), gynecological malignancies, and head and neck tumors¹⁸⁹⁻¹⁹¹; frequency estimates suggest an incidence of 3/1,000,000 individuals in Europe and the US, and a heterozygote carrier frequency of 1/181 to 1/300 in the US and Europe, respectively, with slightly higher rates in some populations, such as the Ashkenazi Jewish population (1/89)^{190,192}. In the appropriate clinical context, germline testing of BRIP1 is recommended.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

On the basis of clinical responses in ovarian cancer⁸⁵ and prostate cancer¹⁷⁴, as well as clinical benefit in breast cancer¹⁷⁵, loss or inactivation of BRIP1 may confer sensitivity to PARP inhibitors.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

CDK12

ALTERATION

R1048*, N474fs*8 - subclonal

HGVS VARIANT

NM_016507.2:c.3142C>T (p.R1048*),

NM_016507.2:c.1421del (p.N474fs*8)

VARIANT CHROMOSOMAL POSITION

chr17:37680973, chr17:37627500-37627501

VARIANT ALLELE FREQUENCY (% VAF)

38.9%, 7.8%

IMPACT trial for patients with CDK12-inactivated mCRPC reported PSA $\geq 50\%$ decline in 14% (4/28) of patients and unconfirmed PSA $\geq 30\%$ decline in 21% (6/28) of patients following treatment with nivolumab plus ipilimumab²⁰². Preclinical data suggest CDK12 inactivating alterations may sensitize cells to PARP inhibitors²⁰³⁻²⁰⁸, and the Phase 3 PROfound study reported numerically improved PFS for patients with CDK12-altered tumors treated with olaparib compared to control androgen deprivation therapy (5.1 vs. 2.2 months)²⁰⁹. However, multiple clinical studies have observed no radiographic responses for patients with CDK12-altered CRPC treated with PARP inhibitors^{174,210}.

in mCRPC than in primary prostate cancer, where CDK12 mutations have been observed in 1.2% of patients¹⁹⁴. Retrospective analyses of prostate cancer have found that CDK12 alterations are associated with high-risk features, including high Gleason scores and de novo metastases, with most patients developing CRPC^{200,212,214}. Retrospective analyses of prostate cancer found that among patients presenting with localized disease, CDK12 mutation was associated with shorter time to metastasis (34.9 vs. 55.6-64.7 months, HR=0.46-0.59) and earlier development of CRPC (32.7 vs. 56.2-72.8 months, HR=0.42-0.51) compared with the non-CDK12 study cohorts²⁰⁰; another retrospective analysis reported that patients with CDK12 alterations experienced earlier development of CRPC prostate-specific antigen PFS on first-line abiraterone or enzalutamide treatment¹⁴⁴. Additional studies report that CDK12 alteration or biallelic loss is associated with shorter OS (64.4 vs. 74.9 months, HR=1.70 and 61.2 vs. 76.8 months, HR=1.65, respectively)^{212,214}. A study of 59 patients with hormone-sensitive prostate cancer identified MSH2, CDK12, RB1, and TP53 alterations as independent risk factors for early progression to castration-resistant prostate cancer²¹⁵.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

CDK12 inactivation in cancer is associated with genomic instability characterized by tandem duplications¹⁹³⁻¹⁹⁷ and has been shown to increase tumor immunogenicity in advanced prostate cancer¹⁹⁴. On the basis of clinical and preclinical evidence in prostate cancer, CDK12 inactivation may predict benefit from immune checkpoint inhibitors^{194,198-199}. Retrospective studies observed prostate-specific antigen (PSA) response rates of 11-50% (2/19-2/4) for patients with CDK12-mutated metastatic prostate cancer on immune checkpoint inhibitors (predominantly anti-PD-1 monotherapy)^{194,198-200}. The prospective NEPTUNES trial of nivolumab plus ipilimumab reported a composite response rate of 14% (1/7) in CDK12-mutated metastatic castration-resistant prostate cancer (mCRPC), with the responder also having high tumor-infiltrating lymphocytes²⁰¹. In a similar treatment population, the prospective

— Nontargeted Approaches —

Cells lacking CDK12 incur spontaneous DNA damage and exhibit heightened sensitivity to DNA-damaging agents²⁰³⁻²⁰⁸. Alterations in DNA repair genes such as BRCA1, BRCA2, ATM, PALB2, FANCA, RAD51D, CHEK2, and CDK12 have been reported to be predictive for sensitivity to platinum agents in castration resistant prostate cancer (CRPC) (NCCN Prostate Cancer Guidelines, v1.2023)¹⁰⁷⁻¹¹⁰. Studies on the impact of CDK12 alterations on response to taxanes for patients with metastatic castration-resistant prostate cancer (mCRPC) are conflicting (NCCN Prostate Cancer Guidelines, v1.2023)^{198,211}.

FREQUENCY & PROGNOSIS

CDK12 mutation has been reported in 4.3-6.9% of patients with metastatic castration-resistant prostate cancer (mCRPC)^{194,212-214}. The frequency of CDK12 mutations has been reported to be higher

FINDING SUMMARY

CDK12 encodes a cyclin-dependent kinase that interacts with cyclin K to regulate the phosphorylation of RNA polymerase II and the expression of genes involved in maintaining genomic stability, including BRCA1 and ATR²¹⁶. Alterations such as seen here may disrupt CDK12 function or expression^{206,217-219}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS
GENE

CHEK1

ALTERATION

T226fs*14

HGVS VARIANT

NM_001274.4:c.676del (p.T226Hfs*14)

VARIANT CHROMOSOMAL POSITION

chr11:125505377-125505378

VARIANT ALLELE FREQUENCY (% VAF)

41.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

In preclinical studies, loss of CHEK1 activity has been shown to sensitize cancer cells to DNA-damaging agents, including PARP inhibitors²²⁰⁻²²⁴. CHEK1 inhibitors have shown clinical benefit for

patients with a variety of solid and hematologic cancers, but it is not known if CHEK1 mutation confers sensitivity to these inhibitors²²⁵⁻²²⁸.

FREQUENCY & PROGNOSIS

CHEK1 mutations have been generally observed at low frequencies in cancer, with greatest prevalence in non-melanoma skin cancer (2.7%), endometrial cancer (1.8%), and melanoma (1.4%)²²⁹. CHEK1 expression was reported to be higher in TNBC and basal-like breast cancer as compared with benign lesions and other types of malignant breast cancer²³⁰⁻²³¹. One study found that CHEK1 was the most upregulated kinase in glioblastoma and was overexpressed in all subtypes compared with healthy cells or to low-grade glioma²³². However, CHEK1 deletion or methylation was detected more frequently in cervical carcinoma than in premalignant cervical lesions (80% vs. 35%, respectively), and CHEK1 expression was also

reduced in these cells as compared with normal cervical tissues²³³. CHEK1 alterations have been associated with reduced survival for patients with cervical cancer²³³. High CHEK1 expression or phosphorylation correlated with aggressive phenotypes and a poorer prognosis in glioma, breast cancer, hepatocellular carcinoma, and non-small cell lung cancer^{232,234-236}.

FINDING SUMMARY

CHEK1 encodes CHK1, a serine/threonine protein kinase that is required for checkpoint-mediated cell cycle arrest in response to DNA damage and maintains genome integrity by activating homologous recombination repair^{223,237-240}. In more than half of cancers, the ATM/CHK2/p53 pathway is compromised, and the ATR/CHK1/CDC25 pathway provides cell-cycle arrest needed for repair of DNA damage²³⁸. Alterations such as seen here may disrupt CHEK1 function or expression²⁴¹⁻²⁴⁴.

GENE

FANCL

ALTERATION

G119*

HGVS VARIANT

NM_018062.3:c.355G>T (p.G119*)

VARIANT CHROMOSOMAL POSITION

chr2:58449096

VARIANT ALLELE FREQUENCY (% VAF)

40.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies that directly address

genomic alterations in FANCL. Clinical evidence in ovarian cancer indicates that FANCL inactivation may confer sensitivity to PARP inhibitors^{70,245}.

FREQUENCY & PROGNOSIS

FANCL mutations are most frequently observed in tumors of the prostate (5.3%) and liver (4.0%), and are seen at lower frequency in other tumor types (COSMIC, May 2023)²⁴⁶. Published data investigating the prognostic implications of FANCL alterations in solid tumors and hematologic malignancies are limited (PubMed, May 2023). In a prospective study of 255 patients with follicular lymphoma, 2p gain, which includes VRK2, FANCL, and LINC01122, was associated with worse PFS and OS in multivariate analysis²⁴⁷.

FINDING SUMMARY

FANCL encodes a member of the Fanconi anemia nuclear complex, a multiprotein structure also including the products of FANCA, FANCC, FANCF and FANCG. The activity of this complex is essential to prevention of chromosome breakage caused by DNA damage²⁴⁸. Germline mutations in FANCL cause Fanconi anemia, a clinically heterogeneous disorder involving various developmental abnormalities as well as predisposition to cancer; underlying these phenotypes are defects in DNA repair²⁴⁹. Alterations such as seen here may disrupt FANCL function or expression²⁵⁰⁻²⁵⁷. Germline mutations in FANCL, such as T367fs*13, have been associated with Fanconi anemia, breast cancer, and ovarian cancer and with an increased risk of esophageal cancer and prostate cancer^{191,258-261}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

EGFR

ALTERATION

G724S

HGVS VARIANT

NM_005228.3:c.2170G>A (p.G724S)

VARIANT CHROMOSOMAL POSITION

chr7:55241722

VARIANT ALLELE FREQUENCY (% VAF)

37.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

For patients with non-small cell lung cancer (NSCLC), EGFR activating mutations may predict sensitivity to EGFR-TKIs, including erlotinib²⁶², gefitinib²⁶³⁻²⁶⁶, afatinib²⁶⁷⁻²⁷⁰, dacomitinib²⁷¹, and osimertinib^{268,272}; however, the data for patients with other tumor types are limited²⁷³⁻²⁷⁸.

— Potential Resistance —

EGFR G724S mediates resistance to osimertinib; notably, virtually all reported evidence for clinical and preclinical resistance of G724S to this agent has been in the context of co-occurrence with activating exon 19 deletions²⁷⁹⁻²⁸⁵ irrespective of the presence of T790M^{279-283,286}. G724S co-occurring with other activating alterations is rare²⁸⁵, and, therefore, it is unclear whether EGFR G724S would reduce sensitivity to osimertinib outside of EGFR exon 19 deletions. G724S also reduces resistance to first-generation EGFR inhibitors, with a retrospective analysis reporting emergence of G724S in 4 patients with non-small cell lung cancer (NSCLC) who progressed on erlotinib or gefitinib; 3 harbored concurrent activating exon 19 deletions and 1 harbored a concurrent L861G activating mutation²⁸⁷. However, in other studies, 2 additional patients, 1 with concurrent S768I activating mutation, experienced SD to these inhibitors²⁸⁸⁻²⁸⁹.

FREQUENCY & PROGNOSIS

EGFR mutation and amplification have been observed in 6.6% and 0.7-4.1% of prostate

carcinoma cases, respectively (cBioPortal, COSMIC, Dec 2022)^{176-177,246}. Many studies have reported EGFR overexpression in prostate cancer, cited in 25-42% of cases²⁹⁰⁻²⁹⁴. One study reports no significant correlation between EGFR expression and PSA levels²⁹³. Higher levels of EGFR have been correlated with poor prognosis and higher Gleason score in prostate cancer patients^{291,293}.

FINDING SUMMARY

EGFR encodes the epidermal growth factor receptor, which belongs to a class of proteins called receptor tyrosine kinases. In response to signals from the environment, EGFR passes biochemical messages to the cell that stimulate it to grow and divide²⁹⁵. The G724S mutation seen here has been characterized as activating²⁹⁶ and confers resistance to first-generation EGFR inhibitors in the context of co-occurring activating EGFR alterations^{287,297-300}. In the context of co-occurring EGFR exon 19 deletions, G724S has demonstrated resistance to osimertinib^{279-283,286}. Limited preclinical evidence for resistance to erlotinib²⁸⁵, gefitinib²⁸¹, and osimertinib²⁸¹ has also been reported for G724S alone.

GENE

APC

ALTERATION

R876*

HGVS VARIANT

NM_000038.4:c.2626C>T (p.R876*)

VARIANT CHROMOSOMAL POSITION

chr5:112173917

VARIANT ALLELE FREQUENCY (% VAF)

39.6%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved drugs targeting APC inactivation in cancer. Loss of APC function leads to accumulation of beta-catenin and upregulation of WNT pathway transcription programs³⁰¹, and potential therapeutic approaches to target this pathway include CBP/beta-catenin antagonists, which interfere with the ability of beta-catenin to interact with transcriptional co-activator

CBP³⁰²⁻³⁰³. In a Phase 1 trial of the CBP/beta-catenin antagonist E7386, 1 patient with APC-mutated small bowel adenocarcinoma achieved a PR with tumor shrinkage of ~69% and response duration of 165 days³⁰⁴; preclinical data support sensitivity of APC-deficient gastric or colorectal cancer models to E7386³⁰⁵⁻³⁰⁶.

FREQUENCY & PROGNOSIS

APC mutations and homozygous deletions were observed in 0.6%-6.7% and 0.6%-4.5% of prostate adenocarcinoma cases, respectively^{56,59,132,307}. APC methylation has been frequently observed in prostate carcinoma³⁰⁸⁻³¹⁰, and dysregulation of the WNT signaling pathway is believed to play a role in a subset of prostate cancers³¹¹⁻³¹². In prostate cancer, APC promoter hypermethylation is associated with shorter time to relapse in both univariate (P = 0.002) and multivariate analyses³¹³. Solid tumors with WNT/beta-catenin pathway alterations, as seen here, were observed to have significantly less T-cell inflammation in one study³¹⁴.

FINDING SUMMARY

APC (adenomatous polyposis coli) encodes a tumor suppressor with critical roles in regulating cell division and adhesion. APC interacts with beta-catenin and controls signaling in the WNT pathway, which regulates embryonic development and cell differentiation³¹⁵. Alterations such as seen here may disrupt APC function or expression³¹⁶⁻³²⁰.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the APC variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with familial adenomatous polyposis (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in APC are found in more than 90% of patients with familial adenomatous polyposis (FAP)³²¹⁻³²³. The prevalence for FAP in the general population is estimated to be 1:8300 from birth³²⁴, and in the appropriate clinical context germline testing of APC is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

PIK3CA

ALTERATION

E970K

HGVS VARIANT

NM_006218.2:c.2908G>A (p.E970K)

VARIANT CHROMOSOMAL POSITION

chr3:178948136

VARIANT ALLELE FREQUENCY (% VAF)

41.5%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical and preclinical data in various tumor types indicate that PIK3CA activating alterations may predict sensitivity to therapies targeting PI3K³²⁵⁻³³², AKT³³³⁻³³⁴, or mTOR³³⁵⁻³⁴². The Phase 2 NCI-MATCH study of copanlisib for patients with refractory solid tumors harboring PIK3CA mutations with or without PTEN loss met its primary endpoint with an ORR of 16% (4/25 PRs); responses (PR or SD >6 months) were seen in patients with ameloblastoma, liposarcoma, and

carcinomas of the endometrium, ovary, esophagus, lung, and prostate³³². However, the Phase 2 study of copanlisib for patients with endometrial carcinoma harboring PIK3CA hotspot mutations failed to report any objective responses (n=11)³³¹. Two other studies of copanlisib for patients with genomically unselected tumors reported 1 CR and 2 PRs (1 unconfirmed) among 16 total patients with PIK3CA-mutated solid tumors with or without PTEN alterations³²⁹⁻³³⁰. In the Phase 2 MATCH trial for patients with PIK3CA-mutated solid tumors, 28% (18/65) of patients experienced PFS lasting at least 6 months after treatment with tasislisib; however, no ORs were observed in this study³⁴³. A separate Phase 1b study of tasislisib in combination with the CDK4/6 inhibitor palbociclib for patients with PIK3CA-mutated solid tumors reported an ORR of 0% (n=12) and a DCR of 17% (2/12)³⁴⁴. In a Phase 1 trial of the dual PI3K/mTOR kinase inhibitor apitolisib, 79% (11/14) of patients with PIK3CA-mutated advanced solid tumors experienced disease control (3 PRs, 8 SDs)³⁴⁵. The PI3K inhibitor alpelisib is approved as a single agent for the treatment of patients with PIK3CA-related overgrowth spectrum (PROS)³⁴⁶, but has shown limited activity as monotherapy for PIK3CA-mutated solid tumors with a Phase 1a

study reporting an ORR of 6.0% (8/134) and a DCR of 58% (78/134)³²⁶.

FREQUENCY & PROGNOSIS

PIK3CA mutations have been reported in up to 4.3% of prostate adenocarcinomas^{54,56,75,347}. PI3K pathway dysregulation in prostate cancer has been reported in numerous studies, noting either PIK3CA mutation, amplification, or overexpression, or PTEN deletion³⁴⁸⁻³⁵⁰. Retrospective analysis of the Prostate Adenocarcinoma TCGA dataset showed a significant association between PIK3CA alteration, including amplification, with decreased survival (HR=0.55), increased regional lymph node metastasis, higher primary tumor category, and higher Gleason grade³⁵¹.

FINDING SUMMARY

PIK3CA encodes p110-alpha, which is the catalytic subunit of phosphatidylinositol 3-kinase (PI3K). The PI3K pathway is involved in cell signaling that regulates a number of critical cellular functions, including cell growth, proliferation, differentiation, motility, and survival³⁵²⁻³⁵³. PIK3CA alterations that have been characterized as activating, such as observed here, are predicted to be oncogenic³⁵⁴⁻³⁷⁵.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

PTEN

ALTERATION

P246L, K267fs*9, R173C

HGVS VARIANT

NM_000314.4:c.737C>T (p.P246L),
NM_000314.4:c.800del (p.K267Rfs*9),
NM_000314.4:c.517C>T (p.R173C)

VARIANT CHROMOSOMAL POSITION

chr10:89717712, chr10:89717769-89717770, chr10:89711899

VARIANT ALLELE FREQUENCY (% VAF)

43.4%, 36.6%, 39.3%

including everolimus and temsirolimus have been terminated due to limited efficacy, reporting brief PFS (1.9-3.6 months)³⁸²⁻³⁸⁶. Preclinical data indicate that PTEN loss or inactivation may predict sensitivity to PARP inhibitors³⁸⁷⁻³⁹¹, and clinical benefit has been observed for patients with PTEN-altered breast cancer including triple negative breast cancer³⁹², ovarian cancer²⁴⁵, uterine leiomyosarcoma³⁹³, and endometrial cancer³⁹¹ treated with PARP inhibitors. However, some studies have reported a lack of association between PTEN mutation and PARP inhibitor sensitivity^{84,394}.

advanced stage, more rapid progression to metastasis, and reduced patient survival³⁹⁶⁻⁴⁰⁰. Concurrent alterations in at least 2 of the TP53, RB1, and PTEN genes, as seen in this sample, molecularly define a subtype of prostate cancer with aggressive clinical course and reduced sensitivity to androgen-deprivation therapies (NCCN Prostate Cancer Guidelines, v1.2023)^{395,401-404}.

FINDING SUMMARY

PTEN encodes an inositol phosphatase that functions as a tumor suppressor by negatively regulating the PI3K-AKT-mTOR pathway; loss of PTEN can lead to uncontrolled cell growth and suppression of apoptosis³⁷⁷. Alterations such as seen here may disrupt PTEN function or expression⁴⁰⁵⁻⁴⁴⁶.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the PTEN variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with hamartoma tumor syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. PTEN mutations underlie several inherited disorders, collectively termed PTEN hamartoma tumor syndrome (PHTS), which include Cowden syndrome (CS) and its variant Lhermitte-Duclos disease (LD), Bannayan-Riley-Ruvalcaba syndrome (BRRS), PTEN-related Proteus syndrome (PS), and Proteus-like syndrome⁴⁴⁷⁻⁴⁴⁸. The mutation rate for PTEN in these disorders ranges from 20 to 85% of patients^{447,449}. The estimated incidence of Cowden syndrome is 1/200,000, which may be an underestimate due to the high variability of this disorder⁴⁴⁷. Given the association between PTEN and these inherited syndromes, in the appropriate clinical context, germline testing for mutations affecting PTEN is recommended.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

PTEN loss or mutation leads to activation of the PI3K-AKT-mTOR pathway and may predict sensitivity to inhibitors of this pathway^{330,376-378}. The Phase 3 IPATential150 trial combining either ipatasertib or placebo with abiraterone and prednisolone for patients with no prior treatment for metastatic castration-resistant prostate cancer (CRPC) reported superior radiographic PFS (18.5 vs. 16.5 months, HR=0.77) and ORR (61% vs. 39%) with ipatasertib for patients with PTEN loss as determined by immunohistochemistry³⁷⁹. A Phase 2 study combining either ipatasertib or placebo with abiraterone and prednisolone for metastatic CRPC reported improved radiographic PFS from the addition of ipatasertib for patients with PTEN loss (11.5 vs. 4.6 months, HR=0.39), but limited benefit for those without PTEN loss (7.5 vs. 5.6 months, HR=0.84)³⁸⁰. A Phase 2 trial combining either the AKT inhibitor capivasertib or placebo with docetaxel and prednisolone for patients with metastatic CRPC reported prolonged OS (31.2 vs. 20.3 months, HR=0.54) with the addition of capivasertib but comparable composite PFS (7.0 vs. 6.7 months, HR= 0.92) between trial arms; OS and PFS were consistent irrespective of PTEN, PI3K, or AKT status³⁸¹. Multiple studies of mTOR inhibitors

— Nontargeted Approaches —

In post-hoc analyses of a Phase 1/2 trial, the addition of carboplatin to cabazitaxel significantly improved PFS (6.0 vs. 2.2 months) and OS (17.4 vs. 9.9 months) for patients with an aggressive variant prostate cancer molecular signature (AVPC-MS, composed of alterations in at least 2 of the genes RB1, TP53, and PTEN); no significant difference in outcomes were observed for patients lacking this signature (NCCN Prostate Cancer Guidelines, v1.2023)³⁹⁵. Post-hoc analysis of a Phase 2 trial in metastatic castration-resistant prostate cancer suggests that patients with aggressive variant prostate cancer (AVPC), molecularly characterized by harboring alterations in at least 2 genes of TP53, RB1, and PTEN, as seen in this sample, may benefit from cabazitaxel combined with carboplatin compared with cabazitaxel alone (median PFS of 5.1 vs. 2.2 months, p=0.03; estimated median OS of 11.2 vs. 10.5 months, p=0.11), whereas patients who were AVPC-negative did not benefit from the more intense chemotherapy combination in this study³⁹⁵.

FREQUENCY & PROGNOSIS

PTEN mutations have been reported in up to 10% of prostate carcinoma cases^{56,59,133,347}, while PTEN loss been reported in 15-36% of cases^{56,59,133}. Reduction or loss of PTEN protein expression has been found in 18-61% of prostate cancer cases and has been found to be associated with more

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

RNF43

ALTERATION

G659fs*41, H683fs*17

HGVS VARIANT

NM_017763.4:c.1976del (p.G659Vfs*41),

NM_017763.4:c.2043del (p.H683fs*17)

VARIANT CHROMOSOMAL POSITION

chr17:56435160-56435161, chr17:56435093-56435094

VARIANT ALLELE FREQUENCY (% VAF)

37.9%, 41.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

RNF43 alterations may lead to WNT activation and therefore may confer sensitivity to WNT pathway

inhibitors such as porcupine (PORCN) inhibitors⁴⁵⁰⁻⁴⁵⁴, but this has not been clinically established⁴⁵⁵. In a Phase 1 basket study of the PORCN inhibitor RXC004, SD was reported for 1/2 patients with RNF43-mutated tumors⁴⁵⁶. RNF43 mutations did not clearly correlate with change in tumor baseline in a Phase 2 study of the addition of the PORCN inhibitor WNT974 to the combination of encorafenib and cetuximab to treat patients with BRAF V600E-mutated metastatic colorectal cancer (CRC)⁴⁵⁵. A meta-analysis of patients with BRAF V600E-mutated CRC suggests that RNF43 mutations may predict improved clinical benefit from BRAF/EGFR combinatorial therapies in the context of microsatellite-stable disease⁴⁵⁷.

FREQUENCY & PROGNOSIS

RNF43 alterations have been reported

predominantly in endometrial (18-27%)⁴⁵⁸⁻⁴⁵⁹ and gastrointestinal cancers such as colorectal cancer (CRC; 19%)^{68,459-460} and gastric cancer (11-18%)⁴⁶¹⁻⁴⁶², where they are associated with MMR deficiency and microsatellite instability (MSI)^{457,459-463}. One study found that patients with Stage 4 RNF43-mutated right-sided CRC had significantly worse OS than patients with RNF43-wildtype right-sided or RNF43-mutated left-sided CRC⁴⁶⁴. Co-occurrence of RNF43 and KRAS activating mutations are associated with numerically poorer outcomes in CRC⁴⁶⁵.

FINDING SUMMARY

RNF43 is a tumor suppressor and encodes a ubiquitin ligase that functions as a negative regulator of WNT signaling^{450-454,465-466}.

GENE

TSC2

ALTERATION

R57H

HGVS VARIANT

NM_000548.3:c.170G>A (p.R57H)

VARIANT CHROMOSOMAL POSITION

chr16:2100432

VARIANT ALLELE FREQUENCY (% VAF)

43.1%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Loss or inactivation of TSC2 can activate mTOR signaling⁴⁶⁷. mTOR inhibitors such as everolimus, temsirolimus, and sirolimus have shown activity against tumors associated with the genetic disease tuberous sclerosis complex (TSC), including subependymal giant cell astrocytomas and renal angiomyolipomas⁴⁶⁸⁻⁴⁷³. Sirolimus and nab-sirolimus have shown activity for patients with TSC2-altered malignant perivascular epithelioid cell tumors⁴⁷⁴⁻⁴⁷⁶. Nab-sirolimus has also shown

limited activity for patients with TSC2-mutated sarcomas as a monotherapy⁴⁷⁷ or in combination with nivolumab⁴⁷⁸. In the context of TSC2-altered malignancies unrelated to TSC, everolimus and temsirolimus activity has been limited⁴⁷⁹⁻⁴⁸¹ with the exception of anecdotal reports across various solid tumors, including anaplastic thyroid cancer⁴⁸², renal cell carcinoma (RCC)⁴⁸³⁻⁴⁸⁴, glioblastoma (GBM)⁴⁸⁵, and central nervous system embryonal tumor⁴⁸⁶, as well as a case of Hodgkin lymphoma⁴⁸⁷. In the prospective NCI-MATCH study, only 6.7% (1/15) of patients with TSC2-mutated solid tumors responded to everolimus, with the single response reported for 1 patient with uterine leiomyosarcoma⁴⁷⁹. Retrospective analyses of the RECORD-3, GRANITE-1, and EVOLVE-1 studies of everolimus to treat patients with RCC, gastric cancer, or hepatocellular carcinoma, respectively, showed no significant association between alterations in MTOR, TSC1, or TSC2 and median PFS⁴⁸⁸.

FREQUENCY & PROGNOSIS

TSC2 mutation and loss have been reported in up to 1.8% and 4.2%, respectively, of prostate carcinoma samples analyzed in large

datasets^{132,214,489-490}. Published data investigating the prognostic implications of TSC2 alterations in prostate carcinoma are limited (PubMed, Feb 2023).

FINDING SUMMARY

The tumor suppressor protein Tuberin (TSC2) binds with Hamartin (TSC1) to inhibit mTOR signaling and cell growth^{467,491}. Alterations such as seen here may disrupt TSC2 function or expression⁴⁹²⁻⁴⁹⁴.

POTENTIAL GERMLINE IMPLICATIONS

Inactivating germline mutations in TSC2 are associated with the autosomal dominant disorder tuberous sclerosis complex, which results in the development of hamartomas in multiple organ systems and an increased risk of developing renal cell carcinoma (RCC)⁴⁹⁵⁻⁴⁹⁷. TSC2 mutations account for approximately 75 to 80% of reported sporadic cases⁴⁹⁸. Prevalence for this disorder in the general population is estimated to be 1/6,000 from birth and 1/12,000 to 1/14,000 in children under 10 years of age⁴⁹⁸. In the appropriate clinical context, germline testing of TSC2 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

ASXL1

ALTERATION

G646fs*12

HGVS VARIANT

NM_015338.5:c.1934dup (p.G646Wfs*12)

VARIANT CHROMOSOMAL POSITION

chr20:31022441

VARIANT ALLELE FREQUENCY (% VAF)

35.1%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in ASXL1.

FREQUENCY & PROGNOSIS

ASXL1 alterations occur infrequently across various solid tumor types²²⁹ and are not known to act as drivers in any specific solid cancer type⁴⁹⁹. Published data investigating the prognostic implications of ASXL1 alterations in solid tumors are limited (PubMed, May 2023). In the context of clonal hematopoiesis, ASXL1 mutations are significantly enriched in current or former smokers⁵⁰⁰.

FINDING SUMMARY

ASXL1 regulates epigenetic marks and transcription through interaction with polycomb complex proteins and various transcription activators and repressors⁵⁰¹⁻⁵⁰³. Alterations such as seen here may disrupt ASXL1 function or expression⁵⁰⁴⁻⁵⁰⁶.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁵⁰⁷⁻⁵¹². CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy⁵⁰⁷⁻⁵⁰⁸. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease⁵¹³. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH^{511,514-515}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

GENE

ATR

ALTERATION

F222fs*11

HGVS VARIANT

NM_001184.3:c.666del (p.F222Lfs*11)

VARIANT CHROMOSOMAL POSITION

chr3:142281577-142281578

VARIANT ALLELE FREQUENCY (% VAF)

36.2%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

A Phase 2 study reported talazoparib led to a SD

lasting over 6 months for a patient with ATR-mutated breast cancer⁵¹⁶. Based on preclinical evidence, ATR-deficient tumors may be sensitive to PARP inhibitors^{224,517}.

FREQUENCY & PROGNOSIS

ATR mutations or loss have been reported in <1% of prostate adenocarcinomas in multiple studies^{54-57,59,133}. ATR inactivation, either by mutation or decreased expression, is associated with increased microsatellite instability (MSI) and chromosome instability (CIN) in a variety of tumor types, including colorectal and endometrial cancers⁵¹⁸⁻⁵²⁰. Published data investigating the prognostic implications of ATR alterations in prostate carcinoma are limited (PubMed, Oct 2022). Other alterations leading to defective DNA repair have been reported in 23-33% of metastatic

castration-resistant prostate cancer cases, and mutations in DNA repair-associated genes have been associated with response to the PARP inhibitor olaparib for patients with prostate cancer^{59,80}; however, alterations in ATR have not been associated with outcomes in these studies.

FINDING SUMMARY

ATR encodes the protein ataxia telangiectasia and RAD3 related, which phosphorylates the tumor suppressor BRCA1, and several cell cycle checkpoint proteins including CHK1; it plays a key role in maintaining genome integrity via regulation of DNA repair and replication⁵²¹⁻⁵²². Alterations such as seen here may disrupt ATR function or expression⁵²³.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

CASP8

ALTERATION

R250Q, Q308*

HGVS VARIANT

NM_001228.4:c.749G>A (p.R250Q),
NM_001228.4:c.922C>T (p.Q308*)

VARIANT CHROMOSOMAL POSITION

chr2:202141587, chr2:202149607

VARIANT ALLELE FREQUENCY (% VAF)

42.2%, 43.8%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted approaches to directly address alterations in CASP8.

FREQUENCY & PROGNOSIS

CASP8 mutations have been observed across solid tumors²²⁹, including 8.2% of head and neck squamous cell⁵²⁴, 3.2% of colorectal⁵²⁵, and 4% of cervical⁵²⁶ carcinoma cases. Reduced expression of CASP8 has been associated with poor prognosis in ovarian cancer⁵²⁷, prostate cancer⁵²⁸, and medulloblastoma⁵²⁹.

FINDING SUMMARY

CASP8 encodes caspase-8, a multifunctional protein that mediates apoptosis⁵³⁰⁻⁵³³, cell motility⁵³⁴⁻⁵³⁵, and cell signaling, including through the NFkB⁵³⁶⁻⁵³⁸ and MAPK⁵³⁹⁻⁵⁴⁰ pathways. The role of CASP8 in cancer is complex and context-dependent, with diverse cancer types exhibiting either overexpression or loss of expression. CASP8 mutations found in the context of cancer tend to be truncating or missense mutations; most of the characterized mutations impair apoptosis⁵⁴¹⁻⁵⁴⁵ and promote NFkB activation⁵⁴⁶.

GENE

CBL

ALTERATION

splice site 1007+2T>C

HGVS VARIANT

NM_005188.2:c.1007+2T>C (p.?)

VARIANT CHROMOSOMAL POSITION

chr11:119146846

VARIANT ALLELE FREQUENCY (% VAF)

38.9%

advanced solid tumors⁵⁵³. Among 8 patients with CBL inactivating alterations in a Phase 1b trial, sitravatinib produced 2 PRs (25% ORR), with 1 NSCLC and 1 melanoma responding for over 4 months, and 4 SD outcomes, with 3 prolonged SDs seen in a patient with NSCLC, a patient with esophageal cancer, and a patient with a pancreatic neuroendocrine tumor⁵⁵³. CBL has been shown to downregulate EGFR⁵⁵⁴⁻⁵⁵⁸ and FLT3⁵⁵⁹⁻⁵⁶¹. Preclinical models of myeloid malignancies have demonstrated that CBL inactivation confers sensitivity to the FLT3-targeting therapies sunitinib⁵⁵⁹, midostaurin⁵⁶¹, and quizartinib⁵⁶², as well as to dasatinib⁵⁶³, although clinical evidence for this approach in solid tumors is lacking.

FREQUENCY & PROGNOSIS

CBL mutations have been reported in 2.4% of prostate adenocarcinoma samples analyzed in the COSMIC database (COSMIC, Jul 2022)²⁴⁶. Preclinical studies have shown that targeted reduction of CBL expression reduced proliferation

and viability of prostate cancer cell lines and tumor growth of prostate cancer xenotransplants⁵⁶⁴⁻⁵⁶⁵. One study identified strong cytoplasmic CBL expression by immunohistochemistry in prostate cancer samples as compared with normal prostate tissue; increased levels of CBL protein were a significant independent predictor of reduced patient survival⁵⁶⁵.

FINDING SUMMARY

CBL encodes an E3 ubiquitin protein ligase that is involved in cell signaling and ubiquitination, targeting proteins such as EGFR, FGFR1, FGFR2, PDGFR-alpha, PDGFR-beta, FLT3, and SRC for degradation by the proteasome⁵⁶⁶⁻⁵⁷⁰. CBL alterations that result in loss or disruption of the tyrosine kinase binding domain, RING finger domain, and/or tail domain, as observed here, are predicted to be inactivating and to promote tumorigenesis⁵⁷¹⁻⁵⁸⁸.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

CBL inactivation may lead to the hyperactivation of various receptor tyrosine kinases (RTKs), including MET⁵⁴⁷, PDGFRA⁵⁴⁸, KIT⁵⁴⁹, VEGFR2⁵⁵⁰, and the TAM (TYRO3, AXL, MER) RTKs⁵⁵¹. These RTKs are targets of the multikinase inhibitor sitravatinib⁵⁵², which has shown activity in CBL-mutated

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

CD79A

ALTERATION

V69I - subclonal

HGVS VARIANT

NM_001783.3:c.205G>A (p.V69I)

VARIANT CHROMOSOMAL POSITION

chr19:42383185

VARIANT ALLELE FREQUENCY (% VAF)

2.7%

affecting the CD79A ITAM domain experienced a CR⁵⁸⁹. Limited preclinical data in ABC-DLBCL cell lines suggest that CD79A mutation may also be associated with sensitivity to the SRC/BCR-ABL kinase inhibitor dasatinib⁵⁹⁰ or to PKC inhibition⁵⁹². However, these approaches have not been evaluated in the context of CD79A mutations in solid tumors. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

CD79a or Ig-alpha, a critical component of the B-cell receptor (BCR) that forms a heterodimer with CD79b, which functions in BCR signal transduction. BCR signaling ultimately leads to the activation of several downstream signaling pathways, including NF-kappaB, PI3K, mitogen-activated protein kinase (MAPK), and RAS pathways, which promote cell proliferation and survival⁵⁹⁴. BCR signaling is considered an important oncogenic pathway in several types of lymphomas, and somatic mutations in CD79A and especially CD79B have been reported to occur frequently in the activated B-cell-like (ABC) subtype of diffuse large B-cell lymphoma (DLBCL) but only rarely or not at all in the other DLBCLs and types of lymphomas studied^{590,594}. Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of limited clinical⁵⁸⁹ and preclinical⁵⁹⁰⁻⁵⁹¹ studies in activated B-cell (ABC)-diffuse large B-cell lymphoma (DLBCL), CD79A activating alterations may indicate sensitivity to the BTK inhibitor ibrutinib. In a Phase 1/2 study of ibrutinib for patients with relapsed or refractory DLBCL, 1 patient with a subclonal 45bp deletion

FREQUENCY & PROGNOSIS

CD79A mutations are rare (0-1.7%) in solid tumors and more commonly found in diffuse large B-cell lymphomas (DLBCL) (COSMIC, Dec 2022)^{246,593}. Published data investigating the prognostic implications of CD79A alteration in solid tumors are limited (PubMed, Dec 2022).

FINDING SUMMARY

CD79A encodes B-cell antigen receptor complex-associated protein alpha chain, also known as

GENE

CDK8

ALTERATION

Q331H - subclonal

HGVS VARIANT

NM_001260.1:c.993G>T (p.Q331H)

VARIANT CHROMOSOMAL POSITION

chr13:26974649

VARIANT ALLELE FREQUENCY (% VAF)

1.5%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies that directly address genomic alterations in CDK8. Investigational inhibitors that selectively target CDK8 and the related CDK19 have shown activity against WNT-dependent tumors in preclinical assays⁵⁹⁵⁻⁵⁹⁹.

FREQUENCY & PROGNOSIS

CDK8 amplification has been reported at low frequencies in solid tumors, including 3% of colorectal (CRC) and <1% of bladder, breast,

sarcoma, and non-small cell lung cancer (NSCLC) samples²²⁹. In CRC, CDK8 expression is associated with reduced survival⁶⁰⁰⁻⁶⁰¹.

FINDING SUMMARY

CDK8 encodes a member of the cyclin-dependent kinase (CDK) family that regulates gene expression and cellular signaling including the WNT/beta-catenin and NOTCH pathways^{595,602-603}. CDK8 acts as an oncogene in colon cancer and melanoma⁶⁰⁴⁻⁶⁰⁶.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS
GENE
CTCF
ALTERATION

T204fs*26

HGVS VARIANT

NM_006565.3:c.610dup (p.T204Nfs*26)

VARIANT CHROMOSOMAL POSITION

chr16:67645338

VARIANT ALLELE FREQUENCY (% VAF)

40.3%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in CTCF.

FREQUENCY & PROGNOSIS

Somatic mutations in CTCF are infrequently reported in most cancers, but have been observed more commonly (24%) in uterine corpus endometrial carcinoma (cBioPortal, 2023)¹⁷⁶⁻¹⁷⁷; nearly half of the observed mutations were truncating, suggesting a tumor suppressor role for CTCF in this disease. In addition, CTCF has been found to act as a tumor suppressor in breast cancer cell line studies⁶⁰⁷⁻⁶⁰⁸.

FINDING SUMMARY

CTCF encodes an 11-zinc-finger protein that is implicated in various regulatory roles, including gene activation and repression, imprinting, insulation, methylation, and X chromosome inactivation⁶⁰⁹. CTCF plays a role in transcriptional regulation of a number of key cancer-associated genes, including the oncogene MYC⁶¹⁰ and tumor suppressor TP53⁶¹¹, via maintenance of local DNA methylation status. Decreased expression levels of CTCF and/or BORIS, another 11-zinc-finger transcriptional regulator, were reported to be closely associated with global DNA methylation variability and decreased OS in epithelial ovarian cancer⁶¹²⁻⁶¹³.

GENE
CTNNA1
ALTERATION

K695fs*27

HGVS VARIANT

NM_001903.2:c.2084del (p.K695Rfs*27)

VARIANT CHROMOSOMAL POSITION

chr5:138266231-138266232

VARIANT ALLELE FREQUENCY (% VAF)

40.7%

highest incidence in endometrial carcinoma (7.0%)⁶¹⁴, cutaneous melanoma (6.4%)⁶¹⁵, colorectal adenocarcinoma (4.9%)⁶¹⁴, and stomach adenocarcinoma (3.90%) TCGA datasets (cBioPortal, 2023)¹⁷⁶⁻¹⁷⁷. CTNNA1 mutations have been observed in patients with hereditary diffuse gastric carcinoma without CDH1 mutations⁶¹⁶⁻⁶¹⁷. Reduced CTNNA1 expression in patients with breast cancer has been correlated with a poor clinical outcome and breast cancer brain metastasis⁶¹⁸⁻⁶¹⁹. Deletion and hypermethylation of CTNNA1 has been observed in up to 22% (18/83) of myelodysplastic syndrome (MDS) cases and associated with poor clinicopathological features⁶²⁰⁻⁶²² and a trend for inferior survival⁶²⁰. Loss of CTNNA1 expression via 5q deletion or hypermethylation has been reported as a frequent event in acute myeloid leukemia and associated with shorter relapse-free survival in one study⁶²²⁻⁶²⁴.

FINDING SUMMARY

CTNNA1 encodes alpha-catenin, a member of the cadherin family that functions in cell adhesion. Alpha-catenin acts as a tumor suppressor, through mechanisms that can vary by tissue⁶²⁵⁻⁶²⁶. Alpha-catenin is one of three catenin proteins that are in complex with E-cadherin to help mediate cell-cell adhesion in epithelial tumor suppression⁶²⁵⁻⁶²⁶; loss of cell adhesion may contribute to cancer cell invasiveness and formation of metastases. In epidermal cells, alpha-catenin acts as a tumor suppressor by inducing YAP1 phosphorylation and cytoplasmic localization^{619,627}. Alpha-catenin also acts as a tumor suppressor by interacting with IKBalpha to influence the NF-KB pathway in E-cadherin-negative basal-like breast cancer cells⁶¹⁹. Loss of alpha-catenin expression is also hypothesized to alter the balance between the cytoplasmic (cell adhesion) and nuclear (cell proliferation) functions of beta-catenin, further contributing to oncogenesis⁶²⁸.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no available targeted therapies to address genomic alterations in CTNNA1.

FREQUENCY & PROGNOSIS

CTNNA1 mutations have been observed with

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

EPHB4

ALTERATION

V420I

HGVS VARIANT

NM_004444.4:c.1258G>A (p.V420I)

VARIANT CHROMOSOMAL POSITION

chr7:100417218

VARIANT ALLELE FREQUENCY (% VAF)

48.6%

demonstrated that sEPHB4-HSA inhibits cell proliferation and xenograft tumor growth, including for cells expressing cancer-associated EPHB4 mutants or overexpressing wild-type EPHB4^{629,631-635}. In addition, small-molecule inhibitors targeting multiple tyrosine kinases including EPHB4, such as JI-101 and XL647, have been under preclinical and clinical investigation⁶³⁶⁻⁶³⁸.

FREQUENCY & PROGNOSIS

Increased EPHB4 mRNA and/or protein expression has been reported in a variety of cancer types, including head and neck squamous cell carcinoma (HNSCC)⁶³⁹⁻⁶⁴², gastric and esophageal⁶⁴³⁻⁶⁴⁷, colorectal carcinoma (CRC)⁶⁴⁸⁻⁶⁵⁴, breast⁶⁵⁵⁻⁶⁵⁹, ovarian⁶⁵⁹⁻⁶⁶¹, endometrial⁶⁶²⁻⁶⁶⁴, thyroid⁶⁶⁵⁻⁶⁶⁷, lung⁶⁶⁸⁻⁶⁶⁹, glioma⁶⁷⁰⁻⁶⁷¹, and other solid tumors^{631,672-679}. In several of these studies, increased EPHB4 expression has been associated with clinicopathologic features, including disease stage^{631,639,655,660-661,669,672,674}, histological grade^{645,655,662,671}, and hormone receptor status^{658,663}. High EPHB4 expression has been

associated with inferior survival in multivariate analyses for patients with CRC treated with bevacizumab [hazard ratio (HR) = 5.95]⁶⁵³, HNSCC (HR = 2.95)⁶⁴², epithelial ovarian cancer (HR = 4.53)⁶⁵⁹, or glioma (HR = 3.21)⁶⁷¹.

FINDING SUMMARY

EPHB4 encodes a member of the EPH family of receptor tyrosine kinases⁶⁸⁰. Ephrin signaling has been implicated in multiple processes, including cell adhesion, cytoskeletal organization, and cell migration⁶⁸¹, and signaling between EPHB4 and its ligand EphrinB2 is particularly important for angiogenesis⁶⁸²⁻⁶⁸³. EPHB receptors, including EPHB4, have been shown to undergo dysregulation (amplification, mutation, under- or overexpression) in a number of different cancer types⁶⁸⁴. EPHB4 amplification has been reported in several solid tumor types^{639-640,645,685-686} and was associated with advanced disease stage in head and neck squamous cell carcinoma (HNSCC)⁶³⁹. Activating missense mutations in or near the tyrosine kinase domain, including G723S, A742V, and P881S, have also been identified in lung cancer⁶³⁵.

GENE

ERRFI1

ALTERATION

F411fs*41

HGVS VARIANT

NM_018948.3:c.1233del (p.F411Lfs*41)

VARIANT CHROMOSOMAL POSITION

chr1:8073425-8073426

VARIANT ALLELE FREQUENCY (% VAF)

41.0%

approved to treat lung or breast cancers. One patient with cholangiocarcinoma and an ERRFI1 loss of function mutation was reported to have a partial response to the EGFR inhibitor erlotinib⁶⁸⁷. In preclinical studies, the EGFR inhibitor gefitinib was reported to cause regression of tumors in ERRFI1 knockout mice⁶⁸⁸ and inhibit signaling downstream EGFR in ERRFI1-depleted human cell lines⁶⁸⁹.

FREQUENCY & PROGNOSIS

ERRFI1 mutation has been reported in 0-2% of metastatic prostate cancer cases^{57,59}. In one study, ERRFI1 was identified as one of 87 genes with significantly altered expression levels in prostate cancer cell lines upon androgen treatment⁶⁹⁰. Published data investigating the prognostic implications of ERRFI1 alterations in prostate carcinoma are limited (PubMed, Nov 2022).

FINDING SUMMARY

The ERRFI1 gene (also known as MIG6 or RALT) encodes a tumor suppressor that negatively regulates the ERBB family of receptors⁶⁹¹⁻⁶⁹⁴, and is transcriptionally activated by RAS-RAF-ERK signaling⁶⁹⁵. ERRFI1 directly binds to the kinase domain of ERBB proteins and consequently antagonizes their oncogenic signaling and cell proliferation⁶⁹¹⁻⁶⁹⁴. ERRFI1 also negatively regulates signaling by promoting EGFR endocytosis and degradation⁶⁹⁶⁻⁶⁹⁷. Knockout or depletion of ERRFI1 was shown to result in EGFR and ERBB2/ERBB3 activation^{688,695,697-698}, and disruption of the ERRFI1 gene promoted tumorigenesis in mice^{688,699}, whereas overexpression of ERRFI1 inhibited EGFR- or ERBB2-mediated signaling and oncogenic transformation in vitro^{697,700-701}. Alterations such as seen here may disrupt ERRFI1 function or expression^{691,693}.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

ERRFI1 loss or inactivating mutations may result in EGFR activation and predict sensitivity to ERBB tyrosine kinase inhibitors such as erlotinib, gefitinib, afatinib, and lapatinib, which are

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

FH

ALTERATION

splice site 1237-1G>A

HGVS VARIANT

NM_000143.3:c.1237-1G>A (p.?)

VARIANT CHROMOSOMAL POSITION

chr1:241663891

VARIANT ALLELE FREQUENCY (% VAF)

38.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

A preclinical study showed that FH-deficient renal cancer cells are dependent on ABL1 activity and sensitive to the multikinase inhibitor vandetanib; treatment with vandetanib inhibited the growth and tumorigenicity of these cells in vitro and in vivo⁷⁰². Tumors with FH loss or inactivation may

therefore be sensitive to vandetanib, which is approved to treat medullary thyroid cancer and is in clinical trials in solid tumors. A Phase 2 trial of bevacizumab and erlotinib reported overall response rate in 60% (12/20) of patients with hereditary leiomyomatosis and renal cell cancer, and 29% (6/21) of patients with sporadic papillary renal cell carcinoma⁷⁰³.

FREQUENCY & PROGNOSIS

FH mutations are rare in solid and hematologic malignancies²²⁹. FH-deficient renal cell carcinoma (RCC) arises in about 20% of families affected by hereditary leiomyomatosis and renal cell cancer (HLRCC) and is associated with aggressive disease and poor prognosis⁷⁰⁴⁻⁷⁰⁶.

FINDING SUMMARY

FH encodes fumarate hydratase, an enzymatic component of the Krebs cycle. FH has been identified as a possible hypoxia inducible factor activating gene⁷⁰⁷. Loss-of-function germline

mutations in FH are associated with hereditary leiomyomatosis and renal cell cancer (HLRCC); tumors arising in FH mutation carriers often demonstrate FH biallelic inactivation^{704,706,708-709}.

POTENTIAL GERMLINE IMPLICATIONS

FH germline inactivating alterations are associated with FH tumor predisposition syndrome, also known as hereditary leiomyomatosis and renal cell cancer (HLRCC), an autosomal-dominant syndrome characterized by cutaneous leiomyomata, uterine fibroids, and aggressive renal cell carcinoma (RCC)⁷⁰⁹. Pheochromocytoma and paraganglioma have also been described at lower frequency⁷¹⁰⁻⁷¹¹. Whereas cutaneous leiomyomata appear at a mean age of 30 years, increasing in size and number with age, the age at diagnosis of uterine fibroids ranges from 18 to 53 years⁷¹¹⁻⁷¹². HLRCC has been associated with a 21% lifetime risk of RCC⁷¹³. In the appropriate clinical context, germline testing of FH is recommended.

GENE

FUBP1

ALTERATION

I301fs*4

HGVS VARIANT

NM_003902.3:c.901dup (p.I301Nfs*4)

VARIANT CHROMOSOMAL POSITION

chr1:78429977

VARIANT ALLELE FREQUENCY (% VAF)

42.1%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Therapies targeting FUBP1 mutation directly or

downstream effectors have not been tested preclinically or clinically in tumors that harbor FUBP1 mutations.

FREQUENCY & PROGNOSIS

FUBP1 alteration has been reported in 1.5% of samples analyzed in COSMIC, with the highest incidences reported in tumors of the endometrium (3.6%), central nervous system (3.2%), large intestine (3.0%), skin (2.8%), stomach (2.7%), and liver (2.6%) (COSMIC, Feb 2023)²⁴⁶. One study reported higher expression of FUBP1 in colorectal carcinoma tissues compared to adenoma and normal colon epithelial tissues⁷¹⁴. A genetic signature defined by concomitant alterations in IDH1, CIC, and FUBP1 is associated with longer survival in patients with glioma⁷¹⁵. FUBP1 has

been shown to activate the expression of MYC⁷¹⁶⁻⁷¹⁹, activate p27KIP1⁷²⁰, and regulate the splicing of MDM2⁷²¹.

FINDING SUMMARY

FUBP1 encodes far upstream element binding protein 1 (also called FBP-1), a DNA-binding protein reported to have roles in transcriptional activation and splicing regulation of target genes. It is believed to act as an oncogene in some tumor types, such as hepatocellular carcinoma and non-small-cell lung cancer⁷²²⁻⁷²³, and as a tumor suppressor in others, particularly oligodendroglioma, for which mutations and/or loss of FUBP1 often co-occur with alterations in CIC or IDH1^{715,724-727}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS
GENE
IRF2
ALTERATION

N167fs*7

HGVS VARIANT

NM_002199.3:c.500del (p.N167Mfs*7)

VARIANT CHROMOSOMAL POSITION

chr4:185329340-185329341

VARIANT ALLELE FREQUENCY (% VAF)

38.5%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no therapies available to directly address genomic alterations in IRF2.

FREQUENCY & PROGNOSIS

Point mutations in the DNA binding domain of IRF2 have been identified in pancreatic cancer and have been shown to be inactivating⁷²⁸. Inactivating mutation or homozygous deletion of IRF2 has also been reported in 4/125 and 2/125 hepatocellular

carcinoma (HCC) tumors, respectively, all 6 of which were hepatitis B virus (HBV)-related⁷²⁹.

FINDING SUMMARY

IRF2 encodes the transcription factor interferon regulatory factor 2, which has been shown to regulate cellular responses to interferons, as well as cell growth and transformation.

GENE
KMT2D (MLL2)
ALTERATION

A221fs*40

HGVS VARIANT

NM_003482.4:c.661del (p.A221Lfs*40)

VARIANT CHROMOSOMAL POSITION

chr12:49447772-49447773

VARIANT ALLELE FREQUENCY (% VAF)

40.6%

FREQUENCY & PROGNOSIS

MLL2 alterations are observed in a number of solid tumor contexts²²⁹, and are especially prevalent in lung squamous cell carcinoma (SCC)⁷³⁰ and small cell lung carcinoma (SCLC)⁷³¹. MLL2 mutation was found to be an independent prognostic factor of poor PFS and OS in non-small cell lung cancer, but not in SCLC⁷³². One study reported that MLL2 truncating mutations were more common in recurrent ovary granulosa cell tumors (GCT) compared with primary GCTs (24% [10/42] vs. 3.0% [1/32])⁷³³. In a study of esophageal SCC, high MLL2 expression positively correlated with tumor stage, differentiation, and size, and negatively correlated with OS⁷³⁴.

FINDING SUMMARY

MLL2 encodes an H3K4-specific histone methyltransferase that is involved in the transcriptional response to progesterone

signaling⁷³⁵. Germline de novo mutations of MLL2 are responsible for the majority of cases of Kabuki syndrome, a complex and phenotypically distinctive developmental disorder⁷³⁶. A significant number of inactivating MLL2 alterations have been observed in multiple tumor types, suggesting a tumor suppressor role⁷³⁷.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁵⁰⁷⁻⁵¹². Comprehensive genomic profiling of solid tumors may detect nontumor alterations that are due to CH^{511,514-515}. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in MLL2.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

MAPK1

ALTERATION

R191H

HGVS VARIANT

NM_002745.4:c.572G>A (p.R191H)

VARIANT CHROMOSOMAL POSITION

chr22:22153338

VARIANT ALLELE FREQUENCY (% VAF)

37.7%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved drugs that directly target ERK2, although ERK1/2 inhibitors are in clinical

trials. In preclinical studies, ERK1/2 inhibitors have been shown to inhibit the activities of ERK2 mutations⁷³⁸⁻⁷⁴⁰ or restore sensitivity to the EGFR inhibitor WZ4002 in a WZ4002-resistant cell line harboring MAPK1 amplification⁷⁴¹. These approaches would not be relevant in the context of inactivating alterations, as seen here.

FREQUENCY & PROGNOSIS

MAPK1 mutations have been reported infrequently in solid and hematological malignancies, with highest incidences observed in liver (4.2%) and cervical (3.4%) cancers, and at lower frequencies in other tumor types (COSMIC, Dec 2022)²⁴⁶. MAPK activation (as assessed by ERK phosphorylation) has been frequently detected in glioblastoma tumors⁷⁴²⁻⁷⁴⁴ and higher levels of activation correlated with increasingly inferior survival (high

levels vs. medium levels vs. low levels)⁷⁴²⁻⁷⁴³. MAPK1-mediated activation of Wnt/beta-catenin signaling through osteopontin has been implicated in cholangiocarcinoma progression⁷⁴⁵. In addition, a MAPK1-associated SNP was correlated with increased risk of high-risk rectal cancer⁷⁴⁶. However, downregulation of MAPK1 was found to significantly associated with poorer disease free survival for HNSCC patients⁷⁴⁷.

FINDING SUMMARY

MAPK1, also known as ERK2, encodes a mitogen-activated protein kinase family member that is involved in the transduction of proliferation and differentiation signals. MAPK1 mutations located in the binding site for ERK and FXFP, such as seen here, have been characterized as inactivating⁷³⁹.

GENE

MLH1

ALTERATION

R127I

HGVS VARIANT

NM_000249.3:c.380G>T (p.R127I)

VARIANT CHROMOSOMAL POSITION

chr3:37045965

VARIANT ALLELE FREQUENCY (% VAF)

40.2%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

MLH1 inactivation leads to MMR defects, high MSI, and increased mutational burden^{26,68,748-749}, which may predict response to the anti-PD-1 immunotherapies pembrolizumab and nivolumab⁶⁻⁸. In a Phase 2 study of pembrolizumab in MSI-high colorectal cancer (CRC), three patients with MLH1 (germline) mutations experienced one partial response and two stable diseases⁷. Pembrolizumab demonstrated a significantly higher objective response rate in MSI-high CRC compared with microsatellite stable CRC (40% vs. 0%)⁷ and its efficacy correlated with high mutational burden in non-small cell lung cancer⁸. Nivolumab achieved a complete response in a patient with MSI-high CRC⁶. Furthermore, MSI status correlates with

higher PD-1 and PD-L1 expression⁷⁵⁰, potential biomarkers of response to anti-PD-1 immunotherapies. These therapies are in clinical trials for various tumor types and may be appropriate, particularly for hypermutant tumors. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

FREQUENCY & PROGNOSIS

MLH1 mutation has been reported in 0-1.6% of prostate carcinoma cases^{54-55,57,59,75}. Approximately 12% (7/60) of prostate cancers were found to be hypermutated, with two of the hypermutated cases having MLH1 mutations¹⁴. MLH1 expression was shown to be reduced in approximately 50% of 39 prostate tumors analyzed in one study⁷⁵¹. In another analysis, MLH1 nuclear expression was detected in 85% (6220/7275) prostate cancers; strong expression of MLH1 was observed in 55.2% of TMPRSS2-ERG-positive cases, but only in 32.9% of TMPRSS2-ERG-negative cases⁷⁵². MLH1 expression was significantly associated with early biochemical recurrence in univariate analyses in one study⁷⁵².

FINDING SUMMARY

MLH1 encodes the protein MutL homolog 1, colon cancer, nonpolyposis type 2, which binds PMS2 to form MutLalpha, a complex involved in DNA mismatch repair (MMR)⁷⁵³. Defective MMR

occurring as a result of mutation(s) in the MMR family (MLH1, MSH2, MSH6, or PMS2) can result in microsatellite instability (MSI), common in colon, endometrium, and stomach cancers²². Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the MLH1 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Lynch syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in MLH1 are associated with an autosomal dominant condition known as Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC), which is characterized by increased risk of a number of cancers²⁷. Approximately 50% of Lynch syndrome-associated mutations have been attributed to alterations in MLH1⁷⁵⁴. Lynch syndrome accounts for 1-7% of all colorectal cancers and has an estimated prevalence in the general population between 1:600 and 1:2000³⁰⁻³². In the appropriate clinical context, germline testing of MLH1 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

MSH2

ALTERATION

loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

MSH2 inactivation leads to MMR defects, MSI, and high mutational burden^{14,18,755-757}, which may predict response to the anti-PD-1 immunotherapies pembrolizumab and nivolumab⁶⁻⁸. In a Phase 2 study of MSI-high cancers, six patients with MSH2 (germline) mutations reported one partial response and two stable diseases⁷. Pembrolizumab therapy resulted in a significantly higher objective response rate in MSI-high CRC compared with microsatellite stable CRC (40% vs. 0%)⁷ and its efficacy correlated with high mutational burden in non-small cell lung cancer⁸. Treatment with nivolumab resulted in a complete response in a patient with MSI-high CRC⁶. Furthermore, MSI status correlates with higher PD-1 and PD-L1 expression⁷⁵⁰, potential biomarkers of response to PD-1 targeted immunotherapies. These therapies are in clinical trials for various tumor types and may be appropriate, particularly in hypermutant tumors.

— Nontargeted Approaches —

Preclinical studies have shown that tumor cells deficient in MSH2 are markedly sensitive to methotrexate in vitro⁷⁵⁸. Low levels of MSH2 have been observed by immunohistochemistry (IHC) in NSCLC and may predict benefit to cisplatin-based adjuvant chemotherapy⁷⁵⁹⁻⁷⁶⁰.

FREQUENCY & PROGNOSIS

MSH2 mutations are rare in prostate carcinoma and have not been detected in this context in multiple datasets^{54-55,57,75}. Deep deletion of MSH2 has been observed in ~1% of prostate carcinoma samples in large datasets^{56,489}. Loss of MSH2 protein expression was detected in 1.2% (14/1176) of primary prostate cancer samples in 1 study; of these 14 samples that could be further analyzed, 12/12 samples had loss of function alterations in MSH2 which included biallelic inactivation or hypermutation⁷⁶¹. A study of 59 patients with hormone-sensitive prostate cancer identified MSH2, CDK12, RB1, and TP53 alterations as independent risk factors for early progression to castration-resistant prostate cancer⁷¹⁵. Multiple studies have correlated low or absent MSH2 expression with a decreased risk of recurrence and increased disease-free survival and OS in patients with prostate cancer⁷⁶²⁻⁷⁶⁵.

FINDING SUMMARY

MSH2 encodes a DNA mismatch repair protein belonging to the mismatch repair (MMR) gene family. Defective MMR occurring as a result of mutation(s) in the MMR family (MLH1, MSH2, MSH6, or PMS2) can result in microsatellite instability (MSI), common in colon, endometrium, and stomach cancers²². Alterations such as seen here may disrupt MSH2 function or expression⁷⁶⁶⁻⁷⁷⁸.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in MSH2 are associated with an autosomal dominant condition known as Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC), which is characterized by increased risk of a number of cancers²⁷. In one large study of Lynch syndrome, endometrial cancer was the most common cancer reported outside the colon, with an incidence of 13.8%²⁸. For family members of patients with newly diagnosed endometrial cancer, the clinical utility of testing the patient for germline mutations in MMR genes is higher for mutations in MLH1 or MSH2 than it is for MSH6 or PMS2 mutations⁷⁷⁹. Therefore, in the appropriate clinical context, germline testing of MSH2 is recommended.

GENE

MSH3

ALTERATION

N385fs*19

HGVS VARIANT

NM_002439.3:c.1148dup (p.N385Qfs*19)

VARIANT CHROMOSOMAL POSITION

chr5:79970914

VARIANT ALLELE FREQUENCY (% VAF)

31.0%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Preclinical studies in the context of MSH3-deficient cancer cells have demonstrated antitumor efficacy of DNA-PKcs inhibitors⁷⁸⁰ and PARP inhibitors such as olaparib⁷⁸¹ as well as increased

chemosensitivity to cisplatin, oxaliplatin, and SN-38⁷⁸¹⁻⁷⁸². However, these remain to be tested in a clinical context.

FREQUENCY & PROGNOSIS

For patients with metastatic disease, MSH3 mutations have been reported with highest prevalence (4-5%) in colorectal (CRC) and endometrial cancers and are rare (<1%) in other tumor types such as prostate, lung, breast, ovarian, or pancreatic⁵⁹³. MSH3 loss has been correlated with the late development and progression of a variety of sporadic cancers including lung, ovarian, bladder, breast, and CRC tumors⁷⁸³⁻⁷⁸⁷. Certain germline polymorphisms in MSH3 have been associated with poor prognosis in CRC⁷⁸⁷, head and neck squamous cell carcinoma (HNSCC)⁷⁸⁸, non-small cell lung cancer (NSCLC)⁷⁸⁹, and pancreatic cancer⁷⁹⁰, while MSH3 loss was associated with improved post-surgery outcome in

1 study of patients with MLH1-deficient CRC⁷⁹¹.

FINDING SUMMARY

MSH3, while in the MutSβ heterodimer complex with MSH2, detects mismatched bases and participates in double-strand break repair by homologous recombination^{780,792-794}. Inactivating MSH3 frameshift mutations may increase microsatellite instability (MSI) and lead to the production of elevated microsatellite alterations at selected tetranucleotide repeats (EMAST)^{793,795}, which have been recognized as a genomic signature in colorectal cancer (CRC) with MSI⁷⁹⁶⁻⁷⁹⁷. Certain germline polymorphisms in MSH3 have been reported to increase the risk of various cancers including CRC⁷⁹⁸⁻⁸⁰³, breast^{798,804}, esophageal⁸⁰⁵, prostate^{798,806-807}, gastric⁸⁰⁸, and head and neck squamous cell carcinoma (HNSCC)⁷⁸⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

MSH6

ALTERATION

loss

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Numerous studies in various cancer types have shown that MSH6 loss or inactivation is associated with MSI and increased mutation burden^{14,18,22,68,756-757}. Clinical studies have shown that MSI is associated with patient responses to anti-programmed death 1 (PD-1) immune checkpoint inhibitors pembrolizumab^{7,809} and nivolumab⁶. Higher mutation burden was also reported to be associated with response to pembrolizumab⁸. Furthermore, MSI status correlates with higher PD-1 and PD-L1 expression⁷⁵⁰, potential biomarkers of response to PD-1 targeted immunotherapies. Therefore, inactivation of MSH6 may confer sensitivity to anti-PD-1 immune checkpoint inhibitors.

FREQUENCY & PROGNOSIS

MSH6 mutations have been reported at low frequencies in prostate adenocarcinoma, in up to 1% of cases across several studies^{54-55,57,75}. In one study, 10% (5/50) of primary advanced prostate cancer samples analyzed were reported to be hypermutated and characterized by MSI; of these hypermutated tumors, 4/5 harbored inactivating mutations in MSH6 and/or MSH2 and loss of MSH6 and/or MSH2 protein expression¹⁴. Published data investigating the prognostic implications of MSH6 alterations in prostate cancer are limited (PubMed, Oct 2022).

FINDING SUMMARY

MSH6 encodes MutS homolog 6 protein, a member of the mismatch repair (MMR) gene family. Defective MMR occurring as a result of mutation(s) in the MMR family (MLH1, MSH2, MSH6, or PMS2) can result in microsatellite instability (MSI), common in colon, endometrium, and stomach cancers²². Alterations such as seen here may disrupt MSH6 function or expression⁸¹⁰⁻⁸¹⁵.

POTENTIAL GERMLINE IMPLICATIONS

Germline mutations in MSH6 are associated with both "typical" and "atypical" forms of autosomal dominant Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC), which accounts for 1-7% of all colorectal cancers³⁰. Approximately 10% of all Lynch syndrome-associated mutations have been attributed to alterations in MSH6⁷⁵⁴. Carriers of mutations in MSH6 have a 60-80% risk of colorectal cancer²⁹. Lynch syndrome has an estimated prevalence in the general population ranging from 1:600 to 1:2000³⁰⁻³². Biallelic germline mutation of MSH6 has been shown to account for 20% of cases of the very rare syndrome Constitutional Mismatch Repair Deficiency (CMMRD), which is characterized by a 95% incidence rate of childhood onset lymphoma, leukemia and brain tumors, followed by early-onset colorectal cancer⁸¹⁶⁻⁸²⁰. Given the association between MSH6 and these inherited syndromes, in the appropriate clinical context, germline testing of MSH6 is recommended.

GENE

NOTCH3

ALTERATION

G463*

HGVS VARIANT

NM_000435.2:c.1387G>T (p.G463*)

VARIANT CHROMOSOMAL POSITION

chr19:15299151

VARIANT ALLELE FREQUENCY (% VAF)

42.8%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Several approaches for inhibiting NOTCH3 signaling are being developed, including neutralizing NOTCH antibodies such as tarextumab (OMP-59R5)⁸²¹, which targets NOTCH2 and NOTCH3, and pan-NOTCH inhibitors, such as

gamma-secretase inhibitors (GSI)⁸²²⁻⁸²⁴. In a Phase 2 study, the GSI AL101 (BMS-906024) elicited PR in 15% (6/39) and SD in 54% (21/39) of patients with metastatic adenoid cystic carcinoma harboring NOTCH activating alterations⁸²⁵. Phase 2 studies have evaluated the efficacy of tarextumab in combination with chemotherapy in metastatic pancreatic cancer or extensive-stage small cell lung cancer, though NOTCH3 expression was not found to be a predictor of OS or PFS in either study⁸²⁶. These approaches would not be relevant in the context of inactivating alterations, as seen here.

FREQUENCY & PROGNOSIS

NOTCH3 mutations have been reported in up to 2% of metastatic prostate cancers^{57,59} but not in two other studies of prostate adenocarcinoma^{54,56}. NOTCH3 amplification or overexpression has been associated with poor clinicopathological features in prostate carcinoma, although the role of NOTCH signaling in this disease is complex⁸²⁷⁻⁸²⁸.

FINDING SUMMARY

NOTCH3 encodes a member of the NOTCH family of receptors, which are involved in cell fate determination and various developmental processes. Upon binding of membrane-bound ligands, NOTCH signaling involves cleavage of the NOTCH intracellular domain (NICD), which subsequently forms part of a transcription factor complex that regulates downstream target genes⁸²⁹⁻⁸³⁰. Alterations that disrupt or remove the transmembrane domain (amino acids 1644-1664), RAM domain (amino acids 1665-1837), and/or ANK repeats region (amino acids 1838-2000), which are necessary for the transcriptional activity of NOTCH family proteins, as well as internal deletions that remove EGF repeats (7-10 and 21-22), have been shown in vitro to negatively affect ligand binding and reduce NOTCH3 transcriptional activity and are predicted to be inactivating⁸³⁰⁻⁸³⁴.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

NTRK2

ALTERATION

A11V

HGVS VARIANT

NM_006180.3:c.32C>T (p.A11V)

VARIANT CHROMOSOMAL POSITION

chr9:87285695

VARIANT ALLELE FREQUENCY (% VAF)

41.4%

as larotrectinib, entrectinib, AZD7451, belizatinib, and PLX7486⁸³⁵⁻⁸⁴⁵. In a Phase 1 study, the activity of larotrectinib was limited in patients who harbored NTRK amplification and not observed in patients with NTRK mutations in the absence of fusion⁸³⁵. A case study of a patient with esophageal carcinoma who harbored NTRK1 amplification reported a short PR⁸⁴⁶. It is not known whether these therapeutic approaches would be relevant in the context of alterations that have not been fully characterized, as seen here.

FREQUENCY & PROGNOSIS

NTRK2 mutation was detected in 8.6% of prostate carcinoma samples in COSMIC (Jun 2023)²⁴⁶. Published data investigating the prognostic implications of NTRK2 alterations in prostate cancer are limited (PubMed, Jan 2023).

FINDING SUMMARY

NTRK2 encodes neurotrophic tyrosine kinase receptor type 2 (TRKB), also known as tropomyosin-related kinase B. TRKB is a receptor for brain-derived neurotrophic factor (BDNF), and neurotrophin-4 and is a member of the insulin receptor subfamily⁸⁴⁷⁻⁸⁴⁸. TRKB activates the RAS-ERK and PI3K-AKT signaling pathways, regulating cell survival, proliferation, and differentiation in normal and neoplastic cells^{847,849-851}. Overexpression of TRKB promotes malignant transformation, cell invasiveness, and tumor metastasis, thereby suggesting a role for NTRK2 as an oncogene⁸⁵². Although alterations such as seen here have not been fully characterized and are of unknown functional significance, similar alterations have been previously reported in the context of cancer, which may indicate biological relevance.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical and preclinical data indicate that NTRK fusions predict sensitivity to TRK inhibitors such

GENE

PMS2

ALTERATION

Y268fs*39

HGVS VARIANT

NM_000535.4:c.802del (p.Y268Tfs*39)

VARIANT CHROMOSOMAL POSITION

chr7:6036957-6036958

VARIANT ALLELE FREQUENCY (% VAF)

46.9%

predict sensitivity to anti-PD-1 immune checkpoint inhibitors. However, this has not been directly demonstrated.

FREQUENCY & PROGNOSIS

PMS2 deletion or amplification have been each detected in <0.5% of analyzed prostate carcinomas, and PMS2 mutations have been detected in up to 1% of prostate cancer samples (cBioPortal, COSMIC, Sep 2022)^{176-177,246}. Alterations in MMR genes, including PMS2, associated with increased risk of certain cancers, including prostate carcinoma⁸⁵⁶, although data regarding the role of PMS2 in this tumor type are conflicting. Reduced PMS2 expression has been associated with poorly differentiated tumors and more aggressive prostate cancer^{751,857-859}. Increased PMS2 expression has been associated with genetic instability and decreased time to recurrence in prostate cancer⁸⁶⁰⁻⁸⁶¹. Restoring PMS2 expression had a protective effect in a PMS2-deficient preclinical model of prostate cancer, primarily through increased apoptosis⁸⁶².

FINDING SUMMARY

PMS2 encodes an endonuclease that has been shown to play a critical role in DNA mismatch repair (MMR) and apoptotic responses to DNA damage⁸⁶³. Both abnormally high levels of PMS2, caused by protein overexpression, and inactivating

mutation leading to loss of PMS2 activity have been shown to result in genomic instability, resistance to genotoxic chemotherapy, and increased tumorigenicity in vivo⁸⁶⁴⁻⁸⁶⁵. Alterations such as seen here may disrupt PMS2 function or expression^{853,866-870}.

POTENTIAL GERMLINE IMPLICATIONS

One or more of the PMS2 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Lynch syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline PMS2 mutations have been associated with autosomal dominant Lynch syndrome and the rarer autosomal recessive Turcot syndrome⁸⁷¹⁻⁸⁷³. Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer or HNPCC) accounts for 1-7% of all colorectal cancers³⁰⁻³¹. One study reported germline PMS2 mutation in 62% (61/99) of patients diagnosed with Lynch syndrome-associated tumors⁸⁶⁷. Turcot syndrome is characterized by concurring primary brain tumors and colon cancers and/or colorectal adenomas in pediatric patients⁸⁷⁴. Therefore, in the appropriate clinical context, germline testing of PMS2 is recommended.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Defective MMR that occurs because of mutation(s) in the MMR family, which includes MLH1, MSH2, MSH6, and PMS2, can result in microsatellite instability (MSI), which is common in colon, endometrial, and stomach cancers²². Alterations in PMS2 can lead to impaired MMR activity⁸⁵³, and selective loss of PMS2 by either mutation or loss of expression has been reported in colorectal cancer and endometrial cancer with MSI-high phenotype^{757,854-855}. Clinical studies have shown that MSI predicts patient responses to the anti-programmed death 1 (PD-1) immune checkpoint inhibitors pembrolizumab^{7,809} and nivolumab⁶, and alterations resulting in PMS2 functional loss may

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS
GENE

PRKAR1A

ALTERATION
 Q275*

HGVS VARIANT

NM_212472.1:c.823C>T (p.Q275*)

VARIANT CHROMOSOMAL POSITION

chr17:66525064

VARIANT ALLELE FREQUENCY (% VAF)

39.3%

genomic alterations in PRKAR1A. Preclinical studies have shown that depletion of PRKAR1A may sensitize cells to mTOR inhibitors such as rapamycin⁸⁷⁵⁻⁸⁷⁶ or TKIs targeting SRC such as dasatinib⁸⁷⁷.

FREQUENCY & PROGNOSIS

PRKAR1A mutations have been reported in less than 1% of solid tumor types and less than 0.5% of hematologic cancer samples analyzed in the COSMIC database (2023)²⁴⁶. Small studies of melanotic schwannoma, a rare tumor that can be sporadic or associated with Carney complex, have identified PRKAR1A mutation in 100% (12/12)⁸⁷⁸ and loss of protein expression in 33% (7/21)⁸⁷⁹ of cases. Published data investigating the prognostic implications of PRKAR1A alterations in cancer are

limited (PubMed, Jan 2023). One small study reported that low tumor PRKAR1A protein expression was associated with reduced OS in patients with lung adenocarcinoma (HR=1.88o)⁸⁸⁰.

FINDING SUMMARY

PRKAR1A encodes a regulatory subunit of cyclic AMP-dependent protein kinase⁸⁸¹. PRKAR1A is thought to be a tumor suppressor, and inactivating germline mutations and deletions are associated with Carney complex, an autosomal dominant syndrome characterized by increased risk of primary pigmented nodular adrenocortical disease, cardiac and other myxomas, endocrine tumors, and schwannomas⁸⁸²⁻⁸⁸⁶. Somatic PRKAR1A mutations have also been reported in cardiac myxoma⁸⁸⁷⁻⁸⁸⁹.

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address

GENE

QKI

ALTERATION

K134fs*14, R124*

HGVS VARIANT

NM_006775.2:c.401del (p.K134Rfs*14),

NM_006775.2:c.370C>T (p.R124*)

VARIANT CHROMOSOMAL POSITION

chr6:163899919-163899920, chr6:163899896

VARIANT ALLELE FREQUENCY (% VAF)

83.2%, 40.1%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies approved or in clinical trials that directly address genomic alterations in QKI.

FREQUENCY & PROGNOSIS

Recurrent deletions of QKI have been reported in astrocytomas and glioblastomas⁸⁹⁰⁻⁸⁹². Decreased expression of quaking has also been reported in colon and gastric cancer, largely due to QKI promoter hypermethylation; in one study, forced

expression of quaking in colon cancer cells inhibited proliferation and tumorigenesis⁸⁹³⁻⁸⁹⁴. However, another study reported that QKI acts as an oncogene in breast cancer by repressing the expression of the tumor suppressor FOXO1⁸⁹⁵.

FINDING SUMMARY

QKI encodes quaking, an RNA-binding protein that plays roles in RNA metabolism and signal transduction and is necessary for myelination and embryonic development⁸⁹⁶⁻⁸⁹⁷. QKI acts as a negative regulator of cell-cycle progression⁸⁹⁸.

GENE

SETD2

ALTERATION

R1407fs*5

HGVS VARIANT

NM_014159.6:c.4219del (p.R1407Gfs*5)

VARIANT CHROMOSOMAL POSITION

chr3:47161906-47161907

VARIANT ALLELE FREQUENCY (% VAF)

37.4%

POTENTIAL TREATMENT STRATEGIES
— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in SETD2.

FREQUENCY & PROGNOSIS

Somatic inactivating alterations of SETD2 are documented to occur at low frequency in a number of solid tumors, most commonly in renal carcinoma⁸⁹⁹. SETD2 has been associated with favorable prognosis in gastric cancer⁹⁰⁰. SETD2 has also been associated with poor prognosis in RCC and MDS⁹⁰¹⁻⁹⁰², while data in other tumor types is limited (PubMed, Jun 2023).

FINDING SUMMARY

SETD2 encodes a histone lysine-36 methyltransferase⁹⁰³ that preferentially interacts with the expanded N-terminal polyglutamine tracts present in mutant huntingtin, implicating it in the pathogenesis of Huntington disease⁹⁰⁴. SETD2 mRNA expression has been observed to be consistently reduced in breast tumors relative to adjacent non-tumor tissue, suggesting a potential tumor suppressor role⁹⁰⁵. SETD2 alterations such as observed here have been shown to be inactivating⁹⁰⁶⁻⁹¹¹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

SMAD4

ALTERATION

G386S - subclonal

HGVS VARIANT

NM_005359.5:c.1156G>A (p.G386S)

VARIANT CHROMOSOMAL POSITION

chr18:48593405

VARIANT ALLELE FREQUENCY (% VAF)

4.9%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in SMAD4. Preclinical studies in colorectal cancer have reported associations of SMAD4 inactivation or loss with sensitivity to inhibitors of Aurora kinase A⁹¹² and the Wnt/beta-catenin pathway⁹¹³.

— Nontargeted Approaches —

Clinical studies have reported associations of SMAD4 loss or low SMAD4 expression with improved responses to chemotherapeutic agents in

patients with pancreatic cancer⁹¹⁴⁻⁹¹⁶ and non-small cell lung cancer (NSCLC)⁹¹⁷. Other clinical studies in pancreatic cancer have reported an association of high SMAD4 expression with better responses to neoadjuvant chemotherapy⁹¹⁸ and adjuvant chemoradiotherapy⁹¹⁹.

FREQUENCY & PROGNOSIS

SMAD4 mutation or homozygous deletion is most frequently observed in pancreatic adenocarcinoma (43%)⁹²⁰, pancreatic acinar cell carcinoma (26%)⁹²¹, cholangiocarcinoma (25%)⁹²², small intestine cancer (20%)⁹²³, appendiceal adenocarcinoma (14-20% mutation; 57% deletion)⁹²⁴⁻⁹²⁵, colorectal adenocarcinoma (CRC; 14%)⁶⁸, esophageal adenocarcinoma (14%)⁹²⁶, and stomach adenocarcinoma (13%)⁴⁶². In preclinical studies, SMAD4 loss of function has been implicated in the development of mucinous neoplasms of the pancreas, including mucinous cystic neoplasms (MCN)⁹²⁷ and intraductal papillary mucinous neoplasms (IPMN)⁹²⁸; in clinical samples, SMAD4 homozygous deletion has been observed in 10% of IPMNs and 8% of MCNs, and mutation was also observed in 5% of IPMNs⁹²⁹. SMAD4 gene alterations have been associated with reduced OS for patients with pancreatic adenocarcinoma⁹³⁰. Reduced SMAD4 expression has been associated

with worse prognosis in various cancer types, including CRC⁹³¹⁻⁹³³, appendiceal mucinous neoplasm⁹³⁴, gastric adenocarcinoma⁹³⁵⁻⁹³⁶, esophageal adenocarcinoma⁹³⁷, esophageal squamous cell carcinoma⁹³⁸, breast cancer⁹³⁹, and prostate cancer⁹⁴⁰.

FINDING SUMMARY

SMAD4, also known as DPC4, encodes a tumor suppressor that regulates transcriptional activity downstream of TGF-beta receptor signaling⁹⁴¹⁻⁹⁴². SMAD4 alterations that result in loss or disruption of the MH1 domain (aa 18-142), MH2 domain (aa 323-552), or SAD domain (aa 275-320) are predicted to be inactivating⁹⁴³⁻⁹⁵⁶.

POTENTIAL GERMLINE IMPLICATIONS

Germline SMAD4 mutations, including those at the R361 hotspot, have been observed in patients with juvenile polyposis syndrome⁹⁵⁷⁻⁹⁵⁹, which is associated with an increased risk of gastrointestinal cancers⁹⁶⁰. The penetrance of deleterious SMAD4 mutations in patients with colon cancer is estimated at 20% by age 35 and 70% by age 65⁹⁶¹. In the appropriate clinical context, germline testing of SMAD4 is recommended.

GENE

SPEN

ALTERATION

K1999fs*23

HGVS VARIANT

NM_015001.2:c.5996del (p.K1999Rfs*23)

VARIANT CHROMOSOMAL POSITION

chr1:16258726-16258727

VARIANT ALLELE FREQUENCY (% VAF)

37.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address SPEN inactivating mutations. Although gamma-secretase inhibitors are in clinical development to target NOTCH activation, it is not known if these therapies would be beneficial in the context of SPEN mutations.

FREQUENCY & PROGNOSIS

Pan-cancer analysis of the MSK-MET dataset identified SPEN alterations in 1.2% of all samples⁵⁹³. SPEN truncating mutations have been reported at a higher frequency in adenoid cystic

carcinoma (ACC; 21%)⁹⁶² and splenic marginal zone lymphoma (SMZL; 5%)⁹⁶³.

FINDING SUMMARY

SPEN (also known as MINT or SHARP) encodes a transcriptional regulator that interacts with histone deacetylase 1 (HDAC1) and the silencing mediator for retinoid and thyroid receptors/nuclear receptor corepressor (SMRT/NcoR)⁹⁶⁴⁻⁹⁶⁵ and represses the transcriptional activity of the NOTCH signaling pathway⁹⁶⁶⁻⁹⁶⁷. SPEN alterations that result in the disruption of functional domains (aa 2804-2816 or aa 3498-3664) are likely to be inactivating⁹⁶⁵⁻⁹⁶⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE
TGFB2

ALTERATION

R497* - subclonal

HGVS VARIANT

NM_003242.5:c.1489C>T (p.R497*)

VARIANT CHROMOSOMAL POSITION

chr3:30729968

VARIANT ALLELE FREQUENCY (% VAF)

6.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies in development or in use for patients with inactivating mutations of TGFB2. TGFB2 activating mutations may predict sensitivity to therapies targeting TGFB2, such as Lucanix and IMC-TR1, which have been in development and are in clinical trials in certain tumor types, but are not currently recruiting patients⁹⁶⁸⁻⁹⁷². These approaches would not be

relevant in the context of inactivating alterations, as seen here.

FREQUENCY & PROGNOSIS

TGFB2 deletions or mutations have been reported in each 0-1% of prostate adenocarcinomas^{54,56,59}. Down-regulation of TGFB2 protein has been observed in 12/20 high-grade prostatic intraepithelial neoplasia and 36/60 prostate cancers⁹⁷³. Published data investigating the prognostic implications of TGFB2 in prostate carcinoma are limited (PubMed, Sep 2022). Conditional knockout of TGFB2 in stromal fibroblastic cells in a transgenic mouse model results in prostatic intraepithelial neoplasia and adenocarcinoma⁹⁷⁴.

FINDING SUMMARY

TGFB2 encodes a type 2 receptor for transforming growth factor-beta (TGF-beta)⁹⁷⁵. Loss of TGF-beta signaling results in insensitivity to the growth inhibitory effects of TGF-beta and ultimately causes an increase in cellular growth and transformation⁹⁷⁶. Constitutive activation of

TGFB2 has been reported to increase the invasive behavior of carcinoma cells⁹⁶⁹. TGFB2 R497* is predicted to be inactivating⁹⁷⁷⁻⁹⁸⁰.

POTENTIAL GERMLINE IMPLICATIONS

The TGFB2 R497* variant observed here has been described in the ClinVar database as a pathogenic germline mutation (by an expert panel or multiple submitters) associated with Loeys-Dietz syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TGFB2 are associated with hereditary nonpolyposis colorectal cancer (HNPCC) type 6⁹⁸¹ and the autosomal dominant disorder Loeys-Dietz syndrome type 2^{980,982-984}. Both disorders are rare, with no clear prevalence data reported; HNPCC type 6 has only been described in one case report⁹⁸¹ and Loeys-Dietz syndrome type 2 is also rarely reported⁹⁸⁵. However, in the appropriate clinical context, germline testing of TGFB2 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

GENE

TP53

ALTERATION

R196*, Y205H - subclonal, R175C

HGVS VARIANT

NM_000546.4:c.586C>T (p.R196*),

NM_000546.4:c.613T>C (p.Y205H),

NM_000546.4:c.523C>T (p.R175C)

VARIANT CHROMOSOMAL POSITION

chr17:7578263, chr17:7578236, chr17:7578407

VARIANT ALLELE FREQUENCY (% VAF)

41.2%, 0.78%, 39.3%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies to address TP53 mutation or loss. However, tumors with TP53 loss of function alterations may be sensitive to the WEE1 inhibitor adavosertib⁹⁸⁶⁻⁹⁸⁹ or p53 gene therapy such as SGT53⁹⁹⁰⁻⁹⁹⁴. In a Phase 1 study, adavosertib in combination with gemcitabine, cisplatin, or carboplatin elicited PRs in 9.7% and SDs in 53% of patients with solid tumors; the response rate was 21% (4/19) for patients with TP53 mutations versus 12% (4/33) for patients who were TP53 wildtype⁹⁹⁵. A Phase 2 trial of adavosertib in combination with chemotherapy (gemcitabine, carboplatin, paclitaxel, or doxorubicin) reported a 32% (30/94, 3 CR) ORR and a 73% (69/94) DCR for patients with platinum-refractory TP53-mutated ovarian, Fallopian tube, or peritoneal cancer⁹⁹⁶. A smaller Phase 2 trial of adavosertib in combination with carboplatin achieved a 43% (9/21, 1 CR) ORR and a 76% (16/21) DCR for patients with platinum-refractory TP53-mutated ovarian cancer⁹⁹⁷. The combination of adavosertib with paclitaxel and carboplatin for patients with TP53-mutated ovarian cancer also significantly increased PFS compared with paclitaxel and carboplatin alone⁹⁹⁸. In the Phase 2 VIKTORY trial, patients with TP53-mutated metastatic and/or recurrent gastric cancer experienced a 24% (6/25) ORR with adavosertib combined with paclitaxel⁹⁹⁹. A Phase 1 trial of neoadjuvant adavosertib in combination with cisplatin and docetaxel for head and neck squamous cell carcinoma (HNSCC) elicited a 71% (5/7) response rate for patients with TP53 alterations¹⁰⁰⁰. The Phase 2 FOCUS4-C trial for patients with TP53- and RAS-mutated colorectal cancer reported improvement in PFS (3.61 vs. 1.87

months, HR=0.35, p=0.0022), but not OS (14.0 vs 12.8 months, p=0.93), following adavosertib treatment compared with active monitoring¹⁰⁰¹. In a Phase 1b clinical trial of SGT-53 in combination with docetaxel for patients with solid tumors, 75% (9/12) of evaluable patients experienced clinical benefit, including 2 confirmed and 1 unconfirmed PRs and 2 instances of SD with significant tumor shrinkage⁹⁹⁴. Missense mutations leading to TP53 inactivation may be sensitive to therapies that reactivate mutated p53 such as eprenetapopt. In a Phase 1b trial for patients with p53-positive high-grade serous ovarian cancer, eprenetapopt combined with carboplatin and pegylated liposomal doxorubicin achieved a 52% (11/21) response rate and 100% DCR¹⁰⁰². A Phase 1 trial of eprenetapopt with pembrolizumab for patients with solid tumors reported an ORR of 10% (3/29)¹⁰⁰³.

— Nontargeted Approaches —

In post-hoc analyses of a Phase 1/2 trial, the addition of carboplatin to cabazitaxel significantly improved PFS (6.0 vs. 2.2 months) and OS (17.4 vs. 9.9 months) for patients with an aggressive variant prostate cancer molecular signature (AVPC-MS, composed of alterations in at least 2 of the genes RB1, TP53, and PTEN); no significant difference in outcomes were observed for patients lacking this signature (NCCN Prostate Cancer Guidelines, v1.2023)³⁹⁵. Post-hoc analysis of a Phase 2 trial in metastatic castration-resistant prostate cancer suggests that patients with aggressive variant prostate cancer (AVPC), molecularly characterized by harboring alterations in at least 2 genes of TP53, RB1, and PTEN, as seen in this sample, may benefit from cabazitaxel combined with carboplatin compared with cabazitaxel alone (median PFS of 5.1 vs. 2.2 months, p=0.03; estimated median OS of 11.2 vs. 10.5 months, p=0.11), whereas patients who were AVPC-negative did not benefit from the more intense chemotherapy combination in this study³⁹⁵.

FREQUENCY & PROGNOSIS

TP53 mutations have been reported in 18-40% of prostate cancers^{350,1004}. A study of 59 patients with hormone-sensitive prostate cancer identified MSH2, CDK12, RB1, and TP53 alterations as independent risk factors for early progression to castration-resistant prostate cancer²¹⁵. Overexpression of p53, which is indicative of TP53 dysregulation, has been reported to be significantly more common in late-stage and hormone-refractory prostate cancers and has been found to

be associated with prostate-specific antigen (PSA) recurrence in low- and intermediate-grade prostate cancer¹⁰⁰⁵. TP53 loss has been found to be associated with prostate cancer-specific mortality in univariate analysis¹⁰⁰⁶. Concurrent alterations in at least 2 of the TP53, RB1, and PTEN genes, as seen in this sample, molecularly define a subtype of prostate cancer with aggressive clinical course and reduced sensitivity to androgen-deprivation therapies (NCCN Prostate Cancer Guidelines, v1.2023)^{395,401-404}.

FINDING SUMMARY

Functional loss of the tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers¹⁰⁰⁷. Alterations such as seen here may disrupt TP53 function or expression¹⁰⁰⁸⁻¹⁰¹².

POTENTIAL GERMLINE IMPLICATIONS

One or more of the TP53 variants observed here has been described in the ClinVar database as a likely pathogenic or pathogenic germline mutation (by an expert panel or multiple submitters) associated with Li-Fraumeni syndrome (ClinVar, Apr 2023)¹⁶³. Follow-up germline testing would be needed to distinguish whether the finding in this patient is somatic or germline. Germline mutations in TP53 are associated with the very rare autosomal dominant disorder Li-Fraumeni syndrome and the early onset of many cancers¹⁰¹³⁻¹⁰¹⁵, including sarcomas¹⁰¹⁶⁻¹⁰¹⁷. Estimates for the prevalence of germline TP53 mutations in the general population range from 1:5,000¹⁰¹⁸ to 1:20,000¹⁰¹⁷. For pathogenic TP53 mutations identified during tumor sequencing, the rate of germline mutations was 1% in the overall population and 6% in tumors arising before age 30¹⁰¹⁹. In the appropriate clinical context, germline testing of TP53 is recommended.

POTENTIAL CLONAL HEMATOPOIESIS IMPLICATIONS

Variants seen in this gene have been reported to occur in clonal hematopoiesis (CH), an age-related process in which hematopoietic stem cells acquire somatic mutations that allow for clonal expansion⁵⁰⁷⁻⁵¹². CH in this gene has been associated with increased mortality, risk of coronary heart disease, risk of ischemic stroke, and risk of secondary hematologic malignancy⁵⁰⁷⁻⁵⁰⁸. Clinical management of patients with CH in this gene may include monitoring for hematologic changes and reduction of controllable risk factors for cardiovascular disease⁵¹³. Comprehensive

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

GENOMIC FINDINGS

genomic profiling of solid tumors detects nontumor alterations that are due to CH^{511,514-515}.

Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if

this alteration is present in tumor or is secondary to CH.

GENE

XRCC2

ALTERATION

L117fs*17

HGVS VARIANT

NM_005431.1:c.350del (p.L117Wfs*17)

VARIANT CHROMOSOMAL POSITION

chr7:152346219-152346220

VARIANT ALLELE FREQUENCY (% VAF)

37.4%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no therapies to directly target XRCC2 inactivation. However, loss of functional XRCC2 is associated with increased sensitivity to DNA cross-linking agents such as mitomycin C and to ionizing radiation¹⁰²⁰⁻¹⁰²².

FREQUENCY & PROGNOSIS

XRCC2 mutation has been seen in 2.4% of colorectal adenocarcinomas⁶⁸, 2.4% of stomach adenocarcinomas⁴⁶², and 1.3% of uterus corpus endometrioid carcinomas⁶⁵. High-level amplification of XRCC2 has been reported in 4.7% of ovarian serous cystadenocarcinomas¹⁰²³, 3.9% of lung adenocarcinomas¹⁰²⁴, and 1.8% of glioblastomas¹⁰²⁵, whereas homozygous deletion of XRCC2 has been seen in 3.1% of acute myeloid leukemias¹⁰²⁶, 3% of esophagus-stomach cancers¹⁰²⁷, and 1.4% of head and neck squamous cell carcinomas⁵²⁴. Alterations in other tumor types are more rarely observed (cBioPortal, Jan 2023)¹⁷⁶⁻¹⁷⁷. XRCC2 polymorphisms, including R188H, have been associated with low penetrance risk of developing breast cancer¹⁰²⁸⁻¹⁰³⁰; however, these findings are controversial, with some studies instead showing a protective effect¹⁰³¹⁻¹⁰³⁶. XRCC2 polymorphisms have also been implicated in decreased survival in the context of pancreatic¹⁰³⁷⁻¹⁰³⁸, lung¹⁰³⁹, and rectal¹⁰⁴⁰ cancers

but reduced risk of bladder¹⁰⁴¹ and ovarian¹⁰³⁶ cancers.

FINDING SUMMARY

XRCC2 encodes a RAD51 homolog involved in homologous recombination; in complex with RAD51 proteins, XRCC2 enables the repair of DNA double-strand breaks caused by genotoxic stresses such as DNA cross-linking agents and ionizing radiation^{1020,1022,1042-1046}. Alterations that disrupt the Walker A (residues 48-55) or Walker B (residues 145-149) motifs, or the C-terminal region of XRCC2 involved in RAD51 binding, disrupt homologous recombination by XRCC2^{1021,1028,1047}. The missense mutations R91W, I95L, C120Y, L133P, R188C, and E207G also impair XRCC2 function, although to a lesser extent than alterations leading to premature protein truncation¹⁰⁴⁷.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT
IN PATIENT'S TUMOR TYPE

Dostarlimab

Assay findings association

CDK12

R1048*, N474fs*8 - subclonal

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Dostarlimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor response. It is FDA approved to treat patients with mismatch repair deficient recurrent or advanced endometrial cancer or solid tumors. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵. On the basis of limited clinical evidence in prostate cancer,

including multiple prostate-specific antigen (PSA) responses, CDK12 inactivating alterations may predict sensitivity to PD-1 inhibitors^{194,1049-1050}. On the basis of prospective clinical data showing efficacy of dostarlimab against various microsatellite instability-high (MSI-H) solid tumors^{9,1051-1053}, MSI-H status may predict sensitivity to dostarlimab.

SUPPORTING DATA

Clinical data on the efficacy of dostarlimab for the treatment of prostate cancer are limited (PubMed, Mar 2023). Dostarlimab has been studied primarily in recurrent and advanced mismatch repair-deficient (dMMR) endometrial and non-endometrial cancers^{9,1052,1054}. In the Phase 1 GARNET trial, single-agent dostarlimab elicited an ORR of 39% (41/106) and an immune-related ORR of 46% (50/110) for patients with non-endometrial dMMR solid tumors^{1052,1055}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Olaparib

Assay findings association

BARD1
R406*

BRCA2
D946fs*14

BRIP1
splice site 918+1G>A - subclonal

CDK12
R1048*, N474fs*8 - subclonal

CHEK1
T226fs*14

FANCL
G119*

AREAS OF THERAPEUTIC USE

The PARP inhibitor olaparib is FDA approved to treat patients with epithelial ovarian, Fallopian tube, or primary peritoneal cancer, patients with deleterious or suspected deleterious gBRCA-mutated pancreatic adenocarcinoma or HER2-negative breast cancer, and patients with prostate cancer and mutations in homologous recombination repair genes. Olaparib is also approved in combination with bevacizumab to treat patients with ovarian, Fallopian tube, or primary peritoneal cancer with deleterious or suspected deleterious somatic or gBRCA mutation and/or genomic instability. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in ovarian^{85,245,1056}, breast¹⁷⁵, endometrial¹⁰⁵⁷, and prostate cancer¹⁷⁴, loss or inactivation of BRIP1 may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in ovarian cancer^{70,245}, FANCL loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in prostate^{80,174,210,1058-1059}, loss or inactivation of genes that are involved in homologous recombination repair may confer sensitivity to PARP inhibitors. CHEK1 inhibition has been associated with sensitivity to PARPi in preclinical studies^{224,1060}. On the basis of extensive clinical evidence in ovarian cancer⁸⁹⁻⁹³ as well as strong clinical evidence in multiple other cancer types^{79-81,89,92,96,1061}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to olaparib. On the basis of preclinical evidence^{71,74} and clinical evidence in ovarian cancer⁷⁰, BARD1 loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of limited clinical benefit in prostate^{209-210,1062} and ovarian¹⁰⁶³ cancer, CDK12 inactivating alterations may confer sensitivity to PARP inhibitors.

SUPPORTING DATA

The Phase 3 PROfound study for patients with metastatic castration-resistant prostate cancer (mCRPC) reported improved radiographic PFS (rPFS; 7.4 vs. 3.6 months, HR=0.34) and median OS (mOS; 19.1 vs. 14.7 months, HR=0.69) with olaparib compared with physician's choice of abiraterone plus prednisone or enzalutamide for patients with BRCA1/2 or ATM alterations^{209,1064}. For

patients with other homologous recombination repair (HRR) gene alterations, PFS (4.8 vs. 3.3 months, HR=0.88) and mOS (14.1 vs. 11.5 months, HR=0.96) were numerically increased with olaparib¹⁰⁶⁴. Other studies, including the Phase 2 TOPARP-A and TOPARP-B studies, reported similar results^{80,210,1065}. In a real-world study of olaparib and/or rucaparib for heavily pretreated prostate cancer, patients with pathogenic BRCA2 mutations experienced greater benefit than patients with other HRR mutations (median PFS of 7.2 vs. 2.8 months, p=0.291; 30% decrease in prostate-specific antigen of 69% vs. 4.0%, p<0.001)¹⁰⁶⁶. In the Phase 3 KEYLYNK-010 study for patients with pretreated mCRPC, treatment with olaparib plus pembrolizumab did not lead to significantly improved rPFS (4.4 months vs. 4.2 months, HR=1.02) or OS (15.8 months vs. 14.6 months, HR=0.94)¹⁰⁶⁷. Benefits were also seen in Phase 1 or 2 studies of olaparib in combination with durvalumab¹⁰⁶⁸, pembrolizumab¹⁰⁶⁹⁻¹⁰⁷⁰, or the ATP inhibitor ceralasertib¹⁰⁷¹ for patients with prostate cancer. PROfound patients with BRCA1/2 or ATM alterations also had improved ORR (33.3% [28/84] vs. 2.3% [1/43], p<0.001) with olaparib compared with physician's choice of enzalutamide or abiraterone/prednisone²⁰⁹. A Phase 2 trial of olaparib for patients with germline BRCA1/2 mutation reported 50% (4/8) PRs and 25% (2/8) SDs for patients with previously treated prostate cancer⁷⁹. In the Phase 3 PROpel study for patients with metastatic castration-resistant prostate cancer, treatment with first-line olaparib plus abiraterone and prednisone led to significantly improved rPFS (not reached vs. 8.4 months, HR=0.23) in patients with BRCA1 and BRCA2 mutations compared with placebo¹⁰⁷². PROfound patients with CDK12 alterations treated with olaparib experienced numerically improved PFS (5.1 vs. 2.2 months) and OS (14.1 vs. 11.5 months, HR=0.97) compared with physician's choice of abiraterone/prednisone or enzalutamide²⁰⁹. The Phase 2 TOPARP-B study reported reduced circulating tumor cell levels for 41.7% (5/12) of patients with CDK12-altered mCRPC treated with olaparib, but no PSA50 or objective responses²¹⁰. In a Phase 2 study of olaparib plus pembrolizumab for advanced solid tumors, patients with BRCA1 or BRCA2 mutations achieved an ORR of 29% (6/21), whereas patients with mutations in other homologous recombination repair genes achieved an ORR of 6.3% (2/32)¹⁰⁷³.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Olaparib + Abiraterone

Assay findings association

BRCA2

D946fs*14

AREAS OF THERAPEUTIC USE

Olaparib is a PARP inhibitor, and abiraterone is an orally available CYP17 inhibitor. These 2 therapies in combination together with prednisone or prednisolone are FDA approved to treat patients with BRCA1/2-mutated metastatic castration-resistant prostate cancer.

GENE ASSOCIATION

On the basis of clinical evidence in prostate cancer¹⁰⁷⁴⁻¹⁰⁷⁸, inactivation of either BRCA1 or BRCA2 may confer sensitivity to the combination of a PARP inhibitor and abiraterone.

SUPPORTING DATA

The Phase 3 PROpel study assessed first-line olaparib plus abiraterone and prednisone for patients with metastatic castration-resistant prostate cancer (mCRPC) and reported improved median radiographic PFS (rPFS) (27.6 vs. 16.4 months, HR=0.61) and improved median OS (42.1 vs. 34.7

months, HR=0.81) compared with abiraterone, prednisone, and placebo; greater benefit was observed among patients with homologous recombination repair (HRR) mutations for both rPFS (NR vs. 13.9 months, HR=0.50) and OS (NR vs. 28.5 months, HR=0.66), than patients without HRR mutations (rPFS: 24.1 vs. 19.0 months, HR=0.76; OS: 42.1 vs. 38.9 months, HR=0.89)¹⁰⁷⁴⁻¹⁰⁷⁵. For patients with mCRPC harboring inactivating germline or somatic alterations in BRCA1, BRCA2, and/or ATM, the triple combination of olaparib with abiraterone and prednisone improved median PFS (not reached vs. 11.3 or 10.4 months, respectively) relative to single-agent olaparib or abiraterone plus prednisone in the Phase 2 BRCA Away trial¹⁰⁷⁶. Another Phase 2 study comparing olaparib or placebo combined with abiraterone and prednisone or prednisolone for patients with mCRPC who had progressed on chemotherapy reported similar results, with median rPFS of 13.8 months and 8.2 months, respectively (HR=0.65, p=0.034)¹⁰⁷⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Pembrolizumab

Assay findings association

CDK12

R1048*, N474fs*8 - subclonal

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Pembrolizumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved for patients with tumor mutational burden-high (≥ 10 Muts/Mb), microsatellite instability-high (MSI-H), or MMR-deficient (dMMR) solid tumors; as monotherapy for PD-L1-positive head and neck squamous cell cancer (HNSCC), cervical cancer, or esophageal cancer; and in combination with chemotherapy for PD-L1-positive triple-negative breast cancer (TNBC) or cervical cancer. It is also approved in various treatment settings as monotherapy for patients with non-small cell lung cancer (NSCLC), melanoma, HNSCC, urothelial carcinoma, hepatocellular carcinoma, Merkel cell carcinoma, cutaneous squamous cell carcinoma, MSI-H or dMMR endometrial carcinoma, classical Hodgkin lymphoma, or primary mediastinal large B-cell lymphoma; and in combination with chemotherapy or targeted therapy for NSCLC, HNSCC, esophageal or gastroesophageal junction cancer, renal cell carcinoma, TNBC, urothelial carcinoma, or endometrial carcinoma that is not MSI-H or dMMR. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of limited clinical evidence in prostate cancer, including multiple prostate-specific antigen (PSA) responses, CDK12 inactivating alterations may predict sensitivity to PD-1 inhibitors^{194,1049-1050}. On the basis of multiple prospective clinical studies showing efficacy of pembrolizumab against MSI-H or mismatch repair-deficient (dMMR) solid tumors^{7,1080-1084}, MSI-H status may predict sensitivity to pembrolizumab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

For patients with prostate cancer enrolled in various clinical trials, Tumor Mutational Burden (TMB) of 10

Muts/Mb or higher was associated with an ORR of 9.1% (1/11) on pembrolizumab, whereas a lower TMB was associated with an ORR of 6% (n=115)¹⁰⁸⁵. A case study described a patient with POLE-mutant prostate cancer and high TMB who experienced progression on prior treatment and exhibited a durable response from pembrolizumab¹⁰⁸⁶. In a retrospective study of patients with CDK12-mutated mCRPC receiving fourth to sixth line pembrolizumab or nivolumab, a median PFS of 5.4 months and 33.3% (3/9) prostate-specific antigen (PSA) response rate were reported¹⁹⁸. In another retrospective study of patients with CDK12-mutated mCRPC, a 10.5% (2/19) rate of PSA decline greater than 50% (PSA50) and mPFS of 2.8 months following immunotherapy (15/19 pembrolizumab) treatment were reported¹⁰⁵⁰. In the Phase 3 KEYNOTE-921 trial, the addition of pembrolizumab to docetaxel did not significantly improve radiographic PFS (rPFS; 8.6 vs. 8.3 months, HR=0.85) or median OS (mOS; 19.6 vs. 19.0 months, HR=0.92) compared with docetaxel for patients with metastatic castration-resistant prostate cancer (mCRPC)¹⁰⁸⁷. In the KEYNOTE-199 and -365 trials, patients with mCRPC treated with pembrolizumab plus enzalutamide experienced a prostate-specific antigen (PSA) decrease $\geq 50\%$ (PSA50) of 8.8-23% and a DCR of 32-51%; an ORR of 12.0-12.3% was reported for patients who were Response Evaluation Criteria in Solid Tumors (RECIST) measurable¹⁰⁸⁸⁻¹⁰⁸⁹. Within the pembrolizumab monotherapy arms of KEYNOTE-199, patients with mCRPC reported a 13-39% DCR and an mOS of 7.9-14.1 months between all cohorts¹⁰⁹⁰. In another cohort of KEYNOTE-365, patients with mCRPC treated with pembrolizumab plus olaparib achieved a confirmed PSA response rate of 15% (15/102), median rPFS of 4.5 months, and mOS of 14 months; an ORR of 8.5% (5/59 PRs) was reported for patients with RECIST-measurable disease and was generally consistent across PD-L1 and homologous recombination repair subgroups¹⁰⁹¹. A Phase 1b study for small cell cancer of the urothelium and neuroendocrine cancer of the prostate (NECP) reported an ORR of 43% (3/7) and a 36% PFS rate at 12 months in the NECP cohort following treatment with pembrolizumab in combination with chemotherapy¹⁰⁹². One case study reported durable PR to pembrolizumab for 1 patient with metastatic platinum-refractory small cell carcinoma of the prostate¹⁰⁹³.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Rucaparib

Assay findings association
BARD1
R406*

BRCA2
D946fs*14

BRIP1
splice site 918+1G>A - subclonal

CDK12
R1048*, N474fs*8 - subclonal

CHEK1
T226fs*14

FANCL
G119*

AREAS OF THERAPEUTIC USE

The PARP inhibitor rucaparib is FDA approved to treat patients with metastatic castration-resistant prostate cancer (mCRPC) and deleterious somatic or germline BRCA mutations. Rucaparib is also approved as a maintenance treatment of patients with recurrent epithelial ovarian, Fallopian tube, or primary peritoneal cancer. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in ovarian cancer^{70,245}, FANCL loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in prostate^{80,174,210,1058-1059}, loss or inactivation of genes that are involved in homologous recombination repair may confer sensitivity to PARP inhibitors. CHEK1 inhibition has been associated with sensitivity to PARPi in preclinical studies^{224,1060}. On the basis of preclinical evidence^{71,74} and clinical evidence in ovarian cancer⁷⁰, BARD1 loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of strong clinical evidence in ovarian cancer^{85-86,996}, as well as clinical data in other cancer types^{86,1094-1095}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to rucaparib. On the basis of clinical evidence in ovarian^{85,245,1056}, breast¹⁷⁵, endometrial¹⁰⁵⁷, and prostate cancer¹⁷⁴, loss or inactivation of BRIP1 may confer sensitivity to PARP inhibitors. On the basis of limited clinical benefit in prostate^{209-210,1062} and ovarian¹⁰⁶³ cancer, CDK12 inactivating alterations may confer sensitivity to PARP inhibitors.

SUPPORTING DATA

The Phase 3 TRITON3 study of rucaparib for patients with chemotherapy-naïve metastatic castration-resistant prostate cancer (mCRPC) and BRCA alterations reported an improved radiographic PFS (10.2 vs. 6.4 months, HR=0.61) and median OS (mOS; 23.6 vs. 20.9 months, HR=0.94) compared to physician's choice docetaxel or second-generation androgen receptor pathway inhibitors (ARPIs). Patients with chemotherapy-naïve mCRPC and ATM alterations reported slightly improved radiographic PFS (8.1 vs. 6.8 months, HR=0.95) compared to physician's choice docetaxel or second-generation ARPI¹⁰⁹⁶. The Phase 2 TRITON2 study of rucaparib for patients with mCRPC and deleterious DNA-repair gene alterations reported an ORR of 44% (27/62; 7 CRs) and radiographic PFS of 9.0 months¹⁰⁹⁷. Objective responses were reported for patients with ATM, BRIP1, CHEK2, FANCA, PALB2, and RAD51B alterations¹⁷⁴. In a real-world study of olaparib and/or rucaparib for heavily pretreated prostate cancer, patients with pathogenic BRCA2 mutations experienced greater benefit than patients with other homologous recombination repair mutations (median PFS 7.2 vs. 2.8 months, p=0.291; PSA30 69.2% vs. 4.0%, p<0.001)¹⁰⁶⁶. The Phase 1b/2 BrUOG360 study of rucaparib combined with copanlisib to treat patients with mCRPC achieved a confirmed prostate-specific antigen (PSA) ≥50% decline for 2 patients (22% [2/9]), 1 of whom had a BRCA2 loss and 1 of whom had a PALB2 alteration¹⁰⁹⁸. A Phase 1b study of rucaparib combined with ipatasertib for patients with mCRPC reported a PSA ≥50% decline rate of 35% (9/26) and an mOS of 13.3 months¹⁰⁹⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Atezolizumab

Assay findings association

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Atezolizumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC) as well as adult and pediatric patients 2 years and older with alveolar soft part sarcoma, depending on treatment setting. Atezolizumab is also approved in combination with other therapies to treat patients with non-squamous NSCLC lacking EGFR or ALK alterations, small cell lung cancer, hepatocellular carcinoma, and BRAF V600-positive melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of emerging clinical data showing efficacy of atezolizumab alone or in combination with antiangiogenic therapy for patients with MSI-H colorectal cancer¹¹⁰⁰ or endometrial cancer¹¹⁰¹, MSI-H status may predict sensitivity to atezolizumab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

A Phase 1 study evaluating atezolizumab monotherapy for patients with heavily pretreated metastatic castration-resistant prostate cancer (mCRPC) reported limited

efficacy, with only 4% (1/25) of patients achieving PR and 8.6% (3/35) of patients with a confirmed prostate-specific antigen (PSA) response, a PFS of 2.7 months, and median OS (mOS) of 14.7 months¹¹⁰². The Phase 3 IMbassador250 study evaluating atezolizumab and enzalutamide versus enzalutamide alone failed to meet its primary endpoint of OS (HR=1.12) for patients with mCRPC previously treated with abiraterone; no differences were observed for radiographic PFS (HR=0.90) or time to PSA progression (HR=1.04) between treatment groups, though an increase in responders was observed in the atezolizumab arm (12% vs. 6.7% PRs)¹¹⁰³. The COSMIC-021 Phase 1b trial of atezolizumab plus cabozantinib for patients with mCRPC reported an ORR of 23% (3 CRs, 28 PRs), DCR of 84%, mOS of 18.4 months, and median PFS of 5.5 months¹¹⁰⁴. Another Phase 1 study investigating combination treatment of atezolizumab and the IDO1-inhibitor navoximod observed 1 PR and 2 PDs for patients with prostate cancer¹¹⁰⁵. In the prospective Phase 2a MyPathway basket study evaluating atezolizumab for patients with TMB-High solid tumors, patients with TMB ≥ 16 Muts/Mb achieved improved ORR (38% [16/42] vs. 2.1% [1/48]), DCR (62% [26/42] vs. 23% [11/48]), mPFS (5.7 vs. 1.8 months, HR 0.34), and mOS (19.8 vs. 11.4, HR 0.53) as compared to those with TMB ≥ 10 and <16 Muts/Mb⁴⁶. In a retrospective analysis of patients with 17 solid tumor types (comprised of 47% NSCLC, 40% urothelial carcinoma, and 13% encompassing 15 other solid tumors), TMB of 16 Muts/Mb or greater was reported to be associated with an improved ORR to atezolizumab compared to chemotherapy (30% vs. 14%)⁴⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Avelumab

Assay findings association

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Avelumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 in order to enhance antitumor immune responses. It is FDA approved to treat patients 12 years and older with Merkel cell carcinoma, or for urothelial carcinoma in various treatment settings. The combination of avelumab and axitinib is FDA approved for patients with renal cell carcinoma (RCC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of emerging clinical data in patients with MSI-H colorectal cancer¹¹⁰⁰, endometrial cancer¹¹⁰¹, or gastric/gastroesophageal junction cancer¹¹⁰⁶, MSI-H status may predict sensitivity to anti-PD-L1 therapies such as avelumab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

A Phase 2 trial of avelumab in combination with radiotherapy for patients with metastatic castration-resistant prostate cancer (mCRPC) reported an ORR of 31% (5/16) with median PFS and median OS of 8.4 months and 14.1 months, respectively¹¹⁰⁷. A trial of avelumab in combination with androgen receptor antagonists for patients with mCRPC who had progressed on previous treatment reported durable SD >24 weeks for 39% (7/18) of patients¹¹⁰⁸. The JAVELIN Phase 1b study has demonstrated clinical benefit from single-agent avelumab in a variety of solid tumor types, including non-small cell lung carcinoma (NSCLC)¹¹⁰⁹, gastric carcinoma and gastroesophageal junction (GEJ) adenocarcinoma¹¹¹⁰, urothelial carcinoma¹¹¹¹, mesothelioma¹¹¹², ovarian carcinoma¹¹¹³, and breast cancer¹¹¹⁴, and from avelumab combined with axitinib in renal cell carcinoma¹¹¹⁵. Emerging clinical data show a positive trend toward the association of tumor cell PD-L1 expression and improved ORR, PFS, or OS in NSCLC in the first-line setting and in ovarian and breast cancer^{1109,1113-1114}. Limited clinical data indicate activity of avelumab in adrenocortical carcinoma, metastatic castration-resistant prostate cancer, and thymic cancer¹¹¹⁶⁻¹¹¹⁸.

Cemiplimab

Assay findings association

CDK12

R1048*, N474fs*8 - subclonal

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Cemiplimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with the ligands PD-L1 and PD-L2 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC), cutaneous squamous cell carcinoma, or basal cell carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of limited clinical evidence in prostate cancer, including multiple prostate-specific antigen (PSA) responses, CDK12 inactivating alterations may predict sensitivity to PD-1 inhibitors^{194,1049-1050}. On the basis of prospective clinical data showing efficacy of anti-PD-1 therapies against various MSI-high (MSI-H) solid tumors^{3,6-7,1080-1083}, MSI-H status may predict sensitivity to cemiplimab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An

association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

Clinical data on the efficacy of cemiplimab for the treatment of prostate carcinoma are limited (PubMed, Mar 2023). Cemiplimab has been studied primarily in advanced cutaneous squamous cell carcinoma (CSCC), where it elicited a combined ORR of 48% (41/85) in Phase 1 and 2 studies¹¹¹⁹. A Phase 2 trial of cemiplimab in patients with basal cell carcinoma (BCC) reported ORRs of 31% (5 CRs and 21 PRs) in patients with locally advanced BCC and 21% (6 PRs) in patients with metastatic BCC¹¹²⁰⁻¹¹²¹. The Phase 3 EMPOWER-Lung 1 trial for advanced non-small cell lung cancer (NSCLC) with PD-L1 expression $\geq 50\%$ reported that cemiplimab is associated with improved PFS (8.2 vs. 5.7 months), OS (not reached vs. 14.2 months), and ORR (37% vs. 21%) compared with chemotherapy¹¹²².

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Durvalumab

Assay findings association
Microsatellite status
MSI-High

Tumor Mutational Burden
350 Muts/Mb

AREAS OF THERAPEUTIC USE

Durvalumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses. It is FDA approved to treat patients with non-small cell lung cancer (NSCLC), small cell lung cancer (SCLC), and biliary tract cancer. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of emerging clinical data in patients with MSI-H colorectal cancer¹¹⁰⁰, endometrial cancer¹¹⁰¹, or gastric/gastroesophageal junction cancer¹¹⁰⁶, MSI-H status may predict sensitivity to anti-PD-L1 therapies such as durvalumab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

A randomized Phase 2 study of durvalumab for patients with metastatic castration-resistant prostate cancer (mCRPC) reported an ORR of 0% (0/13)¹¹²³. Another Phase 2 study of durvalumab in combination with the PARP inhibitor olaparib reported prostate-specific antigen decline $\geq 50\%$ for 53% (9/17) of patients with mCRPC who had progressed on enzalutamide and/or abiraterone; 24% (4/17) of the patients experienced radiographic responses, median radiographic PFS was 16.1 months, and 12-month PFS probability for patients with or without mutations in DNA damage repair genes was 83% and 36%, respectively ($p=0.03$)¹⁰⁶⁸. A Phase 2 study for patients with mCRPC reported a low response rate of 3.4% (1/29), median radiographic PFS of 2.3 months, and median OS of 10.7 months with treatment of durvalumab combined with adenosine 2A receptor inhibitor AZD4635¹¹²⁴. In a case report, a patient with prostate sarcoma achieved a PR on durvalumab¹¹²⁵. A case study reported no responses for 2 patients with prostate neuroendocrine tumor treated with durvalumab combined with cabazitaxel¹¹²⁶.

Durvalumab + Tremelimumab

Assay findings association
Microsatellite status
MSI-High

AREAS OF THERAPEUTIC USE

Durvalumab is a monoclonal antibody that binds to PD-L1 and blocks its interaction with PD-1 to enhance antitumor immune responses; tremelimumab is a cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4)-blocking antibody. These therapies are FDA approved in combination to treat adult patients with unresectable hepatocellular carcinoma and metastatic non-small cell lung cancer. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{10-11,1127-1128}, microsatellite instability high (MSI-H) status may predict sensitivity to combination durvalumab and

tremelimumab.

SUPPORTING DATA

A randomized Phase 2 study of durvalumab in combination with tremelimumab for patients with metastatic castration-resistant prostate cancer (mCRPC) reported an ORR of 16% (6/37), with 38% (5/13) of PD-L1+ ($\geq 1\%$) and 5% (1/19) of PD-L1- disease responding¹¹²³. Another Phase 2 study of this therapy combination reported that 12% (3/25) and 4% (1/25) of patients with mCRPC who were chemotherapy-naïve experienced prostate-specific antigen decline $\geq 50\%$ and $\geq 90\%$, respectively; median radiographic PFS was 3.7 months, and median OS was 28.1 months¹¹²⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Erlotinib

Assay findings association
EGFR
G724S

AREAS OF THERAPEUTIC USE

Erlotinib is a small-molecule inhibitor of EGFR. It is FDA approved as a monotherapy or in combination with ramucirumab for patients with metastatic non-small cell lung cancer (NSCLC) harboring EGFR exon 19 deletions or exon 21 (L858R) mutations. Erlotinib is also FDA approved in combination with gemcitabine as a first-line treatment for advanced pancreatic cancer. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Amplification or activation of EGFR may predict sensitivity to therapies such as erlotinib. For patients with activating mutations in EGFR, treatment with erlotinib has been associated with improved response and lengthened time to progression^{262,1130-1132}. For patients harboring co-occurring EGFR activating alterations, G724S was found to be associated with acquired resistance to first-generation EGFR TKIs in a case series²⁸⁷, although 2 case studies reported SD to such inhibitors²⁸⁸⁻²⁸⁹. First-generation TKIs are therefore

unlikely to benefit patients with G724S and concurrent activating EGFR mutations. Limited preclinical evidence suggests that G724S in the absence of another activating EGFR alteration may reduce sensitivity to erlotinib²⁸⁵ and gefitinib²⁸¹.

SUPPORTING DATA

In the MyPathway Phase 2a basket study for advanced solid tumors, 1 of 9 patients with EGFR activating mutations responded to erlotinib monotherapy; the responding patient had urethral adenocarcinoma¹¹³³. A patient with EGFR-mutated metastatic lacrimal gland adenoid cystic carcinoma experienced clinical benefit from erlotinib treatment that was ongoing at 14 months¹¹³⁴. A Phase 2 trial of erlotinib in patients with metastatic prostate cancer reported clinical benefit in 40% of patients, though no decreases in PSA upon treatment were seen in any of 30 patients¹¹³⁵. Another Phase 2 trial of erlotinib in castration-resistant prostate cancer reported a clinical benefit in 31% of 22 patients and a strong decrease in PSA levels in 2 patients¹¹³⁶.

Gefitinib

Assay findings association
EGFR
G724S

AREAS OF THERAPEUTIC USE

Gefitinib targets the tyrosine kinase EGFR and is FDA approved to treat non-small cell lung cancer (NSCLC) harboring exon 19 deletions or exon 21 (L858R) substitution mutations in EGFR. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Activation of EGFR may predict sensitivity to therapies such as gefitinib. Clinical studies have consistently shown significant improvement in response rates and PFS for patients with EGFR-mutated non-small cell lung cancer (NSCLC) treated with gefitinib compared with chemotherapy^{1132,1137-1142}, and responses have been reported for patients with EGFR-rearranged NSCLC¹¹⁴³⁻¹¹⁴⁴. For patients harboring co-occurring EGFR activating alterations, G724S was found to be associated with acquired resistance to first-generation EGFR TKIs in a case series²⁸⁷, although 2 case studies reported SD to such inhibitors²⁸⁸⁻²⁸⁹. First-generation TKIs are therefore

unlikely to benefit patients with G724S and concurrent activating EGFR mutations. Limited preclinical evidence suggests that G724S in the absence of another activating EGFR alteration may reduce sensitivity to erlotinib²⁸⁵ and gefitinib²⁸¹.

SUPPORTING DATA

A Phase 2 trial of gefitinib in patients with hormone-naïve prostate cancer did not meet its primary objective in PSA stabilization or reduction¹¹⁴⁵. In patients with castration-resistant prostate cancer, the results of a Phase 2 trial of gefitinib as a single agent were not promising, with a response rate of just 2% and toxicities reported¹¹⁴⁶. A Phase 1/2 trial in 42 patients with non-metastatic prostate cancer of gefitinib with concurrent radiotherapy reported clinical activity and an overall survival rate of 87%¹¹⁴⁷. Another Phase 2 trial did not report much increased clinical activity upon addition of gefitinib to prednisone treatment in hormone-refractory prostate cancer¹¹⁴⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Niraparib

Assay findings association
BARD1
R406*

BRCA2
D946fs*14

BRIP1
splice site 918+1G>A - subclonal

CDK12
R1048*, N474fs*8 - subclonal

CHEK1
T226fs*14

FANCL
G119*

AREAS OF THERAPEUTIC USE

The PARP inhibitor niraparib is FDA approved to treat patients with epithelial ovarian, fallopian tube, or primary peritoneal cancer, with or without homologous recombination deficiency (HRD)-positive status. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in ovarian^{85,245,1056}, breast¹⁷⁵, endometrial¹⁰⁵⁷, and prostate cancer¹⁷⁴, loss or inactivation of BRIP1 may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in ovarian and breast cancers^{83-84,1149}, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to PARP inhibitors such as niraparib. On the basis of clinical evidence in ovarian cancer^{70,245}, FANCL loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in prostate^{80,174,210,1058-1059}, loss or inactivation of genes that are involved in homologous recombination repair may confer sensitivity

to PARP inhibitors. CHEK1 inhibition has been associated with sensitivity to PARPi in preclinical studies^{224,1060}. On the basis of preclinical evidence^{71,74} and clinical evidence in ovarian cancer⁷⁰, BARD1 loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of limited clinical benefit in prostate^{209-210,1062} and ovarian¹⁰⁶³ cancer, CDK12 inactivating alterations may confer sensitivity to PARP inhibitors.

SUPPORTING DATA

The Phase 2 GALAHAD study of niraparib for patients with metastatic castration-resistant prostate cancer who had progressed on at least 1 line of AR-targeted therapy in addition to at least 1 line of taxane chemotherapy reported an ORR of 34% (26/76) and a median radiographic PFS of 5.5 months for patients with biallelic BRCA1 or BRCA2 alterations¹¹⁵⁰. Patients in this trial with biallelic alterations in non-BRCA1/2 DNA repair genes experienced an 11% (5/47) ORR¹¹⁵⁰.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Nivolumab

Assay findings association

CDK12

R1048*, N474fs*8 - subclonal

Microsatellite status

MSI-High

Tumor Mutational Burden

350 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response. It is FDA approved as a monotherapy in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), head and neck squamous cell carcinoma (HNSCC), urothelial carcinoma, colorectal cancer (CRC), classical Hodgkin lymphoma (cHL), gastric cancer, gastroesophageal junction cancer, or esophageal adenocarcinoma or squamous cell carcinoma (ESCC). It is also approved in combination with chemotherapy to treat ESCC, in combination with cabozantinib to treat RCC, and in combination with relatlimab to treat melanoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of limited clinical evidence in prostate cancer, including multiple prostate-specific antigen (PSA) responses, CDK12 inactivating alterations may predict sensitivity to PD-1 inhibitors^{194,1049-1050}. On the basis of prospective clinical data showing efficacy of nivolumab for patients with MSI-H CRC^{3,6}, MSI-H status may predict sensitivity to nivolumab. On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵.

SUPPORTING DATA

Patients with CDK12-mutated metastatic castration-

resistant prostate cancer (mCRPC) treated with nivolumab combined with the anti-CTLA-4 immunotherapy ipilimumab experienced a composite response rate of 14% (1/7) in the NEPTUNES trial²⁰¹ and a prostate-specific antigen (PSA) $\geq 50\%$ decline in 14% (4/28) of patients in the IMPACT trial; unconfirmed PSA $\geq 30\%$ decline was additionally observed in 21% (6/28) of patients²⁰². PSA response rates of approximately 11-50% (2/19-2/4) were observed in retrospective studies of patients with CDK12-mutated metastatic prostate cancer treated with immune checkpoint inhibitors, which were predominantly anti-PD-1 monotherapies, such as nivolumab^{194,198-200}. A Phase 2 study of nivolumab and ipilimumab combination therapy for patients with metastatic castration-resistant prostate cancer (mCRPC) reported ORRs of 25% (8/32; 2 CRs) and 10% (3/30; 2 CRs) for chemotherapy-naïve patients who had progressed on androgen deprivation therapy and patients who had progressed on chemotherapy, respectively⁴². Patients with tumor mutational burden (TMB) values above the study median achieved significantly better outcomes compared with patients with lower TMB values in the combined cohorts (ORR of 50% [9/18] vs. 5.3% [1/19], median PFS of 7.4 vs. 2.4 months)⁴². The CheckMate 9KD Phase 2 trial for patients with mCRPC reported ORRs of 37% (7/19; 1 CR, 6 PRs) and a median radiographic PFS of 8.2 months following combination treatment of nivolumab and docetaxel¹¹²². Arm A2 of the same trial observed an ORR of 15% (6/39; all PRs) and median OS (mOS) of 20.2 months following combination treatment of nivolumab and rucaparib, with higher responses observed in a subset of patients with HRD+ tumors (ORR 25% [5/20]; all PRs, mOS of 22.7 months)¹¹⁵¹. A Phase 1 study of nivolumab monotherapy did not report any objective responses for the 17 patients with genomically unselected CRPC¹¹⁵².

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Nivolumab + Ipilimumab

Assay findings association
Microsatellite status
MSI-High

Tumor Mutational Burden
350 Muts/Mb

AREAS OF THERAPEUTIC USE

Nivolumab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor immune response, and ipilimumab is a cytotoxic T-lymphocyte antigen 4 (CTLA-4)-blocking antibody. The combination is FDA approved in various treatment settings for patients with melanoma, renal cell carcinoma (RCC), non-small cell lung cancer (NSCLC), hepatocellular carcinoma (HCC), pleural mesothelioma, and esophageal squamous cell carcinoma (ESCC). Furthermore, nivolumab is approved in combination with ipilimumab to treat patients with mismatch repair-deficient (dMMR) or microsatellite instability-high (MSI-H) colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{37-38,1153}, a TMB score of ≥ 10 Muts/Mb (as measured by this assay) may predict sensitivity to combination nivolumab and

ipilimumab treatment. On the basis of clinical data across solid tumors^{3-5,1154-1158}, microsatellite instability high (MSI-H) status may predict sensitivity to combination nivolumab and ipilimumab.

SUPPORTING DATA

Preliminary results from a Phase 2 study of nivolumab and ipilimumab combination therapy in metastatic castration-resistant prostate cancer (mCRPC) reported ORRs of 25% (8/32, 2 CRs) and 10% (3/30, 2 CRs) for patients who were chemotherapy-naïve and had progressed on androgen deprivation therapy and for patients who had progressed on chemotherapy, respectively⁴². Within the same trial, patients with post-chemotherapy mCRPC reported ORRs of 9-20% (4/43 and 8/41) for cohorts treated with nivolumab and ipilimumab regimens, 5% (1/22) for those receiving ipilimumab alone, and 12% (5/41) for cohorts receiving cabazitaxel; CRs were only observed in cohorts receiving the combination treatment¹¹⁵⁹.

Retifanlimab

Assay findings association
CDK12
R1048*, N474fs*8 - subclonal

Microsatellite status
MSI-High

Tumor Mutational Burden
350 Muts/Mb

AREAS OF THERAPEUTIC USE

Retifanlimab is a monoclonal antibody that binds to the PD-1 receptor and blocks its interaction with PD-L1 and PD-L2, reducing inhibition of the antitumor response. It is FDA approved to treat patients with Merkel cell carcinoma. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical data across solid tumors^{34-36,49,1048}, TMB of ≥ 10 Muts/Mb (based on this assay or others) may predict sensitivity to immune checkpoint inhibitors targeting PD-1 or PD-L1. An association between higher TMB and improved OS, median PFS, and ORR has been observed in large pan-solid tumor studies for patients treated with immune checkpoint inhibitors³⁴⁻³⁵. On the basis of limited clinical evidence in prostate cancer, including multiple prostate-specific antigen (PSA) responses, CDK12 inactivating alterations may predict sensitivity to PD-1 inhibitors^{194,1049-1050}. On the basis of prospective clinical data showing efficacy of anti-PD-1 therapies against various MSI-high (MSI-H) solid tumors^{3,6-7,1080-1083,1160}, MSI-H status may predict sensitivity to retifanlimab.

SUPPORTING DATA

Clinical data on the efficacy of retifanlimab for the treatment of prostate cancer are limited (PubMed, Mar 2023). The efficacy of retifanlimab has been demonstrated

in various treatment settings for multiple advanced solid tumors, including Merkel cell carcinoma¹¹⁶¹, anal squamous cell carcinoma (SCC)¹¹⁶², microsatellite instability-high or deficient MMR endometrial carcinoma¹¹⁶⁰, glioblastoma¹¹⁶³, soft tissue sarcoma¹¹⁶⁴, and gastroesophageal adenocarcinoma¹¹⁶⁵. The Phase 2 POD1UM-201 trial of retifanlimab for patients with chemotherapy-naïve advanced Merkel cell carcinoma reported an ORR of 51% (33/65; 11 CRs, 22 PRs), unreached median duration of response, and median PFS (mPFS) of 13.8 months¹¹⁶¹. In the Phase 2 POD1UM-202 study for patients with previously treated advanced squamous carcinoma of the anal canal, retifanlimab elicited an ORR of 14% (13/94; 1 CR, 12 PRs), mPFS of 2.3 months, and median OS of 10.1 months, with responses observed regardless of PD-L1 expression¹¹⁶². In the Phase 2 POD1UM-203 trial for multiple tumor types, retifanlimab yielded an ORR of 35% (8/23) and mPFS of 4.4 months for patients with treatment-naïve metastatic non-small cell lung cancer (NSCLC) with high PD-L1 expression (TPS $\geq 50\%$), an ORR of 40% (14/35) and mPFS of 3.6 months for patients with unresectable or metastatic melanoma, an ORR of 38% (11/29) and mPFS of 5.7 months for patients with cisplatin-ineligible locally advanced or metastatic urothelial carcinoma with PD-L1 expression (CPS $\geq 10\%$), and an ORR of 24% (8/34; 1 CR, 7 PRs) and mPFS of 5.4 months for patients with treatment-naïve advanced clear cell renal cell carcinoma (RCC)¹¹⁶⁶.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

THERAPIES WITH CLINICAL BENEFIT

IN OTHER TUMOR TYPE

Talazoparib

Assay findings association
BARD1
R406*

BRCA2
D946fs*14

BRIP1
splice site 918+1G>A - subclonal

CDK12
R1048*, N474fs*8 - subclonal

CHEK1
T226fs*14

FANCL
G119*

AREAS OF THERAPEUTIC USE

The PARP inhibitor talazoparib is FDA approved to treat HER2-negative locally advanced or metastatic breast cancer with deleterious or suspected deleterious germline BRCA mutations. Please see the drug label for full prescribing information.

GENE ASSOCIATION

On the basis of clinical evidence in ovarian^{85,245,1056}, breast¹⁷⁵, endometrial¹⁰⁵⁷, and prostate cancer¹⁷⁴, loss or inactivation of BRIP1 may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in ovarian cancer^{70,245}, FANCL loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of clinical evidence in prostate^{80,174,210,1058-1059}, loss or inactivation of genes that are involved in homologous recombination repair may confer sensitivity to PARP inhibitors. CHEK1 inhibition has been associated with sensitivity to PARPi in preclinical studies^{224,1060}. On the basis of preclinical evidence^{71,74} and clinical evidence in ovarian cancer⁷⁰, BARD1 loss or inactivation may confer sensitivity to PARP inhibitors. On the basis of limited clinical benefit in prostate^{209-210,1062} and ovarian¹⁰⁶³ cancer, CDK12 inactivating alterations may confer sensitivity to PARP inhibitors. On the basis of strong clinical data in breast cancer¹¹⁶⁷⁻¹¹⁶⁹ and additional clinical evidence in ovarian, pancreatic, and prostate cancer¹¹⁷⁰⁻¹¹⁷³, loss or inactivation of either BRCA1 or BRCA2 may confer sensitivity to

talazoparib.

SUPPORTING DATA

The Phase 3 TALAPRO-2 trial of talazoparib plus enzalutamide as a first-line treatment for patients with metastatic castration-resistant prostate cancer (mCRPC) reported a significantly improved radiographic PFS (rPFS) (NR vs. 21.9 months, HR=0.63) and ORR (62% vs. 44%), but OS (36.4 months vs. NR, HR=0.89) was not improved compared with placebo plus enzalutamide; greater rPFS benefit was observed among patients with homologous recombination repair (HRR) mutations (27.9 vs. 16.4 months, HR=0.46) than those without HRR mutations or of unknown HRR status (NR vs. 22.5 months, HR=0.70)¹¹⁷⁴. The Phase 2 TALAPRO-1 trial of talazoparib monotherapy for patients with docetaxel-treated mCRPC harboring alterations in DNA repair genes reported a study-wide ORR of 30% with median rPFS of 5.6 months, with an ORR of 46% (28/61) and rPFS of 11.2 months for patients with BRCA1/2 mutations¹¹⁷⁵. A retrospective subgroup analysis found no association between antitumor activity and germline HRR alterations (gHRRm) (ORR: 31% [5/16] vs. 26% [14/54] in patients assigned male at birth with vs. without gHRRm, respectively); ORRs were also similar for patients with gHRRm or with only somatic HRRm (25% [10/40] vs. 19% [4/21], respectively, [p=0.7528])¹¹⁷⁶.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

NOTE Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually

updated and should be investigated by the physician or research staff. This is not a comprehensive list of all available clinical trials. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials listed here may have additional enrollment criteria that

may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see [clinicaltrials.gov](https://www.foundationmedicine.com/genomic-testing#support-services). Or, visit <https://www.foundationmedicine.com/genomic-testing#support-services>.

BIOMARKER

Microsatellite status

RESULT

MSI-High

RATIONALE

High microsatellite instability (MSI) may predict response to anti-PD-1 and anti-PD-L1 immune checkpoint inhibitors (alone or in combination with anti-CTLA-4).

NCT04237649
PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS
 ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Sunto Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT04100018
PHASE 3

A Study of Nivolumab or Placebo in Combination With Docetaxel in Men With Advanced Castration-resistant Prostate Cancer

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Fuzhou (China), Tainan (Taiwan), Niasong (Taiwan), Hangzhou (China), Shanghai (China), Nanchang (China), Nanchang Shi (China), Hong Kong (Hong Kong)

NCT02628067
PHASE 2

Study of Pembrolizumab (MK-3475) in Participants With Advanced Solid Tumors (MK-3475-158/KEYNOTE-158)

TARGETS
 PD-1

LOCATIONS: Taipei (Taiwan), Makati (Philippines), Seoul (Korea, Republic of), North Ryde (Australia), Moscow (Russian Federation), Hod Hasharon (Israel), Drammen (Norway), Glostrup (Denmark), Haar (Germany), Haarlem (Netherlands)

NCT02861573
PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS
 AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04446117
PHASE 3

Study of Cabozantinib in Combination With Atezolizumab Versus Second NHT in Subjects With mCRPC

TARGETS
 AR, PD-L1, MET, ROS1, RET, VEGFRs, CYP17

LOCATIONS: Taoyuan (Taiwan), Taichung (Taiwan), Taichung City (Taiwan), Kaohsiung (Taiwan), Taipei city (Taiwan), Tainan City (Taiwan), Tainan (Taiwan), Hwasun (Korea, Republic of), Gwangju (Korea, Republic of), Fukuoka (Japan)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT04047862
PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS
PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Hefei (China), Guangdong (China), Changsha (China)

NCT05166577
PHASE 1/2

Nanatinostat Plus Valganciclovir in Patients With Advanced EBV+ Solid Tumors, and in Combination With Pembrolizumab in EBV+ RM-NPC

TARGETS
HDAC, PD-1

LOCATIONS: Taipei City (Taiwan), Taipei (Taiwan), Taoyuan City (Taiwan), Sha Tin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Kuching (Malaysia), Kuala Lumpur (Malaysia), Singapore (Singapore), Blacktown (Australia)

NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Melbourne (Australia), Amsterdam (Netherlands), Ravenna (Italy), Meldola (Italy)

NCT03821935
PHASE 1

Study to Determine the Safety, Tolerability, Pharmacokinetics and Recommended Phase 2 Dose (RP2D) of ABBV-151 as a Single Agent and in Combination With ABBV-181 in Participants With Locally Advanced or Metastatic Solid Tumors

TARGETS
PD-1, GARP

LOCATIONS: Taichung City (Taiwan), Taipei City (Taiwan), Seoul (Korea, Republic of), Chuo-ku (Japan), Kashiwa-shi (Japan), South Brisbane (Australia), Camperdown (Australia), Ramat Gan (Israel), Tel Aviv-Yafo (Israel), Haifa (Israel)

NCT04282018
PHASE 1/2

Brief Title: Study of BGB-10188 as Monotherapy, and in Combination With Zanubrutinib, and Tislelizumab

TARGETS
PI3K-delta, PD-1, BTK

LOCATIONS: Fuzhou (China), Zhejiang (China), Shanghai (China), Suzhou (China), Changsha (China), Jining (China), Chengdu (China), West Perth (Australia), Adelaide (Australia), Blacktown (Australia)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
BIOMARKER

Tumor Mutational Burden

RESULT

350 Muts/Mb

RATIONALE

Increased tumor mutational burden may predict response to anti-PD-1 (alone or in combination with anti-CTLA-4) or anti-PD-L1 immune checkpoint inhibitors.

NCT04237649
PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS
ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Sunto Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT04100018
PHASE 3

A Study of Nivolumab or Placebo in Combination With Docetaxel in Men With Advanced Castration-resistant Prostate Cancer

TARGETS
PD-1

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Fuzhou (China), Tainan (Taiwan), Niasong (Taiwan), Hangzhou (China), Shanghai (China), Nanchang (China), Nanchang Shi (China), Hong Kong (Hong Kong)

NCT02861573
PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS
AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS
TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha, RAFs, NRAS

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Shanghai City (China), Shanghai (China), Shatin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Seongnam-si (Korea, Republic of), Xi'an (China)

NCT04446117
PHASE 3

Study of Cabozantinib in Combination With Atezolizumab Versus Second NHT in Subjects With mCRPC

TARGETS
AR, PD-L1, MET, ROS1, RET, VEGFRs, CYP17

LOCATIONS: Taoyuan (Taiwan), Taichung (Taiwan), Taichung City (Taiwan), Kaohsiung (Taiwan), Taipei city (Taiwan), Tainan City (Taiwan), Tainan (Taiwan), Hwasun (Korea, Republic of), Gwangju (Korea, Republic of), Fukuoka (Japan)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT04047862
PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS
PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Hefei (China), Guangdong (China), Changsha (China)

NCT05166577
PHASE 1/2

Nanatinostat Plus Valganciclovir in Patients With Advanced EBV+ Solid Tumors, and in Combination With Pembrolizumab in EBV+ RM-NPC

TARGETS
HDAC, PD-1

LOCATIONS: Taipei City (Taiwan), Taipei (Taiwan), Taoyuan City (Taiwan), Sha Tin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Kuching (Malaysia), Kuala Lumpur (Malaysia), Singapore (Singapore), Blacktown (Australia)

NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Melbourne (Australia), Amsterdam (Netherlands), Ravenna (Italy), Meldola (Italy)

NCT03821935
PHASE 1

Study to Determine the Safety, Tolerability, Pharmacokinetics and Recommended Phase 2 Dose (RP2D) of ABBV-151 as a Single Agent and in Combination With ABBV-181 in Participants With Locally Advanced or Metastatic Solid Tumors

TARGETS
PD-1, GARP

LOCATIONS: Taichung City (Taiwan), Taipei City (Taiwan), Seoul (Korea, Republic of), Chuo-ku (Japan), Kashiwa-shi (Japan), South Brisbane (Australia), Camperdown (Australia), Ramat Gan (Israel), Tel Aviv-Yafo (Israel), Haifa (Israel)

NCT04215978
PHASE 1

Safety and Preliminary Effectiveness of BGB-A445 in Combination With Tislelizumab in Participants With Advanced Solid Tumors

TARGETS
PD-1, OX40

LOCATIONS: Changhua (Taiwan), Taipei (Taiwan), Tianan (Taiwan), Hangzhou (China), Shanghai (China), Changsha (China), Wuhan (China), Linyi (China), Gyeonggi-do (Korea, Republic of), Gyeongju (Korea, Republic of)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

APC

ALTERATION

R876*

RATIONALE

Based on preclinical and limited clinical data, APC inactivation may be associated with sensitivity to CBP/beta-catenin interaction inhibitors.

NCT04008797

PHASE 1

A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor

TARGETS

CBP, Beta-catenin, FGFRs, RET, PDGFRA, VEGFRs, KIT

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Kurume (Japan), Matsuyama (Japan), Seongnamsi Bundang (Korea, Republic of), Songpa-gu (Korea, Republic of), Seodaemun (Korea, Republic of), Seoul (Korea, Republic of), Jongno-gu (Korea, Republic of), Osakasayama (Japan)

NCT03264664

PHASE 1

Study of E7386 in Participants With Selected Advanced Neoplasms

TARGETS

CBP, Beta-catenin

LOCATIONS: Glasgow (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Sutton (United Kingdom), California, Arizona, Minnesota

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

BARD1

ALTERATION

R406*

RATIONALE

Tumors with BARD1 inactivating mutation or loss may be sensitive to PARP inhibitors.

NCT02861573
PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS

AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804
PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS

CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT05489211
PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS

TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT04434482
PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS

PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05405439
PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS

PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05223582
PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
CYP17, PARP

LOCATIONS: Shanghai (China)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05376722
PHASE 2

A Study of Pamiparib Combined With Abiraterone Acetate in Neoadjuvant Treatment of Prostate Cancer

TARGETS
CYP17

LOCATIONS: Nanjing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE
BRCA2

ALTERATION
D946fs*14

RATIONALE
BRCA2 loss or inactivating alterations may predict sensitivity to PARP inhibitors or to ATR inhibitors.

NCT02861573

PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS
AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804

PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS
CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT05489211

PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS
TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT04434482

PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS
PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05457257

PHASE 4

Clinical Study to Assess the Efficacy and Safety of Olaparib in Chinese Patients With Metastatic Castration-Resistant Prostate Cancer Who Have Failed Prior Treatment With a New Hormonal Agent and Have BRCA1/2 Mutations

TARGETS
CYP17, PARP, AR

LOCATIONS: Fuzhou (China), Ningbo (China), Hangzhou (China), Jiaxing (China), Shanghai (China), Nanchang (China), Suzhou (China), Wuxi (China), Nantong (China), Hefei (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05405439
PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS
 PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

NCT05223582
PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
 CYP17, PARP

LOCATIONS: Shanghai (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
 EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
 PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

BRIP1

ALTERATION

splice site 918+1G>A - subclonal

RATIONALE

BRIP1 inactivation may predict sensitivity to inhibitors of other DNA repair pathways, including inhibitors of PARP.

NCT02861573

PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS

AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804

PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS

CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT05489211

PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS

TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT04434482

PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS

PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05405439

PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS

PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05223582
PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
 CYP17, PARP

LOCATIONS: Shanghai (China)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
 PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05376722
PHASE 2

A Study of Pamiparib Combined With Abiraterone Acetate in Neoadjuvant Treatment of Prostate Cancer

TARGETS
 CYP17

LOCATIONS: Nanjing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

CDK12

ALTERATION

R1048*, N474fs*8 - subclonal

RATIONALE

Preclinical and clinical data suggest that tumors with CDK12 mutation or loss may be sensitive to PARP inhibitors. Preclinical and limited clinical

evidence indicate that CDK12 inactivation in prostate cancer may predict benefit from anti-PD-1 immune checkpoint inhibitors.

NCT04237649

PHASE NULL

KAZ954 Alone and With PDR001, NZV930 and NIR178 in Advanced Solid Tumors

TARGETS

ADORA2A, CD73, PD-1

LOCATIONS: Taipei (Taiwan), Shatin, New Territories (Hong Kong), Suntou Gun (Japan), Singapore (Singapore), Milano (Italy), Barcelona (Spain), California, Illinois, Toronto (Canada), Missouri

NCT04100018

PHASE 3

A Study of Nivolumab or Placebo in Combination With Docetaxel in Men With Advanced Castration-resistant Prostate Cancer

TARGETS

PD-1

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Fuzhou (China), Tainan (Taiwan), Niao-sung (Taiwan), Hangzhou (China), Shanghai (China), Nanchang (China), Nanchang Shi (China), Hong Kong (Hong Kong)

NCT02861573

PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS

AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804

PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS

CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT04047862

PHASE 1

Study of BGB-A1217 in Combination With Tislelizumab in Advanced Solid Tumors

TARGETS

PD-1, TIGIT

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Hualien City (Taiwan), Taichung (Taiwan), Fujian (China), Hangzhou (China), Shanghai (China), Hefei (China), Guangdong (China), Changsha (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05166577
PHASE 1/2

Nanatinostat Plus Valganciclovir in Patients With Advanced EBV+ Solid Tumors, and in Combination With Pembrolizumab in EBV+ RM-NPC

TARGETS
HDAC, PD-1

LOCATIONS: Taipei City (Taiwan), Taipei (Taiwan), Taoyuan City (Taiwan), Sha Tin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Kuching (Malaysia), Kuala Lumpur (Malaysia), Singapore (Singapore), Blacktown (Australia)

NCT05489211
PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS
TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT03530397
PHASE 1

A Study to Evaluate MEDI5752 in Subjects With Advanced Solid Tumors

TARGETS
PD-L1, PD-1, CTLA-4

LOCATIONS: Taipei (Taiwan), Tainan (Taiwan), Cheongju-si (Korea, Republic of), Incheon (Korea, Republic of), Seoul (Korea, Republic of), Gyeonggi-do (Korea, Republic of), Melbourne (Australia), Amsterdam (Netherlands), Ravenna (Italy), Meldola (Italy)

NCT04434482
PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS
PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT03821935
PHASE 1

Study to Determine the Safety, Tolerability, Pharmacokinetics and Recommended Phase 2 Dose (RP2D) of ABBV-151 as a Single Agent and in Combination With ABBV-181 in Participants With Locally Advanced or Metastatic Solid Tumors

TARGETS
PD-1, GARP

LOCATIONS: Taichung City (Taiwan), Taipei City (Taiwan), Seoul (Korea, Republic of), Chuo-ku (Japan), Kashiwa-shi (Japan), South Brisbane (Australia), Camperdown (Australia), Ramat Gan (Israel), Tel Aviv-Yafo (Israel), Haifa (Israel)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

CHEK1

ALTERATION

T226fs*14

RATIONALE

On the basis of limited clinical benefit in prostate cancer, CHEK1 alterations may confer sensitivity to PARP inhibitors.

NCT02861573
PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS

AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804
PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS

CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT05489211
PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS

TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT04434482
PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS

PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05405439
PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS

PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05223582
PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
CYP17, PARP

LOCATIONS: Shanghai (China)

NCT04123366
PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05376722
PHASE 2

A Study of Pamiparib Combined With Abiraterone Acetate in Neoadjuvant Treatment of Prostate Cancer

TARGETS
CYP17

LOCATIONS: Nanjing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
GENE
EGFR
ALTERATION
G724S
RATIONALE

EGFR activating mutations, rearrangements, or amplification may predict sensitivity to EGFR-targeted therapies. Strategies to overcome resistance to current agents include next-generation EGFR inhibitors and combination

therapies. On the basis of clinical data, EGFR G724S may reduce response to erlotinib, gefitinib, and osimertinib, as well as other third-generation EGFR TKIs.

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04946968
PHASE 2

Phase-2 Dacomitinib Study on Patients With EGFR-Driven Advanced Solid Tumours With Low EGFR-AS1 lncRNA Expr or Other Novel Emerging Biomarkers

TARGETS

ERBB4, EGFR, ERBB2

LOCATIONS: Singapore (Singapore)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT04720976
PHASE 1/2

JAB-3312 Activity in Adult Patients With Advanced Solid Tumors

TARGETS

MEK, SHP2, PD-1, EGFR, KRAS

LOCATIONS: Utah, California, Arizona, Minnesota, Illinois, Michigan, Oklahoma, Missouri, Indiana, Connecticut

NCT04670679
PHASE 1

A Dose Escalation/Expansion Study of ERAS-601 in Patients With Advanced or Metastatic Solid Tumors

TARGETS

SHP2, EGFR

LOCATIONS: Perth (Australia), Melbourne (Australia), Nevada, California, Missouri, Texas, Massachusetts, New York, Pennsylvania, Tennessee

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
GENE
FANCL
ALTERATION
G119*
RATIONALE

On the basis of clinical evidence in ovarian cancer, FANCL loss or inactivation may confer sensitivity to PARP inhibitors.

NCT02861573
PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS

AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804
PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS

CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT05489211
PHASE 2

Study of Dato-Dxd as Monotherapy and in Combination With Anti-cancer Agents in Patients With Advanced Solid Tumours (TROPION-PanTumor03)

TARGETS

TROP2, PD-L1, PARP1, PD-1

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Liou Ying Township (Taiwan), Shanghai (China), Seoul (Korea, Republic of), Seodaemun-gu (Korea, Republic of), Suita-shi (Japan), Chuo-ku (Japan), Koto-ku (Japan), Kashiwa (Japan)

NCT04434482
PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS

PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05405439
PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS

PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

NCT05223582

PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
CYP17, PARP

LOCATIONS: Shanghai (China)

NCT04123366

PHASE 2

Study of Olaparib (MK-7339) in Combination With Pembrolizumab (MK-3475) in the Treatment of Homologous Recombination Repair Mutation (HRRm) and/or Homologous Recombination Deficiency (HRD)-Positive Advanced Cancer (MK-7339-007/KEYLYNK-007)

TARGETS
PARP, PD-1

LOCATIONS: Fukuoka (Japan), Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Okayama (Japan), Nagoya (Japan), Tokyo (Japan), Kashiwa (Japan), Sapporo (Japan), Nedlands (Australia), Southport (Australia)

NCT03742895

PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

NCT02264678

PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05376722

PHASE 2

A Study of Pamiparib Combined With Abiraterone Acetate in Neoadjuvant Treatment of Prostate Cancer

TARGETS
CYP17

LOCATIONS: Nanjing (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
GENE
PIK3CA
ALTERATION
E970K
RATIONALE

PIK3CA activating mutations may lead to activation of the PI3K-AKT-mTOR pathway and may therefore indicate sensitivity to inhibitors of

this pathway. Strong clinical data support sensitivity of PIK3CA-mutated solid tumors to the PI3K-alpha inhibitor alpelisib.

NCT05348577
PHASE 3

Study of Capivasertib + Docetaxel vs Placebo + Docetaxel as Treatment for Metastatic Castration Resistant Prostate Cancer (mCRPC)

TARGETS
AKTs

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Kaohsiung (Taiwan), Ningbo (China), Hangzhou (China), Jiaxing (China), Shanghai (China), Nanchang (China), Shenzhen (China)

NCT04589845
PHASE 2

Tumor-Agnostic Precision Immuno-Oncology and Somatic Targeting Rational for You (TAPISTRY) Platform Study

TARGETS
TRKB, ALK, TRKC, ROS1, TRKA, RET, PD-L1, AKTs, ERBB2, MDM2, PI3K-alpha, RAFs, NRAS

LOCATIONS: Taipei City (Taiwan), Taoyuan County (Taiwan), Tainan (Taiwan), Shanghai City (China), Shanghai (China), Shatin (Hong Kong), Hong Kong (Hong Kong), Seoul (Korea, Republic of), Seongnam-si (Korea, Republic of), Xi'an (China)

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS
EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS
mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT04526470
PHASE 1/2

Alpelisib and Paclitaxel in PIK3CA-altered Gastric Cancer

TARGETS
PI3K-alpha

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS
mTOR

LOCATIONS: Tianjin (China)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT04551521
PHASE 2

CRAFT: The NCT-PMO-1602 Phase II Trial

TARGETS
PD-L1, AKTs, MEK, BRAF, ALK, RET, ERBB2

LOCATIONS: Lübeck (Germany), Würzburg (Germany), Mainz (Germany), Heidelberg (Germany), Tübingen (Germany)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT03385655
PHASE 2

Prostate Cancer Biomarker Enrichment and Treatment Selection

TARGETS
AR, WEE1, PD-L1, AKTs, PLK4, CTLA-4, MET

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), London (Canada), Hamilton (Canada), Halifax (Canada)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE PTEN

ALTERATION

P246L, K267fs*9, R173C

RATIONALE

PTEN loss or inactivating mutations may lead to increased activation of the PI3K-AKT-mTOR pathway and may indicate sensitivity to inhibitors

of this pathway. PTEN loss or inactivation may also predict sensitivity to PARP inhibitors.

NCT05348577

PHASE 3

Study of Capivasertib + Docetaxel vs Placebo + Docetaxel as Treatment for Metastatic Castration Resistant Prostate Cancer (mCRPC)

TARGETS
AKTs

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Kaohsiung (Taiwan), Ningbo (China), Hangzhou (China), Jiaxing (China), Shanghai (China), Nanchang (China), Shenzhen (China)

NCT02861573

PHASE 1/2

Study of Pembrolizumab (MK-3475) Combination Therapies in Metastatic Castration-Resistant Prostate Cancer (MK-3475-365/KEYNOTE-365)

TARGETS
AR, PD-1, PARP, CYP17

LOCATIONS: Taipei (Taiwan), North Ryde (Australia), Moscow (Russian Federation), Kiev (Ukraine), Espoo (Finland), Stockholm (Sweden), Istanbul (Turkey), Warsaw (Poland), Glostrup (Denmark), Auckland (New Zealand)

NCT04691804

PHASE 3

A Multicenter, Randomized, Double-Blind, Placebo-Controlled Phase III Study of Fuzuloparib Combined With Abiraterone Acetate and Prednisone (AA-P) Versus Placebo Combined With AA-P as First-Line Treatment in Patients With Metastatic Castration-Resistant Prostate Cancer

TARGETS
CYP17, PARP

LOCATIONS: Taipei (Taiwan), Taoyuan (Taiwan), Taichung (Taiwan), Changhua (Taiwan), Fuzhou (China), Tainan (Taiwan), Kaohsiung (Taiwan), Hangzhou (China), Jiaxing (China), Shanghai (China)

NCT04434482

PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS
PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT05405439

PHASE 1/2

To Evaluate the Efficacy of TQB3823 Combined With Abiraterone and Prednisone in Metastatic Castration-resistant Prostate Cancer Patientsprednisone Acetate Tablets in Patients With Metastatic Castration-resistant Prostate Cancer

TARGETS
PARP, CYP17

LOCATIONS: Wenzhou (China), Shanghai (China), Nanjing (China), Guangzhou (China), Qingyuan (China), Changsha (China), Jinan (China), Nanning (China), Chongqing (China), Xi'an (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT05223582
PHASE 2

Fluzoparib and Abiraterone in the preSurgery Treatment of Prostate Cancer: FAST Trial

TARGETS
CYP17, PARP

LOCATIONS: Shanghai (China)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
ATR, PARP, PD-L1

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

NCT05376722
PHASE 2

A Study of Pamiparib Combined With Abiraterone Acetate in Neoadjuvant Treatment of Prostate Cancer

TARGETS
CYP17

LOCATIONS: Nanjing (China)

NCT05406999
PHASE 2

Neoadjuvant Intense Endocrine Therapy for High Risk and Locally Advanced Prostate Cancer

TARGETS
PARP, CYP17

LOCATIONS: Nanjing (China)

NCT05021367
PHASE 1

A Clinical Study of TQB3823 in Patients With Advanced Malignant Tumor

TARGETS
PARP

LOCATIONS: Guangzhou (China)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS

GENE

RNF43

ALTERATION

G659fs*41, H683fs*17

RATIONALE

Based on preclinical evidence, tumors with loss or inactivation of RNF43 may be sensitive to inhibitors of the WNT signaling pathway.

NCT02521844

PHASE 1

A Study to Evaluate the Safety and Tolerability of ETC-1922159 in Advanced Solid Tumours

TARGETS
PORCN

LOCATIONS: Singapore (Singapore), Colorado, Texas, North Carolina

NCT03447470

PHASE 1

Study to Evaluate the Safety and Tolerability of RXC004 in Advanced Malignancies

TARGETS
PORCN

LOCATIONS: Newcastle (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
GENE
TSC2
ALTERATION
R57H
RATIONALE

Inactivating TSC2 alterations may lead to increased mTOR activation and predict sensitivity to mTOR inhibitors.

NCT03239015
PHASE 2

Efficacy and Safety of Targeted Precision Therapy in Refractory Tumor With Druggable Molecular Event

TARGETS

EGFR, ERBB4, ERBB2, PARP, mTOR, MET, ROS1, RET, VEGFRs, BRAF, CDK4, CDK6

LOCATIONS: Shanghai (China)

NCT04803318
PHASE 2

Trametinib Combined With Everolimus and Lenvatinib for Recurrent/Refractory Advanced Solid Tumors

TARGETS

mTOR, FGFRs, RET, PDGFRA, VEGFRs, KIT, MEK

LOCATIONS: Guangzhou (China)

NCT05125523
PHASE 1

A Study of Sirolimus for Injection (Albumin Bound) in Patients With Advanced Solid Tumors

TARGETS

mTOR

LOCATIONS: Tianjin (China)

NCT05103358
PHASE 2

Phase 2 Basket Trial of Nab-sirolimus in Patients With Malignant Solid Tumors With Pathogenic Alterations in TSC1 or TSC2 Genes (PRECISION 1)

TARGETS

mTOR

LOCATIONS: Hawaii, Washington, California, Utah

NCT02693535
PHASE 2

TAPUR: Testing the Use of Food and Drug Administration (FDA) Approved Drugs That Target a Specific Abnormality in a Tumor Gene in People With Advanced Stage Cancer

TARGETS

CDK4, CDK6, FLT3, VEGFRs, CSF1R, KIT, RET, mTOR, ERBB2, MEK, BRAF, PARP, PD-1, CTLA-4, PD-L1, TRKB, ALK, TRKC, ROS1, TRKA, FGFRs

LOCATIONS: Hawaii, Washington, Oregon, California

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

CLINICAL TRIALS
NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS

VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT05036226
PHASE 1/2

COAST Therapy in Advanced Solid Tumors and Prostate Cancer

TARGETS

DDR2, ABL, SRC, KIT, mTOR

LOCATIONS: South Carolina

NCT05661461
PHASE 1

Dose-escalation Study to Assess Safety and Pharmacokinetics of Nab-Sirolimus in Patients With Locally Advanced or Metastatic Solid Tumors and Moderate Liver Impairment

TARGETS

mTOR

LOCATIONS: Utah, Texas

NCT01582191
PHASE 1

A Phase 1 Trial of Vandetanib (a Multi-kinase Inhibitor of EGFR, VEGFR and RET Inhibitor) in Combination With Everolimus (an mTOR Inhibitor) in Advanced Cancer

TARGETS

mTOR, EGFR, SRC, RET, VEGFRs

LOCATIONS: Texas

NCT03203525
PHASE 1

Combination Chemotherapy and Bevacizumab With the NovoTTF-100L(P) System in Treating Participants With Advanced, Recurrent, or Refractory Hepatic Metastatic Cancer

TARGETS

VEGFA, mTOR

LOCATIONS: Texas

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX

Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

ABL1

NM_005157.4: c.3001G>A
(p.A1001T)
chr9:133760678 and
NM_005157.4: c.443G>A
(p.S148N)
chr9:133730377

ALK

NM_004304.4: c.4690G>A
(p.A1564T)
chr2:29416263 and
NM_004304.4: c.4306C>T
(p.R1436C)
chr2:29416647

AMER1 (FAM123B OR WTX)

NM_152424.3: c.1828G>A
(p.A610T)
chrX:63411339 and
NM_152424.3: c.1732C>T
(p.L578F)
chrX:63411435

APC

NM_000038.4: c.2876C>T
(p.S959F)
chr5:112174167 and
NM_000038.4:
c.5465_5467delinsACG
(p.V1822_F1823delinsDV)
chr5:112176756-112176758

AR

NM_000044.2: c.791G>A
(p.R264Q)
chrX:66765779

ARAF

NM_001654.3: c.229G>T
(p.A77S)
chrX:47424224 and
NM_001654.3: c.521G>A
(p.R174H)
chrX:47424713

ARFRP1

NM_003224.3: c.2T>A
(p.M1?)
chr20:62338442

ARID1A

NM_006015.4: c.4973G>A
(p.R1658Q)
chr1:27101691, NM_006015.4:
c.6694C>T (p.R2232W)
chr1:27107083 and
NM_006015.4: c.3869C>T
(p.T1290M)
chr1:27100073

ASXL1

NM_015338.5: c.3442C>T
(p.R1148C)
chr20:31023957

ATM

NM_000051.3: c.1411A>G
(p.N471D)
chr11:108121603,
NM_000051.3: c.7375C>T
(p.R2459C)
chr11:108201008,
NM_000051.3: c.7858G>A
(p.V2620I)
chr11:108203558 and
NM_000051.3: c.2746G>A
(p.V916M)
chr11:108139244

AURKA

NM_003600.2: c.251T>C
(p.V84A)
chr20:54961381

BARD1

NM_000465.2: c.268G>A
(p.A90T)
chr2:215657117

BCL6

NM_001706.4: c.996G>T
(p.Q332H)
chr3:187447197

BRCA2

NM_000059.3: c.2918C>T
(p.S973L)
chr13:32911410 and
NM_000059.3: c.7591G>A
(p.V2531I)
chr13:32930720

CCND1

NM_053056.2: c.733C>T
(p.R245W)
chr11:69465895

CCND2

NM_001759.3: c.445T>C
(p.W149R)
chr12:4387959

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance
CCNE1

 NM_001238.1: c.236A>G
(p.D79G)
chr19:30308099

CD22

 NM_001771.3: c.2029G>A
(p.V677I)
chr19:35832862

CD79B

 NM_000626.2: c.338G>A
(p.R113Q)
chr17:62007526

CDK12

 NM_016507.2: c.2456A>T
(p.D819V)
chr17:37657539 and
NM_016507.2: c.1753C>T
(p.P585S)
chr17:37627838

CDK8

 NM_001260.1: c.1328G>A
(p.S443N)
chr13:26978151

CDKN2A/B

 NM_004936.3: c.20G>A
(p.G7D)
chr9:22008933

CHEK1

 NM_001274.4: c.587T>C
(p.I196T)
chr11:125503220

CIC

 NM_015125.4: c.1852G>A
(p.A618T)
chr19:42794772

CREBBP

 NM_004380.2: c.6431C>T
(p.A2144V)
chr16:3778617,
NM_004380.2: c.2896G>T
(p.A966S)
chr16:3819339 and
NM_004380.2: c.1184T>C
(p.M395T)
chr16:3843419

CSF3R

 NM_156039.3: c.881T>C
(p.L294P)
chr1:36937955

CTCF

 NM_006565.3: c.528G>T
(p.E176D)
chr16:67645263,
NM_006565.3: c.187A>G
(p.M63V)
chr16:67644922 and
NM_006565.3: c.752C>T
(p.P251L)
chr16:67645487

CTNNA1

 NM_001903.2: c.1486C>T
(p.R496C)
chr5:138253527

CTNNB1

 NM_001904.3: c.932G>T
(p.S311I)
chr3:41267348

CXCR4

 NM_001008540.1: c.1021C>T
(p.H341Y)
chr2:136872489

CYP17A1

 NM_000102.3: c.213G>T
(p.K71N)
chr10:104596906

DAXX

 NM_001350.4: c.1986G>T
(p.K662N)
chr6:33286951

DDR1

 NM_001954.4: c.1756dup
(p.E586Gfs*6)
chr6:30864636,
NM_001954.4: c.474del
(p.M159Wfs*16)
chr6:30858800-30858801,
NM_001954.4: c.178C>T
(p.R60C)
chr6:30856777 and
NM_001954.4: c.1514-1G>A
(p.?)
chr6:30864397

DIS3

 NM_001128226.1: c.29del
(p.K10Rfs*7)
chr13:73355941-73355942
and NM_001128226.1:
c.892C>T (p.R298*)
chr13:73349354

DOT1L

 NM_032482.2: c.4504C>T
(p.P1502S)
chr19:2227024,
NM_032482.2: c.1237C>T
(p.R413C)
chr19:2210740 and
NM_032482.2: c.2191T>C
(p.S731P)
chr19:2216547

EGFR

 NM_005228.3: c.162G>A
(p.M54I)
chr7:55210052 and
NM_005228.3: c.2011C>T
(p.R671C)
chr7:55240767

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance
EP300

NM_001429.3: c.6413C>T
(p.A2138V)
chr22:41574128,
NM_001429.3: c.4383G>T
(p.K1461N)
chr22:41566506,
NM_001429.3: c.4411C>T
(p.L1471F)
chr22:41566534 and
NM_001429.3: c.7177C>G
(p.L2393V)
chr22:41574892

EPHA3

NM_005233.4: c.962C>T
(p.A321V)
chr3:89390213,
NM_005233.4: c.860G>A
(p.C287Y)
chr3:89390111 and
NM_005233.4: c.1777C>A
(p.L593I)
chr3:89462305

EPHB1

NM_004441.4: c.2092G>A
(p.E698K)
chr3:134911627

EPHB4

NM_004444.4: c.898G>A
(p.A300T)
chr7:100417829,
NM_004444.4: c.2918A>C
(p.K973T)
chr7:100401129 and
NM_004444.4: c.718C>T
(p.R240C)
chr7:100419983

ERBB4

NM_005235.2: c.3749G>A
(p.R1250Q)
chr2:212248518

ERCC4

NM_005236.2: c.341T>C
(p.L114P)
chr16:14016021

ERG

NM_182918.3: c.1099C>T
(p.R367C)
chr21:39755666

FANCA

NM_000135.2: c.1247C>T
(p.A416V)
chr16:89857923

FANCG

NM_004629.1: c.1235A>G
(p.D412G)
chr9:35075660

FANCL

NM_018062.3: c.203G>A
(p.R68Q)
chr2:58456962

FGF10

NM_004465.1: c.82T>C
(p.F28L)
chr5:44388703 and
NM_004465.1: c.101C>T
(p.P34L)
chr5:44388684

FGF4

NM_002007.2: c.428G>T
(p.G143V)
chr11:69588808

FGFR2

NM_000141.4: c.108A>C
(p.E36D)
chr10:123353224

FGFR3

NM_000142.3: c.1630G>A
(p.A544T)
chr4:1807381, NM_000142.3:
c.1697_1715dup
(p.P573Sfs*98)
chr4:1807527 and
NM_000142.3: c.587G>A
(p.R196H)
chr4:1803235

FGFR4

NM_213647.3: c.1094C>T
(p.A365V)
chr5:176520175 and
NM_213647.3: c.1793G>A
(p.R598Q)
chr5:176522696

FLCN

NM_144997.5: c.1327G>A
(p.A443T)
chr17:17118604

FLT1

NM_002019.4: c.166del
(p.E56Kfs*12)
chr13:29041261-29041262
and NM_002019.4: c.3695del
(p.L1232Yfs*24)
chr13:28883004-28883005

FLT3

NM_004119.2: c.2416T>C
(p.S806P)
chr13:28597489,
NM_004119.2: c.807G>A
(p.W269*)
chr13:28623847 and
NM_004119.2: c.2542-1G>A
(p.?)
chr13:28589839

GABRA6

NM_000811.2: c.825del
(p.F275Lfs*7)
chr5:161117353-161117354,
NM_000811.2: c.340C>T
(p.P114S)
chr5:161116069 and
NM_000811.2: c.248dup
(p.R84Pfs*6)
chr5:161115970

GATA3

NM_001002295.1: c.373G>A
(p.G125S)
chr10:8100399

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance
GATA6

 NM_005257.3: c.493G>A
(p.A165T)
chr18:19751598 and
NM_005257.3: c.323G>T
(p.S108I)
chr18:19751428

GNA11

 NM_002067.2: c.752G>A
(p.S251N)
chr19:3119220

GNAS

 NM_080425.2: c.317C>T
(p.A106V)
chr20:57428637,
NM_016592.2: c.505C>T
(p.R169*)
chr20:57415666 and
NM_000516.4: c.970+1G>A
(p.?)
chr20:57485137

GRM3

 NM_000840.2: c.691G>A
(p.E231K)
chr7:86415799 and
NM_000840.2: c.868C>T
(p.R290C)
chr7:86415976

GSK3B

 NM_002093.3: c.316C>T
(p.H106Y)
chr3:119666165

HDAC1

 NM_004964.2: c.1354G>T
(p.D452Y)
chr1:32797825

HNF1A

 NM_000545.4:
c.864delinsCC
(p.G292Rfs*25)
chr12:121432117 and
NM_000545.4: c.775G>A
(p.V259I)
chr12:121432028

INPP4B

 NM_003866.2: c.2337A>C
(p.E779D)
chr4:143029283 and
NM_003866.2: c.1058G>A
(p.S353N)
chr4:143129592

IRF2

 NM_002199.3: c.745C>T
(p.R249W)
chr4:185310217

IRF4

 NM_002460.3: c.946G>A
(p.V316I)
chr6:401624

IRS2

 NM_003749.2: c.1826C>T
(p.A609V)
chr13:110436575,
NM_003749.2: c.2321C>T
(p.A774V)
chr13:110436080,
NM_003749.2: c.1658G>A
(p.G553D)
chr13:110436743,
NM_003749.2: c.3626C>T
(p.P1209L)
chr13:110434775 and
NM_003749.2: c.2591C>A
(p.P864H)
chr13:110435810

JAK1

 NM_002227.2: c.521G>A
(p.R174Q)
chr1:65335120

JAK2

 NM_004972.3: c.3245dup
(p.L1082Ffs*4)
chr9:5126396

JAK3

 NM_000215.3: c.1820C>T
(p.A607V)
chr19:17946827 and
NM_000215.3: c.309-1G>T
(p.?)
chr19:17954301

KDM5A

 NM_001042603.1: c.3597del
(p.G1200Dfs*9)
chr12:416952-416953,
NM_001042603.1: c.473del
(p.L158*)
chr12:475163-475164,
NM_001042603.1: c.2947A>G
(p.N983D)
chr12:422311 and
NM_001042603.1: c.3091C>T
(p.R1031C)
chr12:420176

KEAP1

 NM_012289.3: c.1510A>G
(p.N504D)
chr19:10600345

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance

KEL NM_000420.2: c.538C>T (p.R180C) chr7:142655048	KMT2A (MLL) NM_005933.3: c.9938C>T (p.A3313V) chr11:118376554 and NM_005933.3: c.11456G>A (p.R3819H) chr11:118391552	KMT2D (MLL2) NM_003482.4: c.592G>A (p.A198T) chr12:49447842, NM_003482.4: c.2120C>T (p.A707V) chr12:49445346, NM_003482.4: c.6139A>G (p.K2047E) chr12:49435744 and NM_003482.4: c.7254G>C (p.Q2418H) chr12:49434299	LTK NM_002344.5: c.1808G>T (p.R603M) chr15:41797618
MAF NM_005360.4: c.905C>T (p.A302V) chr16:79632895	MAPK1 NM_002745.4: c.802C>T (p.P268S) chr22:22142600	MDM2 NM_002392.3: c.1328C>A (p.P443H) chr12:69233463	MEF2B NM_001145785.1: c.407C>T (p.A136V) chr19:19257979, NM_001145785.1: c.861del (p.A288Pfs*116) chr19:19257101-19257102 and NM_001145785.1: c.944dup (p.V316Sfs*6) chr19:19256768
MERTK NM_006343.2: c.2186G>A (p.C729Y) chr2:112777096, NM_006343.2: c.1714G>A (p.E572K) chr2:112760692 and NM_006343.2: c.2782C>T (p.R928W) chr2:112786223	MET NM_000245.2: c.1475T>C (p.V492A) chr7:116380086	MKNK1 NM_003684.4: c.103C>T (p.R35W) chr1:47048933	MST1R NM_002447.2: c.464T>C (p.V155A) chr3:49940579
MTAP NM_002451.3: c.362C>A (p.P121H) chr9:21837921	MTOR NM_004958.3: c.2095G>A (p.A699T) chr1:11298013, NM_004958.3: c.607C>T (p.Q203*) chr1:11316147 and NM_004958.3: c.3781A>G (p.S1261G) chr1:11269389	MYC NM_002467.4: c.650G>A (p.S217N) chr8:128751113	MYCL (MYCL1) NM_005376.4: c.341G>A (p.R114K) chr1:40366856

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance
MYCN

NM_005378.4: c.511G>A
(p.A171T)
chr2:16082697,
NM_005378.4: c.551C>T
(p.A184V)
chr2:16082737 and
NM_005378.4: c.754C>T
(p.L252F)
chr2:16082940

NOTCH2

NM_024408.3: c.1105G>A
(p.A369T)
chr1:120512137 and
NM_024408.3: c.1043C>T
(p.T348I)
chr1:120512199

NTRK2

NM_006180.3: c.1264G>A
(p.V422I)
chr9:87359956

PDCD1 (PD-1)

NM_005018.2: c.852C>A
(p.C284*)
chr2:242793225 and
NM_005018.2: c.755T>C
(p.V252A)
chr2:242793322

NBN

NM_002485.4: c.452T>C
(p.V151A)
chr8:90992990 and
NM_002485.4: c.1249G>A
(p.V417I)
chr8:90967659

NOTCH3

NM_000435.2: c.1739A>G
(p.E580G)
chr19:15298017,
NM_000435.2: c.2413C>A
(p.P805T)
chr19:15295259,
NM_000435.2: c.3725G>A
(p.R1242H)
chr19:15289746 and
NM_000435.2: c.4619G>A
(p.R1540H)
chr19:15284996

NTRK3

NM_002530.2: c.2470C>T
(p.L824F)
chr15:88420174

PDGFRA

NM_006206.4: c.1891C>T
(p.P631S)
chr4:55143659

NF1

NM_001042492.2:
c.8498A>G (p.N2833S)
chr17:29701151,
NM_001042492.2: c.197A>G
(p.N66S)
chr17:29483137 and
NM_001042492.2: c.467G>A
(p.R156H)
chr17:29490382

**NSD2 (WHSC1 OR
MMSET)**

NM_133335.3: c.1862C>T
(p.A621V)
chr4:1941486 and
NM_133335.3: c.1601A>G
(p.D534G)
chr4:1936916

PARP1

NM_001618.3: c.989G>A
(p.R330Q)
chr1:226573227 and
NM_001618.3: c.1063C>T
(p.R355C)
chr1:226570833

PDGFRB

NM_002609.3: c.2278G>A
(p.D760N)
chr5:149501509 and
NM_002609.3: c.1811G>A
(p.R604H)
chr5:149504391

NOTCH1

NM_017617.3: c.6787C>T
(p.R2263W)
chr9:139391404

NTRK1

NM_002529.3: c.922C>T
(p.Q308*)
chr1:156843496

PARP2

NM_005484.3: c.1112C>T
(p.P371L)
chr14:20824162 and
NM_005484.3: c.1320C>A
(p.S440R)
chr14:20824800

PIK3C2G

NM_004570.4: c.310G>A
(p.A104T)
chr12:18435325 and
NM_004570.4: c.1523T>C
(p.V508A)
chr12:18499668

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX
Variants of Unknown Significance
PIK3R1

 NM_181523.3: c.1469C>T
(p.T490I)
chr5:67590407

PIM1

 NM_002648.3: c.480C>A
(p.N160K)
chr6:37139140

POLE

 NM_006231.2: c.3593C>T
(p.A1198V)
chr12:133226465,
NM_006231.2: c.2683G>A
(p.A895T)
chr12:133240613,
NM_006231.2: c.2091dup
(p.F699Vfs*11)
chr12:133245023,
NM_006231.2: c.4472A>G
(p.H1491R)
chr12:133219889 and
NM_006231.2: c.1372T>C
(p.Y458H)
chr12:133249851

PPP2R1A

 NM_014225.5: c.313C>T
(p.R105W)
chr19:52714555

PRKN (PARK2)

 NM_004562.2: c.1381C>A
(p.H461N)
chr6:161771148

PTCH1

 NM_000264.3: c.1715C>T
(p.A572V)
chr9:98238329

PTPRO

 NM_030667.1: c.1544C>A
(p.P515H)
chr12:15668511

RAD51B

 NM_133509.3: c.769A>C
(p.N257H)
chr14:68758613

RAD54L

 NM_003579.3: c.296G>A
(p.G99D)
chr1:46725660 and
NM_003579.3: c.785G>A
(p.R262H)
chr1:46726951

RAF1

 NM_002880.3: c.61G>A
(p.V21M)
chr3:12660160

RARA

 NM_000964.3: c.310G>A
(p.A104T)
chr17:38504699

RET

 NM_020975.4: c.1538C>T
(p.A513V)
chr10:43607562

RICTOR

 NM_152756.3: c.2974G>A
(p.A992T)
chr5:38952451

RNF43

 NM_017763.4: c.2195G>A
(p.R732H)
chr17:56434942 and
NM_017763.4: c.1982C>T
(p.S661F)
chr17:56435155

ROS1

 NM_002944.2: c.4487A>G
(p.D1496G)
chr6:117665260

RPTOR

 NM_020761.2: c.1568C>T
(p.A523V)
chr17:78854273

SETD2

 NM_014159.6: c.4205G>A
(p.G1402E)
chr3:47161921, NM_014159.6:
c.1666A>G (p.K556E)
chr3:47164460,
NM_014159.6: c.1202G>A
(p.R401Q)
chr3:47164924, NM_014159.6:
c.209G>A (p.R70Q)
chr3:47165917 and
NM_014159.6: c.2848C>T
(p.R950C)
chr3:47163278

SF3B1

 NM_012433.2: c.2993A>G
(p.K998R)
chr2:198264799

SMARCA4

 NM_003072.3: c.145C>T
(p.P49S)
chr19:11094972

SMO

 NM_005631.4: c.586C>T
(p.P196S)
chr7:128845092 and
NM_005631.4: c.782G>A
(p.R261H)
chr7:128845485

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX

Variants of Unknown Significance

SPEN

 NM_015001.2: c.8476G>T
(p.G2826C)
chr1:16261211 and
NM_015001.2: c.3065T>G
(p.V1022G)
chr1:16255800

TEK

 NM_000459.3: c.2744G>A
(p.R915H)
chr9:27212762

TIPARP

 NM_015508.4: c.322T>C
(p.S108P)
chr3:156395808 and
NM_015508.4: c.1219T>A
(p.S407T)
chr3:156413786

XPO1

 NM_003400.3:
c.1608_1610del (p.N536del)
chr2:61719572-61719575

SRC

 NM_005417.3: c.248C>T
(p.A83V)
chr20:36012804 and
NM_005417.3: c.1163G>A
(p.R388Q)
chr20:36030884

TENT5C (FAM46C)

 NM_017709.3: c.110G>A
(p.G37E)
chr1:118165600

TSC1

 NM_000368.4: c.1837C>T
(p.P613S)
chr9:135781128

SYK

 NM_003177.5: c.1108C>A
(p.L370M)
chr9:93637058

TET2

 NM_001127208.2: c.2932C>T
(p.H978Y)
chr4:106158031,
NM_001127208.2: c.4909C>A
(p.L1637I)
chr4:106196576,
NM_001127208.2: c.3437C>A
(p.P1146H)
chr4:106162523 and
NM_001127208.2: c.1088C>A
(p.P363H)
chr4:106156187

TSC2

 NM_000548.3: c.4370G>A
(p.R1457Q)
chr16:2134593 and
NM_000548.3: c.1384C>T
(p.R462C)
chr16:2112995

TBX3

 NM_016569.3: c.694G>A
(p.G232R)
chr12:115117741

TGFBR2

 NM_003242.5: c.61G>A
(p.A21T)
chr3:30648436,
NM_003242.5: c.383del
(p.K128Sfs*35)
chr3:30691871-30691872 and
NM_003242.5: c.1484G>A
(p.R495Q)
chr3:30729963

WT1

 NM_024426.4: c.680G>A
(p.S227N)
chr11:32450132

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX

Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFA1	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFBP2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPR2SS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS


Homologous Recombination status
Loss of Heterozygosity (LOH) score
Microsatellite (MS) status
Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium. 

ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® HiSeq platform, hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI) and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is one that the FoundationOne CDx analytical

methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and MSS cut-off thresholds were determined by

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX

About FoundationOne®CDx

- analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage <300X, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score ≥ 16 will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the genome on the FoundationOne CDx test and

extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
- Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal

hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >10%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Brennan Decker, M.D., Ph.D. | 04 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1637928-01

APPENDIX

About FoundationOne®CDx

tumor sequencing is germline or somatic.
 Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D (MLL2)*, *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking

into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.9.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. The median exon coverage for this sample is 874x. Foundation Medicine Inc. does not warrant the suitability of use.

ORDERED TEST # **ORD-1637928-01**
APPENDIX
References

1. Histopathology (2007) PMID: 17204026
2. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
3. Overman MJ, et al. Lancet Oncol. (2017) PMID: 28734759
4. Overman MJ, et al. J. Clin. Oncol. (2018) PMID: 29355075
5. Shitara K, et al. Nature (2022) PMID: 35322232
6. Lipson EJ, et al. Clin. Cancer Res. (2013) PMID: 23169436
7. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
8. Rizvi NA, et al. Science (2015) PMID: 25765070
9. Oaknin A, et al. JAMA Oncol (2020) PMID: 33001143
10. Rubinstein MM, et al. Gynecol Oncol (2022) PMID: 36512912
11. Kanikarla Marie P, et al. Clin Cancer Res (2021) PMID: 33811152
12. Hall et al., 2016; ASCO Abstract 1523
13. Cheng et al., 2016; ASCO Abstract 251
14. Pritchard CC, et al. Nat Commun (2014) PMID: 25255306
15. Schwarzenbach H, et al. BJU Int. (2008) PMID: 18336598
16. Suzuki H, et al. Jpn. J. Cancer Res. (1995) PMID: 7493915
17. Dominguez-Valentin M, et al. BMC Urol (2016) PMID: 27013479
18. Rosty C, et al. Fam. Cancer (2014) PMID: 25117503
19. Ahman AK, et al. BJU Int. (2001) PMID: 11251526
20. Rodrigues DN, et al. J. Clin. Invest. (2018) PMID: 30382943
21. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
22. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
23. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
24. Boland CR, et al. Cancer Res. (1998) PMID: 9823339
25. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
26. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
27. Lynch HT, et al. Clin. Genet. (2009) PMID: 19659756
28. Pande M, et al. Fam. Cancer (2012) PMID: 22714864
29. Kastrinos F, et al. Semin. Oncol. (2007) PMID: 17920897
30. Silva FC, et al. Sao Paulo Med J (2009) PMID: 19466295
31. Sehgal R, et al. Genes (Basel) (2014) PMID: 24978665
32. Fam. Cancer (2005) PMID: 16136383
33. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
34. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
35. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
36. Cristescu R, et al. Science (2018) PMID: 30309915
37. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
38. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
39. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
40. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
41. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
42. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
43. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
44. Ott PA, et al. J. Clin. Oncol. (2019) PMID: 30557521
45. Cristescu R, et al. J Immunother Cancer (2022) PMID: 35101941
46. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
47. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
48. Schenker et al., 2022; AACR Abstract 7845
49. Legrand et al., 2018; ASCO Abstract 12000
50. Graf RP, et al. JAMA Netw Open (2022) PMID: 35357449
51. Chalmers ZR, et al. Genome Med (2017) PMID: 28420421
52. Lawrence MS, et al. Nature (2013) PMID: 23770567
53. Alexandrov LB, et al. Nature (2013) PMID: 23945592
54. Barbieri CE, et al. Nat. Genet. (2012) PMID: 22610119
55. Baca SC, et al. Cell (2013) PMID: 23622249
56. Cell (2015) PMID: 26544944
57. Grasso CS, et al. Nature (2012) PMID: 22722839
58. Gundem G, et al. Nature (2015) PMID: 25830880
59. Robinson D, et al. Cell (2015) PMID: 26000489
60. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
61. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
62. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
63. Johnson BE, et al. Science (2014) PMID: 24336570
64. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
65. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
66. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
67. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
68. Nature (2012) PMID: 22810696
69. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
70. O'Malley et al., 2017; AACR-NCI-EORTC Abstract LB-A12
71. Wu W, et al. Cancer Res. (2015) PMID: 25634209
72. Wang Y, et al. J. Clin. Invest. (2016) PMID: 27454289
73. Ozden O, et al. Sci Rep (2016) PMID: 27197561
74. Zhao W, et al. Nature (2017) PMID: 28976962
75. Taylor BS, et al. Cancer Cell (2010) PMID: 20579941
76. Gynecol. Oncol. (2010) PMID: 19959210
77. Laufer M, et al. J. Biol. Chem. (2007) PMID: 17848578
78. Westermarck UK, et al. Mol. Cell. Biol. (2003) PMID: 14560035
79. Kaufman B, et al. J. Clin. Oncol. (2015) PMID: 25366685
80. Mateo J, et al. N. Engl. J. Med. (2015) PMID: 26510020
81. Tutt A, et al. Lancet (2010) PMID: 20609467
82. Robson M, et al. N. Engl. J. Med. (2017) PMID: 28578601
83. Mirza MR, et al. N. Engl. J. Med. (2016) PMID: 27717299
84. Sandhu SK, et al. Lancet Oncol. (2013) PMID: 23810788
85. Swisher EM, et al. Lancet Oncol. (2017) PMID: 27908594
86. Drew Y, et al. Br. J. Cancer (2016) PMID: 27002934
87. Pujade-Lauraine E, et al. Lancet Oncol. (2017) PMID: 28754483
88. Ledermann JA, et al. Lancet Oncol. (2016) PMID: 27617661
89. Fong PC, et al. N. Engl. J. Med. (2009) PMID: 19553641
90. Audeh MW, et al. Lancet (2010) PMID: 20609468
91. Fong PC, et al. J. Clin. Oncol. (2010) PMID: 20406929
92. Gelmon KA, et al. Lancet Oncol. (2011) PMID: 21862407
93. Kaye SB, et al. J. Clin. Oncol. (2012) PMID: 22203755
94. Domchek SM, et al. Gynecol. Oncol. (2016) PMID: 26723501
95. Moore K, et al. N. Engl. J. Med. (2018) PMID: 30345884
96. Golan T, et al. N. Engl. J. Med. (2019) PMID: 31157963
97. Yap TA, et al. Cancer Discov (2021) PMID: 32988960
98. Thomas A, et al. J. Clin. Oncol. (2018) PMID: 29252124
99. Saito YD, et al. Cancer Treat Res Commun (2018) PMID: 31299005
100. Westin et al., 2021; ASCO Abstract 5505
101. Do K, et al. J. Clin. Oncol. (2015) PMID: 25964244
102. Pouliot GP, et al. PLoS ONE (2019) PMID: 31721781
103. Kim H, et al. Clin. Cancer Res. (2017) PMID: 27993965
104. Jin J, et al. Neoplasia (2018) PMID: 29605721
105. Hussain et al., 2019; ESMO Abstract LBA12PR
106. Yap et al., 2022; AACR Abstract CT007
107. Cheng HH, et al. Eur. Urol. (2016) PMID: 26724258
108. Imyanito EN, et al. Hered Cancer Clin Pract (2011) PMID: 21819606
109. Mota JM, et al. JCO Precis Oncol (2020) PMID: 32856010
110. Pomerantz MM, et al. Cancer (2017) PMID: 28608931
111. Privé BM, et al. Prostate Cancer Prostatic Dis (2021) PMID: 34253846
112. Crumbaker M, et al. JCO Precis Oncol (2019) PMID: 35100671
113. Satapathy S, et al. Clin Nucl Med (2021) PMID: 33630794
114. Kratochwil C, et al. J Nucl Med (2020) PMID: 31601699
115. Ahmadzadehfard H, et al. Clin Nucl Med (2018) PMID: 29762244
116. De Giorgi U, et al. Br J Cancer (2021) PMID: 34333554
117. Castro E, et al. J. Clin. Oncol. (2019) PMID: 30625039
118. Eur. Urol. (2016) PMID: 26708042
119. Isakoff SJ, et al. J. Clin. Oncol. (2015) PMID: 25847936
120. Dann RB, et al. Gynecol. Oncol. (2012) PMID: 22406760
121. Evers B, et al. Clin. Cancer Res. (2008) PMID: 18559613
122. Soares DG, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17656556
123. Miolo G, et al. Cancer Biol. Ther. (2016) PMID: 27561088
124. Ghouadni A, et al. Breast (2017) PMID: 28467918
125. Case Rep Oncol () PMID: 28626402
126. Monk BJ, et al. Gynecol. Oncol. (2020) PMID: 31924332
127. Harnicek D, et al. Int. J. Cancer (2016) PMID: 26933761
128. Cruz C, et al. J. Clin. Oncol. (2018) PMID: 30240327
129. Poveda A, et al. Ann. Oncol. (2017) PMID: 28368437
130. Chung JH, et al. JCO Precis Oncol (2019) PMID: 31218271
131. Pritchard CC, et al. N. Engl. J. Med. (2016) PMID: 27433846
132. Abida W, et al. Proc. Natl. Acad. Sci. U.S.A. (2019) PMID: 31061129
133. Kumar A, et al. Nat. Med. (2016) PMID: 26928463
134. Mitra AV, et al. BJU Int. (2011) PMID: 20840664
135. Castro E, et al. J. Clin. Oncol. (2013) PMID: 23569316
136. Mitra A, et al. Br J Cancer (2008) PMID: 18182994
137. Na R, et al. Eur Urol (2017) PMID: 27989354
138. Narod SA, et al. Br J Cancer (2008) PMID: 18577985
139. Page EC, et al. Eur Urol (2019) PMID: 31537406
140. Thorne H, et al. Cancer Prev Res (Phila) (2011) PMID: 21733824
141. Wei Y, et al. Oncologist (2020) PMID: 32190957
142. Darst et al., 2022; AACR Abstract 688
143. Castro E, et al. Eur. Urol. (2015) PMID: 25454609
144. Warner E, et al. Clin Cancer Res (2021) PMID: 33414135
145. Yang H, et al. Science (2002) PMID: 12228710
146. Nat. Struct. Mol. Biol. (2011) PMID: 21731065
147. Biswas K, et al. Blood (2011) PMID: 21719596
148. Biswas K, et al. Hum. Mol. Genet. (2012) PMID: 22678057
149. Claes K, et al. Br. J. Cancer (2004) PMID: 15026808
150. Cote S, et al. Breast Cancer Res. Treat. (2012) PMID: 21947752
151. Fackenthal JD, et al. Am. J. Hum. Genet. (2002) PMID: 12145750
152. Farrugia DJ, et al. Cancer Res. (2008) PMID: 18451181
153. Gómez García EB, et al. Breast Cancer Res. (2009) PMID: 19200354
154. Guidugli L, et al. Cancer Res. (2013) PMID: 23108138
155. Kluska A, et al. BMC Med Genomics (2015) PMID: 25948282

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX References

156. Kuznetsov SG, et al. Nat. Med. (2008) PMID: 18607349
157. Roy R, et al. Nat. Rev. Cancer (2011) PMID: 22193408
158. Siaud N, et al. PLoS Genet. (2011) PMID: 22194698
159. Thomassen M, et al. Genet. Test. (2006) PMID: 16792514
160. Xia B, et al. Mol. Cell (2006) PMID: 16793542
161. Guidugli L, et al. Am. J. Hum. Genet. (2018) PMID: 29394989
162. Fraile-Bethencourt E, et al. PLoS Genet. (2017) PMID: 28339459
163. Landrum MJ, et al. Nucleic Acids Res. (2018) PMID: 29165669
164. Miki Y, et al. Science (1994) PMID: 7545954
165. Wooster R, et al. Nature () PMID: 8524414
166. King MC, et al. Science (2003) PMID: 14576434
167. MedGenMed (2005) PMID: 16369438
168. Ford D, et al. Lancet (1994) PMID: 7907678
169. Whittemore AS, et al. Am. J. Hum. Genet. (1997) PMID: 9042908
170. Claus EB, et al. Cancer (1996) PMID: 8635102
171. Struwing JP, et al. N. Engl. J. Med. (1997) PMID: 9145676
172. Oddoux C, et al. Nat. Genet. (1996) PMID: 8841192
173. Hall MJ, et al. Cancer (2009) PMID: 19241424
174. Abida W, et al. Clin. Cancer Res. (2020) PMID: 32086346
175. Dhawan et al., 2017; ASCO Abstract 2527
176. Cerami E, et al. Cancer Discov (2012) PMID: 22588877
177. Gao J, et al. Sci Signal (2013) PMID: 23550210
178. Cantor SB, et al. Cell (2001) PMID: 11301010
179. Litman R, et al. Cancer Cell (2005) PMID: 16153896
180. Bridge WL, et al. Nat. Genet. (2005) PMID: 16116421
181. Cantor SB, et al. Future Oncol (2011) PMID: 21345144
182. Gupta R, et al. Nucleic Acids Res. (2006) PMID: 17145708
183. Cantor S, et al. Proc. Natl. Acad. Sci. U.S.A. (2004) PMID: 14983014
184. Filippini SE, et al. Front Biosci (Landmark Ed) (2013) PMID: 23747889
185. Ren LP, et al. Genet. Mol. Res. (2013) PMID: 24301948
186. Seal S, et al. Nat. Genet. (2006) PMID: 17033622
187. Rafnar T, et al. Nat. Genet. (2011) PMID: 21964575
188. Ma XD, et al. Biochem. Biophys. Res. Commun. (2013) PMID: 23473757
189. Hematology Am Soc Hematol Educ Program (2007) PMID: 18024606
190. Khincha PP, et al. Semin. Hematol. (2013) PMID: 24246701
191. Chandrasekharappa SC, et al. Blood (2013) PMID: 23613520
192. Tischkowitz MD, et al. J. Med. Genet. (2003) PMID: 12525534
193. Popova T, et al. Cancer Res. (2016) PMID: 26787835
194. Wu YM, et al. Cell (2018) PMID: 29906450
195. Viswanathan SR, et al. Cell (2018) PMID: 29909985
196. Menghi F, et al. Cancer Cell (2018) PMID: 30017478
197. Sokol ES, et al. Oncologist (2019) PMID: 31292271
198. Antonarakis ES, et al. JCO Precis Oncol (2020) PMID: 32462107
199. Schweizer MT, et al. JCO Precis Oncol (2020) PMID: 32671317
200. Reimers MA, et al. Eur. Urol. (2019) PMID: 31640893
201. Linch et al., 2021; AACR Abstract LB004
202. Alva et al., 2022; ASCO GU Abstract 103
203. Joshi PM, et al. J. Biol. Chem. (2014) PMID: 24554720
204. Natrajan R, et al. J. Pathol. (2014) PMID: 24395524
205. Bajrami I, et al. Cancer Res. (2014) PMID: 24240700
206. Ekumi KM, et al. Nucleic Acids Res. (2015) PMID: 25712099
207. Johnson SF, et al. Cell Rep (2016) PMID: 27880910
208. AlHilli MM, et al. Gynecol. Oncol. (2016) PMID: 27614696
209. de Bono J, et al. N. Engl. J. Med. (2020) PMID: 32343890
210. Mateo J, et al. Lancet Oncol. (2019) PMID: 31806540
211. Dong B, et al. J Natl Compr Canc Netw (2021) PMID: 33990090
212. Rescigno P, et al. Clin Cancer Res (2021) PMID: 32988971
213. Graf RP, et al. Eur Urol (2021) PMID: 34716049
214. Nguyen B, et al. Eur. Urol. (2020) PMID: 32317181
215. Wang Z, et al. Prostate Cancer Prostatic Dis (2022) PMID: 36401126
216. Blazek D, et al. Genes Dev. (2011) PMID: 22012619
217. Böskén CA, et al. Nat Commun (2014) PMID: 24662513
218. Dixon-Clarke SE, et al. Sci Rep (2015) PMID: 26597175
219. Zhang X, et al. Mol. Cell Proteomics (2019) PMID: 30617155
220. Wang Q, et al. Cell Growth Differ. (1995) PMID: 8547221
221. Zabloudoff SD, et al. Mol. Cancer Ther. (2008) PMID: 18790776
222. Wang L, et al. Int. J. Oncol. (2018) PMID: 30015873
223. Lopes JL, et al. Cancer Genet (2019) PMID: 31078449
224. McCabe N, et al. Cancer Res. (2006) PMID: 16912188
225. Italiano A, et al. Ann. Oncol. (2018) PMID: 29788155
226. Calvo E, et al. Oncology (2016) PMID: 27598338
227. Doi T, et al. Anticancer Drugs (2015) PMID: 26288133
228. Chamoun K, et al. Expert Opin Invest Drugs (2018) PMID: 30084282
229. Zehir A, et al. Nat. Med. (2017) PMID: 28481359
230. Albiges L, et al. Breast (2014) PMID: 24636978
231. Lee HJ, et al. Mol. Cancer Ther. (2017) PMID: 28138032
232. Bai X, et al. Transl Oncol (2018) PMID: 29287241
233. Mazumder Indra D, et al. Int. J. Cancer (2011) PMID: 21154811
234. Grabauskiene S, et al. J. Surg. Res. (2014) PMID: 24418519
235. Abdel-Fatah TM, et al. Mol Oncol (2015) PMID: 25468710
236. Xie Y, et al. Med. Oncol. (2014) PMID: 24464213
237. Chini CC, et al. J. Biol. Chem. (2006) PMID: 16963448
238. Goto H, et al. Cancer Sci. (2012) PMID: 22435685
239. Bahassi EM, et al. Oncogene (2008) PMID: 18317453
240. Sørensen CS, et al. Nat. Cell Biol. (2005) PMID: 15665856
241. Capasso H, et al. J. Cell. Sci. (2002) PMID: 12415000
242. Zhao H, et al. Mol. Cell. Biol. (2001) PMID: 11390642
243. Ng CP, et al. J. Biol. Chem. (2004) PMID: 14681223
244. Gong EY, et al. Sci Rep (2018) PMID: 30510197
245. Dougherty et al., 2014; ASCO Abstract 5536
246. Tate JG, et al. Nucleic Acids Res. (2019) PMID: 30371878
247. Qu X, et al. Blood (2019) PMID: 30446494
248. Pace P, et al. EMBO J. (2002) PMID: 12093742
249. Deakyné JS, et al. Biochemistry Mosc. (2011) PMID: 21568838
250. Alpi A, et al. Mol. Cell. Biol. (2007) PMID: 17938197
251. Alpi AF, et al. Mol. Cell (2018) PMID: 19111657
252. Meetei AR, et al. Nat. Genet. (2003) PMID: 12973351
253. Machida YJ, et al. Mol. Cell (2006) PMID: 16916645
254. Hodson C, et al. J. Biol. Chem. (2011) PMID: 21775430
255. Hodson C, et al. Structure (2014) PMID: 24389026
256. Seki S, et al. Genes Cells (2007) PMID: 17352736
257. Ali AM, et al. Hum. Mutat. (2009) PMID: 19405097
258. Fostira F, et al. Breast Cancer Res. Treat. (2018) PMID: 29335925
259. Hart SN, et al. BMJ Open (2016) PMID: 27084275
260. Del Valle J, et al. Cancers (Basel) (2020) PMID: 32235514
261. Dicks E, et al. Oncotarget (2017) PMID: 28881617
262. Rosell R, et al. Lancet Oncol. (2012) PMID: 22285168
263. Douillard JY, et al. Br. J. Cancer (2014) PMID: 24263064
264. Hayashi T, et al. Hum Pathol (2020) PMID: 32673682
265. Cao L, et al. Onco Targets Ther (2018) PMID: 29780256
266. Yang TY, et al. J. Clin. Oncol. (2011) PMID: 21422421
267. Sequist LV, et al. J. Clin. Oncol. (2013) PMID: 23816960
268. Qin BD, et al. Onco Targets Ther (2018) PMID: 30127622
269. Frega S, et al. J Thorac Oncol (2016) PMID: 27131295
270. Long X, et al. Onco Targets Ther (2020) PMID: 33116645
271. Mok TS, et al. J. Clin. Oncol. (2018) PMID: 29864379
272. Jänne PA, et al. N. Engl. J. Med. (2015) PMID: 25923549
273. Hong MH, et al. Cancer (2020) PMID: 32749686
274. Kim HS, et al. Oncotarget (2015) PMID: 26462025
275. Kim HS, et al. Clin. Cancer Res. (2015) PMID: 25424851
276. Mondal G, et al. Acta Neuropathol (2020) PMID: 32303840
277. Cavalieri S, et al. Eur. J. Cancer (2018) PMID: 29734047
278. Chi AS, et al. JCO Precis Oncol (2020) PMID: 32923886
279. Oztan A, et al. Lung Cancer (2017) PMID: 28838405
280. Le X, et al. Clin. Cancer Res. (2018) PMID: 30228210
281. Fassunke J, et al. Nat Commun (2018) PMID: 30405134
282. Cortiura F, et al. Ann Transl Med (2019) PMID: 30788361
283. Zhang Y, et al. Onco Targets Ther (2019) PMID: 30588029
284. Rosen GM, et al. J. Med. Chem. (1988) PMID: 2828624
285. Brown BP, et al. Clin. Cancer Res. (2019) PMID: 30796031
286. Peled N, et al. J Thorac Oncol (2017) PMID: 28286242
287. Li J, et al. Lung Cancer (2018) PMID: 29502892
288. Cheng C, et al. Sci Rep (2015) PMID: 26354324
289. Ogino S, et al. Clin. Cancer Res. (2005) PMID: 16166444
290. de Muga S, et al. Mod. Pathol. (2010) PMID: 20208477
291. Peraldo-Neia C, et al. BMC Cancer (2011) PMID: 21266046
292. Baek KH, et al. Cancer Res Treat (2012) PMID: 22500161
293. Weber DC, et al. Radiat Oncol (2012) PMID: 22546016
294. Carlsson J, et al. Oncol Lett (2013) PMID: 23255921
295. Ciardiello F, et al. N. Engl. J. Med. (2008) PMID: 18337605
296. Cho J, et al. Mol. Cancer (2014) PMID: 24894453
297. Lu S, et al. J Thorac Oncol (2019) PMID: 30610926
298. de Wit M, et al. EBioMedicine (2020) PMID: 32512509
299. Wei Y, et al. JTO Clin Res Rep (2021) PMID: 34590038
300. Yu N, et al. Oncologist (2017) PMID: 34396638
301. Zhan T, et al. Oncogene (2021) PMID: 27617575
302. Jung YS, et al. Exp Mol Med (2020) PMID: 32037398
303. Krishnamurthy N, et al. Cancer Treat Rev (2018) PMID: 29169144
304. Kawazoe et al., 2021; ESMO Abstract 473P
305. Yamada K, et al. Cancer Res (2021) PMID: 33408116
306. Kanda Y, et al. Biochem Biophys Res Commun (2022) PMID: 34837838
307. Fraser M, et al. Nature (2017) PMID: 28068672
308. Liu L, et al. Int. J. Cancer (2011) PMID: 21207416
309. Yoon HY, et al. J Biomol Screen (2012) PMID: 22511308
310. Brock BJ, et al. BJU Int. (2012) PMID: 22077694
311. Gerstein AV, et al. Genes Chromosomes Cancer (2002) PMID: 11921277

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX References

312. Yardy GW, et al. Eur. Urol. (2009) PMID: 18514389
313. Henrique R, et al. Clin Cancer Res (2007) PMID: 17947477
314. Luke JJ, et al. Clin Cancer Res (2019) PMID: 30635339
315. Logan CY, et al. Annu. Rev. Cell Dev. Biol. (2004) PMID: 15473860
316. Eklof Spink K, et al. EMBO J. (2001) PMID: 11707392
317. Liu J, et al. J. Mol. Biol. (2006) PMID: 16753179
318. Dikovskaya D, et al. J. Cell. Sci. (2010) PMID: 20144988
319. Murphy SJ, et al. Dig. Dis. Sci. (2007) PMID: 17410430
320. Aretz S, et al. Hum. Mutat. (2004) PMID: 15459959
321. Kerr SE, et al. J Mol Diagn (2013) PMID: 23159591
322. Annu Rev Pathol (2011) PMID: 21090969
323. Kastritis E, et al. Int. J. Cancer (2009) PMID: 18844223
324. Half E, et al. Orphanet J Rare Dis (2009) PMID: 19822006
325. Fritsch C, et al. Mol. Cancer Ther. (2014) PMID: 24608574
326. Juric D, et al. J. Clin. Oncol. (2018) PMID: 29401002
327. Gallant JN, et al. NPJ Precis Oncol (2019) PMID: 30793038
328. Delestre F, et al. Sci Transl Med (2021) PMID: 34613809
329. Morschhauser F, et al. Mol Cancer Ther (2020) PMID: 31619463
330. Patnaik A, et al. Ann. Oncol. (2016) PMID: 27672108
331. Santin AD, et al. Gynecol Oncol Rep (2020) PMID: 31934607
332. Damodaran S, et al. J Clin Oncol (2022) PMID: 35133871
333. André F, et al. N. Engl. J. Med. (2019) PMID: 31091374
334. Smyth LM, et al. NPJ Breast Cancer (2021) PMID: 33863913
335. Varnier R, et al. Eur J Cancer (2019) PMID: 31351267
336. Basse C, et al. JCO Precis Oncol (2018) PMID: 32914004
337. Sultova E, et al. Arch Gynecol Obstet (2021) PMID: 33277683
338. Mackay HJ, et al. Cancer (2014) PMID: 24166148
339. Myers AP, et al. Gynecol. Oncol. (2016) PMID: 27016228
340. Dhani J, et al. Cold Spring Harb Mol Case Stud (2018) PMID: 29588307
341. Harris EJ, et al. Front Oncol (2019) PMID: 30863722
342. Hanna GJ, et al. Clin Cancer Res (2018) PMID: 29301825
343. Krop et al., 2018; ASCO Abstract 101
344. Pascual J, et al. Cancer Discov (2021) PMID: 32958578
345. Dolly SO, et al. Clin. Cancer Res. (2016) PMID: 26787751
346. Canaud et al., 2021; ESMO Abstract LBA23
347. Stopsack KH, et al. Clin. Cancer Res. (2020) PMID: 32220891
348. Sun X, et al. Anticancer Res. (2009) PMID: 19443396
349. Agell L, et al. Mod. Pathol. (2011) PMID: 21113138
350. Beltran H, et al. Eur. Urol. (2013) PMID: 22981675
351. Pearson HB, et al. Cancer Discov (2018) PMID: 29581176
352. Samuels Y, et al. Cancer Cell (2005) PMID: 15950905
353. Nat. Rev. Cancer (2009) PMID: 19629070
354. Kang S, et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PMID: 15647370
355. Ikenoue T, et al. Cancer Res. (2005) PMID: 15930273
356. Gymnopoulos M, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17376864
357. Horn S, et al. Oncogene (2008) PMID: 18317450
358. Rudd ML, et al. Clin. Cancer Res. (2011) PMID: 21266528
359. Hon WC, et al. Oncogene (2012) PMID: 22120714
360. Burke JE, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22949682
361. Wu H, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19915146
362. Laurenti R, et al. Rev Saude Publica (1990) PMID: 2103068
363. Dan S, et al. Cancer Res. (2010) PMID: 20530683
364. Oda K, et al. Cancer Res. (2008) PMID: 18829572
365. Zhao L, et al. Oncogene (2008) PMID: 18794883
366. Lui VW, et al. Cancer Discov (2013) PMID: 23619167
367. Ross RL, et al. Oncogene (2013) PMID: 22430209
368. Rivière JB, et al. Nat. Genet. (2012) PMID: 22729224
369. Shibata T, et al. Cancer Lett. (2009) PMID: 19394761
370. Dogruluk T, et al. Cancer Res. (2015) PMID: 26627007
371. Croessmann S, et al. Clin. Cancer Res. (2018) PMID: 29284706
372. Ng PK, et al. Cancer Cell (2018) PMID: 29533785
373. Spangle JM, et al. (2020) PMID: 32929011
374. Chen L, et al. Nat Commun (2018) PMID: 29636477
375. Jin N, et al. J Clin Invest (2021) PMID: 34779417
376. Courtney KD, et al. J. Clin. Oncol. (2010) PMID: 20085938
377. Simpson L, et al. Exp. Cell Res. (2001) PMID: 11237521
378. Milella M, et al. Sci Rep (2017) PMID: 28220839
379. Sweeney C, et al. Lancet (2021) PMID: 34246347
380. de Bono JS, et al. Clin. Cancer Res. (2019) PMID: 30037818
381. Crabb SJ, et al. J Clin Oncol (2021) PMID: 33326257
382. George DJ, et al. Urol Oncol (2020) PMID: 31522863
383. Eule CJ, et al. Prostate Cancer Prostatic Dis (2022) PMID: 36402814
384. Templeton AJ, et al. Eur. Urol. (2013) PMID: 23582881
385. Koshkin VS, et al. Invest New Drugs (2019) PMID: 31037562
386. Armstrong AJ, et al. Clin Genitourin Cancer (2013) PMID: 23830964
387. Mendes-Pereira AM, et al. EMBO Mol Med (2009) PMID: 20049735
388. Shen Y, et al. Clin. Cancer Res. (2013) PMID: 23881923
389. Chatterjee P, et al. PLoS ONE (2013) PMID: 23565244
390. McCormick A, et al. Int. J. Gynecol. Cancer (2016) PMID: 26905328
391. Forster MD, et al. Nat Rev Clin Oncol (2011) PMID: 21468130
392. Eikesdal HP, et al. Ann Oncol (2021) PMID: 33242536
393. Pan M, et al. Perm J (2021) PMID: 33970096
394. Romero I, et al. Gynecol Oncol (2020) PMID: 32988624
395. Corn PG, et al. Lancet Oncol (2019) PMID: 31515154
396. Lotan TL, et al. Clin. Cancer Res. (2011) PMID: 21878536
397. Reid AH, et al. Br. J. Cancer (2010) PMID: 20104229
398. Antonarakis ES, et al. Cancer (2012) PMID: 22674438
399. Zu K, et al. Cancer Epidemiol. Biomarkers Prev. (2013) PMID: 23983239
400. Cuzick J, et al. Br. J. Cancer (2013) PMID: 23695019
401. Tsaur I, et al. Cancer Treat Rev (2019) PMID: 30875581
402. Spetsieris N, et al. Cancers (Basel) (2020) PMID: 33339136
403. Aparicio AM, et al. Clin Cancer Res (2013) PMID: 23649003
404. Aparicio AM, et al. Clin Cancer Res (2016) PMID: 26546618
405. Campbell RB, et al. J. Biol. Chem. (2003) PMID: 12857747
406. Rodríguez-Escudero I, et al. Hum. Mol. Genet. (2011) PMID: 21828076
407. He X, et al. Cancer Res. (2013) PMID: 23475934
408. Han SY, et al. Cancer Res. (2000) PMID: 10866302
409. Myers MP, et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PMID: 9811831
410. Pradella LM, et al. BMC Cancer (2014) PMID: 24498881
411. Kim JS, et al. Mol. Cell. Biol. (2011) PMID: 21536651
412. Denning G, et al. Oncogene (2007) PMID: 17213812
413. Hlobilkova A, et al. Anticancer Res. (2016) PMID: 16619501
414. Redfern RE, et al. Protein Sci. (2010) PMID: 20718038
415. Shenoy S, et al. PLoS ONE (2012) PMID: 22505997
416. Wang Y, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19329485
417. Okumura K, et al. J. Biol. Chem. (2006) PMID: 16829519
418. Lee JO, et al. Cell (1999) PMID: 10555148
419. Maxwell GL, et al. Cancer Res. (1998) PMID: 9635567
420. Risinger JL, et al. Clin. Cancer Res. (1998) PMID: 9865913
421. Kato H, et al. Clin. Cancer Res. (2000) PMID: 11051241
422. Fenton TR, et al. Proc. Natl. Acad. Sci. U.S.A. (2012) PMID: 22891331
423. Ngeow J, et al. J. Clin. Endocrinol. Metab. (2012) PMID: 23066114
424. Lobo GP, et al. Hum. Mol. Genet. (2009) PMID: 19457929
425. Liu J, et al. Oncogene (2014) PMID: 23995781
426. Maehama T, et al. Annu. Rev. Biochem. (2001) PMID: 11395408
427. De Vivo I, et al. J. Med. Genet. (2000) PMID: 10807691
428. Ramaswamy S, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10051603
429. Liu JL, et al. Mol. Cell. Biol. (2005) PMID: 15988030
430. Karoui M, et al. Br. J. Cancer (2004) PMID: 15026806
431. Gil A, et al. PLoS ONE (2015) PMID: 25875300
432. Furnari FB, et al. Cancer Res. (1998) PMID: 9823298
433. Spinelli L, et al. J. Med. Genet. (2015) PMID: 25527629
434. Mingo J, et al. Eur. J. Hum. Genet. (2018) PMID: 29706633
435. Wang Q, et al. J. Mol. Graph. Model. (2010) PMID: 20538496
436. Andrés-Pons A, et al. Cancer Res. (2007) PMID: 17942903
437. Butler MG, et al. J. Med. Genet. (2005) PMID: 15805158
438. Georgescu MM, et al. Proc. Natl. Acad. Sci. U.S.A. (1999) PMID: 10468583
439. Staal FJ, et al. Br. J. Cancer (2002) PMID: 12085208
440. Nguyen HN, et al. Oncogene (2014) PMID: 24292679
441. Rahdar M, et al. Proc. Natl. Acad. Sci. U.S.A. (2009) PMID: 19114656
442. Das S, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12808147
443. Wang X, et al. Biochem. J. (2008) PMID: 18498243
444. Valiente M, et al. J. Biol. Chem. (2005) PMID: 15951562
445. Nguyen HN, et al. Oncogene (2015) PMID: 25263454
446. Shan L, et al. Cell Discov (2020) PMID: 32704382
447. Blumenthal GM, et al. Eur. J. Hum. Genet. (2008) PMID: 18781191
448. Orloff MS, et al. Oncogene (2008) PMID: 18794875
449. Zbuk KM, et al. Nat. Rev. Cancer (2007) PMID: 17167516
450. Hao HX, et al. Nature (2012) PMID: 22575959
451. Koo BK, et al. Nature (2012) PMID: 22895187
452. Jiang X, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 23847203
453. Koo BK, et al. Proc. Natl. Acad. Sci. U.S.A. (2015) PMID: 26023187
454. Tsukiyama T, et al. Mol. Cell. Biol. (2015) PMID: 25825523
455. Tabernero J, et al. Oncologist (2023) PMID: 36811382
456. Cook et al., 2021; ESMO Abstract 517MO
457. Elez E, et al. Nat Med (2022) PMID: 36097219
458. Kinde I, et al. Sci Transl Med (2013) PMID: 23303603
459. Giannakis M, et al. Nat. Genet. (2014) PMID: 25344691
460. Seeber A, et al. Clin Cancer Res (2022) PMID: 35254413
461. Wang K, et al. Nat. Genet. (2014) PMID: 24816253

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX

References

462. Nature (2014) PMID: 25079317
463. Quintanilha JCF, et al. Oncologist (2023) PMID: 36779536
464. Matsumoto A, et al. Oncol Rep (2020) PMID: 32236609
465. Tsukiyama T, et al. Nat Commun (2020) PMID: 32934222
466. Sugiyama T, et al. Exp. Cell Res. (2008) PMID: 18313049
467. Tee AR, et al. Curr. Biol. (2003) PMID: 12906785
468. Kwiatkowski DJ, et al. Eur J Hum Genet (2015) PMID: 25782670
469. Luo C, et al. Orphanet J Rare Dis (2021) PMID: 34217357
470. Wang T, et al. Cancer Biol Ther (2020) PMID: 31597506
471. Guo G, et al. Front Oncol (2020) PMID: 33575217
472. Espinosa M, et al. BMC Cancer (2018) PMID: 29764404
473. Chuang CK, et al. Int Urol Nephrol (2017) PMID: 28547571
474. Wagner AJ, et al. J. Clin. Oncol. (2010) PMID: 20048174
475. Dickson MA, et al. Int. J. Cancer (2013) PMID: 22927055
476. Wagner AJ, et al. J Clin Oncol (2021) PMID: 34637337
477. Dickson et al., 2021; ASCO Abstract 3111
478. Gordon EM, et al. Anticancer Res (2023) PMID: 37097693
479. Adib E, et al. Clin Cancer Res (2021) PMID: 33727259
480. Nassar AH, et al. Mol Cancer Ther (2020) PMID: 31653662
481. De S, et al. Gegenbaurs Morphol Jahrb (1986) PMID: 3032730
482. Wagle N, et al. N. Engl. J. Med. (2014) PMID: 25295501
483. Tannir NM, et al. Eur. Urol. (2016) PMID: 26626617
484. Maroto P, et al. J Natl Compr Canc Netw (2018) PMID: 29632054
485. Zureick AH, et al. BMJ Case Rep (2019) PMID: 31154346
486. Hu W, et al. Front Oncol (2020) PMID: 33344249
487. Perini GF, et al. Blood Cancer J (2016) PMID: 27176796
488. Voss MH, et al. Clin. Cancer Res. (2018) PMID: 30327302
489. Armenia J, et al. Nat. Genet. (2018) PMID: 29610475
490. Abida W, et al. JCO Precis Oncol (2017) PMID: 28825054
491. Inoki K, et al. Genes Dev. (2003) PMID: 12869586
492. Hodges AK, et al. Hum. Mol. Genet. (2001) PMID: 11741833
493. Int. J. Cancer (2006) PMID: 16206276
494. Li Y, et al. Mol. Cell. Biol. (2004) PMID: 15340059
495. Ann. N. Y. Acad. Sci. (1991) PMID: 2039135
496. Kandt RS, et al. Nat. Genet. (1992) PMID: 1303246
497. Cell (1993) PMID: 8269512
498. Curatolo P, et al. Lancet (2008) PMID: 18722871
499. Bailey MH, et al. Cell (2018) PMID: 29625053
500. Bolton KL, et al. Nat Genet (2020) PMID: 33106634
501. Schuermann JC, et al. Nature (2010) PMID: 20436459
502. Cho YS, et al. J. Biol. Chem. (2006) PMID: 16606617
503. Park UH, et al. J. Biol. Chem. (2011) PMID: 21047783
504. Inoue D, et al. J. Clin. Invest. (2013) PMID: 24216483
505. Abdel-Wahab O, et al. Cancer Cell (2012) PMID: 22897849
506. Br. J. Cancer (2013) PMID: 23736028
507. Jaiswal S, et al. N. Engl. J. Med. (2014) PMID: 25426837
508. Genovese G, et al. N. Engl. J. Med. (2014) PMID: 25426838
509. Xie M, et al. Nat. Med. (2014) PMID: 25326804
510. Acuna-Hidalgo R, et al. Am. J. Hum. Genet. (2017) PMID: 28669404
511. Severson EA, et al. Blood (2018) PMID: 29678827
512. Fuster JJ, et al. Circ. Res. (2018) PMID: 29420212
513. Hematology Am Soc Hematol Educ Program (2018) PMID: 30504320
514. Chabon JJ, et al. Nature (2020) PMID: 32269342
515. Razavi P, et al. Nat. Med. (2019) PMID: 31768066
516. Gruber et al., 2019; ASCO Abstract 3006
517. Peasland A, et al. Br. J. Cancer (2011) PMID: 21730979
518. Vassileva V, et al. Cancer Res. (2002) PMID: 12124347
519. Miquel C, et al. Oncogene (2007) PMID: 17384679
520. Jardim MJ, et al. Mol. Biol. Cell (2009) PMID: 19570909
521. Ha K, et al. Mol. Cancer Ther. (2011) PMID: 21566061
522. Liang Y, et al. World J Surg (2009) PMID: 19034564
523. Mordes DA, et al. Genes Dev. (2008) PMID: 18519640
524. Nature (2015) PMID: 25631445
525. Yaeger R, et al. Cancer Cell (2018) PMID: 29316426
526. Cancer Genome Atlas Research Network, et al. Nature (2017) PMID: 28112728
527. Hernandez L, et al. Cell Death Discov (2015) PMID: 28179987
528. Rodriguez-Berriguete G, et al. BMC Cancer (2015) PMID: 26507126
529. Pingoud-Meier C, et al. Clin. Cancer Res. (2003) PMID: 14695141
530. Galluzzi L, et al. Immunity (2016) PMID: 26885855
531. Tummers B, et al. Immunol. Rev. (2017) PMID: 28462525
532. Crowder RN, et al. Exp. Oncol. (2012) PMID: 23070000
533. Liu J, et al. Mol. Cancer (2011) PMID: 21801448
534. Graf RP, et al. Curr. Mol. Med. (2014) PMID: 24467204
535. Barbero S, et al. Cancer Res. (2009) PMID: 19383910
536. Zhang L, et al. Cell. Signal. (2015) PMID: 25446254
537. Su H, et al. Science (2005) PMID: 15746428
538. Henry CM, et al. Mol. Cell (2017) PMID: 28212752
539. Kober AM, et al. Cell Death Dis (2011) PMID: 21975294
540. Sun BK, et al. Oncol. Rep. (2011) PMID: 21152872
541. Soung YH, et al. Cancer Res. (2005) PMID: 15705878
542. Soung YH, et al. Oncogene (2005) PMID: 15531912
543. Mandruzzato S, et al. J. Exp. Med. (1997) PMID: 9271594
544. Kim HS, et al. Gastroenterology (2003) PMID: 12949717
545. Li C, et al. Mol Oncol (2014) PMID: 24816188
546. Ando M, et al. Cancer Sci. (2013) PMID: 23659359
547. Mancini A, et al. J. Biol. Chem. (2002) PMID: 11847211
548. Miyake S, et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PMID: 9653117
549. Masson K, et al. Biochem. J. (2006) PMID: 16780420
550. Singh AJ, et al. Proc. Natl. Acad. Sci. U.S.A. (2007) PMID: 17372230
551. Paolino M, et al. Nature (2014) PMID: 24553136
552. Patwardhan PP, et al. Oncotarget (2016) PMID: 26675259
553. Bazhenova et al., 2018; ESMO Abstract 4080
554. Shtiegman K, et al. Oncogene (2007) PMID: 17486068
555. Padrón D, et al. Cancer Res. (2007) PMID: 17699773
556. Hosaka T, et al. Anticancer Res. () PMID: 17695511
557. Han W, et al. Cancer Biol. Ther. (2006) PMID: 16969069
558. Yang S, et al. Cancer Res. (2006) PMID: 16849543
559. Sargin B, et al. Blood (2007) PMID: 17446348
560. Oshikawa G, et al. J. Biol. Chem. (2011) PMID: 21768087
561. Reindl C, et al. Clin. Cancer Res. (2009) PMID: 19276253
562. Taylor SJ, et al. Blood (2012) PMID: 22990016
563. Makishima H, et al. Leukemia (2012) PMID: 22246246
564. Kim SY, et al. Cancer Gene Ther. (2013) PMID: 23306613
565. Knight JF, et al. Br. J. Cancer (2008) PMID: 19002168
566. Bacher U, et al. Ann. Hematol. (2010) PMID: 20195608
567. Miyake S, et al. J. Biol. Chem. (1999) PMID: 10347229
568. Polzer H, et al. Exp. Hematol. (2013) PMID: 23127761
569. Levkowitz G, et al. Genes Dev. (1998) PMID: 9851973
570. Bunda S, et al. Cancer Res. (2013) PMID: 23400592
571. Andoniou CE, et al. EMBO J. (1994) PMID: 7925293
572. Aranaz P, et al. Haematologica (2012) PMID: 22315494
573. Fernandes MS, et al. J. Biol. Chem. (2010) PMID: 20622007
574. Grand FH, et al. Blood (2009) PMID: 19387008
575. Javadi M, et al. J. Biol. Chem. (2013) PMID: 23696637
576. Kassenbrock CK, et al. J. Biol. Chem. (2004) PMID: 15117950
577. Levkowitz G, et al. Mol. Cell (1999) PMID: 10635327
578. Loh ML, et al. Blood (2009) PMID: 19571318
579. Martinelli S, et al. Am. J. Hum. Genet. (2010) PMID: 20619386
580. Saito Y, et al. Leuk. Res. (2012) PMID: 22591685
581. Sanada M, et al. Nature (2009) PMID: 19620960
582. Score J, et al. Blood (2012) PMID: 22053108
583. Shiba N, et al. Leukemia (2011) PMID: 21494262
584. Standaert ML, et al. Biochemistry (2004) PMID: 15581361
585. Tan YH, et al. PLoS ONE (2010) PMID: 20126411
586. Thien CB, et al. Mol. Cell (2001) PMID: 11239464
587. Visser GD, et al. Exp. Cell Res. (2005) PMID: 16246327
588. Li M, et al. Cancer Res. (2016) PMID: 26676746
589. Wilson WH, et al. Nat. Med. (2015) PMID: 26193343
590. Davis RE, et al. Nature (2010) PMID: 20054396
591. Nagel D, et al. Oncotarget (2015) PMID: 26540570
592. Naylor TL, et al. Cancer Res. (2011) PMID: 21324920
593. Nguyen B, et al. Cell (2022) PMID: 35120664
594. Young RM, et al. Nat Rev Drug Discov (2013) PMID: 23449308
595. Rzymiski T, et al. Biochim. Biophys. Acta (2015) PMID: 26006748
596. Dale T, et al. Nat. Chem. Biol. (2015) PMID: 26502155
597. Mallinger A, et al. ACS Med Chem Lett (2016) PMID: 27326329
598. Koehler MF, et al. ACS Med Chem Lett (2016) PMID: 26985305
599. Schiemann K, et al. Bioorg. Med. Chem. Lett. (2016) PMID: 26852363
600. Menzl I, et al. Pharmaceuticals (Basel) (2019) PMID: 31248103
601. Firestein R, et al. Int. J. Cancer (2010) PMID: 19790197
602. Li YC, et al. Sci Adv (2021) PMID: 33523904
603. Bian J, et al. Cells (2020) PMID: 32961708
604. Firestein R, et al. Nature (2008) PMID: 18794900
605. Morris EJ, et al. Nature (2008) PMID: 18794899
606. Adler AS, et al. Cancer Res. (2012) PMID: 22345154
607. Méndez-Catalá CF, et al. Neoplasia (2013) PMID: 23908591
608. Tiffen JC, et al. Int. J. Cancer (2013) PMID: 23553099
609. Phillips JE, et al. Cell (2009) PMID: 19563753
610. Gombert WM, et al. PLoS ONE (2009) PMID: 19568426
611. Soto-Reyes E, et al. Oncogene (2010) PMID: 20101205
612. Woloszyńska-Read A, et al. Clin. Cancer Res. (2011) PMID: 21296871
613. Kemp CJ, et al. Cell Rep (2014) PMID: 24794443
614. Hadley KA, et al. Cell (2018) PMID: 29625048
615. Van Allen EM, et al. Cancer Discov (2014) PMID: 24265153
616. Hansford S, et al. JAMA Oncol (2015) PMID: 26182300
617. Majewski IJ, et al. J. Pathol. (2013) PMID: 23208944
618. Ding L, et al. Nature (2010) PMID: 20393555
619. Piao HL, et al. Nat. Cell Biol. (2014) PMID: 24509793
620. Qian J, et al. Clin. Chem. Lab. Med. (2014) PMID: 25153418
621. Ye Y, et al. Cancer Res. (2009) PMID: 19826047
622. Liu TX, et al. Nat. Med. (2007) PMID: 17159988

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX **References**

623. Chen XX, et al. Leuk. Res. (2014) PMID: 24685333
624. Li M, et al. Oncotarget (2016) PMID: 27129146
625. Kobiela A, et al. Nat. Rev. Mol. Cell Biol. (2004) PMID: 15366705
626. Bajpai S, et al. J. Biol. Chem. (2009) PMID: 19458087
627. Silvis MR, et al. Sci Signal (2011) PMID: 21610251
628. Inge LJ, et al. Mol. Cancer Ther. (2008) PMID: 18566211
629. Kertesz N, et al. Blood (2006) PMID: 16322467
630. Shi S, et al. J Pharm Sci (2012) PMID: 22411527
631. Li X, et al. PLoS ONE (2014) PMID: 25148033
632. Liu R, et al. BMC Cancer (2013) PMID: 23721559
633. Bhatia S, et al. Sci Rep (2016) PMID: 27941840
634. Scheinet JS, et al. Blood (2009) PMID: 18836096
635. Ferguson BD, et al. Sci Rep (2015) PMID: 26073592
636. Werner TL, et al. Invest New Drugs (2015) PMID: 26365907
637. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22011666
638. Pietanza MC, et al. J Thorac Oncol (2012) PMID: 22722787
639. Sinha UK, et al. Arch. Otolaryngol. Head Neck Surg. (2006) PMID: 17043250
640. Masood R, et al. Int. J. Cancer (2006) PMID: 16615113
641. Ferguson BD, et al. Growth Factors (2014) PMID: 25391996
642. Yavrouian EJ, et al. Arch. Otolaryngol. Head Neck Surg. (2008) PMID: 18794445
643. Liersch-Löhn B, et al. Int. J. Cancer (2016) PMID: 26414866
644. Hu F, et al. Tumour Biol. (2014) PMID: 24771266
645. Hasina R, et al. Cancer Res. (2013) PMID: 23100466
646. Li M, et al. Dig. Dis. Sci. (2011) PMID: 20686847
647. Yin J, et al. Anticancer Res. (2017) PMID: 28739744
648. Stephenson SA, et al. BMC Mol. Biol. (2001) PMID: 11801186
649. Liu W, et al. Cancer (2002) PMID: 11920461
650. McCall JL, et al. Mol. Cell. Biol. (2016) PMID: 27273865
651. Stremtizer S, et al. Mol. Cancer Ther. (2016) PMID: 27535973
652. Lv J, et al. Exp. Mol. Pathol. (2016) PMID: 27072105
653. Gujjarro-Muñoz I, et al. Med. Oncol. (2013) PMID: 23579861
654. Kumar SR, et al. Cancer Res. (2009) PMID: 19366806
655. Wu Q, et al. Pathol. Oncol. Res. (2004) PMID: 15029258
656. Berclaz G, et al. Oncol. Rep. () PMID: 12168060
657. Brantley-Sieders DM, et al. PLoS ONE (2011) PMID: 21935409
658. Huang G, et al. Int J Clin Exp Pathol (2015) PMID: 26191333
659. Pradeep S, et al. Cancer Cell (2015) PMID: 26481148
660. Alam SM, et al. Br. J. Cancer (2008) PMID: 18231102
661. Kumar SR, et al. Br. J. Cancer (2007) PMID: 17353927
662. Takai N, et al. Oncol. Rep. () PMID: 11295082
663. Dong LD, et al. Oncol Lett (2017) PMID: 28454369
664. Berclaz G, et al. Ann. Oncol. (2003) PMID: 12562648
665. Sharma GK, et al. Head Neck (2015) PMID: 24634162
666. Giaginis C, et al. Pathol. Oncol. Res. (2016) PMID: 26220827
667. Xuqing W, et al. Tumour Biol. (2012) PMID: 22528941
668. Ferguson BD, et al. PLoS ONE (2013) PMID: 23844053
669. Zheng MF, et al. Mol Med Rep (2012) PMID: 22684742
670. Chen T, et al. Tumour Biol. (2013) PMID: 23138393
671. Tu Y, et al. Clin Transl Oncol (2012) PMID: 22374425
672. Li M, et al. Mol. Biol. Rep. (2013) PMID: 23079712
673. Xia G, et al. Cancer Res. (2005) PMID: 15930280
674. Alam SM, et al. Gynecol. Oncol. (2009) PMID: 19356789
675. Özgür E, et al. Urol. Oncol. () PMID: 19272799
676. Pierscianek D, et al. Neuropathology (2017) PMID: 27388534
677. Pierscianek D, et al. Brain Tumor Pathol (2016) PMID: 26951238
678. Becerikli M, et al. Int. J. Cancer (2015) PMID: 25274141
679. Xia G, et al. Clin. Cancer Res. (2005) PMID: 15958611
680. Noren NK, et al. Cancer Res. (2007) PMID: 17483308
681. Cell (2008) PMID: 18394988
682. Salvucci O, et al. Adv. Cancer Res. (2012) PMID: 22588055
683. Pitulescu ME, et al. Genes Dev. (2010) PMID: 21078817
684. Nat. Rev. Cancer (2010) PMID: 20179713
685. Boberg DR, et al. Chem. Biol. Interact. (2013) PMID: 23063927
686. Cromer A, et al. Oncogene (2004) PMID: 14676830
687. Borad MJ, et al. PLoS Genet. (2014) PMID: 24550739
688. Ferby I, et al. Nat. Med. (2006) PMID: 16648858
689. Walsh AM, et al. J. Cell. Sci. (2013) PMID: 23868981
690. Romanuk TL, et al. BMC Genomics (2009) PMID: 19832994
691. Anastasi S, et al. Oncogene (2003) PMID: 12833145
692. Anastasi S, et al. Oncogene (2007) PMID: 17599051
693. Zhang X, et al. Nature (2007) PMID: 18046415
694. Anastasi S, et al. Oncogene (2005) PMID: 15856022
695. Fiorini M, et al. Oncogene (2002) PMID: 12226756
696. Frosi Y, et al. J. Cell Biol. (2010) PMID: 20421427
697. Ying H, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20351267
698. Reschke M, et al. Hepatology (2010) PMID: 20044804
699. Zhang YW, et al. Oncogene (2007) PMID: 16819504
700. Hackel PO, et al. Biol. Chem. (2001) PMID: 11843178
701. Fiorentino L, et al. Mol. Cell. Biol. (2000) PMID: 11003669
702. Sourbier C, et al. Cancer Cell (2014) PMID: 25490448
703. Srinivasan et al., 2014; EORTC-NCI-AACR Abstract 5
704. Schmidt C, et al. Semin Cell Dev Biol (2020) PMID: 31085323
705. Lau HD, et al. Am J Surg Pathol (2020) PMID: 31524643
706. Gardie B, et al. J Med Genet (2011) PMID: 21398687
707. Morris MR, et al. J. Clin. Pathol. (2004) PMID: 15220362
708. Lehtonen HJ, et al. J Med Genet (2006) PMID: 16155190
709. Tomlinson IP, et al. Nat. Genet. (2002) PMID: 11865300
710. Castro-Vega LJ, et al. Hum Mol Genet (2014) PMID: 24334767
711. Muller M, et al. Clin Genet (2017) PMID: 28300276
712. Fam Cancer (2011) PMID: 21404119
713. Forde C, et al. Eur Urol Oncol (2019) PMID: 31831373
714. Jia MY, et al. Neoplasia (2012) PMID: 25030436
715. Jiao Y, et al. Oncotarget (2012) PMID: 22869205
716. Duncan R, et al. Genes Dev. (1994) PMID: 8125259
717. Kim MJ, et al. Nat. Genet. (2003) PMID: 12819782
718. Weber A, et al. BMC Cancer (2008) PMID: 19087307
719. Jang M, et al. Oncogene (2009) PMID: 19219071
720. Zheng Y, et al. Int. J. Biochem. Cell Biol. (2011) PMID: 21855647
721. Jacob AG, et al. J. Biol. Chem. (2014) PMID: 24798327
722. Rabenhorst U, et al. Hepatology (2009) PMID: 19637194
723. Singer S, et al. Cancer Res. (2009) PMID: 19258502
724. Sahm F, et al. Acta Neuropathol. (2012) PMID: 22588899
725. Bettgowda C, et al. Science (2011) PMID: 21817013
726. Baumgarten P, et al. Neuropathol. Appl. Neurobiol. (2014) PMID: 24117486
727. Chan AK, et al. Mod. Pathol. (2014) PMID: 24030748
728. Xi H, et al. Int. J. Cancer (2000) PMID: 10956389
729. Guichard C, et al. Nat. Genet. (2012) PMID: 22561517
730. Nature (2012) PMID: 22960745
731. Augert A, et al. J Thorac Oncol (2017) PMID: 28007623
732. Ardeshir-Larijani F, et al. Clin Lung Cancer (2018) PMID: 29627316
733. Hillman RT, et al. Nat Commun (2018) PMID: 29950560
734. Abudurehman A, et al. J. Cancer Res. Clin. Oncol. (2018) PMID: 29532228
735. Vicent GP, et al. Genes Dev. (2011) PMID: 21447625
736. Hannibal MC, et al. Am. J. Med. Genet. A (2011) PMID: 21671394
737. Fagan RJ, et al. Cancer Lett. (2019) PMID: 31128216
738. Chakraborty R, et al. Oncotarget (2017) PMID: 28512266
739. Brennan L, et al. Cell Rep (2016) PMID: 27760319
740. Goetz EM, et al. Cancer Res. (2014) PMID: 25320010
741. Ercan D, et al. Cancer Discov (2012) PMID: 22961667
742. Patil CG, et al. Neuro-oncology (2013) PMID: 23115159
743. Pelloski CE, et al. Clin. Cancer Res. (2006) PMID: 16818690
744. Feldkamp MM, et al. Neurosurgery (1999) PMID: 10598712
745. Zheng Y, et al. Cell Death Dis (2018) PMID: 29415992
746. Slattery ML, et al. Carcinogenesis (2012) PMID: 23027623
747. Ngan et al., 2017; AACR 5712
748. Bronner CE, et al. Nature (1994) PMID: 8145827
749. Gazzoli I, et al. Cancer Res. (2002) PMID: 12124320
750. Gatalica Z, et al. Cancer Epidemiol. Biomarkers Prev. (2014) PMID: 25392179
751. Chen Y, et al. Int. J. Oncol. (2003) PMID: 12684669
752. Wilczak W, et al. Carcinogenesis (2017) PMID: 27803051
753. Kadyrov FA, et al. Cell (2006) PMID: 16873062
754. Raevaara TE, et al. Gastroenterology (2005) PMID: 16083711
755. Mensenkamp AR, et al. Gastroenterology (2014) PMID: 24333619
756. Joly MO, et al. Hum. Mutat. (2015) PMID: 25504677
757. McConechy MK, et al. Gynecol. Oncol. (2015) PMID: 25636458
758. Martin SA, et al. EMBO Mol Med (2009) PMID: 20049736
759. Kamal NS, et al. Clin. Cancer Res. (2010) PMID: 20145178
760. Pierceall WE, et al. Ann. Oncol. (2012) PMID: 22269178
761. Guedes LB, et al. Clin Cancer Res (2017) PMID: 28790115
762. Prtalo A, et al. J. Urol. (2005) PMID: 16217293
763. Velasco A, et al. Cancer Biol. Ther. () PMID: 12432247
764. Burger M, et al. J. Mol. Med. (2006) PMID: 16924473
765. Velasco A, et al. Cancer (2002) PMID: 11857301
766. Christensen LL, et al. Fam. Cancer (2009) PMID: 19697156
767. Drost M, et al. Hum. Mutat. (2012) PMID: 22102614
768. Gammie AE, et al. Genetics (2007) PMID: 17720936
769. Houleberghs H, et al. Proc. Natl. Acad. Sci. U.S.A. (2016) PMID: 26951660
770. Cell Res. (2008) PMID: 18157157
771. Lützen A, et al. Mutat. Res. (2008) PMID: 18822302
772. Martinez SL, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20176959
773. Mastrocola AS, et al. Hum. Mutat. (2010) PMID: 20672385
774. Ollila S, et al. Gastroenterology (2006) PMID: 17101317
775. Ollila S, et al. Hum. Mutat. (2008) PMID: 18951462
776. Petersen SM, et al. BMC Med. Genet. (2013) PMID: 24090359

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1637928-01

APPENDIX
References

777. Ward R, et al. J. Cancer Res. Clin. Oncol. (2002) PMID: 12200596
778. Wielders EA, et al. J. Med. Genet. (2014) PMID: 24501230
779. PLoS Curr (2013) PMID: 24056992
780. Dietlein F, et al. Cancer Discov (2014) PMID: 24556366
781. Takahashi M, et al. J. Biol. Chem. (2011) PMID: 21285347
782. Park JM, et al. PLoS ONE (2013) PMID: 23724141
783. Plaschke J, et al. Int J Colorectal Dis (2012) PMID: 22249440
784. Benachene N, et al. Int. J. Cancer (1998) PMID: 9650548
785. Kawakami T, et al. Biochem. Biophys. Res. Commun. (2004) PMID: 15541380
786. Benachene N, et al. Br. J. Cancer (1999) PMID: 10098729
787. Plaschke J, et al. Cancer Res. (2004) PMID: 14871813
788. Nogueira GA, et al. Int. J. Cancer (2015) PMID: 25598504
789. Xu XL, et al. Genet. Mol. Res. (2015) PMID: 25966119
790. Dong X, et al. Oncologist (2011) PMID: 21212431
791. Laghi L, et al. Clin. Cancer Res. (2012) PMID: 22496206
792. De' Angelis GL, et al. Acta Biomed (2018) PMID: 30561401
793. Haugen AC, et al. Cancer Res. (2008) PMID: 18922920
794. van Oers JM, et al. Oncogene (2014) PMID: 24013230
795. Lee SY, et al. Gastroenterology (2010) PMID: 20708618
796. Watson MM, et al. Br. J. Cancer (2014) PMID: 24691426
797. Campregher C, et al. PLoS One (2012) PMID: 23209772
798. Miao HK, et al. Int J Clin Exp Pathol (2015) PMID: 26617824
799. Morak M, et al. Fam. Cancer (2017) PMID: 28528517
800. Duraturo F, et al. Int. J. Cancer (2011) PMID: 21128252
801. Reeves SG, et al. Cancer Epidemiol (2012) PMID: 21974800
802. Berndt SI, et al. Int. J. Cancer (2007) PMID: 17205513
803. Adam R, et al. Am. J. Hum. Genet. (2016) PMID: 27476653
804. Yang X, et al. PLoS ONE (2015) PMID: 25927356
805. Vogelsang M, et al. PLoS ONE (2012) PMID: 22623965
806. Jafary F, et al. Asian Pac. J. Cancer Prev. (2012) PMID: 23464402
807. Hirata H, et al. J. Urol. (2008) PMID: 18355840
808. Ohmiya N, et al. Gene (2001) PMID: 11470537
809. Le et al., 2015; ASCO Abstract LBA100
810. Li F, et al. Cell (2013) PMID: 23622243
811. Wu H, et al. PLoS ONE (2011) PMID: 21720545
812. Edellbrock MA, et al. Mutat. Res. (2012) PMID: 23391514
813. Berends MJ, et al. Am. J. Hum. Genet. (2002) PMID: 11709755
814. Warren JJ, et al. Mol. Cell (2007) PMID: 17531815
815. Geng H, et al. J. Biol. Chem. (2012) PMID: 22277660
816. Wimmer K, et al. Hum. Genet. (2008) PMID: 18709565
817. Wimmer K, et al. J. Med. Genet. (2014) PMID: 24737826
818. Scott RH, et al. Nat Clin Pract Oncol (2007) PMID: 17259933
819. Ripberger T, et al. Haematologica (2010) PMID: 20015892
820. Baris HN, et al. Pediatr Blood Cancer (2016) PMID: 26544533
821. Yen WC, et al. Clin. Cancer Res. (2015) PMID: 25934888
822. Hu W, et al. Cancer Res. (2014) PMID: 24743243
823. Konishi J, et al. Cancer Res. (2007) PMID: 17804716
824. Xiao Y, et al. Oncogene (2011) PMID: 20838375
825. Ferrarotto et al., 2020; ESMO Abstract 919MO
826. Hu Zl, et al. Cancer Med (2019) PMID: 31347292
827. Carvalho FL, et al. Prostate (2014) PMID: 24737393
828. Kim AR, et al. Int J Clin Exp Pathol (2019) PMID: 31934201
829. Penton AL, et al. Semin. Cell Dev. Biol. (2012) PMID: 22306179
830. Kopan R, et al. Cell (2009) PMID: 19379690
831. Deregowski V, et al. J. Bone Miner. Res. (2006) PMID: 16869730
832. Beà S, et al. Proc. Natl. Acad. Sci. U.S.A. (2013) PMID: 24145436
833. McDaniel R, et al. Am. J. Hum. Genet. (2006) PMID: 16773578
834. Lin L, et al. Cancer Res. (2010) PMID: 20068176
835. Hong DS, et al. Ann. Oncol. (2019) PMID: 30624546
836. Vaishnavi A, et al. Nat. Med. (2013) PMID: 24162815
837. Doebele RC, et al. Cancer Discov (2015) PMID: 26216294
838. Tatematsu T, et al. Mol Clin Oncol (2014) PMID: 25054037
839. Wong V, et al. J. Natl. Cancer Inst. (2016) PMID: 26563356
840. Mody RJ, et al. JAMA (2015) PMID: 26325560
841. Sartore-Bianchi A, et al. J. Natl. Cancer Inst. (2016) PMID: 26563355
842. Farago AF, et al. J Thorac Oncol (2015) PMID: 26565381
843. Drilon A, et al. Cancer Discov (2017) PMID: 28183697
844. Drilon A, et al. N. Engl. J. Med. (2018) PMID: 29466156
845. Laetsch TW, et al. Lancet Oncol. (2018) PMID: 29606586
846. Hempel D, et al. Oncologist (2020) PMID: 32323889
847. Philos. Trans. R. Soc. Lond., B, Biol. Sci. (2006) PMID: 16939974
848. Guiton M, et al. J. Biol. Chem. (1994) PMID: 7982951
849. Massa SM, et al. J. Clin. Invest. (2010) PMID: 20407211
850. Huang EJ, et al. Annu. Rev. Biochem. (2003) PMID: 12676795
851. Schramm A, et al. Cancer Lett. (2005) PMID: 15921851
852. Geiger TR, et al. Cancer Res. (2005) PMID: 16103047
853. Drost M, et al. Hum. Mutat. (2013) PMID: 24027009
854. Gill S, et al. Clin. Cancer Res. (2005) PMID: 16166421
855. Batte BA, et al. Gynecol. Oncol. (2014) PMID: 24933100
856. Win AK, et al. J. Natl. Cancer Inst. (2012) PMID: 22933731
857. Soni A, et al. OMICS (2011) PMID: 21348638
858. Bansal A, et al. Indian J. Med. Res. (2012) PMID: 23168703
859. Chen Y, et al. Cancer Res. (2001) PMID: 11358834
860. Norris AM, et al. Prostate (2007) PMID: 17044039
861. Norris AM, et al. Cancer Epidemiol. Biomarkers Prev. (2009) PMID: 19124481
862. Fukuhara S, et al. Oncotarget (2015) PMID: 26036629
863. Shimodaira H, et al. Proc. Natl. Acad. Sci. U.S.A. (2003) PMID: 12601175
864. Gibson SL, et al. Cancer Lett. (2006) PMID: 16426742
865. van Oers JM, et al. Proc. Natl. Acad. Sci. U.S.A. (2010) PMID: 20624957
866. Nakagawa H, et al. Cancer Res. (2004) PMID: 15256438
867. Senter L, et al. Gastroenterology (2008) PMID: 18602922
868. Wang Z, et al. Cancer Lett. (2014) PMID: 23981578
869. Clendenning M, et al. J. Med. Genet. (2008) PMID: 18178629
870. Hendriks YM, et al. Gastroenterology (2006) PMID: 16472587
871. Ramchander NC, et al. BMC Med. Genet. (2017) PMID: 28381238
872. Fam. Cancer (2005) PMID: 16136382
873. Hegde M, et al. Genet. Med. (2014) PMID: 24310308
874. De Rosa M, et al. Oncogene (2000) PMID: 10763829
875. Mavrikis M, et al. Hum. Mol. Genet. (2006) PMID: 16963469
876. de Jossineau C, et al. Hum. Mol. Genet. (2014) PMID: 24865460
877. Beristain AG, et al. Oncogene (2015) PMID: 24662820
878. Wang L, et al. Genes Chromosomes Cancer (2015) PMID: 26031761
879. Torres-Mora J, et al. Am. J. Surg. Pathol. (2014) PMID: 24145644
880. Wang S, et al. Sci Rep (2016) PMID: 27995993
881. Bossis I, et al. Endocrinology (2004) PMID: 15331577
882. Kirschner LS, et al. Nat. Genet. (2000) PMID: 10973256
883. Carney JA, et al. Am. J. Surg. Pathol. (2014) PMID: 24805858
884. Bertherat J, et al. J Clin Endocrinol Metab (2009) PMID: 19293268
885. Horvath A, et al. Clin Cancer Res (2008) PMID: 18223213
886. Salpea P, et al. J Clin Endocrinol Metab (2014) PMID: 24170103
887. Maleszewski JJ, et al. Am. J. Surg. Pathol. (2014) PMID: 24618615
888. He J, et al. Oncotarget (2017) PMID: 29262613
889. Roque A, et al. J Med Case Rep (2019) PMID: 31874650
890. Ichimura K, et al. Oncogene (2006) PMID: 16205629
891. Mulholland PJ, et al. Cell Cycle (2006) PMID: 16582634
892. Yin D, et al. Mol. Cancer Res. (2009) PMID: 19435819
893. Yang G, et al. Gastroenterology (2010) PMID: 19686745
894. Bian Y, et al. Biochem. Biophys. Res. Commun. (2012) PMID: 22569043
895. Yu F, et al. Oncol. Rep. (2014) PMID: 24398626
896. Ebersole TA, et al. Nat. Genet. (1996) PMID: 8589716
897. Hardy RJ, et al. J. Neurosci. (1996) PMID: 8987822
898. Yang G, et al. Cell Cycle (2011) PMID: 21768773
899. Varela L, et al. Nature (2011) PMID: 21248752
900. Chen Z, et al. Biochem Biophys Res Commun (2018) PMID: 29522714
901. Chen BY, et al. Blood (2020) PMID: 32202636
902. Liu W, et al. Medicine (Baltimore) (2015) PMID: 26559293
903. Sun XJ, et al. J. Biol. Chem. (2005) PMID: 16118227
904. Faber PW, et al. Hum. Mol. Genet. (1998) PMID: 9700202
905. Al Sarakbi W, et al. BMC Cancer (2009) PMID: 19698110
906. Parker H, et al. Leukemia (2016) PMID: 27282254
907. Zhang J, et al. Nature (2012) PMID: 22237106
908. McKinney M, et al. Cancer Discov (2017) PMID: 28122867
909. Moffitt AB, et al. J. Exp. Med. (2017) PMID: 28424246
910. Zhu X, et al. Nat. Genet. (2014) PMID: 24509477
911. Lu C, et al. Science (2016) PMID: 27174990
912. Shi C, et al. Oncogene (2022) PMID: 35393542
913. Park JW, et al. Cancer Med (2022) PMID: 35274815
914. Ormanns S, et al. Int J Mol Sci (2017) PMID: 28534865
915. Fei N, et al. Clin Transl Sci (2021) PMID: 34002944
916. Bachet JB, et al. Ann. Oncol. (2012) PMID: 22377565
917. Ziemke M, et al. Lung Cancer (2017) PMID: 28577946
918. Kassardjian A, et al. Pancreas (2020) PMID: 32897998
919. Pen SL, et al. Radiother Oncol (2021) PMID: 33667587
920. Witkiewicz AK, et al. Nat Commun (2015) PMID: 25855536
921. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
922. Churi CR, et al. PLoS ONE (2014) PMID: 25536104
923. Takeda et al., 2022; ASCO GI Abstract 642
924. Liu X, et al. Clin. Chem. (2014) PMID: 24821835
925. Maru D, et al. Oncogene (2004) PMID: 14647445

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1637928-01**
APPENDIX
References

926. Wang K, et al. *Oncologist* (2015) PMID: 26336083
927. Izeradjene K, et al. *Cancer Cell* (2007) PMID: 17349581
928. Bardeesy N, et al. *Genes Dev.* (2006) PMID: 17114584
929. Springer S, et al. *Gastroenterology* (2015) PMID: 26253305
930. Blackford A, et al. *Clin. Cancer Res.* (2009) PMID: 19584151
931. Yan P, et al. *Clin. Cancer Res.* (2016) PMID: 26861460
932. Kozak MM, et al. *J. Clin. Pathol.* (2015) PMID: 25681512
933. Roth AD, et al. *J. Natl. Cancer Inst.* (2012) PMID: 23104212
934. Davison JM, et al. *Am. J. Surg. Pathol.* (2014) PMID: 24618609
935. Kim YH, et al. *Ann. Oncol.* (2004) PMID: 15033661
936. Xiangming C, et al. *Clin. Cancer Res.* (2001) PMID: 11234879
937. Singhi AD, et al. *Am. J. Surg. Pathol.* (2015) PMID: 25634752
938. Natsugoe S, et al. *Clin. Cancer Res.* (2002) PMID: 12060625
939. de Kruijf EM, et al. *Ann. Oncol.* (2013) PMID: 23022998
940. Shipitsin M, et al. *Br. J. Cancer* (2014) PMID: 25032733
941. *Nat. Rev. Mol. Cell Biol.* (2012) PMID: 22992590
942. *Cell* (2008) PMID: 18662538
943. Massagué J, et al. *Genes Dev.* (2005) PMID: 16322555
944. Morén A, et al. *Oncogene* (2000) PMID: 10980615
945. Xu J, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2000) PMID: 10781087
946. Luo K, et al. *Genes Dev.* (1999) PMID: 10485843
947. Jones JB, et al. *Nucleic Acids Res.* (2000) PMID: 10871368
948. Fink SP, et al. *Cancer Res.* (2001) PMID: 11196171
949. De Bosscher K, et al. *Biochem. J.* (2004) PMID: 14715079
950. Shi Y, et al. *Nature* (1997) PMID: 9214508
951. Miyaki M, et al. *Oncogene* (1999) PMID: 10340381
952. Prokova V, et al. *Biochemistry* (2007) PMID: 17994767
953. Wu JW, et al. *J. Biol. Chem.* (2001) PMID: 11274206
954. Ding L, et al. *J. Clin. Invest.* (2009) PMID: 19139564
955. Kuang C, et al. *Oncogene* (2004) PMID: 14647410
956. Watanabe M, et al. *EMBO Rep.* (2000) PMID: 11265759
957. Houlston R, et al. *Hum. Mol. Genet.* (1998) PMID: 9811934
958. Woodford-Richens K, et al. *Gut* (2000) PMID: 10764709
959. Howe JR, et al. *J. Med. Genet.* (2004) PMID: 15235019
960. Brosens LA, et al. *World J. Gastroenterol.* (2011) PMID: 22171123
961. Kalia SS, et al. *Genet. Med.* (2017) PMID: 27854360
962. Stephens PJ, et al. *J. Clin. Invest.* (2013) PMID: 23778141
963. Rossi D, et al. *J. Exp. Med.* (2012) PMID: 22891273
964. Shi Y, et al. *Genes Dev.* (2001) PMID: 11331609
965. Ariyoshi M, et al. *Genes Dev.* (2003) PMID: 12897056
966. Kuroda K, et al. *Immunity* (2003) PMID: 12594956
967. Oswald F, et al. *EMBO J.* (2002) PMID: 12374742
968. Yingling JM, et al. *Nat Rev Drug Discov* (2004) PMID: 15573100
969. Bharathy S, et al. *Cancer Res.* (2008) PMID: 18339844
970. Bogdahn U, et al. *Neuro-oncology* (2011) PMID: 20980335
971. Strimpakos AS, et al. *JOP* (2012) PMID: 22797386
972. Akhurst RJ, et al. *Nat Rev Drug Discov* (2012) PMID: 23000686
973. Yamashita S, et al. *Cancer Res.* (2008) PMID: 18381416
974. Jackson RS, et al. *Neoplasia* (2012) PMID: 22904682
975. Markowitz S, et al. *Science* (1995) PMID: 7761852
976. Lücke CD, et al. *Cancer Res.* (2001) PMID: 11212236
977. Attias D, et al. *Circulation* (2009) PMID: 19996017
978. Mizuguchi T, et al. *Nat. Genet.* (2004) PMID: 15235604
979. Horbelt D, et al. *J. Cell. Sci.* (2010) PMID: 21098638
980. Loeys BL, et al. *Nat. Genet.* (2005) PMID: 15731757
981. Lu SL, et al. *Nat. Genet.* (1998) PMID: 9590282
982. Loeys BL, et al. *N. Engl. J. Med.* (2006) PMID: 16928994
983. Watanabe Y, et al. *Am. J. Med. Genet. A* (2008) PMID: 19006214
984. Togashi Y, et al. *Reshio Ketsueki* (2008) PMID: 18800617
985. Arslan-Kirchner M, et al. *Eur. J. Hum. Genet.* (2011) PMID: 21522183
986. Hirai H, et al. *Cancer Biol. Ther.* (2010) PMID: 20107315
987. Bridges KA, et al. *Clin. Cancer Res.* (2011) PMID: 21799033
988. Rajeshkumar NV, et al. *Clin. Cancer Res.* (2011) PMID: 21389100
989. Osman AA, et al. *Mol. Cancer Ther.* (2015) PMID: 25504633
990. Xu L, et al. *Mol. Cancer Ther.* (2002) PMID: 12489850
991. Xu L, et al. *Mol. Med.* (2001) PMID: 11713371
992. Camp ER, et al. *Cancer Gene Ther.* (2013) PMID: 23470564
993. Kim SS, et al. *Nanomedicine* (2015) PMID: 25240597
994. Pirolo KF, et al. *Mol. Ther.* (2016) PMID: 27357628
995. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27601554
996. Moore et al., 2019; ASCO Abstract 5513
997. Leijen S, et al. *J. Clin. Oncol.* (2016) PMID: 27998224
998. Oza et al., 2015; ASCO Abstract 5506
999. Lee J, et al. *Cancer Discov* (2019) PMID: 31315834
1000. Méndez E, et al. *Clin. Cancer Res.* (2018) PMID: 29535125
1001. Seligmann JF, et al. *J Clin Oncol* (2021) PMID: 34538072
1002. Gourley et al., 2016; ASCO Abstract 5571
1003. Park H, et al. *ESMO Open* (2022) PMID: 36084396
1004. Kumar A, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2011) PMID: 21949389
1005. Schlomm T, et al. *Mod. Pathol.* (2008) PMID: 18552821
1006. Liu W, et al. *Cancer* (2013) PMID: 23609948
1007. Brown CJ, et al. *Nat. Rev. Cancer* (2009) PMID: 19935675
1008. Joerger AC, et al. *Annu. Rev. Biochem.* (2008) PMID: 18410249
1009. Kato S, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2003) PMID: 12826609
1010. Kamada R, et al. *J. Biol. Chem.* (2011) PMID: 20978130
1011. Zerdoumi Y, et al. *Hum. Mol. Genet.* (2017) PMID: 28472496
1012. Yamada H, et al. *Carcinogenesis* (2007) PMID: 17690113
1013. Bougeard G, et al. *J. Clin. Oncol.* (2015) PMID: 26014290
1014. Sorrell AD, et al. *Mol Diagn Ther* (2013) PMID: 23355100
1015. Nichols KE, et al. *Cancer Epidemiol. Biomarkers Prev.* (2001) PMID: 11219776
1016. Kleihues P, et al. *Am. J. Pathol.* (1997) PMID: 9006316
1017. Gonzalez KD, et al. *J. Clin. Oncol.* (2009) PMID: 19204208
1018. Lalloo F, et al. *Lancet* (2003) PMID: 12672316
1019. Mandelker D, et al. *Ann. Oncol.* (2019) PMID: 31050713
1020. Cui X, et al. *Mutat. Res.* (1999) PMID: 10422536
1021. O'Regan P, et al. *J. Biol. Chem.* (2001) PMID: 11301337
1022. Deans B, et al. *Cancer Res.* (2003) PMID: 14678973
1023. *Nature* (2011) PMID: 21720365
1024. *Nature* (2014) PMID: 25079552
1025. Brennan CW, et al. *Cell* (2013) PMID: 24120142
1026. Cancer Genome Atlas Research Network, et al. *N. Engl. J. Med.* (2013) PMID: 23634996
1027. Cancer Genome Atlas Research Network, et al. *Nature* (2017) PMID: 28052061
1028. Rafii S, et al. *Hum. Mol. Genet.* (2002) PMID: 12023985
1029. Lin WY, et al. *J. Med. Genet.* (2011) PMID: 21632523
1030. Park DJ, et al. *Am. J. Hum. Genet.* (2012) PMID: 22464251
1031. Pooley KA, et al. *Cancer Epidemiol. Biomarkers Prev.* (2008) PMID: 19064565
1032. Loizidou MA, et al. *Breast Cancer Res. Treat.* (2008) PMID: 18188695
1033. Brooks J, et al. *Cancer Epidemiol. Biomarkers Prev.* (2008) PMID: 18398049
1034. Silva SN, et al. *Cancer Epidemiol* (2010) PMID: 20004634
1035. Yu KD, et al. *Breast Cancer Res. Treat.* (2010) PMID: 20127279
1036. He Y, et al. *PLoS ONE* (2014) PMID: 24621646
1037. Li D, et al. *Cancer Res.* (2006) PMID: 16540687
1038. Jiao L, et al. *Am. J. Gastroenterol.* (2008) PMID: 17986315
1039. Yin M, et al. *PLoS ONE* (2011) PMID: 21647442
1040. Curtin K, et al. *Cancer Epidemiol. Biomarkers Prev.* (2009) PMID: 19690184
1041. Figueroa JD, et al. *Carcinogenesis* (2007) PMID: 17557904
1042. Cartwright R, et al. *Nucleic Acids Res.* (1998) PMID: 9628903
1043. Johnson RD, et al. *Nature* (1999) PMID: 10517641
1044. Griffin CS, et al. *Nat. Cell Biol.* (2000) PMID: 11025669
1045. Sale JE, et al. *Nature* (2001) PMID: 11528482
1046. Miller KA, et al. *J. Biol. Chem.* (2002) PMID: 11744692
1047. Hilbers FS, et al. *Hum. Mutat.* (2016) PMID: 27233470
1048. Marabelle et al., 2019; ESMO Abstract 11920
1049. Antonarakis et al., 2020; DOI: 10.1200/PO.19.00399
1050. Schweizer et al., 2020; DOI: 10.1200/PO.19.00383
1051. Subramanian et al., 2020; ESMO Abstract 1399P
1052. Andre et al., 2021; ASCO GI Abstract 9
1053. Gabrail et al., 2019; ASCO Abstract 2560
1054. Berton et al., 2021; ASCO Abstract 2564
1055. Andre et al., 2021; ESMO GI Abstract SO-9
1056. Piha-Paul et al., 2018; AACR Abstract A096
1057. Nakamura et al., 2020; DOI: 10.1200/PO.19.00368
1058. de Bono et al., 2020; ASCO GU Abstract 119
1059. Smith et al., 2019; ESMO Abstract LBA50
1060. Sen T, et al. *Cancer Res.* (2017) PMID: 28490518
1061. Del Conte G, et al. *Br. J. Cancer* (2014) PMID: 25025963
1062. Anderson BJ, et al. *Diabetes Educ* () PMID: 3208634
1063. Kondrashova O, et al. *Cancer Discov* (2017) PMID: 28588062
1064. Hussain M, et al. (2020) PMID: 32955174
1065. Antonarakis et al., 2019; ASCO Abstract 5045
1066. Price MJ, et al. *JCO Precis Oncol* (2022) PMID: 35476551
1067. Yu et al., 2022; ESMO Abstract 1362MO
1068. Karzai F, et al. *J Immunother Cancer* (2018) PMID: 30514390
1069. Yu et al., 2019; ASCO GU Abstract 145
1070. Yu et al., 2020; ASCO GU Abstract 100
1071. Reichert et al., 2022; ASCO GU Abstract 88
1072. Saad et al., 2022; ESMO Abstract 13570
1073. Maio et al., 2021; AACR Abstract CT178
1074. Clarke et al., 2023; ASCO GU Abstract LBA16
1075. Clarke et al., 2022; DOI: 10.1056/EVIDoa2200043
1076. Hussain et al., 2022; ASCO Abstract 5018
1077. Efsthathiou et al., 2023; ASCO GU Abstract 170
1078. Sandhu et al., 2022; ASCO Abstract 5020
1079. Clarke N, et al. *Lancet Oncol.* (2018) PMID: 29880291
1080. Ayers et al., 2016; ASCO-SITC Abstract P60

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1637928-01**
APPENDIX
References

1081. Diaz et al., 2016; ASCO Abstract 3003
1082. Fader et al., 2016; SGO Abstract 3
1083. Le et al., 2016; ASCO GI Abstract 195
1084. Chao et al., 2020; ASCO GI Abstract 430
1085. Marcus L, et al. Clin Cancer Res (2021) pmid: 34083238
1086. Lee et al., 2018; doi/full/10.1200/PO.17.00097
1087. Petrylak et al., 2023; ASCO GU Abstract 19
1088. Graff et al., 2020; ASCO GU Abstract 15
1089. Berry et al., 2020; ASCO GU Abstract 102
1090. Antonarakis ES, et al. J. Clin. Oncol. (2020) pmid: 31774688
1091. Yu EY, et al. Eur Urol (2022) pmid: 36055895
1092. Chin et al., 2022; ASCO Abstract 510
1093. Salhab et al., 2018; DOI: 10.15761/HMO.1000169
1094. Kristeleit et al., 2014; ASCO Abstract 2573
1095. Domcheck et al., 2016; ASCO Abstract 4110
1096. Bryce et al., 2023; ASCO Genitourinary Abstract 18
1097. Abida W, et al. J. Clin. Oncol. (2020) pmid: 32795228
1098. Carneiro et al., 2022; ASCO GU Abstract 128
1099. Pook et al., 2022; ASCO GU Abstract 95
1100. Hochster et al., 2017; ASCO Abstract 673
1101. Fleming et al., 2018; ASCO Abstract 5585
1102. Petrylak DP, et al. Clin Cancer Res (2021) pmid: 33568344
1103. Powles T, et al. Nat Med (2022) pmid: 35013615
1104. Agarwal N, et al. Lancet Oncol (2022) pmid: 35690072
1105. Jung KH, et al. Clin. Cancer Res. (2019) pmid: 30770348
1106. Bang et al., 2018; ASCO Abstract 92
1107. Kwan EM, et al. Eur Urol (2021) pmid: 34493414
1108. Fakhrejahani et al., 2017; ASCO Abstract 5037
1109. Verschraegen et al., 2016; ASCO Abstract 9036
1110. Chung et al., 2016; ASCO Abstract 4009
1111. Patel et al., 2016; ESMO Abstract 777PD
1112. Hassan et al., 2016; ASCO Abstract 8503
1113. Disis et al., 2016; ASCO Abstract 5533
1114. Dirix et al., 2016; SABCS Abstract 51-04
1115. Larkin et al., 2016; ESMO Abstract 775PD
1116. Le Tourneau et al., 2016; ASCO Abstract 4516
1117. Fakhrejahani et al., 2017; ASCO GU Abstract 159
1118. Rajan et al., 2016; ASCO Abstract e20106
1119. Migden MR, et al. N. Engl. J. Med. (2018) pmid: 29863979
1120. Stratigos et al., 2020; EMSO Abstract LBA47
1121. Lewis et al. 2020; doi: 10.1136/jitc-2020-SITC2020.0428
1122. Fizazi et al., 2019; ESMO Abstract LBA52
1123. Hotte et al., 2019; ESMO Abstract LBA51
1124. Lim EA, et al. Clin Cancer Res (2022) pmid: 36044531
1125. Wang L, et al. Transl Androl Urol (2021) pmid: 34984194
1126. Pokhrel A, et al. J Investig Med High Impact Case Rep (2022) pmid: 35473437
1127. Pietrantonio et al., 2023; ASCO GI Abstract 358
1128. Lee et al., 2023; ASCO GI abstract 401
1129. Subudhi SK, et al. J Immunother Cancer (2021) pmid: 34663638
1130. Cappuzzo F, et al. Lancet Oncol. (2010) pmid: 20493771
1131. Zhong WZ, et al. J. Clin. Oncol. (2019) pmid: 31194613
1132. Petrelli F, et al. Clin Lung Cancer (2012) pmid: 22056888
1133. Hainsworth JD, et al. J. Clin. Oncol. (2018) pmid: 29320312
1134. Nie KK, et al. Chin Med J (Engl) (2018) pmid: 29998897
1135. Gravis G, et al. Ann. Oncol. (2008) pmid: 18467313
1136. Nabhan C, et al. Urology (2009) pmid: 19616281
1137. Han JY, et al. J. Clin. Oncol. (2012) pmid: 22370314
1138. Maemondo M, et al. N. Engl. J. Med. (2010) pmid: 20573926
1139. Mitsudomi T, et al. Lancet Oncol. (2010) pmid: 20022809
1140. Mok TS, et al. N. Engl. J. Med. (2009) pmid: 19692680
1141. Qi WX, et al. Curr Med Res Opin (2015) pmid: 25329826
1142. Zhao H, et al. J Thorac Oncol (2015) pmid: 25546556
1143. Wang J, et al. Int. J. Cancer (2019) pmid: 30255937
1144. Baik CS, et al. J Thorac Oncol (2015) pmid: 26398831
1145. Joensuu G, et al. Acta Oncol (2012) pmid: 22150168
1146. Pezaro C, et al. Am. J. Clin. Oncol. (2009) pmid: 19363437
1147. Joensuu G, et al. Int. J. Radiat. Oncol. Biol. Phys. (2010) pmid: 20004525
1148. Boccardo F, et al. Oncology (2008) pmid: 18714171
1149. Konstantinopoulos et al., 2018; ASCO Abstract 106
1150. Smith MR, et al. Lancet Oncol (2022) pmid: 35131040
1151. Petrylak et al., 2021; ESMO Abstract 579MO
1152. Topalian SL, et al. N. Engl. J. Med. (2012) pmid: 22658127
1153. Hodi et al., 2019; AACR abstract CT037
1154. Overman et al., 2019; ASCO Abstract 635
1155. Klein O, et al. Oncologist (2021) pmid: 33856094
1156. Andre et al., 2022; ASCO GI Abstract 244
1157. Klein O, et al. Oncoimmunology (2021) pmid: 33889439
1158. Noor et al., 2021; DOI: 10.1200/JCO.2021.39.3_suppl.415
1159. Sharma et al., 2023; ASCO GU Abstract 22
1160. Berton et al., 2021; SITC Abstract 956
1161. Grignani et al., 2021; SITC Abstract 545
1162. Rao S, et al. ESMO Open (2022) pmid: 35816951
1163. Bagley et al., 2022; SNO Abstract CTIM-35
1164. Rosenbaum et al., 2022; ASCO Abstract 11516
1165. Catenacci DVT, et al. ESMO Open (2022) pmid: 36029651
1166. Maio et al., 2021; ASCO Abstract 2571
1167. Turner et al., 2017; ASCO Abstract 1007
1168. Litton JK, et al. N. Engl. J. Med. (2018) pmid: 30110579
1169. Ettl J, et al. Ann. Oncol. (2018) pmid: 30124753
1170. Meehan et al., 2017; AACR Abstract 4687
1171. de Bono J, et al. Cancer Discov (2017) pmid: 28242752
1172. Lu E, et al. J Natl Compr Canc Netw (2018) pmid: 30099369
1173. De Bono et al., 2020; ASCO Abstract 5566
1174. Agarwal et al., 2023; ASCO Abstract LBA17
1175. de Bono JS, et al. Lancet Oncol (2021) pmid: 34388386
1176. Dorff et al., 2022; ASCO GU Abstract 157

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.