



Sample Information

Patient Name: 洪秀鸞  
Gender: Female  
ID No.: K201544825  
History No.: 20712565  
Age: 89  
  
Ordering Doctor: DOC3016D 江起陸  
Ordering REQ.: C32DJ58  
Signing in Date: 2023/12/21

Path No.: M112-00333  
MP No.: F23095  
Assay: Oncomine Focus Assay  
Sample Type: FFPE  
Block No.: S112-62172A  
Percentage of tumor cells: 80%

Reporting Doctor: DOC5466K 葉奕成 (Phone: 8#5466)

Note:

Sample Cancer Type: Non-Small Cell Lung Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	1 Relevant Biomarkers
Biomarker Descriptions	2	2 Therapies Available
Relevant Therapy Summary	3	8 Clinical Trials
Relevant Therapy Details	5	
Clinical Trials Summary	8	
Alert Details	8	

Relevant Non-Small Cell Lung Cancer Variants

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	ERBB2 exon 20 insertion	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	ERBB2 exon 20 insertion erb-b2 receptor tyrosine kinase 2 Allele Frequency: 45.55%	trastuzumab deruxtecan <sup>1</sup>	trastuzumab	8

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO  
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

### Prevalent cancer biomarkers without relevant evidence based on included data sources

CTNNB1 p.(I35S) c.104T>G

## Variant Details

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
CTNNB1	p.(I35S)	c.104T>G	COSM5674	chr3:41266107	32.80%	NM_001904.4	missense	1701
ERBB2	p.(Y772_A775dup)	c.2324_2325insATAC GTGATGGC	COSM20959	chr17:37880981	45.55%	NM_004448.3	nonframeshift Insertion	1978
PDGFRA	p.(G313=)	c.939T>G	.	chr4:55133726	50.08%	NM_006206.6	synonymous	1997
PDGFRA	p.(V824=)	c.2472C>T	.	chr4:55152040	49.92%	NM_006206.6	synonymous	1997
FGFR4	p.(P136L)	c.407C>T	.	chr5:176517797	99.60%	NM_213647.3	missense	1984
FGFR4	p.(I197T)	c.590T>C	.	chr5:176518092	16.06%	NM_213647.3	missense	1407
MET	p.(N375S)	c.1124A>G	.	chr7:116340262	39.10%	NM_001127500.3	missense	2000

## Biomarker Descriptions

### CTNNB1 (catenin beta 1)

**Background:** The CTNNB1 gene encodes catenin beta-1 (β-catenin), an integral component of cadherin-based adherens junctions involved in maintaining adhesion and regulating the growth of epithelial cell layers<sup>1</sup>. CTNNB1 binds to the APC protein in the cytoplasm and also interacts with TCF and LEF transcription factors in the nucleus to regulate WNT signaling<sup>2</sup>. Steady state levels of CTNNB1 are regulated by ubiquitin-dependent proteolysis<sup>3,4,5</sup>.

**Alterations and prevalence:** Recurrent somatic mutations leading to CTNNB1 activation are common in cancer. The most prevalent alterations include missense mutations in exon 3 at codons S33, S37, T41, and S45 that block phosphorylation by GSK-β and inhibit CTNNB1 degradation<sup>6,7,8,9</sup>. These activating mutations are observed in diverse solid tumors and have a prevalence of 20-30% in hepatocellular carcinoma, 20% of uterine carcinoma, and 15% of adrenocortical carcinoma<sup>10,11,12,13,14,15,16</sup>.

**Potential relevance:** Currently, no therapies have been approved for CTNNB1 aberrations. CTNNB1 alterations have been proposed to promote cancer progression and limit the response to EGFR tyrosine kinase inhibitors in EGFR positive lung cancer<sup>17</sup>. Mutation of CTNNB1 is considered useful as an ancillary diagnostic biomarker for desmoid fibromatosis<sup>18</sup>.

### ERBB2 (erb-b2 receptor tyrosine kinase 2)

**Background:** The ERBB2 gene encodes the erb-b2 receptor tyrosine kinase 2, a member of the human epidermal growth factor receptor (HER) family. Along with ERBB2/HER2, EGFR/ERBB1/HER1, ERBB3/HER3, and ERBB4/HER4 make up the HER protein family<sup>19</sup>. All ERBB/HER proteins encode transmembrane receptor tyrosine kinases. However, ERBB2/HER2 is an orphan receptor with no known ligand. ERBB2 preferentially binds other ligand bound ERBB/HER family members to form hetero-dimers resulting in the activation of ERBB2 tyrosine kinase activity and subsequent activation of the PI3K/AKT/MTOR and RAS/RAF/MAPK/ERK signaling pathways which promote cell proliferation, differentiation, and survival<sup>20</sup>. Recurrent focal amplification of the ERBB2 gene leads to increased expression

## Biomarker Descriptions (continued)

in several cancer types. ERBB2 overexpression in immortalized cell lines is oncogenic and leads to ERBB2 homo-dimerization and activation without ligand binding<sup>21,22,23</sup>.

**Alterations and prevalence:** ERBB2 gene amplification occurs in 10-20% of breast, esophageal, and gastric cancers, 5-10% of bladder, cervical, pancreas, and uterine cancers, and 1-5% of colorectal, lung, and ovarian cancers<sup>15,16,24,25,26,27,28,29</sup>. Recurrent somatic activating mutations in ERBB2/HER2 occur at low frequencies (<1%) in diverse cancer types<sup>16,30,31</sup>. In breast, bladder, and colorectal cancers, the most common recurrent ERBB2 activating mutations include kinase domain mutations L755S and V777L and the extracellular domain mutation S310F. In lung cancer, the most common recurrent ERBB2 activating mutations include in-frame exon 20 insertions, particularly Y772\_A775dup.

**Potential relevance:** The discovery of ERBB2/HER2 as an important driver of breast cancer in 1987 led to the development of trastuzumab, a humanized monoclonal antibody with specificity to the extracellular domain of HER2<sup>32,33</sup>. Trastuzumab<sup>34</sup> was FDA approved for the treatment of HER2 positive breast cancer in 1998, and subsequently in HER2 positive metastatic gastric and gastroesophageal junction adenocarcinoma in 2010. Additional monoclonal antibody therapies have been approved by the FDA for HER2-positive breast cancer including pertuzumab<sup>35</sup> (2012), a humanized monoclonal antibody that inhibits HER2 dimerization, and ado-trastuzumab emtansine<sup>36</sup> (2013), a conjugate of trastuzumab and a potent antimicrotubule agent. The combination of pertuzumab, trastuzumab, and a taxane is the preferred front-line regimen for HER2-positive metastatic breast cancer<sup>37</sup>. In addition to monoclonal antibodies, the small molecule inhibitor lapatinib<sup>38</sup>, with specificity for both EGFR and ERBB2, was FDA approved (2007) for the treatment of patients with advanced HER2-positive breast cancer who have received prior therapy including trastuzumab. In 2017, the FDA approved the use of neratinib<sup>39</sup>, an irreversible kinase inhibitor of EGFR, ERBB2/HER2, and ERBB4, for the extended adjuvant treatment of adult patients with early stage HER2-positive breast cancer. In 2020, the FDA approved neratinib<sup>39</sup> in combination with capecitabine for HER2-positive advanced or metastatic patients after two or more prior HER2-directed therapies. Also in 2020, the TKI irbininib<sup>40</sup> was FDA approved for HER2 overexpressing or amplified breast cancer in combination with trastuzumab and capecitabine. In 2021, the PD-1 blocking antibody, pembrolizumab, in combination with trastuzumab, fluoropyrimidine- and platinum-based chemotherapy, was approved for HER2 amplified gastric or gastroesophageal (GEJ) adenocarcinoma in the first line<sup>41</sup>. The vaccine, nelipecimut-S<sup>42</sup>, was granted fast-track designation by the FDA (2016) in patients with low to intermediate HER2 expressing (IHC score 1+ or 2+) breast cancer. In 2018 fast-track designation was granted to the monoclonal antibody margetuximab<sup>43</sup> in patients with ERBB2 positive breast cancer previously treated with an anti-HER2 therapy. In 2019, fast track designation was granted to the HER2-targeting antibody drug conjugate, amcenestrant<sup>44</sup>, for HER2-positive advanced or metastatic breast cancer after one or more prior anti-HER2 based regimens. Additionally, in 2019, the novel bispecific antibody, zanidatamab<sup>45</sup>, received fast-track designation in combination with standard chemotherapy for patients with HER2-overexpressing gastroesophageal adenocarcinoma (GEA) and breakthrough therapy designation (2020) as a monotherapy for patients with HER2-amplified biliary tract cancer<sup>46</sup>. In 2020, BDTX-189<sup>47</sup> received fast-track designation for adult patients with solid tumors harboring an allosteric human ERBB2 mutation or exon 20 insertion, and the humanized anti-HER2 antibody drug conjugate disitamab vedotin received breakthrough designation for adult patients with HER2-positive urothelial cancer after previous platinum-chemotherapy treatment<sup>48</sup>. In 2021, the antibody-drug conjugate ARX788<sup>49</sup> received fast-track designation as a monotherapy for advanced or metastatic HER2-positive breast cancer that have progressed on one or more anti-HER2 regimens. Additionally, in 2021, fast track designation was granted to HER2 targeted chimeric antigen receptor macrophage (CAR-M), CT-0508<sup>50</sup>, for HER2-overexpressing solid tumors. Certain activating mutations have been observed to impart sensitivity to neratinib, afatinib, lapatinib, and trastuzumab, or dacomitinib in early and ongoing clinical studies<sup>51,52,53,54,55</sup>. ERBB2 kinase domain mutations R896G and V659E both showed response to afatinib in two NSCLC case studies<sup>56,57</sup>. Additionally, acquired HER2 mutations in estrogen receptor-positive (ER+) breast cancer have been shown to confer resistance to hormone therapy<sup>58</sup>. However, this was shown to be overcome by neratinib in combination with therapies targeting ER<sup>58</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### ERBB2 exon 20 insertion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
trastuzumab deruxtecan	●	●	×	●	×
trastuzumab	×	×	×	○	×
trastuzumab deruxtecan, pembrolizumab, chemotherapy	×	×	×	×	● (III)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types    ☒ No evidence

### ERBB2 exon 20 insertion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
ado-trastuzumab emtansine	×	×	×	×	● (II)
ABT-101	×	×	×	×	● (I/II)
sunvozertinib	×	×	×	×	● (I/II)
BAY-2927088	×	×	×	×	● (I)
ELVN-002, trastuzumab deruxtecan	×	×	×	×	● (I)
SAR-443216	×	×	×	×	● (I)
SHR-A1811	×	×	×	×	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

FDA information is current as of 2023-09-13. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### ERBB2 exon 20 insertion

##### ☒ trastuzumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer

Label as of: 2022-11-04

Variant class: ERBB2 Y772\_A775dup mutation

##### Indications and usage:

ENHERTU® is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of:

- adult patients with unresectable or metastatic HER2-positive breast cancer who have received a prior anti-HER2-based regimen either:
  - in the metastatic setting, or
  - in the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy.
- adult patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer, as determined by an FDA-approved test, who have received a prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy.
- adult patients with unresectable or metastatic non-small cell lung cancer (NSCLC) whose tumors have activating HER2 (ERBB2) mutations, as detected by an FDA-approved test, and who have received a prior systemic therapy.
  - This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.
- adult patients with locally advanced or metastatic HER2-positive gastric or gastroesophageal junction adenocarcinoma who have received a prior trastuzumab-based regimen.

##### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/761139s024lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/761139s024lbl.pdf)

## Current NCCN Information

☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

NCCN information is current as of 2023-09-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

### ERBB2 exon 20 insertion

#### ☒ trastuzumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 Y772\_A775dup mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Adenocarcinoma, Large Cell, Not otherwise specified (NOS), Squamous Cell; Advanced, Metastatic, Progression (Subsequent therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2023]

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

ESMO information is current as of 2023-09-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### ERBB2 exon 20 insertion

#### ☒ trastuzumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer

Variant class: ERBB2 Y772\_A775dup mutation

ESMO Level of Evidence/Grade of Recommendation: III / B

Population segment (Line of therapy):

- Advanced, Metastatic, Unresectable (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Oncogene-addicted Metastatic Non-Small-Cell Lung Cancer [Annals of Oncology (2023), doi: <https://doi.org/10.1016/j.annonc.2022.12.009> (Published)]

#### ☐ trastuzumab

Cancer type: Cholangiocarcinoma, Gallbladder Carcinoma

Variant class: ERBB2 mutation

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- Progression (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Biliary Cancer [Ann Oncol (2023), doi: <https://doi.org/10.1016/j.annonc.2022.10.506>]

## Clinical Trials in Taiwan region:

### Clinical Trials Summary

#### ERBB2 exon 20 insertion

NCT ID	Title	Phase
<a href="#">NCT05532696</a>	A Phase Ib/II, Open-Label, Multicenter Study to Evaluate Safety, Tolerability, Pharmacokinetics, and Antitumor Activity of ABT-101 in Patients With Advanced Solid Tumors and HER2 Exon 20 Insertions Mutated Non- Small Cell Lung Cancer	I/II
<a href="#">NCT03974022</a>	A Phase I/II, Open-Label, Multicenter Study to Assess the Safety, Tolerability, Pharmacokinetics and Anti-tumor Efficacy of DZD9008 in Patients With Advanced Non-Small Cell Lung Cancer (NSCLC) with EGFR or HER2 Mutation	I/II
<a href="#">NCT05099172</a>	An Open Label, First-in-human Study of BAY 2927088 in Participants With Advanced Non-small Cell Lung Cancer (NSCLC) Harboring an EGFR and/or HER2 Mutation	I
<a href="#">NCT05048797</a>	An Open-label, Randomized, Multicenter, Phase III Study to Assess the Efficacy and Safety of Trastuzumab Deruxtecan as First-line Treatment of Unresectable, Locally Advanced, or Metastatic NSCLC Harboring HER2 Exon 19 or 20 Mutations (DESTINY-Lung04)	III
<a href="#">NCT05650879</a>	A Phase Ia/Ib Study of ELVN-002 for the Treatment of Patients With HER2 Mutant Non-Small Cell Lung Cancer	I
<a href="#">NCT05013554</a>	A Phase I/Ib Open-label, First-in-human, Single Agent, Dose Escalation and Expansion Study for the Evaluation of Safety, Pharmacokinetics, Pharmacodynamics, and Anti-tumor Activity of SAR443216 in Participants with Relapsed/Refractory HER2 Expressing Solid Tumors.	I
<a href="#">NCT04589845</a>	Tumor-Agnostic Precision Immunooncology and Somatic Targeting Rational for You (TAPISTRY) Phase II Platform Trial	II
<a href="#">NCT04446260</a>	A Phase I Multi-Country, Multi-Center, Open-Label Study to Evaluate the Safety, Tolerability, Pharmacokinetics and Efficacy of SHR-A1811 in HER2 Expressing or Mutated Advanced Malignant Solid Tumor Subjects	I

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2023-09-13. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### ERBB2 exon 20 insertion

### BDTX-189

Cancer type: Solid Tumor

Variant class: ERBB2 exon 20 insertion

#### Supporting Statement:

The FDA has granted Fast Track Designation to BDTX-189 for solid tumors harboring a HER2 mutation or an EGFR or HER2 exon 20 insertion after progression on prior therapy.

#### Reference:

<https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>



## References

1. Valenta et al. The many faces and functions of  $\beta$ -catenin. *EMBO J.* 2012 Jun 13;31(12):2714-36. PMID: 22617422
2. Korinek et al. Constitutive transcriptional activation by a beta-catenin-Tcf complex in APC-/- colon carcinoma. *Science.* 1997 Mar 21;275(5307):1784-7. PMID: 9065401
3. Aberle et al. beta-catenin is a target for the ubiquitin-proteasome pathway. *EMBO J.* 1997 Jul 1;16(13):3797-804. PMID: 9233789
4. Winston et al. The SCFbeta-TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in IkappaBalpha and beta-catenin and stimulates IkappaBalpha ubiquitination in vitro. *Genes Dev.* 1999 Feb 1;13(3):270-83. PMID: 9990852
5. Kitagawa et al. An F-box protein, FWD1, mediates ubiquitin-dependent proteolysis of beta-catenin. *EMBO J.* 1999 May 4;18(9):2401-10. PMID: 10228155
6. Liu et al. Control of beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell.* 2002 Mar 22;108(6):837-47. PMID: 11955436
7. Miyoshi et al. Activation of the beta-catenin gene in primary hepatocellular carcinomas by somatic alterations involving exon 3. *Cancer Res.* 1998 Jun 15;58(12):2524-7. PMID: 9635572
8. Gao et al. Exon 3 mutations of CTNNB1 drive tumorigenesis: a review. *Oncotarget.* 2018 Jan 12;9(4):5492-5508. PMID: 29435196
9. Morin et al. Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC. *Science.* 1997 Mar 21;275(5307):1787-90. PMID: 9065402
10. Schulze et al. Exome sequencing of hepatocellular carcinomas identifies new mutational signatures and potential therapeutic targets. *Nat. Genet.* 2015 May;47(5):505-511. PMID: 25822088
11. Ahn et al. Genomic portrait of resectable hepatocellular carcinomas: implications of RB1 and FGF19 aberrations for patient stratification. *Hepatology.* 2014 Dec;60(6):1972-82. PMID: 24798001
12. Harding et al. Prospective Genotyping of Hepatocellular Carcinoma: Clinical Implications of Next-Generation Sequencing for Matching Patients to Targeted and Immune Therapies. *Clin. Cancer Res.* 2018 Oct 29. PMID: 30373752
13. Cancer et al. Integrated genomic characterization of endometrial carcinoma. *Nature.* 2013 May 2;497(7447):67-73. PMID: 23636398
14. Soumerai et al. Clinical Utility of Prospective Molecular Characterization in Advanced Endometrial Cancer. *Clin. Cancer Res.* 2018 Dec 1;24(23):5939-5947. PMID: 30068706
15. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
16. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
17. Blakely et al. Evolution and clinical impact of co-occurring genetic alterations in advanced-stage EGFR-mutant lung cancers. *Nat. Genet.* 2017 Dec;49(12):1693-1704. PMID: 29106415
18. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2023]
19. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science.* 1985 Sep 6;229(4717):974-6. PMID: 2992089
20. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
21. Di et al. erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science.* 1987 Jul 10;237(4811):178-82. PMID: 2885917
22. Hudziak et al. Increased expression of the putative growth factor receptor p185HER2 causes transformation and tumorigenesis of NIH 3T3 cells. *Proc. Natl. Acad. Sci. U.S.A.* 1987 Oct;84(20):7159-63. PMID: 2890160
23. Lonardo et al. The normal erbB-2 product is an atypical receptor-like tyrosine kinase with constitutive activity in the absence of ligand. *New Biol.* 1990 Nov;2(11):992-1003. PMID: 1983208
24. Ciriello et al. Comprehensive Molecular Portraits of Invasive Lobular Breast Cancer. *Cell.* 2015 Oct 8;163(2):506-19. PMID: 26451490
25. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
26. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature.* 2014 Mar 20;507(7492):315-22. doi: 10.1038/nature12965. Epub 2014 Jan 29. PMID: 24476821
27. Donna et al. Comprehensive molecular characterization of human colon and rectal cancer. *Nature.* 2012 Jul 18;487(7407):330-7. PMID: 22810696

## References (continued)

28. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
29. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011 Jun 29;474(7353):609-15. PMID: 21720365
30. Petrelli et al. Clinical and pathological characterization of HER2 mutations in human breast cancer: a systematic review of the literature. *Breast Cancer Res. Treat.* 2017 Nov;166(2):339-349. PMID: 28762010
31. Bose et al. Activating HER2 mutations in HER2 gene amplification negative breast cancer. *Cancer Discov.* 2013 Feb;3(2):224-37. doi: 10.1158/2159-8290.CD-12-0349. Epub 2012 Dec 7. PMID: 23220880
32. Hudis. Trastuzumab—mechanism of action and use in clinical practice. *N. Engl. J. Med.* 2007 Jul 5;357(1):39-51. PMID: 17611206
33. Slamon et al. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/neu oncogene. *Science*. 1987 Jan 9;235(4785):177-82. PMID: 3798106
34. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2018/103792s5345lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2018/103792s5345lbl.pdf)
35. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/125409s124lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/125409s124lbl.pdf)
36. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/125427s111lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/125427s111lbl.pdf)
37. NCCN Guidelines® - NCCN-Breast Cancer [Version 4.2023]
38. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/022059s031lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/022059s031lbl.pdf)
39. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/208051s009lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/208051s009lbl.pdf)
40. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/213411s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/213411s004lbl.pdf)
41. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/125514s135lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/125514s135lbl.pdf)
42. <https://www.globenewswire.com/news-release/2016/06/01/845166/0/en/Galena-Biopharma-Receives-Fast-Track-Designation-for-NeuVax-nelipepimut-S-PRESENT-Clinical-Trial.html>
43. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/761150s005lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/761150s005lbl.pdf)
44. <https://www.prnewswire.com/news-releases/fda-grants-arx788-fast-track-designation-for-her2-positive-metastatic-breast-cancer-301199951.html>
45. <https://www.targetedonc.com/view/her2targeted-antibody-zw25-earns-fda-fast-track-designation-in-gea>
46. <https://www.targetedonc.com/view/fda-grants-breakthrough-designation-to-zanidatamab-for-her2-amplified-biliary-tract-cancer>
47. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>
48. <https://www.prnewswire.com/news-releases/remegen-announces-us-fda-has-granted-breakthrough-therapy-designation-for-disitamab-vedotin-rc48-in-urothelial-cancer-301138315.html>
49. <https://ir.ambrx.com/news/news-details/2023/ACE-Breast-02-Pivotal-Phase-3-Study-of-Ambrxs-ARX788-for-the-Treatment-of-HER2-Positive-Metastatic-Breast-Cancer-Achieves-Positive-Results/default.aspx>
50. <https://www.prnewswire.com/news-releases/carisma-therapeutics-announces-us-food-and-drug-administration-grants-fast-track-designation-to-ct-0508-for-the-treatment-of-patients-with-solid-tumors-301381843.html>
51. Ma et al. Neratinib Efficacy and Circulating Tumor DNA Detection of HER2 Mutations in HER2 Nonamplified Metastatic Breast Cancer. *Clin. Cancer Res.* 2017 Oct 1;23(19):5687-5695. PMID: 28679771
52. De et al. Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu. *Lung Cancer*. 2012 Apr;76(1):123-7. PMID: 22325357
53. Kris et al. Targeting HER2 aberrations as actionable drivers in lung cancers: phase II trial of the pan-HER tyrosine kinase inhibitor dacomitinib in patients with HER2-mutant or amplified tumors. *Ann. Oncol.* 2015 Jul;26(7):1421-7. PMID: 25899785
54. Falchook et al. Non-small-cell lung cancer with HER2 exon 20 mutation: regression with dual HER2 inhibition and anti-VEGF combination treatment. *J Thorac Oncol.* 2013 Feb;8(2):e19-20. PMID: 23328556
55. David et al. Neratinib in HER2- or HER3-mutant solid tumors: SUMMIT, a global, multi-histology, open-label, phase 2 'basket' study. AACR 2017. Abstract CT001
56. Lin et al. Response to Afatinib in a Patient with Non-Small Cell Lung Cancer Harboring HER2 R896G Mutation: A Case Report. *Onco Targets Ther.* 2019;12:10897-10902. PMID: 31849493
57. Chang et al. Sustained Partial Response to Afatinib in a Patient With Lung Adenocarcinoma Harboring HER2V659E Mutation. *JCO Precis Oncol.* 2020 Aug; 912-915. DOI: 10.1200/PO.20.00114
58. Nayar et al. Acquired HER2 mutations in ER+ metastatic breast cancer confer resistance to estrogen receptor-directed therapies. *Nat. Genet.* 2019 Feb;51(2):207-216. PMID: 30531871