

ABOUT THE TEST FoundationOne®CDx is a next-generation sequencing (NGS) based assay that identifies genomic findings within hundreds of cancer-related genes.

PATIENT	DISEASE Colon adenocarcinoma (CRC)	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN SITE Ovary
	NAME Huang, Chien Chung		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN ID S112-66168G (PF23084)
	DATE OF BIRTH 04 January 1951		ADDITIONAL RECIPIENT None		SPECIMEN TYPE Slide Deck
	SEX Female		MEDICAL FACILITY ID 205872		DATE OF COLLECTION 21 March 2023
	MEDICAL RECORD # 17051911 PF23084		PATHOLOGIST Not Provided		SPECIMEN RECEIVED 05 July 2023

Sample qualified for low tumor purity. Sensitivity for detecting copy-number alterations (including in *ERBB2*), other genomic alterations, and signatures may be reduced and TMB may be underreported. Refer to the appendix for limitations statements.

Biomarker Findings

Microsatellite status - Cannot Be Determined ^α
Tumor Mutational Burden - Cannot Be Determined

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

KRAS wildtype
NRAS wildtype
SMAD4 G386D

4 Disease relevant genes with no reportable alterations: **BRAF, ERBB2, KRAS, NRAS**

^α Patients with Microsatellite status of Cannot Be Determined should be re-tested with an orthogonal (alternative) method.

Report Highlights

- Targeted therapies with NCCN categories of evidence in this tumor type: Cetuximab (p. 7), Panitumumab (p. 7)

BIOMARKER FINDINGS

Microsatellite status -
Cannot Be Determined

Tumor Mutational Burden -
Cannot Be Determined

GENOMIC FINDINGS

KRAS - wildtype

0 Trials

NRAS - wildtype

0 Trials

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

No therapies or clinical trials. See Biomarker Findings section

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)		THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)
Cetuximab	2A	none
Panitumumab	2A	
Cetuximab	2A	none
Panitumumab	2A	

 NCCN category

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

SMAD4 - G386D **p. 6**

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the agents listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and exhaustive. Neither the therapeutic agents nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies.

Therapies contained in this report may have been approved by the US FDA.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

BIOMARKER FINDINGS

BIOMARKER

Microsatellite status

RESULT

Cannot Be Determined

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in multiple solid tumor types, microsatellite instability (MSI) and associated increased tumor mutational burden (TMB)¹⁻² may predict sensitivity to immune checkpoint inhibitors, including the approved PD-1-targeting agents cemiplimab, dostarlimab, nivolumab (alone or in combination with ipilimumab), retifanlimab, and pembrolizumab³⁻⁹, as well as PD-L1-targeting agents atezolizumab, avelumab, and durvalumab (alone or in combination with tremelimumab)¹⁰⁻¹¹. Pembrolizumab therapy resulted in a significantly higher ORR in MSI-H CRC compared with MSS CRC (40% vs. 0%)⁷. Similarly, a clinical study of nivolumab, alone or in combination with ipilimumab, in patients with CRC reported a significantly higher response rate in patients with tumors with high MSI than those without³⁻⁴. An earlier case study reported that nivolumab therapy resulted in a complete response in a patient with MSI-H CRC⁶. A Phase 1b trial of atezolizumab

combined with bevacizumab reported PRs for 40% (4/10) of patients with MSI-H CRC¹². As the MSI status of this tumor is unknown, the relevance of these therapeutic approaches is unclear.

— Nontargeted Approaches —

MSI has not been found to be a predictive biomarker for combination chemotherapy regimens, including FOLFOX¹³⁻¹⁴ and FOLFIRI¹⁵⁻¹⁶. Patients with MSS CRC are more likely to benefit from postsurgical fluorouracil (FU)-based adjuvant therapy¹⁷⁻¹⁸ but less likely to benefit from irinotecan chemotherapy¹⁹.

FREQUENCY & PROGNOSIS

Approximately 10-15% of colorectal cancers (CRCs) are MSI-high (MSI-H), 8-16% are MSI-low (MSI-L), and the remaining majority are microsatellite-stable (MSS)^{2,20-24}. Multiple studies have shown that MSI-H CRCs have a better prognosis than MSI-low (MSI-L) or microsatellite stable (MSS) tumors^{21,25-31}. MSI-H CRCs are associated with certain pathologic and molecular features, including poor differentiation, right-sided and mucinous tumors, increased numbers of tumor infiltrating lymphocytes, diploidy, and a relatively high frequency of BRAF mutations^{22-23,32}. The prognostic implications of MS-Equivocal status for patients with colorectal cancer (CRC) have not been evaluated in published studies. One study reported

that microsatellite instability-low (MSI-L) tumors occur more frequently in early-stage CRC than advanced CRC³³; in contrast, another study found that MSI-L status is associated with advanced stage and worse OS for patients with CRC³⁴. MSI-L CRCs resemble MS-Stable tumors in most clinicopathologic features^{1-2,22,32} but harbor a significantly higher frequency of KRAS mutations than MS-Stable tumors^{20,33,35}. MSS colorectal cancers are molecularly heterogeneous, driven by diverse mechanisms such as extensive DNA methylation, oncogenic mutations in KRAS or BRAF, or chromosomal instability¹.

FINDING SUMMARY

Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor²³. Defective MMR and consequent MSI occur as a result of genetic or epigenetic inactivation of one of the MMR pathway proteins, primarily MLH1, MSH2, MSH6, or PMS2^{23,36-37}. The level of MSI in this sample could not be determined with confidence. Depending on the clinical context, MSI testing of an alternate sample or by another methodology could be considered.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

BIOMARKER FINDINGS

BIOMARKER

Tumor Mutational Burden

RESULT

Cannot Be Determined

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased TMB may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1³⁸⁻⁴⁰, anti-PD-1 therapies³⁸⁻⁴¹, and combination nivolumab and ipilimumab⁴²⁻⁴⁷. In multiple pan-tumor studies, increased tissue tumor mutational burden (TMB) was associated with sensitivity to immune checkpoint inhibitors^{38-41,48-52}. In the KEYNOTE 158 trial of pembrolizumab monotherapy for patients with solid tumors, significant improvement in ORR was observed for patients with TMB ≥ 10 Muts/Mb (as measured by this assay) compared with those with TMB < 10 Muts/Mb in a large cohort that included multiple tumor types⁴⁸; similar findings were observed in the KEYNOTE 028 and 012 trials⁴¹. At the same TMB cutpoint, retrospective analysis of patients with solid tumors treated with any checkpoint inhibitor identified that tissue TMB scores ≥ 10 Muts/Mb were associated with prolonged time to treatment failure compared with scores < 10 Muts/Mb (HR=0.68)⁵². For patients with solid tumors treated with nivolumab plus ipilimumab in the CheckMate 848 trial, improved responses were observed in patients with a tissue TMB ≥ 10 Muts/Mb independent of blood TMB at any cutpoint in matched samples⁵³. However, support for higher TMB thresholds and efficacy was observed in the prospective Phase 2 MyPathway trial of atezolizumab for patients with pan-solid tumors, where improved ORR and DCR was seen in patients with TMB ≥ 16 Muts/Mb than those with TMB ≥ 10 and < 16 Muts/Mb⁵¹.

Similarly, analyses across several solid tumor types reported that patients with higher TMB (defined as ≥ 16 -20 Muts/Mb) achieved greater clinical benefit from PD-1 or PD-L1-targeting monotherapy compared with patients with higher TMB treated with chemotherapy⁵⁴ or those with lower TMB treated with PD-1 or PD-L1-targeting agents³⁹. In CRC specifically, a retrospective analysis of immune checkpoint inhibitor efficacy reported significantly improved OS for patients with tumors harboring TMB ≥ 9.8 Muts/Mb compared with those with tumors with TMB < 9.8 Muts/Mb (~equivalency < 12 Muts/Mb as measured by this assay)³⁸. Another retrospective study reported that a TMB ≥ 12 Muts/Mb cutoff identifies $> 99\%$ of MSI-High CRC cases but only 3% of MSS cases, indicating the utility of this cutoff for identification of patients with CRC likely to benefit from treatment with immune checkpoint inhibitors⁵⁵. As the TMB status of this tumor cannot be determined with confidence, the benefit of these therapeutic approaches is unclear.

FREQUENCY & PROGNOSIS

Elevated tumor mutational burden (TMB) has been reported in 8-25% of colorectal cancer (CRC) samples^{24,56-57}. Multiple studies have reported that up to 90% of hypermutated CRC cases exhibit high levels of microsatellite instability (MSI-H) and mismatch repair deficiency (MMR-D)^{24,56}. Increased TMB is significantly associated with MSI-H and MMR-D, with studies reporting that 100% of MSI-H CRCs harbor elevated TMB and conversely that 100% of tumors with low TMB harbor intact MMR⁵⁶. A subset of CRCs that harbor increased TMB but not MSI-H are driven by mutations in POLE, which leads to an "ultramutated" phenotype with especially high TMB^{24,56}. Tumors with increased TMB harbor BRAF V600E mutations more frequently than those with low TMB^{24,56}, whereas TMB-low tumors more frequently harbor mutations in TP53 and APC²⁴. The prognostic value of tumor mutational burden (TMB) in colorectal cancer (CRC) is context- and therapy-dependent. A study

of tissue TMB (tTMB) in 145 CRC samples showed longer OS in TMB-high samples compared with TMB-low ones⁵⁸. Similarly, for patients with metastatic CRC treated with first-line chemotherapy combined with bevacizumab or cetuximab, high tissue TMB (tTMB-H) was associated with longer OS⁵⁹. For patients treated with adjuvant chemotherapy, tTMB-H was associated with better 5-year relapse-free survival⁶⁰. However, for patients with EGFR/BRAF-inhibitor-treated, BRAF-mutated microsatellite stable (MSS) metastatic CRC, intermediate tTMB was associated with significantly poorer PFS and OS compared with TMB-low status; patients with primary resistance to EGFR/BRAF blockage had higher TMB than those sensitive to these therapies⁶¹. In a study for 61 patients with metastatic, MSS CRC treated with best standard of care, plasma TMB scores ≥ 28 Muts/Mb (approximately 14 Muts/Mb as measured by this assay) were associated with reduced OS compared with plasma TMB scores < 28 Muts/Mb (3.0 vs. 5.3 months, HR=0.76, p=0.007), whereas tTMB was not found to be prognostic in this population⁶².

FINDING SUMMARY

Tumor mutation burden (TMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations occurring in a tumor specimen. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma⁶³⁻⁶⁴ and cigarette smoke in lung cancer^{8,65}, treatment with temozolomide-based chemotherapy in glioma⁶⁶⁻⁶⁷, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{24,68-71}, and microsatellite instability (MSI)^{24,68,71}. Elevated TMB has been reported to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors in multiple solid tumor types^{39-41,48}. However, the TMB level in this sample could not be determined with confidence.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

GENOMIC FINDINGS

GENE
KRAS

ALTERATION
wildtype

with clinical benefit of treatment with EGFR-targeting antibodies cetuximab⁷²⁻⁷⁵ or panitumumab⁷⁶⁻⁷⁸ for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

Numerous studies have reported that KRAS wild-type status is associated with decreased metastasis, better clinicopathological features, and longer survival of patients with CRC^{81-84,88-89}.

FINDING SUMMARY

KRAS encodes a member of the RAS family of small GTPases. Activating mutations in RAS genes can cause uncontrolled cell proliferation and tumor formation⁹⁰⁻⁹¹. No alterations in KRAS were identified in this case.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated

FREQUENCY & PROGNOSIS

Approximately 50-65% of colorectal cancers (CRCs) have been reported to lack KRAS mutations⁷⁹⁻⁸⁷.

GENE
NRAS

ALTERATION
wildtype

targeting antibodies cetuximab⁷²⁻⁷⁵ or panitumumab⁷⁶⁻⁷⁸ for patients with CRC. Therefore, these agents are indicated to treat patients with CRC lacking such mutations (NCCN Colon Cancer Guidelines, v3.2022, Rectal Cancer Guidelines, v4.2022).

of metastasis⁸⁷ and longer survival⁹⁷⁻⁹⁸ of patients with CRC.

FINDING SUMMARY

NRAS encodes a member of the RAS family of small GTPases that mediate transduction of growth signals. Activation of RAS signaling causes cell growth, differentiation, and survival by activating the RAF-MAPK-ERK, PI3K, and other pathways⁹⁰. No alterations in NRAS were identified in this case.

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Lack of mutations in KRAS or NRAS is associated with clinical benefit of treatment with EGFR-

FREQUENCY & PROGNOSIS

The majority of colorectal cancers (CRCs) (91-98%) have been reported to lack NRAS mutations^{24,87,92-97}. NRAS wild-type status has been reported to be associated with decreased frequency

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

GENOMIC FINDINGS

GENE

SMAD4

ALTERATION

G386D

HGVS VARIANT

NM_005359.5:c.1157G>A (p.G386D)

VARIANT CHROMOSOMAL POSITION

chr18:48593406

VARIANT ALLELE FREQUENCY (% VAF)

1.7%

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no targeted therapies available to address genomic alterations in SMAD4. Preclinical studies in colorectal cancer have reported associations of SMAD4 inactivation or loss with sensitivity to inhibitors of Aurora kinase A⁹⁹ and the Wnt/beta-catenin pathway¹⁰⁰.

— Nontargeted Approaches —

Clinical studies have reported associations of SMAD4 loss or low SMAD4 expression with improved responses to chemotherapeutic agents in

patients with pancreatic cancer¹⁰¹⁻¹⁰³ and non-small cell lung cancer (NSCLC)¹⁰⁴. Other clinical studies in pancreatic cancer have reported an association of high SMAD4 expression with better responses to neoadjuvant chemotherapy¹⁰⁵ and adjuvant chemoradiotherapy¹⁰⁶.

FREQUENCY & PROGNOSIS

SMAD4 mutation or homozygous deletion is most frequently observed in pancreatic adenocarcinoma (43%)¹⁰⁷, pancreatic acinar cell carcinoma (26%)¹⁰⁸, cholangiocarcinoma (25%)¹⁰⁹, small intestine cancer (20%)¹¹⁰, appendiceal adenocarcinoma (14-20% mutation; 57% deletion)¹¹¹⁻¹¹², colorectal adenocarcinoma (CRC; 14%)²⁴, esophageal adenocarcinoma (14%)¹¹³, and stomach adenocarcinoma (13%)¹¹⁴. In preclinical studies, SMAD4 loss of function has been implicated in the development of mucinous neoplasms of the pancreas, including mucinous cystic neoplasms (MCN)¹¹⁵ and intraductal papillary mucinous neoplasms (IPMN)¹¹⁶; in clinical samples, SMAD4 homozygous deletion has been observed in 10% of IPMNs and 8% of MCNs, and mutation was also observed in 5% of IPMNs¹¹⁷. SMAD4 gene alterations have been associated with reduced OS for patients with pancreatic adenocarcinoma¹¹⁸. Reduced SMAD4 expression has been associated

with worse prognosis in various cancer types, including CRC¹¹⁹⁻¹²¹, appendiceal mucinous neoplasm¹²², gastric adenocarcinoma¹²³⁻¹²⁴, esophageal adenocarcinoma¹²⁵, esophageal squamous cell carcinoma¹²⁶, breast cancer¹²⁷, and prostate cancer¹²⁸.

FINDING SUMMARY

SMAD4, also known as DPC4, encodes a tumor suppressor that regulates transcriptional activity downstream of TGF-beta receptor signaling¹²⁹⁻¹³⁰. SMAD4 alterations that result in loss or disruption of the MH1 domain (aa 18-142), MH2 domain (aa 323-552), or SAD domain (aa 275-320) are predicted to be inactivating¹³¹⁻¹⁴⁴.

POTENTIAL GERMLINE IMPLICATIONS

Germline SMAD4 mutations, including those at the R361 hotspot, have been observed in patients with juvenile polyposis syndrome¹⁴⁵⁻¹⁴⁷, which is associated with an increased risk of gastrointestinal cancers¹⁴⁸. The penetrance of deleterious SMAD4 mutations in patients with colon cancer is estimated at 20% by age 35 and 70% by age 65¹⁴⁹. In the appropriate clinical context, germline testing of SMAD4 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

THERAPIES WITH CLINICAL BENEFIT

IN PATIENT'S TUMOR TYPE

Cetuximab

Assay findings association

KRAS
wildtype

NRAS
wildtype

AREAS OF THERAPEUTIC USE

Cetuximab is a monoclonal antibody that targets EGFR. It is FDA approved for the treatment of head and neck squamous cell carcinoma (HNSCC) and KRAS-wild-type, EGFR-expressing metastatic colorectal cancer (CRC). Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including cetuximab, have been shown to have significant clinical activity for patients with CRC^{72-75,150-151}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of cetuximab in metastatic CRC (NCCN Colon Cancer Guidelines, v3.2022, NCCN Rectal Cancer Guidelines, v4.2022).

SUPPORTING DATA

Cetuximab has been shown to improve OS, PFS, and response rate for patients with KRAS-wildtype colorectal cancer (CRC), both in combination with FOLFIRI, FOLFOX₄, or irinotecan^{72-73,150-152} and as monotherapy for chemotherapy-refractory patients^{75,153}. The Phase 3 study STRATEGIC-1 reported a similar duration of disease control (DDC) for patients with unresectable metastatic CRC (mCRC) and KRAS-, NRAS-, and BRAF-wildtype status treated with mFOLFOX-bevacizumab alternated

with a cetuximab regimen in first or second line, respectively (overall DDC 22.5 vs. 23.5 months); in addition, the study reported similar OS (37.8 vs. 34.4 months) and higher numerical ORR for patients treated with cetuximab in the first line followed by mFOLFOX-bevacizumab compared with those receiving EGFR-directed antibodies in the second or third line¹⁵⁴. A prospective study of cetuximab monotherapy for patients with KRAS-, NRAS-, and BRAF-wildtype mCRC reported 11% (2/19) PRs and 58% (11/19) SDs¹⁵⁵. The Phase 2 AVETUX trial of cetuximab combined with avelumab and mFOLFOX6 for patients with RAS- and BRAF-wildtype mCRC resulted in an ORR of 81% (4 CR and 27 PRs, n=37) and a DCR of 89%¹⁵⁶. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)¹⁵⁷. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)¹⁵⁸.

Panitumumab

Assay findings association

KRAS
wildtype

NRAS
wildtype

AREAS OF THERAPEUTIC USE

Panitumumab is a monoclonal antibody that targets EGFR. It is FDA approved to treat KRAS wild-type and NRAS wild-type metastatic colorectal cancer (CRC) combined with chemotherapy or as monotherapy for patients who have progressed on prior chemotherapy. Please see the drug label for full prescribing information.

GENE ASSOCIATION

Therapies targeting EGFR, including panitumumab, have been shown to have significant clinical activity for patients with CRC^{76,157,159}; wild-type KRAS and NRAS are predictive biomarkers for the efficacy of panitumumab in metastatic CRC (NCCN Colon Cancer Guidelines v3.2022)(NCCN Rectal Cancers Guidelines, v4.2022).

SUPPORTING DATA

Panitumumab has been shown to improve OS, PFS, and ORR for patients with KRAS-wildtype colorectal cancer (CRC), both in combination with FOLFOX₄, FOLFIRI, irinotecan, or best supportive care^{76,160-163}, and as monotherapy for chemotherapy-refractory patients^{157,159,164}. The Phase 3 PARADIGM trial comparing panitumumab

plus mFOLFOX6 versus bevacizumab plus mFOLFOX6 as first-line treatment for patients with RAS-wildtype left-sided metastatic CRC demonstrated that treatment with panitumumab significantly improved median OS (mOS; 36.2 months vs. 31.3 months) compared with bevacizumab¹⁶⁵. A Phase 2 trial reported that, for patients with unresectable RAS-wildtype colorectal adenocarcinoma treated with panitumumab plus FOLFOX₄, maintenance with a combination of panitumumab plus fluorouracil and leucovorin was superior to panitumumab monotherapy (10-month PFS OF 59% vs. 49%)¹⁶⁶. In the Phase 3 ASPECCT study, panitumumab was found to be non-inferior to cetuximab with respect to median OS (10.4 vs. 10.0 months, HR=0.97) for patients with previously treated KRAS exon 2 wildtype metastatic colorectal cancer; median PFS was also similar between the two treatment groups (4.4 vs. 4.1 months, HR=1.00)¹⁵⁷. In a similar patient population, a Phase 2 study of combination panitumumab and irinotecan versus combination cetuximab and irinotecan also demonstrated non-inferiority with respect to median PFS (5.4 vs. 4.3 months, HR = 0.64) and median OS (14.9 vs. 11.5 months, HR=0.66)¹⁵⁸.

NOTE Genomic alterations detected may be associated with activity of certain FDA approved drugs, however, the agents listed in this report may have varied evidence in the patient's tumor type.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

MSH6

 NM_000179.2:
c.4068_4071dup
(p.K1358Dfs*2)
chr2:48033981

TSC1

 NM_000368.4: c.1960C>G
(p.Q654E)
chr9:135781005

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Erik Williams, M.D. | 14 July 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

APPENDIX
Genes Assayed in FoundationOne®CDx

FoundationOne CDx is designed to include genes known to be somatically altered in human solid tumors that are validated targets for therapy, either approved or in clinical trials, and/or that are unambiguous drivers of oncogenesis based on current knowledge. The current assay interrogates 324 genes as well as introns of 36 genes involved in rearrangements. The assay will be updated periodically to reflect new knowledge about cancer biology.

DNA GENE LIST: ENTIRE CODING SEQUENCE FOR THE DETECTION OF BASE SUBSTITUTIONS, INSERTION/DELETIONS, AND COPY NUMBER ALTERATIONS

ABL1	ACVR1B	AKT1	AKT2	AKT3	ALK	ALOX12B	AMER1 (FAM123B or WTX)	
APC	AR	ARAF	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX
AURKA	AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2
BCL6	BCOR	BCORL1	BRAF	BRCA1	BRCA2	BRD4	BRIP1	BTG1
BTG2	BTK	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD274 (PD-L1)	CD70	CD79A	CD79B	CDC73	CDH1
CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B	CDKN2C
CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R	CTCF
CTNNA1	CTNNB1	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1	DDR2
DIS3	DNMT3A	DOT1L	EED	EGFR	EMSY (C11orf30)	EP300	EPHA3	EPHB1
EPHB4	ERBB2	ERBB3	ERBB4	ERCC4	ERG	ERRF1	ESR1	EZH2
FANCA	FANCC	FANCG	FANCL	FAS	FBXW7	FGF10	FGF12	FGF14
FGF19	FGF23	FGF3	FGF4	FGF6	FGFR1	FGFR2	FGFR3	FGFR4
FH	FLCN	FLT1	FLT3	FOXL2	FUBP1	GABRA6	GATA3	GATA4
GATA6	GID4 (C17orf39)	GNA11	GNA13	GNAQ	GNAS	GRM3	GSK3B	H3-3A (H3F3A)
HDAC1	HGF	HNFA1	HRAS	HSD3B1	ID3	IDH1	IDH2	IGF1R
IKBKE	IKZF1	INPP4B	IRF2	IRF4	IRS2	JAK1	JAK2	JAK3
JUN	KDM5A	KDM5C	KDM6A	KDR	KEAP1	KEL	KIT	KLHL6
KMT2A (MLL)	KMT2D (MLL2)	KRAS	LTK	LYN	MAF	MAP2K1 (MEK1)	MAP2K2 (MEK2)	MAP2K4
MAP3K1	MAP3K13	MAPK1	MCL1	MDM2	MDM4	MED12	MEF2B	MEN1
MERTK	MET	MITF	MKNK1	MLH1	MPL	MRE11 (MRE11A)	MSH2	MSH3
MSH6	MST1R	MTAP	MTOR	MUTYH	MYC	MYCL (MYCL1)	MYCN	MYD88
NBN	NF1	NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2	NOTCH3
NPM1	NRAS	NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1	NTRK2	NTRK3	NTRK3
P2RY8	PALB2	PARP1	PARP2	PARP3	PAX5	PBRM1	PDCCD1 (PD-1)	PDCCD1LG2 (PD-L2)
PDGFRA	PDGFRB	PDK1	PIK3C2B	PIK3C2G	PIK3CA	PIK3CB	PIK3R1	PIM1
PMS2	POLD1	POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI
PRKN (PARK2)	PTCH1	PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51
RAD51B	RAD51C	RAD51D	RAD52	RAD54L	RAF1	RARA	RB1	RBM10
REL	RET	RICTOR	RNF43	ROS1	RPTOR	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SMAD2	SMAD4	SMARCA4	SMARCB1	SMO
SNCAIP	SOC1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2	STAT3
STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TET2	TET2	TGFB2
TIPARP	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2	TYRO3	U2AF1	VEGFA
VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703			

DNA GENE LIST: FOR THE DETECTION OF SELECT REARRANGEMENTS

ALK	BCL2	BCR	BRAF	BRCA1	BRCA2	CD74	EGFR	ETV4
ETV5	ETV6	EWSR1	EZR	FGFR1	FGFR2	FGFR3	KIT	KMT2A (MLL)
MSH2	MYB	MYC	NOTCH2	NTRK1	NTRK2	NUTM1	PDGFRA	RAF1
RARA	RET	ROS1	RSP02	SDC4	SLC34A2	TERC*	TERT**	TPRSS2

*TERC is an NCRNA

**Promoter region of TERT is interrogated

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Homologous Recombination status

Loss of Heterozygosity (LOH) score

Microsatellite (MS) status


Tumor Mutational Burden (TMB)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1667994-01

APPENDIX

About FoundationOne®CDx

FoundationOne CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Cipalstraat 3, 2440 Geel, Belgium. 

ABOUT FOUNDATIONONE CDx

FoundationOne CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details:
www.rochefoundationmedicine.com/f1cdxtech.

INTENDED USE

FoundationOne®CDx (F1CDx) is a next generation sequencing based in vitro diagnostic device for detection of substitutions, insertion and deletion alterations (indels), and copy number alterations (CNAs) in 324 genes and select gene rearrangements, as well as genomic signatures including microsatellite instability (MSI), tumor mutational burden (TMB), and for selected forms of ovarian cancer, loss of heterozygosity (LOH) score, using DNA isolated from formalin-fixed, paraffin-embedded (FFPE) tumor tissue specimens. The test is intended as a companion diagnostic to identify patients who may benefit from treatment with therapies in accordance with approved therapeutic product labeling. Additionally, F1CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with solid malignant neoplasms.

TEST PRINCIPLE

FoundationOne CDx will be performed exclusively as a laboratory service using DNA extracted from formalin-fixed, paraffin-embedded (FFPE) tumor samples. The proposed assay will employ a single DNA extraction method from routine FFPE biopsy or surgical resection specimens, 50-1000 ng of which will undergo whole-genome shotgun library construction and hybridization-based capture of all coding exons from 309 cancer-related genes, one promoter region, one non-coding (ncRNA), and select intronic regions from 34 commonly rearranged genes, 21 of which also include the coding exons. The assay therefore includes

detection of alterations in a total of 324 genes.

Using an Illumina® Sequencing platform (HiSeq 4000 or NovaSeq 6000), hybrid capture-selected libraries will be sequenced to high uniform depth (targeting >500X median coverage with >99% of exons at coverage >100X). Sequence data will be processed using a customized analysis pipeline designed to accurately detect all classes of genomic alterations, including base substitutions, indels, focal copy number amplifications, homozygous gene deletions, and selected genomic rearrangements (e.g., gene fusions). Additionally, genomic signatures including loss of heterozygosity (LOH), microsatellite instability (MSI), and tumor mutational burden (TMB) will be reported.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. The F1CDx report may be used as an aid to inform molecular eligibility for clinical trials. Note: A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

Diagnostic Significance

FoundationOne CDx identifies alterations to select cancer-associated genes or portions of genes (biomarkers). In some cases, the Report also highlights selected negative test results regarding biomarkers of clinical significance.

Qualified Alteration Calls (Equivocal and Subclonal)

An alteration denoted as "amplification – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence that the copy number of a gene exceeds the threshold for identifying copy number amplification. The threshold used in FoundationOne CDx for identifying a copy number amplification is four (4) for *ERBB2* and six (6) for all other genes. Conversely, an alteration denoted as "loss – equivocal" implies that the FoundationOne CDx assay data provide some, but not unambiguous, evidence for homozygous deletion of the gene in question. An alteration denoted as "subclonal" is

one that the FoundationOne CDx analytical methodology has identified as being present in <10% of the assayed tumor DNA.

Ranking of Therapies and Clinical Trials

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

Limitations

1. In the fraction-based MSI algorithm, a tumor specimen will be categorized as MSI-H, MSS, or MS-Equivocal according to the fraction of microsatellite loci determined to be altered or unstable (i.e., the fraction unstable loci score). In the F1CDx assay, MSI is evaluated based on a genome-wide analysis across >2000 microsatellite loci. For a given microsatellite locus, non-somatic alleles are discarded, and the microsatellite is categorized as unstable if remaining alleles differ from the reference genome. The final fraction unstable loci score is calculated as the number of unstable microsatellite loci divided by the number of evaluable microsatellite loci. The MSI-H and

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531

ORDERED TEST # ORD-1667994-01

APPENDIX

About FoundationOne®CDx

- MSS cut-off thresholds were determined by analytical concordance to a PCR comparator assay using a pan-tumor FFPE tissue sample set. Patients with results categorized as "MS-Stable" with median exon coverage $<300\times$, "MS-Equivocal," or "Cannot Be Determined" should receive confirmatory testing using a validated orthogonal (alternative) method.
- TMB by F1CDx is determined by counting all synonymous and non-synonymous variants present at 5% allele frequency or greater (after filtering) and the total number is reported as mutations per megabase (mut/Mb) unit. Observed TMB is dependent on characteristics of the specific tumor focus tested for a patient (e.g., primary vs. metastatic, tumor content) and the testing platform used for the detection; therefore, observed TMB results may vary between different specimens for the same patient and between detection methodologies employed on the same sample. The TMB calculation may differ from TMB calculations used by other assays depending on variables such as the amount of genome interrogated, percentage of tumor, assay limit of detection (LoD), filtering of alterations included in the score, and the read depth and other bioinformatic test specifications. Refer to the SSED for a detailed description of these variables in FMI's TMB calculation https://www.accessdata.fda.gov/cdrh_docs/pdf17/P170019B.pdf. The clinical validity of TMB defined by this panel has been established for TMB as a qualitative output for a cut-off of 10 mutations per megabase but has not been established for TMB as a quantitative score.
 - Homologous Recombination status may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas (Coleman et al., 2017; 28916367). Samples with deleterious *BRCA1/2* alteration and/or Loss of Heterozygosity (LOH) score $\geq 16\%$ will be reported as "HRD Positive" and samples with absence of these findings will be reported as "HRD Not Detected," agnostic of potential secondary *BRCA1/2* reversion alterations. Certain potentially deleterious missense or small in-frame deletions in *BRCA1/2* may not be classified as deleterious and, in the absence of an elevated LOH profile, samples with such mutations may be classified as "HRD Not Detected." A result of "HRD Not Detected" does not rule out the presence of a *BRCA1/2* alteration or an elevated LOH profile outside the assay performance characteristic limitations.
 - The LOH score is determined by analyzing SNPs spaced at 1Mb intervals across the

genome on the FoundationOne CDx test and extrapolating an LOH profile, excluding arm- and chromosome-wide LOH segments. Detection of LOH has been verified only for ovarian cancer patients, and the LOH score result may be reported for epithelial ovarian, peritoneal, or Fallopian tube carcinomas. The LOH score will be reported as "Cannot Be Determined" if the sample is not of sufficient quality to confidently determine LOH. Performance of the LOH classification has not been established for samples below 35% tumor content. There may be potential interference of ethanol with LOH detection. The interfering effects of xylene, hemoglobin, and triglycerides on the LOH score have not been demonstrated.

- Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. The test does not provide information about susceptibility.
- Biopsy may pose a risk to the patient when archival tissue is not available for use with the assay. The patient's physician should determine whether the patient is a candidate for biopsy.
- Reflex testing to an alternative FDA approved companion diagnostic should be performed for patients who have an *ERBB2* amplification result detected with copy number equal to 4 (baseline ploidy of tumor +2) for confirmatory testing. While this result is considered negative by FoundationOne®CDx (F1CDx), in a clinical concordance study with an FDA approved FISH test, 70% (7 out of 10 samples) were positive, and 30% (3 out of 10 samples) were negative by the FISH test with an average ratio of 2.3. The frequency of *ERBB2* copy number 4 in breast cancer is estimated to be approximately 2%. Multiple references listed in <https://www.mycancergenome.org/content/disease/breast-cancer/ERBB2/238/> report the frequency of *HER2* overexpression as 20% in breast cancer. Based on the F1CDx *HER2* CDx concordance study, approximately 10% of *HER2* amplified samples had copy number 4. Thus, total frequency is conservatively estimated to be approximately 2%.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic,

nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANT ALLELE FREQUENCY

Variant Allele Frequency (VAF) represents the fraction of sequencing reads in which the variant is observed. This attribute is not taken into account for therapy inclusion, clinical trial matching, or interpretive content. Caution is recommended in interpreting VAF to indicate the potential germline or somatic origin of an alteration, recognizing that tumor fraction and tumor ploidy of samples may vary.

Precision of VAF for base substitutions and indels

BASE SUBSTITUTIONS	%CV*
Repeatability	5.11 - 10.40
Reproducibility	5.95 - 12.31
INDELS	%CV*
Repeatability	6.29 - 10.00
Reproducibility	7.33 - 11.71

*Interquartile Range = 1st Quartile to 3rd Quartile

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of $>10\%$, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *PALB2*, *PMS2*, *POLE*, *RAD51C*, *RAD51D*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *TSC2*, and *VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1667994-01

APPENDIX

About FoundationOne®CDx

distinguish whether a finding in this patient's tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1*, *CBL*, *DNMT3A*, *IDH2*, *JAK2*, *KMT2D* (*MLL2*), *MPL*, *MYD88*, *SF3B1*, *TET2*, and *U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This Report makes no promises or guarantees that a particular drug will be effective in the treatment of disease in any patient. This Report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne CDx.

TREATMENT DECISIONS ARE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent

medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this Test, or the information contained in this Report. Certain sample or variant characteristics may result in reduced sensitivity. FoundationOne CDx is performed using DNA derived from tumor, and as such germline events may not be reported.

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
mut/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.10.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of the data. Please refer to the original report for the suitability of use.

The median exon coverage for this sample is 940x

ORDERED TEST # ORD-1667994-01

APPENDIX
References

1. Histopathology (2007) PMID: 17204026
2. Lal N, et al. Oncoimmunology (2015) PMID: 25949894
3. Overman MJ, et al. Lancet Oncol. (2017) PMID: 28734759
4. Overman MJ, et al. J. Clin. Oncol. (2018) PMID: 29355075
5. Shitara K, et al. Nature (2022) PMID: 35322232
6. Lipson EJ, et al. Clin. Cancer Res. (2013) PMID: 23169436
7. Le DT, et al. N. Engl. J. Med. (2015) PMID: 26028255
8. Rizvi NA, et al. Science (2015) PMID: 25765070
9. Oaknin A, et al. JAMA Oncol (2020) PMID: 33001143
10. Rubinstein MM, et al. Gynecol Oncol (2022) PMID: 36512912
11. Kanikarla Marie P, et al. Clin Cancer Res (2021) PMID: 33811152
12. Hochster et al., 2017; ASCO Abstract 673
13. Sinicrope FA, et al. J. Clin. Oncol. (2013) PMID: 24019539
14. Gavin PG, et al. Clin. Cancer Res. (2012) PMID: 23045248
15. Bertagnolli MM, et al. J. Clin. Oncol. (2009) PMID: 19273709
16. Van Cutsem E, et al. J. Clin. Oncol. (2009) PMID: 19451425
17. Ribic CM, et al. N. Engl. J. Med. (2003) PMID: 12867608
18. Sargent DJ, et al. J. Clin. Oncol. (2010) PMID: 20498393
19. Fallik D, et al. Cancer Res. (2003) PMID: 14522894
20. Jass JR, et al. J. Clin. Pathol. (1999) PMID: 10562815
21. Guastadisegni C, et al. Eur. J. Cancer (2010) PMID: 20627535
22. Pawlik TM, et al. Dis. Markers (2004) PMID: 15528785
23. Kocarnik JM, et al. Gastroenterol Rep (Oxf) (2015) PMID: 26337942
24. Nature (2012) PMID: 22810696
25. Samowitz WS, et al. Cancer Epidemiol. Biomarkers Prev. (2001) PMID: 11535541
26. Elsaleh H, et al. Clin Colorectal Cancer (2001) PMID: 12445368
27. Brueckl WM, et al. Anticancer Res. () PMID: 12820457
28. Guidoboni M, et al. Am. J. Pathol. (2001) PMID: 11438476
29. Gryfe R, et al. N. Engl. J. Med. (2000) PMID: 10631274
30. Sinicrope FA, et al. Gastroenterology (2006) PMID: 16952542
31. Laghi L, et al. Dig Dis (2012) PMID: 22722556
32. Boland CR, et al. Gastroenterology (2010) PMID: 20420947
33. Kambara T, et al. Cancer Res. (2001) PMID: 11691787
34. Torshizi Esfahani A, et al. J Cell Physiol (2019) PMID: 30549036
35. Konishi M, et al. Gastroenterology (1996) PMID: 8690195
36. You JF, et al. Br. J. Cancer (2010) PMID: 21081928
37. Bairwa NK, et al. Methods Mol. Biol. (2014) PMID: 24623249
38. Samstein RM, et al. Nat. Genet. (2019) PMID: 30643254
39. Goodman AM, et al. Mol. Cancer Ther. (2017) PMID: 28835386
40. Goodman AM, et al. Cancer Immunol Res (2019) PMID: 31405947
41. Cristescu R, et al. Science (2018) PMID: 30309915
42. Ready N, et al. J. Clin. Oncol. (2019) PMID: 30785829
43. Hellmann MD, et al. N. Engl. J. Med. (2018) PMID: 29658845
44. Hellmann MD, et al. Cancer Cell (2018) PMID: 29657128
45. Hellmann MD, et al. Cancer Cell (2018) PMID: 29731394
46. Rozeman EA, et al. Nat Med (2021) PMID: 33558721
47. Sharma P, et al. Cancer Cell (2020) PMID: 32916128
48. Marabelle A, et al. Lancet Oncol. (2020) PMID: 32919526
49. Ott PA, et al. J. Clin. Oncol. (2019) PMID: 30557521
50. Cristescu R, et al. J Immunother Cancer (2022) PMID: 35101941
51. Friedman CF, et al. Cancer Discov (2022) PMID: 34876409
52. Sturgill EG, et al. Oncologist (2022) PMID: 35274716
53. Schenker et al., 2022; AACR Abstract 7845
54. Legrand et al., 2018; ASCO Abstract 12000
55. Fabrizio DA, et al. J Gastrointest Oncol (2018) PMID: 30151257
56. Stadler ZK, et al. J. Clin. Oncol. (2016) PMID: 27022117
57. Shao C, et al. JAMA Netw Open (2020) PMID: 33191110
58. Schwartz et al., 2018; ASCO Abstract 572
59. Innocenti F, et al. J Clin Oncol (2019) PMID: 30865548
60. Lee DW, et al. Clin Cancer Res (2019) PMID: 31285374
61. Randon G, et al. Eur J Cancer (2022) PMID: 34933155
62. Chen EX, et al. JAMA Oncol (2020) PMID: 32379280
63. Pfeifer GP, et al. Mutat. Res. (2005) PMID: 15748635
64. Hill VK, et al. Annu Rev Genomics Hum Genet (2013) PMID: 23875803
65. Pfeifer GP, et al. Oncogene (2002) PMID: 12379884
66. Johnson BE, et al. Science (2014) PMID: 24336570
67. Choi S, et al. Neuro-oncology (2018) PMID: 29452419
68. Cancer Genome Atlas Research Network, et al. Nature (2013) PMID: 23636398
69. Briggs S, et al. J. Pathol. (2013) PMID: 23447401
70. Heitzer E, et al. Curr. Opin. Genet. Dev. (2014) PMID: 24583393
71. Roberts SA, et al. Nat. Rev. Cancer (2014) PMID: 25568919
72. Van Cutsem E, et al. J. Clin. Oncol. (2011) PMID: 21502544
73. Bokemeyer C, et al. Ann. Oncol. (2011) PMID: 21228335
74. Karapetis CS, et al. N. Engl. J. Med. (2008) PMID: 18946061
75. De Roock W, et al. Ann. Oncol. (2008) PMID: 17998284
76. Douillard JY, et al. Ann. Oncol. (2014) PMID: 24718886
77. Douillard JY, et al. N. Engl. J. Med. (2013) PMID: 24024839
78. Amado RG, et al. J. Clin. Oncol. (2008) PMID: 18316791
79. Lièvre A, et al. Cancer Res. (2006) PMID: 16618717
80. De Roock W, et al. Lancet Oncol. (2011) PMID: 21163703
81. Chen J, et al. BMC Cancer (2014) PMID: 25367198
82. Li W, et al. BMC Cancer (2015) PMID: 25929517
83. Hu J, et al. Medicine (Baltimore) (2016) PMID: 27977612
84. Zekri J, et al. Genet. Mol. Res. (2017) PMID: 28218784
85. Staudacher JJ, et al. Clin Transl Gastroenterol (2017) PMID: 29048416
86. Wang Y, et al. Virchows Arch. (2018) PMID: 29705968
87. Guo F, et al. Sci Rep (2018) PMID: 29666387
88. Mármol I, et al. Int J Mol Sci (2017) PMID: 28106826
89. Kwak MS, et al. Medicine (Baltimore) (2017) PMID: 28858102
90. Pylyayeva-Gupta Y, et al. Nat. Rev. Cancer (2011) PMID: 21993244
91. Kahn S, et al. Anticancer Res. () PMID: 3310850
92. Pentheroudakis G, et al. BMC Cancer (2013) PMID: 23374602
93. Vaughn CP, et al. Genes Chromosomes Cancer (2011) PMID: 21305640
94. Janku F, et al. Target Oncol (2013) PMID: 23400451
95. De Roock W, et al. Lancet Oncol. (2010) PMID: 20619739
96. Irahara N, et al. Diagn. Mol. Pathol. (2010) PMID: 20736745
97. Schirripa M, et al. Int. J. Cancer (2015) PMID: 24806288
98. Cercek A, et al. Clin. Cancer Res. (2017) PMID: 28446505
99. Shi C, et al. Oncogene (2022) PMID: 35393542
100. Park JW, et al. Cancer Med (2022) PMID: 35274815
101. Ormanns S, et al. Int J Mol Sci (2017) PMID: 28534865
102. Fei N, et al. Clin Transl Sci (2021) PMID: 34002944
103. Bachet JB, et al. Ann. Oncol. (2012) PMID: 22377565
104. Ziemke M, et al. Lung Cancer (2017) PMID: 28577946
105. Kassardjian A, et al. Pancreas (2020) PMID: 32897998
106. Pen SL, et al. Radiother Oncol (2021) PMID: 33667587
107. Witkiewicz AK, et al. Nat Commun (2015) PMID: 25855536
108. Jiao Y, et al. J. Pathol. (2014) PMID: 24293293
109. Churi CR, et al. PLoS ONE (2014) PMID: 25536104
110. Takeda et al., 2022; ASCO GI Abstract 642
111. Liu X, et al. Clin. Chem. (2014) PMID: 24821835
112. Maru D, et al. Oncogene (2004) PMID: 14647445
113. Wang K, et al. Oncologist (2015) PMID: 26336083
114. Nature (2014) PMID: 25079317
115. Izeradjene K, et al. Cancer Cell (2007) PMID: 17349581
116. Bardeesy N, et al. Genes Dev. (2006) PMID: 17114584
117. Springer S, et al. Gastroenterology (2015) PMID: 26253305
118. Blackford A, et al. Clin. Cancer Res. (2009) PMID: 19584151
119. Yan P, et al. Clin. Cancer Res. (2016) PMID: 26861460
120. Kozak MM, et al. J. Clin. Pathol. (2015) PMID: 25681512
121. Roth AD, et al. J. Natl. Cancer Inst. (2012) PMID: 23104212
122. Davison JM, et al. Am. J. Surg. Pathol. (2014) PMID: 24618609
123. Kim YH, et al. Ann. Oncol. (2004) PMID: 15033661
124. Xiangming C, et al. Clin. Cancer Res. (2001) PMID: 11234879
125. Singhi AD, et al. Am. J. Surg. Pathol. (2015) PMID: 25634752
126. Natsugoe S, et al. Clin. Cancer Res. (2002) PMID: 12060625
127. de Kruif EM, et al. Ann. Oncol. (2013) PMID: 23022998
128. Shipitsin M, et al. Br. J. Cancer (2014) PMID: 25032733
129. Nat. Rev. Mol. Cell Biol. (2012) PMID: 22992590
130. Cell (2008) PMID: 18662538
131. Massagué J, et al. Genes Dev. (2005) PMID: 16322555
132. Morén A, et al. Oncogene (2000) PMID: 10980615
133. Xu J, et al. Proc. Natl. Acad. Sci. U.S.A. (2000) PMID: 10781087
134. Luo K, et al. Genes Dev. (1999) PMID: 10485843
135. Jones JB, et al. Nucleic Acids Res. (2000) PMID: 10871368
136. Fink SP, et al. Cancer Res. (2001) PMID: 11196171
137. De Bosscher K, et al. Biochem. J. (2004) PMID: 14715079
138. Shi Y, et al. Nature (1997) PMID: 9214508
139. Miyaki M, et al. Oncogene (1999) PMID: 10340381
140. Prokova V, et al. Biochemistry (2007) PMID: 17994767
141. Wu JW, et al. J. Biol. Chem. (2001) PMID: 11274206
142. Ding L, et al. J. Clin. Invest. (2009) PMID: 19139564
143. Kuang C, et al. Oncogene (2004) PMID: 14647410
144. Watanabe M, et al. EMBO Rep. (2000) PMID: 11265759
145. Houlston R, et al. Hum. Mol. Genet. (1998) PMID: 9811934
146. Woodford-Richens K, et al. Gut (2000) PMID: 10764709
147. Howe JR, et al. J. Med. Genet. (2004) PMID: 15235019
148. Brosens LA, et al. World J. Gastroenterol. (2011) PMID: 22171123

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1667994-01

APPENDIX
References

- | | | |
|---|--|---|
| <p>149. Kalia SS, et al. Genet. Med. (2017) PMID: 27854360</p> <p>150. Cunningham D, et al. N. Engl. J. Med. (2004) PMID: 15269313</p> <p>151. Jonker DJ, et al. N. Engl. J. Med. (2007) PMID: 18003960</p> <p>152. Papamichael D, et al. Eur J Cancer (2022) PMID: 35033994</p> <p>153. Karapetis CS, et al. Clin. Cancer Res. (2014) PMID: 24218517</p> <p>154. Chibaudel et al., 2022; ASCO Abstract 3504</p> <p>155. Moiseyenko VM, et al. Clin Drug Investig (2018) PMID: 29470838</p> | <p>156. Stein A, et al. J Immunother Cancer (2021) PMID: 34315821</p> <p>157. Price TJ, et al. Lancet Oncol. (2014) PMID: 24739896</p> <p>158. Sakai D, et al. Eur J Cancer (2020) PMID: 32526634</p> <p>159. Van Cutsem E, et al. J. Clin. Oncol. (2007) PMID: 17470858</p> <p>160. Peeters M, et al. Clin. Cancer Res. (2015) PMID: 26341920</p> <p>161. Watanabe J, et al. Int J Cancer (2022) PMID: 35723084</p> | <p>162. Kim TW, et al. Clin Colorectal Cancer (2018) PMID: 29703606</p> <p>163. Shitara K, et al. Cancer Sci (2016) PMID: 27712015</p> <p>164. Peeters M, et al. Clin. Cancer Res. (2013) PMID: 23325582</p> <p>165. Yoshino et al., 2022; ASCO Abstract LBA1</p> <p>166. Pietrantonio F, et al. JAMA Oncol (2019) PMID: 31268481</p> |
|---|--|---|

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Erik Williams, M.D. | 14 July 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531